Humana. Claims Payment Policy

Subject: Laboratory – Biochemical Markers of Alzheimer's Disease and Dementia

Application: Medicare Advantage Products
Policy number: LAB2023026
Published date: 06/2023
Revision date: 04/2026

Disclaimer: *The intended audience of this medical claims payment policy is healthcare providers who treat Humana members. This policy is made available to provide information on certain Humana claims payment policies. This policy is a guideline only and does not constitute medical advice, guarantee of payment, plan pre-authorization, an explanation of benefits, or a contract. This policy does not govern whether a procedure is covered under a specific member plan or policy, nor is it intended to address every claim situation. Claims may be affected by other factors, such as: state and federal laws and regulations, provider contract terms, and our professional judgment. This policy is subject to change or termination by Humana. Humana has full and final discretionary authority for its interpretation and application. No part of this policy may be reproduced, stored in a retrieval system or transmitted, in any shape or form or by any means, electronic, mechanical, photocopying or otherwise without express written permission from Humana. When printed, this version becomes uncontrolled. For the most current information, always refer to the following website:* ***Humana.com/ClaimPaymentPolicies***

Medicare Advantage Payment Policy

As a Medicare Advantage plan, we have developed payment policies to explain coverage more fully for services. The criteria outlined will apply to Medicare Advantage plans except when the procedure(s) has established criteria through the Original Medicare National Coverage Determinations (NCDs) and Local Coverage Determinations (LCDs) as outlined below.

In addition to the policy, claims payments are subject to other plan requirements for the processing and payment of claims, including, but not limited to, requirements of medical necessity and reasonableness and applicable referral or authorization requirements.

- For individuals with a diagnosis of Alzheimer's disease, mild cognitive impairment or dementia, measurement of the following biomarkers **IS REIMBURSABLE:**
  - Amyloid beta peptides (CPT Code 82233, CPT code 82234, CPT code 0358U)
  - The ratio (CPT code 0459U) of total tau (t-tau) (CPT code 84394) to amyloid beta 1-42 (Aβ42) (CPT code 82234)
  - The ratio (CPT code 0445U) of phosphorylated tau 181 (p-tau181) (CPT code 84393) to Aβ42 (CPT code 82234)
  - The ratio of p-tau 217 (p-tau217) (CPT code 84393, CPT code 0479U) to Aβ42 (CPT code 82234)
- When not ordered in conjunction with Aβ42 (CPT code 82234), measurement of the following biomarkers **IS NOT REIMBURSABLE:**
  - t-tau (CPT code 84394)
  - p-tau181 (CPT code 84393)
  - p-tau217 (CPT code 84393, CPT code 0479U)
- For individuals with a diagnosis of Alzheimer's disease, mild cognitive impairment or dementia, measurement of other biomarkers (CPT code 83520, CPT code 83884, CPT code 0547U) **IS NOT REIMBURSABLE.**
- For individuals with a diagnosis of Alzheimer's disease, mild cognitive impairment or dementia, measurement of all other cerebral spinal fluid biomarkers (CPT code 0393U, CPT code 0443U, CPT code 0548U) or plasma biomarkers (CPT code 0479U, CPT code 0551U) **IS NOT REIMBURSABLE.**
- The use of DISCERN™ (CPT code 0206U*, CPT code 0207U*) **IS NOT REIMBURSABLE.**
- The use of other multianalyte assays (CPT code 0412U*, CPT code 0503U*, CPT code 0568U*, CPT code 0596U) for Alzheimer's disease or dementia **IS NOT REIMBURSABLE.**
- Gene expression profiling to predict development of Alzheimer's disease (CPT code 0289U**) **IS NOT REIMBURSABLE.**

Applicable procedures for Medicare Advantage plans

Procedure code	Code description
82233	Beta-amyloid; 1-40 (Abeta 40)
82234	Beta-amyloid; 1-42 (Abeta 42)
83520	Immunoassay for analyte other than infectious agent antibody or infectious agent antigen; quantitative, not otherwise specified
83884	Neurofilament light chain (NfL)
84393	Tau, phosphorylated (eg, pTau 181, pTau 217), each
84394	Tau, total (tTau)
0206U*	Neurology (Alzheimer's disease); cell aggregation using morphometric imaging and protein kinase C-epsilon (PKCe) concentration in response to amylospheroid treatment by ELISA, cultured skin fibroblasts, each reported as positive or negative for Alzheimer's disease Proprietary test: DISCERN™ Lab/Manufacturer: NeuroDiagnostics
0207U*	Neurology (Alzheimer's disease); Quantitative imaging of phosphorylated ERK1 and ERK2 in response to bradykinin treatment by in situ immunofluorescence, using cultured skin fibroblasts, reported as a probability index for Alzheimer's disease (List separately in addition to code for primary procedure) Proprietary test: DISCERN™ Lab/Manufacturer: NeuroDiagnostics
0289U**	Neurology (Alzheimer's disease), mRNA, gene expression profiling by RNA sequencing of 24 genes, whole blood, algorithm reported as predictive risk score Proprietary test: MindX Blood Test™ - Memory/Alzheimer's Lab/Manufacturer: MindX Sciences™ Laboratory/MindX Sciences™ Inc
0358U	Neurology (mild cognitive impairment), analysis of B-amyloid 1- and 1-40, chemiluminescence enzyme immunoassay, cerebral spinal fluid, reported as positive, likely positive, or negative Proprietary test: Lumipulse® G βAmyloid Ratio (1–42/1–40) Test Lab/Manufacturer: Fujirebio Diagnostics
0393U	Neurology (eg, Parkinson disease, dementia with Lewy bodies), cerebrospinal fluid (CSF), detection of misfolded a-synuclein protein by seed amplification assay, qualitative Proprietary test: SAAmplify™ -aSYN Lab/Manufacturer: Amprion Clinical Laboratory
0412U*	Beta amyloid, AB42/40 ratio, immunoprecipitation with quantitation by liquid chromatography with tandem mass spectrometry (LC-MS/MS) and qualitative ApoE isoform-specific proteotyping, plasma combined with age, algorithm reported as presence or absence of brain amyloid pathology Proprietary test: PrecivityAD® blood test Lab/Manufacturer: C2N Diagnostics LLC
0443U	Neurofilament light chain (NfL), ultra-sensitive immunoassay, serum or cerebrospinal fluid Proprietary test: Neurofilament Light Chain (NfL) Lab/Manufacturer: Neuromuscular Clinical Laboratory at Washington University in St. Louis School of Medicine
0445U	B-amyloid (Abeta42) and phospho tau (181P) (pTau181), electrochemiluminescent immunoassay (ECLIA), cerebral spinal fluid, ratio reported as positive or negative for amyloid pathology Proprietary test: Elecsys® Phospho-Tau (181P) CSF (pTau181) and B-Amyloid (1-42) CSF II (Abeta 42) Ratio Lab/Manufacturer: Roche Diagnostics Operations, Inc (US owner/operator)
0459U	B-amyloid (Abeta42) and total tau (tTau), electrochemiluminescent immunoassay (ECLIA), cerebral spinal fluid, ratio reported as positive or negative for amyloid pathology Proprietary test: Elecsys® Total Tau CSF (tTau) and B-Amyloid (1-42) CSF II (Abeta 42) Ratio, Lab/Manufacturer: Roche Diagnostics Operations, Inc (US owner/operator)
0479U	Tau, phosphorylated, pTau217 Proprietary test: ALZpath pTau217, Neurocode USA, Inc,

Procedure code	Code description
	Lab/Manufacturer: Quanterix/ALZpath
0503U*	Neurology (Alzheimer's disease), beta amyloid (AB40, AB42, AB42/40 ratio) and tau-protein (ptau217, np-tau217, ptau217/np-tau217 ratio), blood, immunoprecipitation with quantitation by liquid chromatography with tandem mass spectrometry (LC-MS/MS), algorithm score reported as likelihood of positive or negative for amyloid plaques Proprietary test: PrecivityAD2TM Lab Manufacturer: C2N Diagnostics, LLC
0547U	Neurofilament light chain (NfL), chemiluminescent enzyme immunoassay, plasma, quantitative Proprietary test: Neurofilament Light Blood Test Lab Manufacturer: Neurocode USA, Inc, Fujirebio Diagnostics, Inc
0548U	Glial fibrillary acidic protein (GFAP), chemiluminescent enzyme immunoassay, using plasma Proprietary test: Glial Fibrillary Acidic Protein Blood Test Lab Manufacturer: Neurocode USA, Inc, Fujirebio Diagnostics, Inc
0568U*	Neurology (dementia), beta amyloid (AB40, AB42, AB42/40 ratio), tau-protein phosphorylated at residue (eg, pTau217), neurofilament light chain (NfL), and glial fibrillary acidic protein (GFAP), by ultra-high sensitivity molecule array detection, plasma, algorithm reported as positive, intermediate, or negative for Alzheimer's pathology Proprietary test: LucentAD™ Complete Lab Manufacturer: Quanterix Corporation, Quanterix Corporation
0596U	Neurology (Alzheimer's disease), plasma, 3 distinct isoform-specific peptides (APOE2, APOE3, and APOE4) by liquid chromatography with tandem mass spectrometry (LC-MS/MS), reported as an APOE proteotype Proprietary test: Precivity-ApoE™ Lab Manufacturer: C2N Diagnostics, LLC

** The payment criteria for 0206U, 0207U, 0412U, 0503U and 0568U applies to all Medicare jurisdictions except those states awarded to CGS Administrators, LLC; Noridian Healthcare Solutions, LLC; Palmetto GBA; and WPS Insurance Corporation.*
***The payment criteria for 0289U applies to all Medicare Jurisdictions except those states awarded to CGS Administrators; Noridian Healthcare Solutions, LLC; Palmetto GBA; and WPS Insurance Corporation; First Coast Service Options, Inc. and Novitas Solutions, Inc.*

Original Medicare Local Coverage Determinations (LCDs)

Contractor name	Policy title	Policy number	Procedure code
CGS Administrators, LLC	MolDX: Molecular Diagnostic Tests (MDT)	L36021	0206U 0207U 0289U 0412U 0503U 0568U
CGS Administrators, LLC	MolDX: Repeat Germline Testing	L38288	0289U
Noridian Healthcare Solutions, Inc	MolDX: Molecular Diagnostic Tests (MDT)	L35160	0206U 0207U 0289U 0412U 0503U 0568U
Noridian Healthcare Solutions, Inc	MolDX: Molecular Diagnostic Tests (MDT)	L36256	0206U 0207U 0289U 0412U 0503U 0568U

Contractor name	Policy title	Policy number	Procedure code
Noridian Healthcare Solutions, Inc	MolDX: Repeat Germline Testing	L38351	0289U
Noridian Healthcare Solutions, Inc	MolDX: Repeat Germline Testing	L38353	0289U
Palmetto GBA	MolDX: Molecular Diagnostic Tests (MDT)	L35025	0206U 0207U 0289U 0412U 0503U 0568U
Palmetto GBA	MolDX: Repeat Germline Testing	L38274	0289U
WPS Insurance Corporation	MolDX: Molecular Diagnostic Tests (MDT)	L36807	0206U 0207U 0289U 0412U 0503U 0568U
WPS Insurance Corporation	MolDX: Repeat Germline Testing	L38429	0289U

References

Albert MS, DeKosky ST, Dickson D, et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2011;7(3):270-9. doi:10.1016/j.jalz.2011.03.008

Alzheimer's Association. Lecanemab Approved for Treatment of Early Alzheimer's Disease. https://www.alz.org/Alzheimer'ss-dementia/treatments/lecanemab-leqembi

Arranz AM, De Strooper B. The role of astroglia in Alzheimer's disease: pathophysiology and clinical implications. *Lancet Neurol*. 2019;18(4):406-414. doi:10.1016/s1474-4422(18)30490-3

Bennett DA, Wilson RS, Schneider JA, et al. Natural history of mild cognitive impairment in older persons. *Neurology*. 2002;59(2):198-205.

Blennow K, Zetterberg H, Fagan AM. Fluid biomarkers in Alzheimer's disease. *Cold Spring Harbor perspectives in medicine*. 2012;2(9):a006221. doi:10.1101/cshperspect.a006221

BusinessWire. Quanterix Launches LucentAD Biomarker Blood Test to Aid Physician Diagnosis of Alzheimer's Disease in Patients. https://www.businesswire.com/news/home/20230706742971/en/

C2N. SYN-ONE TEST®. https://c2n.com/news-releases/2019/01/29/2019-1-24-c2n-diagnostics-receives-breakthrough-device-designation-from-us-fda-for-blood-test-to-screen-for-Alzheimer'ss-disease-risk

Chen YR, Liang CS, Chu H, et al. Diagnostic accuracy of blood biomarkers for Alzheimer's disease and amnestic mild cognitive impairment: A meta-analysis. *Ageing Res Rev*. 2021;71:101446. doi:10.1016/j.arr.2021.101446

CND Life Sciences. SYN-ONE TEST®. https://cndlifesciences.com/syn-one-test/

Cordell CB, Borson S, Boustani M, et al. Alzheimer's Association recommendations for operationalizing the detection of cognitive impairment during the Medicare Annual Wellness Visit in a primary care setting. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2013;9(2):141-50. doi:10.1016/j.jalz.2012.09.011

Dage JL, Wennberg AMV, Airey DC, et al. Levels of tau protein in plasma are associated with neurodegeneration and cognitive function in a population-based elderly cohort. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2016;12(12):1226-1234. doi:10.1016/j.jalz.2016.06.001

Donadio V, Doppler K, Incensi A, et al. Abnormal alpha-synuclein deposits in skin nerves: intra- and inter-laboratory reproducibility. *Eur J Neurol*. 2019;26(10):1245-1251. doi:10.1111/ene.13939

Dubois B, Feldman HH, Jacova C, et al. Advancing research diagnostic criteria for Alzheimer's disease: the IWG-2 criteria. *Lancet Neurol*. 2014;13(6):614-29. doi:10.1016/s1474-4422(14)70090-0

Dubois B, Villain N, Frisoni GB, et al. Clinical diagnosis of Alzheimer's disease: recommendations of the International Working Group. *Lancet Neurol*. 2021;20(6):484-496. doi:10.1016/s1474-4422(21)00066-1

Farrer LA, Cupples LA, Haines JL, et al. Effects of age, sex, and ethnicity on the association between apolipoprotein E genotype and Alzheimer's disease. A meta-analysis. APOE and Alzheimer's Disease Meta Analysis Consortium. *Jama*. 1997;278(16):1349-56.

FDA. Early Alzheimer's Disease: Developing Drugs for Treatment. U.S. Food and Drug Administration Center for Drug Evaluation and Research Center for Biologics Evaluation and Research. https://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM596728.pdf

FDA. FDA Permits Marketing for New Test to Improve Diagnosis of Alzheimer's Disease. https://www.prnewswire.com/news-releases/fda-permits-marketing-for-new-test-to-improve-diagnosis-of-Alzheimer'ss-disease-301540093.html

FDA. K242706. https://www.accessdata.fda.gov/cdrh_docs/pdf24/K242706.pdf

Fossati S, Ramos Cejudo J, Debure L, et al. Plasma tau complements CSF tau and P-tau in the diagnosis of Alzheimer's disease. *Alzheimer's & dementia (Amsterdam, Netherlands)*. 2019;11:483-492. doi:10.1016/j.dadm.2019.05.001

François M, Bull CF, Fenech MF, Leifert WR. Current State of Saliva Biomarkers for Aging and Alzheimer's Disease. *Curr Alzheimer's Res*. 2019;16(1):56-66. doi:10.2174/1567205015666181022094924

Fujirebio. Lumipulse® G β-Amyloid Ratio (1-42/1-40). https://www.fujirebio.com/en-us/products-solutions/lumipulse-g-beta-amyloid-ratio-142-140

Gatz M, Reynolds CA, Fratiglioni L, et al. Role of genes and environments for explaining Alzheimer's disease. *Archives of general psychiatry*. 2006;63(2):168-74. doi:10.1001/archpsyc.63.2.168

GBSC. Biomarker Consortium. Global Biomarker Standardization Consortium (GBSC); Alzheimer's Association. https://www.alz.org/research/for_researchers/partnerships/biomarker_consortium

Goldman JS, Hahn SE, Catania JW, et al. Genetic counseling and testing for Alzheimer's disease: Joint practice guidelines of the American College of Medical Genetics and the National Society of Genetic Counselors. *Genet Med*. 2011;13(6):597-605. doi:10.1097/GIM.0b013e31821d69b8

Gottlieb S. Statement from FDA Commissioner Scott Gottlieb, M.D. on advancing the development of novel treatments for neurological conditions; part of broader effort on modernizing FDA's new drug review programs. U.S. Food and Drug Administration. https://www.americanpharmaceuticalreview.com/1315-News/347185-Statement-from-FDA-Commissioner-on-Advancing-the-Development-of-Novel-Treatments-for-Neurological-Conditions/

Hansson O, Edelmayer RM, Boxer AL, et al. The Alzheimer's Association appropriate use recommendations for blood biomarkers in Alzheimer's disease. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2022;18(12):2669-2686. doi:10.1002/alz.12756

Hansson O, Lehmann S, Otto M, Zetterberg H, Lewczuk P. Advantages and disadvantages of the use of the CSF Amyloid β (Aβ) 42/40 ratio in the diagnosis of Alzheimer's Disease. *Alzheimer'ss Res Ther*. 2019;11(1):34. doi:10.1186/s13195-019-0485-0

Herukka SK, Simonsen AH, Andreasen N, et al. Recommendations for cerebrospinal fluid Alzheimer's disease biomarkers in the diagnostic evaluation of mild cognitive impairment. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2017;13(3):285-295. doi:10.1016/j.jalz.2016.09.009

Huan T, Tran T, Zheng J, et al. Metabolomics Analyses of Saliva Detect Novel Biomarkers of Alzheimer's Disease. *Journal of Alzheimer's disease : JAD*. 2018;65(4):1401-1416. doi:10.3233/jad-180711

Ismail Z, Black SE, Camicioli R, et al. Recommendations of the 5th Canadian Consensus Conference on the diagnosis and treatment of dementia. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2020;16(8):1182-1195. doi:10.1002/alz.12105

Jack CR, Jr., Andrews SJ, Beach TG, et al. Revised criteria for the diagnosis and staging of Alzheimer's disease. *Nat Med*. Aug 2024;30(8):2121-2124. doi:10.1038/s41591-024-02988-7

Jack CR, Jr., Bennett DA, Blennow K, et al. NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2018;14(4):535-562. doi:10.1016/j.jalz.2018.02.018

Jiang L, Dong H, Cao H, Ji X, Luan S, Liu J. Exosomes in Pathogenesis, Diagnosis, and Treatment of Alzheimer's Disease. *Med Sci Monit*. 2019;25:3329-3335. doi:10.12659/msm.914027

Karch CM, Cruchaga C, Goate AM. Alzheimer's disease genetics: from the bench to the clinic. *Neuron*. 2014;83(1):11-26. doi:10.1016/j.neuron.2014.05.041

Kim JY, Illigens BM, McCormick MP, Wang N, Gibbons CH. Alpha-Synuclein in Skin Nerve Fibers as a Biomarker for Alpha-Synucleinopathies. *J Clin Neurol*. 2019;15(2):135-142. doi:10.3988/jcn.2019.15.2.135

Kim K, Kim M-J, Kim DW, Kim SY, Park S, Park CB. Clinically accurate diagnosis of Alzheimer's disease via multiplexed sensing of core biomarkers in human plasma. *Nature Communications*. 2020;11(1):119. doi:10.1038/s41467-019-13901-z

Knopman DS, DeKosky ST, Cummings JL, et al. Practice parameter: diagnosis of dementia (an evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology*. 2001;56(9):1143-53. doi:10.1212/wnl.56.9.1143

Kramarow E, Tejada-Vera B. Adjusted Death Rates* from Dementia,† by Sex, Race, and Hispanic Origin — National Vital Statistics System, United States, 2017. *MMWR Morb Mortal Wkly Rep*. 2019;68:670.

Levine TD, Bellaire B, Gibbons C, Freeman R. Cutaneous alpha-synuclein deposition in postural tachycardia patients. *Ann Clin Transl Neurol*. 2021;8(4):908-917. doi:10.1002/acn3.51347

Lewczuk P, Matzen A, Blennow K, et al. Cerebrospinal Fluid Abeta42/40 Corresponds Better than Abeta42 to Amyloid PET in Alzheimer's Disease. *Journal of Alzheimer's disease : JAD*. 2017;55(2):813-822. doi:10.3233/jad-160722

Lewczuk P, Riederer P, O'Bryant SE, et al. Cerebrospinal fluid and blood biomarkers for neurodegenerative dementias: An update of the Consensus of the Task Force on Biological Markers in Psychiatry of the World Federation of Societies of Biological Psychiatry. *The world journal of biological psychiatry : the official journal of the World Federation of Societies of Biological Psychiatry*. 2018;19(4):244-328. doi:10.1080/15622975.2017.1375556

Liu Y, Wei M, Yue K, et al. Study on Urine Metabolic Profile of Abeta25-35-Induced Alzheimer's Disease Using UHPLC-Q-TOF-MS. *Neuroscience*. 2018;394:30-43. doi:10.1016/j.neuroscience.2018.10.001

Lonneborg A. Biomarkers for Alzheimer's disease in cerebrospinal fluid, urine, and blood. *Molecular diagnosis & therapy*. 2008;12(5):307-20.

Mattsson N, Andreasson U, Persson S, et al. The Alzheimer's Association external quality control program for cerebrospinal fluid biomarkers. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2011;7(4):386-395 e6. doi:10.1016/j.jalz.2011.05.2243

McKeith IG, Boeve BF, Dickson DW, et al. Diagnosis and management of dementia with Lewy bodies: Fourth consensus report of the DLB Consortium. *Neurology*. 2017;89(1):88-100. doi:10.1212/wnl.0000000000004058

McKhann GM, Knopman DS, Chertkow H, et al. The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2011;7(3):263-9. doi:10.1016/j.jalz.2011.03.005

Morris JC, Blennow K, Froelich L, et al. Harmonized diagnostic criteria for Alzheimer's disease: recommendations. *Journal of internal medicine*. 2014;275(3):204-13. doi:10.1111/joim.12199

NICE. Dementia: assessment, management and support for people living with dementia and their carers. https://www.nice.org.uk/guidance/ng97/chapter/Recommendations

Nojima H, Ito S, Kushida A, et al. Clinical utility of cerebrospinal fluid biomarkers measured by LUMIPULSE(®) system. *Ann Clin Transl Neurol*. 2022;9(12):1898-1909. doi:10.1002/acn3.51681

Owens DK, Davidson KW, Krist AH, et al. Screening for Cognitive Impairment in Older Adults: US Preventive Services Task Force Recommendation Statement. *Jama*. 2020;323(8):757-763. doi:10.1001/jama.2020.0435

Palmqvist S, Janelidze S, Stomrud E, et al. Performance of Fully Automated Plasma Assays as Screening Tests for Alzheimer's Disease-Related β-Amyloid Status. *JAMA Neurol*. 2019;76(9):1060-9. doi:10.1001/jamaneurol.2019.1632

Patel S, Shah RJ, Coleman P, Sabbagh M. Potential peripheral biomarkers for the diagnosis of Alzheimer's disease. *International journal of Alzheimer's disease*. 2011;2011:572495. doi:10.4061/2011/572495

Peterson RC. Mild cognitive impairment: Prognosis and treatment. Updated June 7, 2024. https://www.uptodate.com/contents/mild-cognitive-impairment-prognosis-and-treatment

Qu Y, Ma YH, Huang YY, et al. Blood biomarkers for the diagnosis of amnestic mild cognitive impairment and Alzheimer's disease: A systematic review and meta-analysis. *Neurosci Biobehav Rev*. 2021;128:479-486. doi:10.1016/j.neubiorev.2021.07.007

Rabinovici GD, Knopman DS, Arbizu J, et al. Updated Appropriate Use Criteria for Amyloid and Tau PET: A Report from the Alzheimer's Association and Society for Nuclear Medicine and Molecular Imaging Workgroup. *J Nucl Med*. Jun 6 2025;66(Suppl 2):S5-s31. doi:10.2967/jnumed.124.268756

Roche. Roche Alzheimer's disease Cerebrospinal Fluid (CSF) assays receive FDA clearance, supporting more accurate and timely diagnosis. https://www.globenewswire.com/news-release/2022/12/08/2569879/0/en/Roche-Alzheimer's-s-disease-Cerebrospinal-Fluid-CSF-assays-receive-FDA-clearance-supporting-more-accurate-and-timely-diagnosis.html

Roche. Roche receives FDA clearance for additional Alzheimer's disease Cerebrospinal Fluid (CSF) assays, supporting timely diagnosis and treatment decision-making. https://diagnostics.roche.com/us/en/news-listing/2023/roche-fda-clearance-additional-Alzheimer'ss-disease-cerebrospinal-fluid-ttau.html

Sala Frigerio C, De Strooper B. Alzheimer's Disease Mechanisms and Emerging Roads to Novel Therapeutics. *Annual Review of Neuroscience*. 2016;39(1):57-79. doi:10.1146/annurev-neuro-070815-014015

Schaffer C, Sarad N, DeCrumpe A, et al. Biomarkers in the Diagnosis and Prognosis of Alzheimer's Disease. *Journal of laboratory automation*. 2015;20(5):589-600. doi:10.1177/2211068214559979

Schindler SE, Petersen KK, Saef B, et al. Head-to-head comparison of leading blood tests for Alzheimer's disease pathology. *medRxiv*. Jul 3 2024;doi:10.1101/2024.06.12.24308839

Shaw LM, Arias J, Blennow K, et al. Appropriate use criteria for lumbar puncture and cerebrospinal fluid testing in the diagnosis of Alzheimer's disease. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2018;14(11):1505-1521. doi:10.1016/j.jalz.2018.07.220

Shea YF, Chu LW, Chan AO, Ha J, Li Y, Song YQ. A systematic review of familial Alzheimer's disease: Differences in presentation of clinical features among three mutated genes and potential ethnic differences. *Journal of the Formosan Medical Association = Taiwan yi zhi*. 2016;115(2):67-75. doi:10.1016/j.jfma.2015.08.004

Shen XN, Niu LD, Wang YJ, et al. Inflammatory markers in Alzheimer's disease and mild cognitive impairment: a meta-analysis and systematic review of 170 studies. *J Neurol Neurosurg Psychiatry*. 2019;90(5):590-598. doi:10.1136/jnnp-2018-319148

Simonsen AH, Herukka SK, Andreasen N, et al. Recommendations for CSF AD biomarkers in the diagnostic evaluation of dementia. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2017;13(3):274-284. doi:10.1016/j.jalz.2016.09.008

Simrén J, Leuzy A, Karikari TK, et al. The diagnostic and prognostic capabilities of plasma biomarkers in Alzheimer's disease. *Alzheimer's & Dementia*. 2021;17(7):1145-1156. doi:10.1002/alz.12283

Sorbi S, Hort J, Erkinjuntti T, et al. EFNS-ENS Guidelines on the diagnosis and management of disorders associated with dementia. *Eur J Neurol*. 2012;19(9):1159-79. doi:10.1111/j.1468-1331.2012.03784.x

Talwar P, Sinha J, Grover S, et al. Meta-analysis of apolipoprotein E levels in the cerebrospinal fluid of patients with Alzheimer's disease. *Journal of the neurological sciences*. 2016;360:179-87. doi:10.1016/j.jns.2015.12.004

Tatebe H, Kasai T, Ohmichi T, et al. Quantification of plasma phosphorylated tau to use as a biomarker for brain Alzheimer's pathology: pilot case-control studies including patients with Alzheimer's disease and down syndrome. *Molecular neurodegeneration*. 2017;12(1):63. doi:10.1186/s13024-017-0206-8

Van Cauwenberghe C, Van Broeckhoven C, Sleegers K. The genetic landscape of Alzheimer's disease: clinical implications and perspectives. *Genet Med*. 2016;18(5):421-30. doi:10.1038/gim.2015.117

Wang H, Stewart T, Toledo JB, et al. A Longitudinal Study of Total and Phosphorylated alpha-Synuclein with Other Biomarkers in Cerebrospinal Fluid of Alzheimer's Disease and Mild Cognitive Impairment. *Journal of Alzheimer's disease : JAD*. 2018;61(4):1541-1553. doi:10.3233/jad-171013

Wang S, Kojima K, Mobley JA, West AB. Proteomic analysis of urinary extracellular vesicles reveal biomarkers for neurologic disease. *EBioMedicine*. 2019;45:351-361. doi:10.1016/j.ebiom.2019.06.021

Waqar S, Khan H, Zulfiqar SK, Ahmad A. Skin Biopsy as a Diagnostic Tool for Synucleinopathies. *Cureus*. 2023;15(10):e47179. doi:10.7759/cureus.47179

WHO. Dementia. https://www.who.int/news-room/fact-sheets/detail/dementia

WHO. Preferred product characteristics of blood-based biomarker diagnostics for Alzheimer's disease. https://www.who.int/publications/i/item/9789240099067

Wolk DA, Dickerson BC. Clinical features and diagnosis of Alzheimer's disease. Updated September 30, 2024. https://www.uptodate.com/contents/clinical-features-and-diagnosis-of-Alzheimer's-disease

Yoong SQ, Lu J, Xing H, Gyanwali B, Tan YQ, Wu XV. The prognostic utility of CSF neurogranin in predicting future cognitive decline in the Alzheimer's disease continuum: A systematic review and meta-analysis with narrative synthesis. *Ageing Res Rev*. 2021;72:101491. doi:10.1016/j.arr.2021.101491

Zhang J, Zhang CH, Li RJ, et al. Accuracy of urinary AD7c-NTP for diagnosing Alzheimer's disease: a systematic review and meta-analysis. *Journal of Alzheimer's disease : JAD*. 2014;40(1):153-9. doi:10.3233/jad-131445