Humana® Claims Payment Policy

Subject: Laboratory – In Vitro Chemo-Resistance and Chemo-Sensitivities Assay

Application: Medicare Advantage Products
Policy number: LAB2023012
Published date: 04/2023
Revision date: 04/2026

Disclaimer: *The intended audience of this medical claims payment policy is healthcare providers who treat Humana members. This policy is made available to provide information on certain Humana claims payment policies. This policy is a guideline only and does not constitute medical advice, guarantee of payment, plan pre-authorization, an explanation of benefits, or a contract. This policy does not govern whether a procedure is covered under a specific member plan or policy, nor is it intended to address every claim situation. Claims may be affected by other factors, such as: state and federal laws and regulations, provider contract terms, and our professional judgment. This policy is subject to change or termination by Humana. Humana has full and final discretionary authority for its interpretation and application. No part of this policy may be reproduced, stored in a retrieval system or transmitted, in any shape or form or by any means, electronic, mechanical, photocopying or otherwise without express written permission from Humana. When printed, this version becomes uncontrolled. For the most current information, always refer to the following website:* ***Humana.com/ClaimPaymentPolicies***

Medicare Advantage Payment Policy

As a Medicare Advantage plan, we have developed payment policies to explain coverage more fully for services. The criteria outlined will apply to Medicare Advantage plans except when the procedure(s) has established criteria through the Original Medicare National Coverage Determinations (NCDs) and Local Coverage Determinations (LCDs) as outlined below.

In addition to the policy, claims payments are subject to other plan requirements for the processing and payment of claims, including, but not limited to, requirements of medical necessity and reasonableness and applicable referral or authorization requirements.

- In vitro chemotherapy sensitivity and resistance assays (CPT code 81535**, CPT code 81536**, CPT code 0083U^, CPT code 0248U^^, CPT code 0249U*, CPT code 0285U*, CPT code 0435U^^) **ARE NOT REIMBURSABLE.**

Applicable procedures for Medicare Advantage plans

Procedure code	Code description
81535**	Oncology (gynecologic), live tumor cell culture and chemotherapeutic response by DAPI stain and morphology, predictive algorithm reported as a drug response score; first single drug or drug combination Proprietary test: ChemoFX® Lab/manufacturer: Helomics, Corp
81536**	Oncology (gynecologic), live tumor cell culture and chemotherapeutic response by DAPI stain and morphology, predictive algorithm reported as a drug response score; each additional single drug or drug combination (List separately in addition to code for primary procedure) Proprietary test: ChemoFX® Lab/manufacturer: Helomics, Corp
0083U^	Oncology, response to chemotherapy drugs using motility contrast tomography, fresh or frozen tissue, reported as likelihood of sensitivity or resistance to drugs or drug combinations Proprietary test: Onco4D™ Lab/Manufacturer: Animated Dynamics, Inc
0248U^^	Oncology, spheroid cell culture in a 3D microenvironment, 12 drug panel, brain- or brain metastasis-response prediction for each drug Proprietary test: 3D Predict Glioma Lab/Manufacturer: KIYATEC®, Inc
0249U*	Oncology (breast), semiquantitative analysis of 32 phosphoproteins and protein analytes, includes laser capture microdissection, with algorithmic analysis and interpretative report Proprietary test: Theralink® Reverse Phase Protein Array (RPPA) Lab/Manufacturer: Theralink® Technologies, Inc
0285U*	Oncology, disease progression and response monitoring to radiation, chemotherapy, or other systematic cancer treatments, cell-free DNA, quantitative branched chain DNA amplification, plasma, reported in ng/mL Proprietary test: RadTox™ cfDNA test Lab/Manufacturer: DiaCarta Clinical Lab/DiaCarta Inc

Procedure code	Code description
0435U^^	Oncology, chemotherapeutic drug cytotoxicity assay of cancer stem cells (CSCs), from cultured CSCs and primary tumor cells, categorical drug response reported based on cytotoxicity percentage observed, minimum of 14 drugs or drug combinations Proprietary test: ChemoID® Lab/Manufacturer: Cordgenics, LLC

**The payment criterion for 0249U and 0285U applies to all Medicare jurisdictions except for states awarded to CGS Administrators, LLC., Noridian Healthcare Solutions, LLC., Palmetto GBA and Wisconsin Physician Service Insurance Corporation.*
***The payment criterion for 81535 and 81536 applies to all Medicare jurisdictions except for states awarded to National Government Services, Inc.*
^The payment criterion for 0083U applies to all Medicare jurisdictions except for states awarded to Noridian Healthcare Solutions, LLC. and Palmetto GBA.
^^The payment criterion for 0248U and 0435U applies to all Medicare jurisdictions, except for states awarded to Palmetto GBA.

Original Medicare Local Coverage Determinations (LCDs)

Contractor name	Policy title	Policy number	Procedure code
CGS Administrators, LLC	MolDX: Molecular Diagnostic Tests (MDT)	L36021	0249U 0285U
National Government Services, Inc.	Molecular Pathology Procedures	L35000	81535 81536
Noridian Healthcare Solutions, LLC	MolDX: Molecular Diagnostic Tests (MDT)	L35160	0249U 0285U
Noridian Healthcare Solutions, LLC	MolDX: Molecular Diagnostic Tests (MDT)	L36256	0249U 0285U
Noridian Healthcare Solutions, LLC	In Vitro Chemosensitivity & Chemoresistance Assays	L37628	0083U
Palmetto GBA	In Vitro Chemosensitivity & Chemoresistance Assays	L34554	0083U 0248U 0435U
Palmetto GBA	MolDX: Molecular Diagnostic Tests (MDT)	L35025	0249U 0285U
Wisconsin Physicians Service Insurance Corporation	MolDX: Molecular Diagnostic Tests (MDT)	L36807	0249U 0285U

References

Brower, S. L., Fensterer, J. E., & Bush, J. E. (2008). The ChemoFx® Assay: An Ex Vivo Chemosensitivity and Resistance Assay for Predicting Patient Response to Cancer Chemotherapy. In G. Mor & A. B. Alvero (Eds.), *Apoptosis and Cancer: Methods and Protocols* (pp. 57-78). Humana Press. https://doi.org/10.1007/978-1-59745-339-4_6

Burstein, H. J., Mangu, P. B., Somerfield, M. R., Schrag, D., Samson, D., Holt, L., Zelman, D., & Ajani, J. A. (2011). American Society of Clinical Oncology clinical practice guideline update on the use of chemotherapy sensitivity and resistance assays. *J Clin Oncol*, *29*(24), 3328-3330. https://doi.org/10.1200/jco.2011.36.0354

Chen, Z., Zhang, S., Ma, S., Li, C., Xu, C., Shen, Y., Zhao, J., & Miao, L. (2018). Evaluation of the in vitro Chemosensitivity and Correlation with Clinical Outcomes in Lung Cancer using the ATP-TCA. *Anticancer Agents Med Chem*, *18*(1), 139-145. https://doi.org/10.2174/1871520617666170419123713

Grendys, E. C., Jr., Fiorica, J. V., Orr, J. W., Jr., Holloway, R., Wang, D., Tian, C., Chan, J. K., & Herzog, T. J. (2014). Overview of a chemoresponse assay in ovarian cancer. *Clin Transl Oncol*, *16*(9), 761-769. https://doi.org/10.1007/s12094-014-1192-8

Hatok, J., Babusikova, E., Matakova, T., Mistuna, D., Dobrota, D., & Racay, P. (2009). In vitro assays for the evaluation of drug resistance in tumor cells. *Clin Exp Med*, *9*(1), 1-7. https://doi.org/10.1007/s10238-008-0011-3

Hoffman, R. M. (2018). Clinical Correlation of the Histoculture Drug Response Assay in Gastrointestinal Cancer. *Methods Mol Biol*, *1760*, 61-72. https://doi.org/10.1007/978-1-4939-7745-1_7

Howard, C. M., Valluri, J., Alberico, A., Julien, T., Mazagri, R., Marsh, R., Alastair, H., Cortese, A., Griswold, M., Wang, W., Denning, K., Brown, L., & Claudio, P. P. (2017). Analysis of Chemopredictive Assay for Targeting Cancer Stem Cells in Glioblastoma Patients. *Transl Oncol*, *10*(2), 241-254. https://doi.org/10.1016/j.tranon.2017.01.008

Karam, A. K., Chiang, J. W., Fung, E., Nossov, V., & Karlan, B. Y. (2009). Extreme drug resistance assay results do not influence survival in women with epithelial ovarian cancer. *Gynecologic Oncology*, *114*(2), 246-252. https://doi.org/10.1016/j.ygyno.2009.02.022

Kiyatec. (2023). 3D Predict Platform. https://kiyatec.com/healthcare-professionals/3d-predict-platform/

Krivak, T. C., Lele, S., Richard, S., Secord, A. A., Leath, C. A., 3rd, Brower, S. L., Tian, C., & Moore, R. G. (2014). A chemoresponse assay for prediction of platinum resistance in primary ovarian cancer. *Am J Obstet Gynecol*, *211*(1), 68.e61-68. https://doi.org/10.1016/j.ajog.2014.02.009

Kwon, H. Y., Kim, I. K., Kang, J., Sohn, S. K., & Lee, K. Y. (2016). In Vitro Adenosine Triphosphate-Based Chemotherapy Response Assay as a Predictor of Clinical Response to Fluorouracil-Based Adjuvant Chemotherapy in Stage II Colorectal Cancer. *Cancer Res Treat*, *48*(3), 970-977. https://doi.org/10.4143/crt.2015.140

NCCN. (2024a). *NCCN Clinical Practice Guidelines in Oncology*. https://www.nccn.org/professionals/physician_gls/default.aspx

NCCN. (2024b). *NCCN Clinical Practice Guidelines in Oncology; Ovarian Cancer v 3.2024* https://www.nccn.org/professionals/physician_gls/pdf/ovarian.pdf

Pierian. (2023). Products: ChemoINTEL™. https://pierianbio.com/project/chemo-intel/

RGCC. (2023). *Onconomics RGCC*. https://www.rgcc-group.com/tests/onconomics-plus-rgcc/

Rutherford, T., Orr, J., Jr., Grendys, E., Jr., Edwards, R., Krivak, T. C., Holloway, R., Moore, R. G., Puls, L., Tillmanns, T., Schink, J. C., Brower, S. L., Tian, C., & Herzog, T. J. (2013). A prospective study evaluating the clinical relevance of a chemoresponse assay for treatment of patients with persistent or recurrent ovarian cancer. *Gynecol Oncol*, *131*(2), 362-367. https://doi.org/10.1016/j.ygyno.2013.08.009

Schrag, D., Garewal, H. S., Burstein, H. J., Samson, D. J., Von Hoff, D. D., & Somerfield, M. R. (2004). American Society of Clinical Oncology Technology Assessment: chemotherapy sensitivity and resistance assays. *J Clin Oncol*, *22*(17), 3631-3638. https://doi.org/10.1200/jco.2004.05.065

Shuford, S., Lipinski, L., Abad, A., Smith, A. M., Rayner, M., O'Donnell, L., Stuart, J., Mechtler, L. L., Fabiano, A. J., Edenfield, J., Kanos, C., Gardner, S., Hodge, P., Lynn, M., Butowski, N. A., Han, S. J., Redjal, N., Crosswell, H. E., Vibat, C. R. T., . . . DesRochers, T. M. (2021). Prospective prediction of clinical drug response in high-grade gliomas using an ex vivo 3D cell culture assay. *Neurooncol Adv*, *3*(1), vdab065. https://doi.org/10.1093/noajnl/vdab065

Strickland, S. A., Raptis, A., Hallquist, A., Rutledge, J., Chernick, M., Perree, M., Talbott, M. S., & Presant, C. A. (2013). Correlation of the microculture-kinetic drug-induced apoptosis assay with patient outcomes in initial treatment of adult acute myelocytic leukemia. *Leuk Lymphoma*, *54*(3), 528-534. https://doi.org/10.3109/10428194.2012.722217

Tatar, B., Boyraz, G., Selçuk, İ., Doğan, A. K., Usubütün, A., & Tuncer, Z. S. (2016). In vitro chemosensitivity in ovarian carcinoma: Comparison of three leading assays. In *J Turk Ger Gynecol Assoc* (Vol. 17, pp. 35-40). https://doi.org/10.5152/jtgga.2016.16017

Theralink. (2023). Theralink: Precision Medicine for Life. https://theralink.com/theralink