Humana. Claims Payment Policy

Subject: Laboratory – Serum Tumor Markers for Malignancies

Application: Medicare Advantage Products
Policy number: LAB2023051
Published date: 07/2023
Revision date: 04/2026

Disclaimer: *The intended audience of this medical claims payment policy is healthcare providers who treat Humana members. This policy is made available to provide information on certain Humana claims payment policies. This policy is a guideline only and does not constitute medical advice, guarantee of payment, plan pre-authorization, an explanation of benefits, or a contract. This policy does not govern whether a procedure is covered under a specific member plan or policy, nor is it intended to address every claim situation. Claims may be affected by other factors, such as: state and federal laws and regulations, provider contract terms, and our professional judgment. This policy is subject to change or termination by Humana. Humana has full and final discretionary authority for its interpretation and application. No part of this policy may be reproduced, stored in a retrieval system or transmitted, in any shape or form or by any means, electronic, mechanical, photocopying or otherwise without express written permission from Humana. When printed, this version becomes uncontrolled. For the most current information, always refer to the following website:* ***Humana.com/ClaimPaymentPolicies***

Medicare Advantage Payment Policy

As a Medicare Advantage plan, we have developed payment policies to explain coverage more fully for services. The criteria outlined will apply to Medicare Advantage plans except when the procedure(s) has established criteria through the Original Medicare National Coverage Determinations (NCDs) and Local Coverage Determinations (LCDs) as outlined below.

In addition to the policy, claims payments are subject to other plan requirements for the processing and payment of claims, including, but not limited to, requirements of medical necessity and reasonableness and applicable referral or authorization requirements.

- Serum tumor markers **ARE REIMBURSABLE** for the following indications:

  – Alkaline Phosphatase (ALP) (CPT code 84078, CPT code 84080) every 90 days for:
    - Metastatic and primary bone neoplasms
    - Systemic light chain amyloidosis
    - Uveal melanoma

  – B-type natriuretic peptide (BNP) or N-terminal fragment of B-type natriuretic peptide (NT-proBNP) (CPT code 83880*) every 90 days for:
    - Multiple myeloma

  – Beta human chorionic gonadotropin (beta-HCG) (CPT code 84703, CPT code 84704) every 60 days for:
    - Gestational trophoblastic neoplasia
    - Occult primary mass of the mediastinum or retroperitoneum
    - Ovarian cancer, fallopian tube cancer or primary peritoneal cancer
      - Borderline epithelial tumors
      - Malignant germ cell tumors
      - Malignant sex cord stromal tumors
    - Testicular cancer
    - Thymoma

  – Beta-2 microglobulin (B2M) (CPT code 82232) every 90 days for:
    - B-cell lymphoma or Castleman disease
    - Chronic lymphocytic leukemia or small lymphocytic lymphoma
    - Multiple myeloma
    - Systemic light chain amyloidosis
    - Waldenström's macroglobulinemia or lymphoplasmacytic lymphoma

  – Calcitonin (CALCA) expression (CPT code 82308) every 60 days for:
    - Adenocarcinoma or anaplastic epithelial tumors

  - Multiple endocrine neoplasia, type 2
  - Occult primary adenocarcinoma, anaplastic or undifferentiated tumors of the head and neck
  - Thyroid carcinoma or medullary carcinoma

- Circulating serum kappa or lambda free light chains (CPT code 83521) every 90 days for:
  - Castleman disease
  - Multiple myeloma
  - Systemic light chain amyloidosis

- Human epididymis protein 4 (HE4) expression (CPT code 86305) every 60 days for:
  - Ovarian cancer, fallopian tube cancer or primary peritoneal cancer
    - Borderline epithelial tumors
    - Carcinosarcoma, also known as malignant mixed Müllerian tumors
    - Clear cell carcinoma of the ovary
    - Grade 1 endometrioid carcinoma
    - Low-grade serous carcinoma
    - Mucinous neoplasms of the ovary
    - Malignant germ cell tumors
    - Malignant sex cord stromal tumors

- Inhibin (INHA) expression (CPT code 86336) every 60 days for:
  - Ovarian cancer, fallopian tube cancer or primary peritoneal cancer
    - Borderline epithelial tumors
    - Carcinosarcoma, also known as malignant mix mullerian tumors
    - Clear cell carcinoma of the ovary
    - Grade 1 endometrioid carcinoma
    - Low-grade serous carcinoma
    - Mucinous neoplasms of the ovary
    - Malignant germ cell tumors
    - Malignant sex cord stromal tumors
  - Unspecified occult primary adenocarcinoma or carcinoma

- Troponin T (CPT code 84484) for systemic light chain amyloidosis

- All other applications of serum tumor markers (CPT code 82107, CPT code 83950, CPT code 83951, CPT code 86316**) **ARE NOT REIMBURSABLE**.

- The following tests **ARE NOT REIMBURSABLE**:

  - BeScreened™-CRC (CPT code 0163U)
  - Divitum® TKa (CPT code 0404U)
  - IGoCheck™ (CPT code 0558U)
  - MammoCheck™ Baseline (CPT code 0559U)
  - Overa OVA1 Next Generation (CPT code 0003U***)
  - PancreaSure™ (CPT code 0599U****)
  - REVEAL Lung Nodule Characterization (CPT code 0092U****)

- Analysis of proteomic patterns in serum (CPT code 83789) for screening and detection of cancer **IS NOT REIMBURSABLE**.

Applicable procedures for Medicare Advantage plans

Procedure code	Code description
82107	Alpha-fetoprotein (AFP); AFP-L3 fraction isoform and total AFP (including ratio)
82232	Beta-2 microglobulin
82308	Calcitonin
83521	Immunoglobulin light chains (ie, kappa, lambda), free, each
83789	Mass spectrometry and tandem mass spectrometry (e.g., MS, MS/MS, MALDI, MS-TOF, QTOF), non-drug analyte(s) not elsewhere specified, qualitative or quantitative, each specimen
83880*	Natriuretic peptide
83950	Oncoprotein; HER-2/neu
83951	Oncoprotein; des-gamma-carboxy-prothrombin (DCP)
84078	Phosphatase, alkaline; heat stable (total not included)
84080	Phosphatase, alkaline; isoenzymes
84484	Troponin, quantitative
84703	Gonadotropin, chorionic (hCG); qualitative
84704	Gonadotropin, chorionic (hCG); free beta chain
86305	Human epididymis protein 4 (HE4)
86316**	Immunoassay for tumor antigen, other antigen, quantitative (e.g., CA 50, 72-4, 549), each
86336	Inhibin A
0003U***	Oncology (ovarian) biochemical assays of five proteins (apolipoprotein A-1, CA 125 II, follicle stimulating hormone, human epididymis protein 4, transferrin), utilizing serum, algorithm reported as a likelihood score Proprietary test: Overa™ (OVA1 Next Generation) Lab/manufacturer: Aspira Labs, Inc, Vermillion, Inc
0092U****	Oncology (lung), three protein biomarkers, immunoassay using magnetic nanosensor technology, plasma, algorithm reported as risk score for likelihood of malignancy Proprietary test: REVEAL Lung Nodule Characterization Lab/manufacturer: MagArray, Inc
0163U	Oncology (colorectal) screening, biochemical enzyme-linked immunosorbent assay (ELISA) of 3 plasma or serum proteins (teratocarcinoma derived growth factor-1 [TDGF-1, Cripto-1], carcinoembryonic antigen [CEA], extracellular matrix protein [ECM]), with demographic data (age, gender, CRC-screening compliance) using a proprietary algorithm and reported as likelihood of CRC or advanced adenomas Proprietary test: BeScreened™-CRC Lab/Manufacturer: Beacon Biomedical Inc
0404U	Oncology (breast), semiquantitative measurement of thymidine kinase activity by immunoassay, serum, results reported as risk of disease progression Proprietary test: DiviTum®TKa Lab/Manufacturer: Biovica Inc, Biovica International AB
0558U	Oncology (colorectal), quantitative enzyme-linked immunosorbent assay (ELISA) for secreted colorectal cancer protein marker (BF7 antigen), using serum, result reported as indicative of response/no response to therapy or disease progression/regression Proprietary test: IGoCheck™ Lab/Manufacturer: Milagen, Inc, Milagen, Inc
0559U	Oncology (breast), quantitative enzyme-linked immunosorbent assay (ELISA) for secreted breast cancer protein marker (BF9 antigen), serum, result reported as indicative of response/no response to therapy or disease progression/regression Proprietary test: MammoCheck™ (Blood-Based Breast Cancer Test) Lab/Manufacturer: Milagen, Inc, Milagen, Inc
0599U****	Oncology (pancreatic cancer), multiplex immunoassay of ICAM1, TIMP1, CTSD, THBS1, and CA 19-9, serum, diagnostic algorithm reported as positive or negative

Procedure code	Code description
	Proprietary test: PancreaSure™ Lab/Manufacturer: Immunovia, Inc, Immunovia, Inc

**The payment criterion for 83880 applies to all Medicare jurisdictions, except those states awarded to CGS Administrators, First Coast Service Options, National Government Services, Noridian Healthcare Solutions, Palmetto GBA, and Wisconsin Physicians Service Insurance Corporation.*
***The payment criterion for 86316 applies to all Medicare jurisdictions, except those states awarded to CGS Administrators, Noridian Healthcare Solutions, and Palmetto GBA.*
****The payment criterion for 0003U applies to all Medicare jurisdictions, except those states awarded to CGS Administrators, National Government Services, Noridian Healthcare Solutions, Palmetto GBA, and Wisconsin Physicians Service Insurance Corporation.*
*****The payment criterion for 0092U and 0599U applies to all Medicare jurisdictions, except those states awarded to CGS Administrators, Noridian Healthcare Solutions, Palmetto GBA, and Wisconsin Physicians Service Insurance Corporation.*

Original Medicare Local Coverage Determinations (LCDs)

Contractor name	Policy title	Policy number	Procedure code
CGS Administrators, LLC	Bladder/Urothelial Tumor Markers	L36975	86316
CGS Administrators, LLC	B-type Natriuretic Peptide (BNP) Testing	L33943	83880
First Coast Service Options, Inc.	B-Type Natriuretic Peptide (BNP)	L33267	83880
National Government Services, Inc.	B-type Natriuretic Peptide (BNP) Testing	L33573	83880
Noridian Healthcare Solutions, LLC	B-type Natriuretic Peptide (BNP) Testing	L34038	83880
Noridian Healthcare Solutions, LLC	B-type Natriuretic Peptide (BNP) Testing	L35526	83880
Noridian Healthcare Solutions, LLC	Lab: Bladder/Urothelial Tumor Markers	L36678	86316
Palmetto GBA	Brain Natriuretic Peptide (BNP) Level	L33422	83880
Palmetto GBA	B-type Natriuretic Peptide (BNP) Testing	L34410	83880
Palmetto GBA	Lab: Bladder/Urothelial Tumor Markers	L33420	86316
Wisconsin Physicians Service Insurance Corporation	MolDX: Biomarkers in Cardiovascular Risk Assessment	L36523	83880
CGS Administrators, LLC	MolDX: Molecular Diagnostic Tests (MDT)	L36021	0003U 0092U 0599U
Noridian Healthcare Solutions, LLC	MolDX: Molecular Diagnostic Tests (MDT)	L35160	0003U 0092U 0599U
Noridian Healthcare Solutions, LLC	MolDX: Molecular Diagnostic Tests (MDT)	L36256	0003U 0092U 0599U
Palmetto GBA	MolDX: Molecular Diagnostic Tests (MDT)	L35025	0003U 0092U 0599U
Wisconsin Physicians Service Insurance Corporation	MolDX: Molecular Diagnostic Tests (MDT)	L36807	0003U 0092U 0599U

Contractor name	Policy title	Policy number	Procedure code
National Government Services, Inc.	Multimarker Serum Tests Related to Ovarian Cancer Testing	L38371	0003U

References

AAAAI. Systemic Mastocytosis. https://www.aaaai.org/conditions-treatments/related-conditions/systemic-mastocytosis

ACS. What Is Multiple Myeloma? https://www.cancer.org/cancer/multiple-myeloma/about/what-is-multiple-myeloma.html

ACS. What Is Waldenstrom Macroglobulinemia? https://www.cancer.org/cancer/waldenstrom-macroglobulinemia/about/what-is-wm.html

ADLM. Human Chorionic Gonadotropin (hCG). https://myadlm.org/cln/articles/2021/april/using-human-chorionic-gonadotropin-as-a-tumor-marker

ADLM. Serum Free Light Chains: Optimal Testing Recommendations. https://www.myadlm.org/advocacy-and-outreach/optimal-testing-guide-to-lab-test-utilization/g-s/serum-free-light-chains

Akar H, Seldin DC, Magnani B, et al. Quantitative serum free light chain assay in the diagnostic evaluation of AL amyloidosis. *Amyloid*. Dec 2005;12(4):210-5. doi:10.1080/13506120500352339

Aspira Health. Ova1Plus®. https://aspirawh.com/ova1plus/

ASPIRA. OvaWatch. https://aspirawh.com/ovawatch/

ATA. Revised ATA Management Guidelines for MTC. https://www.thyroid.org/wp-content/uploads/2017/03/revised-ata-management-guidelines-for-MTC.pdf

Berrebi A, Shvidel L, Arditti FD, Bassous L, Haran M, Shtalrid M. The Significance of Elevated Beta 2-Microglobulin (b2-m) in B-CLL: Evidence of in Vitro b2-m Secretion Following Activation of B-CLL Cells. *Blood*. 2009;114(22):4380.

BeScreened. BeScreened. https://bescreened.com/

Bhole MV, Sadler R, Ramasamy K. Serum-free light-chain assay: clinical utility and limitations. *Ann Clin Biochem*. Sep 2014;51(Pt 5):528-42. doi:10.1177/0004563213518758

Bind MK, Mishra RR, Kumar V, Misra V, Singh PA. Serum CA 19-9 and CA 125 as a diagnostic marker in carcinoma of gallbladder. *Indian J Pathol Microbiol*. Jan-Mar 2021;64(1):65-68.

Bowlus CL, Arrive L, Bergquist A, et al. AASLD practice guidance on primary sclerosing cholangitis and cholangiocarcinoma. *Hepatology*. Feb 1 2023;77(2):659-702. doi:10.1002/hep.32771

Cautha S, Gupta S, Hanif A, Moirangthem V, Jain K. Lymphoplasmacytic Lymphoma with Only Lambda Light Chain Monoclonal Paraprotein Expression. *Eur J Case Rep Intern Med*. 2022;9(2):003106. doi:10.12890/2022_003106

Caviglia GP, Abate ML, Petrini E, Gaia S, Rizzetto M, Smedile A. Highly sensitive alpha-fetoprotein, Lens culinaris agglutinin-reactive fraction of alpha-fetoprotein and des-gamma-carboxyprothrombin for hepatocellular carcinoma detection. *Hepatology research : the official journal of the Japan Society of Hepatology*. Mar 2016;46(3):E130-5. doi:10.1111/hepr.12544

Chappuis PO, Dieterich B, Sciretta V, et al. Functional evaluation of plasmin formation in primary breast cancer. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. May 15 2001;19(10):2731-8. doi:10.1200/jco.2001.19.10.2731

Chaulin AM. Biology of Cardiac Troponins: Emphasis on Metabolism. *Biology (Basel)*. Mar 11 2022;11(3)doi:10.3390/biology11030429

Chen F, Shen J, Wang J, Cai P, Huang Y. Clinical analysis of four serum tumor markers in 458 patients with ovarian tumors: diagnostic value of the combined use of HE4, CA125, CA19-9, and CEA in ovarian tumors. *Cancer Manag Res*. 2018;10:1313-1318. doi:10.2147/cmar.S155693

Chen Y, Xie Y, Xu L, et al. Protein content and functional characteristics of serum-purified exosomes from patients with colorectal cancer revealed by quantitative proteomics. *International journal of cancer*. 2017/02/15 2017;140(4):900-913. doi:10.1002/ijc.30496

Cheng J, Wang W, Zhang Y, et al. Prognostic role of pre-treatment serum AFP-L3% in hepatocellular carcinoma: systematic review and meta-analysis. *PloS one*. 2014;9(1):e87011. doi:10.1371/journal.pone.0087011

Di Castelnuovo A, Veronesi G, Costanzo S, et al. NT-proBNP (N-Terminal Pro-B-Type Natriuretic Peptide) and the Risk of Stroke. *Stroke*. Mar 2019;50(3):610-617. doi:10.1161/STROKEAHA.118.023218

Dispenzieri A. Clinical presentation, laboratory manifestations, and diagnosis of immunoglobulin light chain (AL) amyloidosis. Updated February 19, 2025. https://www.uptodate.com/contents/clinical-presentation-laboratory-manifestations-and-diagnosis-of-immunoglobulin-light-chain-al-amyloidosis

Dorigo O, Berek JS. Personalizing CA125 levels for ovarian cancer screening. *Cancer prevention research (Philadelphia, Pa)*. Sep 2011;4(9):1356-9. doi:10.1158/1940-6207.Capr-11-0378

Duffy MJ. Carcinoembryonic Antigen as a Marker for Colorectal Cancer: Is It Clinically Useful? *Clinical Chemistry*. 2001;47(4):624. doi:10.1093/clinchem/47.4.624

Farkkila A, Koskela S, Bryk S, et al. The clinical utility of serum anti-Mullerian hormone in the follow-up of ovarian adult-type granulosa cell tumors--A comparative study with inhibin B. *International journal of cancer*. Oct 1 2015;137(7):1661-71. doi:10.1002/ijc.29532

Febbo PG, Ladanyi M, Aldape KD, et al. NCCN Task Force report: Evaluating the clinical utility of tumor markers in oncology. *Journal of the National Comprehensive Cancer Network : JNCCN*. Nov 2011;9 Suppl 5:S1-32; quiz S33. doi:10.6004/jnccn.2011.0137

Feng F, Tian Y, Xu G, et al. Diagnostic and prognostic value of CEA, CA19-9, AFP and CA125 for early gastric cancer. *BMC cancer*. Nov 9 2017;17(1):737. doi:10.1186/s12885-017-3738-y

Foekens JA, Peters HA, Look MP, et al. The urokinase system of plasminogen activation and prognosis in 2780 breast cancer patients. *Cancer Res*. Feb 01 2000;60(3):636-43.

Foukakis T, Bergh J. Prognostic and predictive factors in early, nonmetastatic breast cancer - UpToDate. Updated March 17, 2025. https://www.uptodate.com/contents/prognostic-and-predictive-factors-in-early-non-metastatic-breast-cancer

Gershenson D. Sex cord-stromal tumors of the ovary: Epidemiology, clinical features, and diagnosis in adults. Updated July 8, 2024. https://www.uptodate.com/contents/sex-cord-stromal-tumors-of-the-ovary-epidemiology-clinical-features-and-diagnosis-in-adults

Gilligan TD, Seidenfeld J, Basch EM, et al. American Society of Clinical Oncology Clinical Practice Guideline on uses of serum tumor markers in adult males with germ cell tumors. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. Jul 10 2010;28(20):3388-404. doi:10.1200/jco.2009.26.4481

Harris LN, Ismaila N, McShane LM, et al. Use of Biomarkers to Guide Decisions on Adjuvant Systemic Therapy for Women With Early-Stage Invasive Breast Cancer: American Society of Clinical Oncology Clinical Practice Guideline. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. Apr 01 2016;34(10):1134-50. doi:10.1200/jco.2015.65.2289

Harvey RA. Human chorionic gonadotropin: Biochemistry and measurement in pregnancy and disease. Updated June 23, 2023. https://www.uptodate.com/contents/human-chorionic-gonadotropin-biochemistry-and-measurement-in-pregnancy-and-disease

Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid : official journal of the American Thyroid Association*. Jan 2016;26(1):1-133. doi:10.1089/thy.2015.0020

Hotakainen K, Ljungberg B, Paju A, Rasmuson T, Alfthan H, Stenman UH. The free beta-subunit of human chorionic gonadotropin as a prognostic factor in renal cell carcinoma. *British journal of cancer*. Jan 21 2002;86(2):185-9. doi:10.1038/sj.bjc.6600050

Hottinger A, Hormigo A. Serum Biomarkers. *Encyclopedia of Cancer*. Springer Berlin Heidelberg; 2011. http://link.springer.com/referenceworkentry/10.1007/978-3-642-16483-5_5269

Isaksson S, Jönsson P, Monsef N, et al. CA 19-9 and CA 125 as potential predictors of disease recurrence in resectable lung adenocarcinoma. *PloS one*. 2017;12(10):e0186284. doi:10.1371/journal.pone.0186284

Katou H, Kanno T, Hoshino M, et al. The role of disulfide bond in the amyloidogenic state of beta(2)-microglobulin studied by heteronuclear NMR. *Protein science : a publication of the Protein Society*. Sep 2002;11(9):2218-29. doi:10.1110/ps.0213202

Katzmann JA, Clark RJ, Abraham RS, et al. Serum reference intervals and diagnostic ranges for free kappa and free lambda immunoglobulin light chains: relative sensitivity for detection of monoclonal light chains. *Clin Chem*. Sep 2002;48(9):1437-44.

Kim NH, Lee MY, Park JH, et al. Serum CEA and CA 19-9 Levels are Associated with the Presence and Severity of Colorectal Neoplasia. *Yonsei medical journal*. Sep 1 2017;58(5):918-924. doi:10.3349/ymj.2017.58.5.918

Kindler HL, Ismaila N, Armato SG, et al. Treatment of Malignant Pleural Mesothelioma: American Society of Clinical Oncology Clinical Practice Guideline. *Journal of Clinical Oncology*. 2018/05/01 2018;36(13):1343-1373. doi:10.1200/JCO.2017.76.6394

Kumar S, Dispenzieri A, Katzmann JA, et al. Serum immunoglobulin free light-chain measurement in primary amyloidosis: prognostic value and correlations with clinical features. *Blood*. Dec 9 2010;116(24):5126-9. doi:10.1182/blood-2010-06-290668

Kyrtsonis MC KE, Bartzis V, Pessah I, Nikolaou E, Karalis V, Maltezas D, Panayiotidis P, Harding S. Monoclonal Immunoglobulin. *Multiple Myeloma - A Quick Reflection on the Fast Progress*. 2012.

Leru PM. Evaluation and Classification of Mast Cell Disorders: A Difficult to Manage Pathology in Clinical Practice. *Cureus*. Feb 2022;14(2):e22177. doi:10.7759/cureus.22177

Li AJ. Serum biomarkers for evaluation of an adnexal mass for epithelial carcinoma of the ovary, fallopian tube, or peritoneum. Updated October 31, 2024. https://www.uptodate.com/contents/serum-biomarkers-for-evaluation-of-an-adnexal-mass-for-epithelial-carcinoma-of-the-ovary-fallopian-tube-or-peritoneum

Li J, Yin M, Song W, et al. B Subunit of Human Chorionic Gonadotropin Promotes Tumor Invasion and Predicts Poor Prognosis of Early-Stage Colorectal Cancer. *Cellular physiology and biochemistry : international journal of experimental cellular physiology, biochemistry, and pharmacology*. 2018;45(1):237-249. doi:10.1159/000486770

Lucarelli G, Ditonno P, Bettocchi C, et al. Diagnostic and prognostic role of preoperative circulating CA 15-3, CA 125, and beta-2 microglobulin in renal cell carcinoma. *Disease markers*. 2014;2014:689795. doi:10.1155/2014/689795

Magnani JL. The discovery, biology, and drug development of sialyl Lea and sialyl Lex. *Archives of Biochemistry and Biophysics*. 2004/06/15/ 2004;426(2):122-131. doi:10.1016/j.abb.2004.04.008

Malmstrom P, Bendahl PO, Boiesen P, Brunner N, Idvall I, Ferno M. S-phase fraction and urokinase plasminogen activator are better markers for distant recurrences than Nottingham Prognostic Index and histologic grade in a prospective study of premenopausal lymph node-negative breast cancer. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. Apr 01 2001;19(7):2010-9. doi:10.1200/jco.2001.19.7.2010

Marcillac I, Troalen F, Bidart J-M, et al. Free Human Chorionic Gonadotropin β Subunit in Gonadal and Nongonadal Neoplasms. *Cancer Research*. 1992;52(14):3901.

Marcinko TM, Dong J, LeBlanc R, Daborowski KV, Vachet RW. Small molecule-mediated inhibition of β-2-microglobulin-based amyloid fibril formation. *The Journal of biological chemistry*. Jun 23 2017;292(25):10630-10638. doi:10.1074/jbc.M116.774083

Merlini G, Wechalekar AD, Palladini G. Systemic light chain amyloidosis: an update for treating physicians. *Blood*. Jun 27 2013;121(26):5124-30. doi:10.1182/blood-2013-01-453001

Moreau AS LX, Manning R, Coiteux V, Darre S, Hatjiharisi E, Hunter Z, Jia X, Ngo H, O'Sullivan G, Santos D, Treon S, Facon T, Anderson K, Ghobrial I. Serum Free Light Chain in Waldenstrom Macroglobulinemia. 2006;doi:10.1182/blood.V108.11.2420.2420

NANETS. NANETS 2024 Compendium. 2025. https://nanets.net/images/NANETS_2024_Symposium_Guidelines_Compendium.pdf

NCCN. Biomarkers Compendium. https://www.nccn.org/compendia-templates/compendia/biomarkers-compendium

NCCN. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) Systemic Light Chain Amyloidosis Version Version 2.2025. https://www.nccn.org/professionals/physician_gls/pdf/amyloidosis.pdf

NCCN. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) Multiple Myeloma Version 1.2025. https://www.nccn.org/professionals/physician_gls/pdf/myeloma.pdf

NCCN. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) Ovarian Cancer Version 2.2025. https://www.nccn.org/professionals/physician_gls/pdf/ovarian.pdf

NCI. Tumor Markers. https://www.cancer.gov/about-cancer/diagnosis-staging/diagnosis/tumor-markers-fact-sheet

Oyaert M, Boone E, De Ceuninck L, et al. Clonal multicentric Castleman's disease with increased free Kappa light chains in a patient with systemic lupus erythematosus. *Ann Hematol*. Jul 2014;93(7):1255-7. doi:10.1007/s00277-013-1962-3

Park SJ, Jang JY, Jeong SW, et al. Usefulness of AFP, AFP-L3, and PIVKA-II, and their combinations in diagnosing hepatocellular carcinoma. *Medicine*. Mar 2017;96(11):e5811. doi:10.1097/md.0000000000005811

Payne V, Kam PC. Mast cell tryptase: a review of its physiology and clinical significance. *Anaesthesia*. Jul 2004;59(7):695-703. doi:10.1111/j.1365-2044.2004.03757.x

Pejler G, Ronnberg E, Waern I, Wernersson S. Mast cell proteases: multifaceted regulators of inflammatory disease. *Blood*. Jun 17 2010;115(24):4981-90. doi:10.1182/blood-2010-01-257287

Perfetto F, Bergesio F, Emdin M, Cappelli F. Troponins in cardiac amyloidosis: multipurpose markers. *Nat Rev Cardiol*. Mar 2014;11(3):179. doi:10.1038/nrcardio.2013.129-c1

Pinzani P, D'Argenio V, Del Re M, et al. Updates on liquid biopsy: current trends and future perspectives for clinical application in solid tumors. *Clin Chem Lab Med*. Jun 25 2021;59(7):1181-1200. doi:10.1515/cclm-2020-1685

Qin J, Yang Q, Ye H, et al. Using Serological Proteome Analysis to Identify and Evaluate Anti-GRP78 Autoantibody as Biomarker in the Detection of Gastric Cancer. *J Oncol*. 2020;2020:9430737. doi:10.1155/2020/9430737

Raby B. Personalized medicine. Updated September 6, 2023. https://www.uptodate.com/contents/personalized-medicine

Ryu T, Takami Y, Wada Y, et al. Double- and Triple-Positive Tumor Markers Predict Early Recurrence and Poor Survival in Patients with Hepatocellular Carcinoma within the Milan Criteria and Child-Pugh Class A. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract*. Jun 2017;21(6):957-966. doi:10.1007/s11605-017-3394-1

Santos Schraiber Ld, de Mattos AA, Zanotelli ML, et al. Alpha-fetoprotein Level Predicts Recurrence After Transplantation in Hepatocellular Carcinoma. *Medicine*. Jan 2016;95(3):e2478. doi:10.1097/md.0000000000002478

Schefer H, Mattmann S, Joss RA. Hereditary persistence of α-fetoproteinCase report and review of the literature. *Annals of Oncology*. 1998;9(6):667-672. doi:10.1023/A:1008243311122

Seo S, Hong JY, Yoon S, et al. Prognostic significance of serum beta-2 microglobulin in patients with diffuse large B-cell lymphoma in the rituximab era. *Oncotarget*. Nov 22 2016;7(47):76934-76943. doi:10.18632/oncotarget.12734

Sharma S, Jackson PG, Makan J. Cardiac troponins. *J Clin Pathol*. Oct 2004;57(10):1025-6. doi:10.1136/jcp.2003.015420

Sharma U, Pal D, Prasad R. Alkaline phosphatase: an overview. *Indian J Clin Biochem*. Jul 2014;29(3):269-78. doi:10.1007/s12291-013-0408-y

Singal AG, Llovet JM, Yarchoan M, et al. AASLD Practice Guidance on prevention, diagnosis, and treatment of hepatocellular carcinoma. *Hepatology*. May 22 2023;doi:10.1097/HEP.0000000000000466

Stankowski-Drengler T, Gertz MA, Katzmann JA, et al. Serum immunoglobulin free light chain measurements and heavy chain isotype usage provide insight into disease biology in patients with POEMS syndrome. *Am J Hematol*. Jun 2010;85(6):431-4. doi:10.1002/ajh.21707

Stephens RW, Brunner N, Janicke F, Schmitt M. The urokinase plasminogen activator system as a target for prognostic studies in breast cancer. *Breast cancer research and treatment*. 1998;52(1-3):99-111. doi:10.1007/978-1-4615-5195-9_15

Stoffel EM, McKernin SE, Brand R, et al. Evaluating Susceptibility to Pancreatic Cancer: ASCO Provisional Clinical Opinion. *Journal of Clinical Oncology*. 2019/01/10 2018;37(2):153-164. doi:10.1200/JCO.18.01489

Strosberg J. Diagnosis of carcinoid syndrome and tumor localization. Updated March 19, 2025. https://www.uptodate.com/contents/diagnosis-of-the-carcinoid-syndrome-and-tumor-localization

Sturgeon CM, Duffy MJ, Hofmann BR, et al. National Academy of Clinical Biochemistry Laboratory Medicine Practice Guidelines for use of tumor markers in liver, bladder, cervical, and gastric cancers. *Clin Chem*. Jun 2010;56(6):e1-48. doi:10.1373/clinchem.2009.133124

Sturgeon CM, Duffy MJ, Stenman UH, et al. National Academy of Clinical Biochemistry laboratory medicine practice guidelines for use of tumor markers in testicular, prostate, colorectal, breast, and ovarian cancers. *Clin Chem*. Dec 2008;54(12):e11-79. doi:10.1373/clinchem.2008.105601

Sturgeon CM, Hoffman BR, Chan DW, et al. National Academy of Clinical Biochemistry Laboratory Medicine Practice Guidelines for Use of Tumor Markers in Clinical Practice: Quality Requirements. *Clinical Chemistry*. 2008;54(8):e1. doi:10.1373/clinchem.2007.094144

Szulc P, Bauer DC, Dempster DW, Luckey M, Cauley JA. Osteoporosis. 2013;1doi:10.1016/B978-0-12-415853-5.00067-4

Thio Q, Karhade AV, Notman E, et al. Serum alkaline phosphatase is a prognostic marker in bone metastatic disease of the extremity. *J Orthop*. Nov-Dec 2020;22:346-351. doi:10.1016/j.jor.2020.08.008

Tian T, Gao J, Li N, et al. Circulating Chromogranin A as A Marker for Monitoring Clinical Response in Advanced Gastroenteropancreatic Neuroendocrine Tumors. *PloS one*. 2016;11(5):e0154679. doi:10.1371/journal.pone.0154679

Tormey WP, Byrne B, Hill AD, Sherlock M, Thompson CJ. Should serum calcitonin be routinely measured in patients presenting with thyroid nodule? *Minerva endocrinologica*. Dec 2017;42(4):306-310. doi:10.23736/s0391-1977.17.02566-4

Tosi P, Tomassetti S, Merli A, Polli V. Serum free light-chain assay for the detection and monitoring of multiple myeloma and related conditions. *Ther Adv Hematol*. Feb 2013;4(1):37-41. doi:10.1177/2040620712466863

Tuttle RM. Medullary thyroid cancer: Clinical manifestations, diagnosis, and staging. Updated July 29, 2024. https://www.uptodate.com/contents/medullary-thyroid-cancer-clinical-manifestations-diagnosis-and-staging

Van Poznak C, Somerfield MR, Bast RC, et al. Use of Biomarkers to Guide Decisions on Systemic Therapy for Women With Metastatic Breast Cancer: American Society of Clinical Oncology Clinical Practice Guideline. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. Aug 20 2015;33(24):2695-704. doi:10.1200/jco.2015.61.1459

Venner CP. AL amyloidosis cardiac staging updated using BNP. *Blood*. Jan 17 2019;133(3):184-185. doi:10.1182/blood-2018-10-882159

Walentowicz P, Krintus M, Sadlecki P, et al. Serum inhibin A and inhibin B levels in epithelial ovarian cancer patients. *PloS one*. 2014;9(3):e90575. doi:10.1371/journal.pone.0090575

Weber M, Hamm C. Role of B-type natriuretic peptide (BNP) and NT-proBNP in clinical routine. *Heart*. Jun 2006;92(6):843-9. doi:10.1136/hrt.2005.071233

Wells SA, Jr., Asa SL, Dralle H, et al. Revised American Thyroid Association guidelines for the management of medullary thyroid carcinoma. *Thyroid : official journal of the American Thyroid Association*. Jun 2015;25(6):567-610. doi:10.1089/thy.2014.0335

Wu D, Lim MS, Jaffe ES. Pathology of Castleman Disease. *Hematol Oncol Clin North Am*. Feb 2018;32(1):37-52. doi:10.1016/j.hoc.2017.09.004

Wu M, Liu H, Liu Z, Liu C, Zhang A, Li N. Analysis of serum alpha-fetoprotein (AFP) and AFP-L3 levels by protein microarray. *The Journal of international medical research*. Oct 2018;46(10):4297-4305. doi:10.1177/0300060518789304

Yang X, Yang Y, Li Z, et al. Diagnostic value of circulating chromogranin a for neuroendocrine tumors: a systematic review and meta-analysis. *PloS one*. 2015;10(4):e0124884. doi:10.1371/journal.pone.0124884