Humana. Claims Payment Policy

Subject: Laboratory – Testing of Homocysteine Metabolism-Related Conditions

Application: Medicare Advantage Products
Policy number: LAB2023054
Published date: 07/2023
Revision date: 05/2026

Disclaimer: *The intended audience of this medical claims payment policy is healthcare providers who treat Humana members. This policy is made available to provide information on certain Humana claims payment policies. This policy is a guideline only and does not constitute medical advice, guarantee of payment, plan pre-authorization, an explanation of benefits, or a contract. This policy does not govern whether a procedure is covered under a specific member plan or policy, nor is it intended to address every claim situation. Claims may be affected by other factors, such as: state and federal laws and regulations, provider contract terms, and our professional judgment. This policy is subject to change or termination by Humana. Humana has full and final discretionary authority for its interpretation and application. No part of this policy may be reproduced, stored in a retrieval system or transmitted, in any shape or form or by any means, electronic, mechanical, photocopying or otherwise without express written permission from Humana. When printed, this version becomes uncontrolled. For the most current information, always refer to the following website:* ***Humana.com/ClaimPaymentPolicies***

Medicare Advantage Payment Policy

As a Medicare Advantage plan, we have developed payment policies to explain coverage more fully for services. The criteria outlined will apply to Medicare Advantage plans except when the procedure(s) has established criteria through the Original Medicare National Coverage Determinations (NCDs) and Local Coverage Determinations (LCDs) as outlined below.

In addition to the policy, claims payments are subject to other plan requirements for the processing and payment of claims, including, but not limited to, requirements of medical necessity and reasonableness and applicable referral or authorization requirements.

- Total homocysteine testing (CPT code 83090*) in plasma **IS REIMBURSABLE** in patients with homocystinuria or vitamin B12 deficiency.

Applicable procedures for Medicare Advantage plans

Procedure code	Code description
83090*	Homocysteine

**The payment criterion for 83090 applies to all Medicare jurisdictions, except those states awarded to CGS Administrators, Noridian Healthcare Solutions, Novitas Solutions, Palmetto GBA, and Wisconsin Physicians Service Insurance Corp.*

Original Medicare Local Coverage Determinations (LCDs)

Contractor name	Policy title	Policy number	Procedure code
CGS Administrators, LLC	MolDX: Biomarkers in Cardiovascular Risk Assessment	L36139	83090
Noridian Healthcare Solutions, LLC	MolDX: Biomarkers in Cardiovascular Risk Assessment	L36358	83090
Novitas Solutions, Inc.	Assays for Vitamins and Metabolic Function	L34914	83090
Palmetto GBA	Homocysteine Level, Serum	L34419	83090
Palmetto GBA	MolDX: Biomarkers in Cardiovascular Risk Assessment	L36129	83090
Wisconsin Physicians Service Insurance Corporation	MolDX: Biomarkers in Cardiovascular Risk Assessment	L36523	83090

References

Abbott, M. H., Folstein, S. E., Abbey, H., & Pyeritz, R. E. (1987). Psychiatric manifestations of homocystinuria due to cystathionine beta-synthase deficiency: prevalence, natural history, and relationship to neurologic impairment and vitamin B6-responsiveness. *Am J Med Genet, 26*(4), 959-969. doi:10.1002/ajmg.1320260427

ACMG. (2009). Methionine Elevated or Decreased. Retrieved from http://www.acmg.net/PDFLibrary/Methionine-Algorithm.pdf

ACMG. (2010). Newborn Screening ACT Sheet [Increased Methionine] Homocystinuria (CBS Deficiency) Retrieved from file:///C:/Users/AHCS6886/OneDrive%20-%20AVALON%20HEALTH%20SERVICES,%20LLC/Downloads/Methionine.pdf

Al-Sadeq, D. W., & Nasrallah, G. K. (2020). The Spectrum of Mutations of Homocystinuria in the MENA Region. *Genes (Basel), 11*(3). doi:10.3390/genes11030330

Belkhiria, M. N., Ducros, V., Harzallah, K., Jarraya, F., Cordonnier, D., Favier, A., & Achour, A. (2007). [Evaluation of plasmatic homocysteine determination by gas chromatography-mass spectrometry]. *Ann Biol Clin (Paris), 65*(4), 393-398.

Collison, F. T., Xie, Y. A., Gambin, T., Jhangiani, S., Muzny, D., Gibbs, R., . . . Allikmets, R. (2015). Whole Exome Sequencing Identifies an Adult-Onset Case of Methylmalonic Aciduria and Homocystinuria Type C (cblC) with Non-Syndromic Bull's Eye Maculopathy. *Ophthalmic Genet, 36*(3), 270-275. doi:10.3109/13816810.2015.1010736

Concepción-Alvarez, A., Camayd-Viera, I., & Nuevas-Paz, L. (2016). Validation of an HPLC method for total homocysteine quantification in plasma. *Revista del Laboratorio Clinico, 9*(2), 40-47. Retrieved from https://www.sciencedirect.com/science/article/pii/S1888400816300022#!

de Sain-van der Velden, M. G. M., van der Ham, M., Jans, J. J., Visser, G., van Hasselt, P. M., Prinsen, H., & Verhoeven-Duif, N. M. (2015). Suitability of methylmalonic acid and total homocysteine analysis in dried bloodspots. *Anal Chim Acta, 853*, 435-441. doi:10.1016/j.aca.2014.10.043

FDA. (2007, October 11, 2007). Nanosphere 510(k) Summary. Retrieved from https://www.accessdata.fda.gov/cdrh_docs/pdf7/K070597.pdf

FDA. (2010, March 24, 2010). eSensor Thrombophila Risk Test on XT-8 System. Retrieved from https://www.accessdata.fda.gov/cdrh_docs/pdf9/K093974.pdf

FDA. (2011a, May 13, 2011). Invader MTHFR 677 510(k) Summary. Retrieved from https://www.accessdata.fda.gov/cdrh_docs/pdf10/K100987.pdf

FDA. (2011b, April 25, 2011). Invader MTHFR 1298 510(k) Summary. Retrieved from https://www.accessdata.fda.gov/cdrh_docs/pdf10/K100496.pdf

Froese, D. S., Kopec, J., Fitzpatrick, F., Schuller, M., McCorvie, T. J., Chalk, R., . . . Yue, W. W. (2015). Structural Insights into the MMACHC-MMADHC Protein Complex Involved in Vitamin B12 Trafficking. *J Biol Chem, 290*(49), 29167-29177. doi:10.1074/jbc.M115.683268

Frosst, P., Blom, H. J., Milos, R., Goyette, P., Sheppard, C. A., Matthews, R. G., . . . et al. (1995). A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. *Nat Genet, 10*(1), 111-113. doi:10.1038/ng0595-111

Gales, A., Masingue, M., Millecamps, S., Giraudier, S., Grosliere, L., Adam, C., . . . Nadjar, Y. (2018). Adolescence/adult onset MTHFR deficiency may manifest as isolated and treatable distinct neuro-psychiatric syndromes. *Orphanet J Rare Dis, 13*(1), 29. doi:10.1186/s13023-018-0767-9

Gaughan, D. J., Kluijtmans, L. A., Barbaux, S., McMaster, D., Young, I. S., Yarnell, J. W., . . . Whitehead, A. S. (2001). The methionine synthase reductase (MTRR) A66G polymorphism is a novel genetic determinant of plasma homocysteine concentrations. *Atherosclerosis, 157*(2), 451-456.

Gaustadnes, M., Ingerslev, J., & Rütiger, N. (1999). Prevalence of Congenital Homocystinuria in Denmark. *New England Journal of Medicine, 340*(19), 1513-1513. doi:10.1056/NEJM199905133401915

Gonzales, M. C., Yu, P., & Shiao, S. P. K. (2017). MTHFR Gene Polymorphism-Mutations and Air Pollution as Risk Factors for Breast Cancer: A Metaprediction Study. *Nursing research, 66*(2), 152-163. doi:10.1097/NNR.0000000000000206

Harmon, D. L., Shields, D. C., Woodside, J. V., McMaster, D., Yarnell, J. W., Young, I. S., . . . Whitehead, A. S. (1999). Methionine synthase D919G polymorphism is a significant but modest determinant of circulating homocysteine concentrations. *Genet Epidemiol, 17*(4), 298-309. doi:10.1002/(sici)1098-2272(199911)17:4<298::Aid-gepi5>3.0.Co;2-v

HHS. (2018, 02/2019). Recommended Uniform Screening Panel. Retrieved from https://www.hrsa.gov/advisory-committees/heritable-disorders/rusp/index.html

Huemer, M., Diodato, D., Schwahn, B., Schiff, M., Bandeira, A., Benoist, J. F., . . . Dionisi-Vici, C. (2017). Guidelines for diagnosis and management of the cobalamin-related remethylation disorders cblC, cblD, cblE, cblF, cblG, cblJ and MTHFR deficiency. *J Inherit Metab Dis, 40*(1), 21-48. doi:10.1007/s10545-016-9991-4

Huemer, M., Kožich, V., Rinaldo, P., Baumgartner, M. R., Merinero, B., Pasquini, E., . . . Blom, H. J. (2015). Newborn screening for homocystinurias and methylation disorders: systematic review and proposed guidelines. *Journal of inherited metabolic disease, 38*(6), 1007-1019. doi:10.1007/s10545-015-9830-z

Ismayilova, N., MacKinnon, A. D., Mundy, H., & Fallon, P. (2019). Reversible Cerebral White Matter Abnormalities in Homocystinuria. *JIMD Rep, 44*, 115-119. doi:10.1007/8904_2018_135

Janosík, M., Sokolová, J., Janosíková, B., Krijt, J., Klatovská, V., & Kozich, V. (2009). Birth prevalence of homocystinuria in Central Europe: frequency and pathogenicity of mutation c.1105C>T (p.R369C) in the cystathionine beta-synthase gene. *The Journal of pediatrics, 154*(3), 431-437. doi:10.1016/j.jpeds.2008.09.015

Kluijtmans, L. A., Young, I. S., Boreham, C. A., Murray, L., McMaster, D., McNulty, H., . . . Whitehead, A. S. (2003). Genetic and nutritional factors contributing to hyperhomocysteinemia in young adults. *Blood, 101*(7), 2483-2488. doi:10.1182/blood.V101.7.2483

Li, Barshop, Feigenbaum, & Khanna. (2018). Brain Magnetic Resonance Imaging Findings in Poorly Controlled Homocystinuria. *J Radiol Case Rep*. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/29875981

Long, S., & Goldblatt, J. (2016). MTHFR genetic testing: Controversy and clinical implications. *Australian Family Physician, 45*, 237-240. Retrieved from http://www.racgp.org.au/afp/2016/april/mthfr-genetic-testing-controversy-and-clinical-implications/

Mazaheri, Mostofizadeh, & Hashemipour. (2017). Homocystinuria with Stroke and Positive Familial History. *Adv Biomed Res*. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/29279830

Morris, A. A. M., Kožich, V., Santra, S., Andria, G., Ben-Omran, T. I. M., Chakrapani, A. B., . . . Chapman, K. A. J. J. o. I. M. D. (2017). Guidelines for the diagnosis and management of cystathionine beta-synthase deficiency. *40*(1), 49-74. doi:10.1007/s10545-016-9979-0

Mudd, S. H., Finkelstein, J. D., Refsum, H., Ueland, P. M., Malinow, M. R., Lentz, S. R., . . . Rosenberg, I. H. (2000). Homocysteine and its disulfide derivatives: a suggested consensus terminology. *Arterioscler Thromb Vasc Biol, 20*(7), 1704-1706.

Mudd, S. H., Skovby, F., Levy, H. L., Pettigrew, K. D., Wilcken, B., Pyeritz, R. E., . . . et al. (1985). The natural history of homocystinuria due to cystathionine beta-synthase deficiency. *Am J Hum Genet, 37*(1), 1-31.

Nelson, B. C., Pfeiffer, C. M., Sniegoski, L. T., & Satterfield, M. B. (2003). Development and evaluation of an isotope dilution LC/MS method for the determination of total homocysteine in human plasma. *Anal Chem, 75*(4), 775-784.

Nexo, E., Engbaek, F., Ueland, P. M., Westby, C., O'Gorman, P., Johnston, C., . . . Refsum, H. (2000). Evaluation of Novel Assays in Clinical Chemistry: Quantification of Plasma Total Homocysteine. *Clinical Chemistry, 46*(8), 1150. Retrieved from http://clinchem.aaccjnls.org/content/46/8/1150.abstract

NIH. (2016, 3/1/2016). Homocystinuria. *Genetics Home Reference.* Retrieved from https://ghr.nlm.nih.gov/condition/homocystinuria#definition

NIH. (2018a, 10/23/2018). Homocystinuria. *Genetics Home Reference.* Retrieved from https://ghr.nlm.nih.gov/condition/homocystinuria#definition

NIH. (2018b). Homocystinuria due to CBS deficiency. Retrieved from https://rarediseases.info.nih.gov/diseases/6667/homocystinuria-due-to-cbs-deficiency

Okun, J. G., Gan-Schreier, H., Ben-Omran, T., Schmidt, K. V., Fang-Hoffmann, J., Gramer, G., . . . Hoffmann, G. F. (2017). Newborn Screening for Vitamin B6 Non-responsive Classical Homocystinuria: Systematical Evaluation of a Two-Tier Strategy. *JIMD Rep, 32*, 87-94. doi:10.1007/8904_2016_556

Rose, N. C., & Dolan, S. M. (2012). Newborn screening and the obstetrician. *Obstetrics and gynecology, 120*(4), 908-917. doi:10.1097/AOG.0b013e31826b2f03

Rosenson, R. S., Smith, C. C., & Bauer, K. A. (2020, October 26). Overview of homocysteine. Retrieved from https://www.uptodate.com/contents/overview-of-homocysteine

Sacharow, S. J., Picker, J. D., & Levy, H. L. (2004). Homocystinuria Caused by Cystathionine Beta-Synthase Deficiency. In M. P. Adam, H. H. Ardinger, R. A. Pagon, S. E. Wallace, L. J. H. Bean, K. Stephens, & A. Amemiya (Eds.), *GeneReviews((R))*. Seattle (WA): University of Washington, Seattle

University of Washington, Seattle. GeneReviews is a registered trademark of the University of Washington, Seattle. All rights reserved.

Wang, W., Jiao, X. H., Wang, X. P., Sun, X. Y., & Dong, C. (2016). MTR, MTRR, and MTHFR Gene Polymorphisms and Susceptibility to Nonsyndromic Cleft Lip With or Without Cleft Palate. *Genet Test Mol Biomarkers, 20*(6), 297-303. doi:10.1089/gtmb.2015.0186

Weisberg, I., Tran, P., Christensen, B., Sibani, S., & Rozen, R. (1998). A second genetic polymorphism in methylenetetrahydrofolate reductase (MTHFR) associated with decreased enzyme activity. *Mol Genet Metab, 64*(3), 169-172. doi:10.1006/mgme.1998.2714

Zhu, H., Blake, S., Chan, K. T., Pearson, R. B., & Kang, J. (2018). Cystathionine beta-Synthase in Physiology and Cancer. Biomed Res Int, 2018, 3205125. doi:10.1155/2018/3205125