

PEDIATRIC DOSING GUIDE

BLUE: 19-23kg/42-51lbs

Normal Vital Signs

HR asleep	HR awake	Respiratory Rate	Systolic BP	Diastolic BP	MAP
65-100	80-120	20-28	89-112	46-72	58-69

NS Fluid Bolus: 10ml/kg	i-gel Airway	Blade for foreign body removal
420ml	2.0	2

DEFIBRILLATION: 2, 4J/kg	1st: 40J	2nd: 80J	CARDIOVERSION: 1, 2J/kg	1st: 20J	2nd: 40J
--------------------------	----------	----------	-------------------------	----------	----------

Medication	Concentration	Dose	Dose in mg	Dose in ml	Details
ACETAMINOPHEN	160mg/5ml (32mg/ml)	15mg/kg PO Max dose: 345mg	315mg	10ml	
ADENOSINE	6mg/2ml (3mg/ml)	0.1mg/kg RIVP Max 1st dose: 6mg Max 2nd dose: 12mg	1st: 2.1mg 2nd: 4.2mg	1st: 0.7ml 2nd: 1.4ml	RIVP w/ 10ml NS flush MR x1 double the dose (0.2mg/kg)
ALBUTEROL	2.5mg/3ml	2.5mg/3ml HHN	2.5mg	3ml	MR x1
AMIODARONE (Pulseless arrest)	150mg/3ml (50mg/ml)	5mg/kg IV/IO Max single dose: 300mg	105mg	2.1ml	Follow with or dilute in 20ml NS flush Give after 3rd shock
ATROPINE (Bradycardia)	1mg/10ml (0.1mg/ml)	0.02mg/kg IV/IO Min dose: 0.1mg Max single dose: 0.5mg	0.4mg	4ml	MR x1 in 3-5 min
ATROPINE (Organophosphate poisoning)	Preload: 1mg/10ml (0.1mg/ml) Vial: 0.4mg/ml	0.05mg/kg IV/IO	1mg	Preload: 10ml Vial: 2.6ml	MR q5-10 min until symptoms resolve
DEXTROSE	10%	5ml/kg IV/IO Max dose: 125ml		105ml	Give over 10 min
DIPHENHYDRAMINE <i>Benadryl</i>	50mg/ml	1mg/kg IM/IV/IO Max dose: 50mg	21mg	0.4ml	
EPINEPHRINE (Cardiac arrest/ Bradycardia)	1mg/10ml (0.1mg/ml)	0.01mg/kg IV/IO	0.2mg	2ml	MR q3-5 min
EPINEPHRINE (Allergic reaction/ Asthma)	1mg/ml	0.01mg/kg IM Total max dose: 0.6mg	0.2mg	0.2ml	MR x1 in 5 min
EPINEPHRINE (Upper airway/Stridor)	1mg/ml	5mg HHN	5mg	5ml	

BLUE: 19-23kg/42-51lbs

Medication	Concentration	Dose	Dose in mg	Dose in ml	Details
FENTANYL (Pain)	50mcg/ml	1mcg/kg IV/IO/IM/IN Max dose: 3mcg/kg	21mcg	0.42ml	MR q5 min For IN: split dose equally in each nostril
GLUCAGON (Hypoglycemia/Beta blocker OD)	1mg/ml	0.03mg/kg IM Max dose: 1mg	0.6mg	0.6ml	MR x2 q15 min if no IV established
IPRATROPIUM <i>Atrovent</i>	500mcg/2.5ml	500mcg/2.5ml HHN	500mcg	2.5ml	
LIDOCAINE 2% preservative free (IO insertion)	20mg/ml	0.5mg/kg slow IO Max dose: 40mg	<u>1st:</u> 10mg <u>2nd:</u> 5mg	<u>1st:</u> 0.5ml <u>2nd:</u> 0.3ml	MR x1 half initial dose (0.25mg/kg)
MIDAZOLAM <i>Versed</i> (Seizure)	5mg/ml	0.2mg/kg IM Max single dose 5mg	4.2mg	0.8ml	MR x1 in 10 min
MIDAZOLAM <i>Versed</i> (Seizure)	5mg/ml	0.2mg/kg IN Max dose 5mg	4.2mg	0.8ml	Split dose equally in each nostril
MIDAZOLAM <i>Versed</i> (Seizure)	2mg/2ml (1mg/ml)	0.05mg/kg slow IV/IO Total max dose: 5mg	1mg	1ml	MR x2 q15 min
MIDAZOLAM <i>Versed</i> (Cardioversion)	2mg/2ml (1mg/ml)	0.05mg/kg slow IV/IO Max dose: 1mg	1mg	1ml	
MORPHINE (Pain/burns)	10mg/ml	0.1mg/kg IV/IO/IM	2.1mg	0.2ml	MR x2 q15 min (IV/IO) MR in 30min (IM)
NALOXONE <i>Narcan</i>	2mg/2ml (1mg/ml)	0.1mg/kg IV/IO/IM/IN	2mg	2ml	MR q5 min For IN: split dose equally in each nostril
ONDANSETRON <i>Zofran</i>	4mg tab 4mg/2ml	4mg ODT/slow IV	4mg	2ml 1 tab	Slow IV over 30 sec
SODIUM BICARBONATE	1mEq/ml	1mEq/kg IV/IO	21mEq	21ml	