

## DAA Assignment - 1

Ques 1

Asymptotic notations are the mathematical notations used to describe the running time of an algorithm when the Input tends toward a particular value or a limiting value.

Eg - In bubble sort, when the Input array is already sorted, the time taken by algorithm is linear i.e. the best case ( $\Omega$  notation)  
(omega notation)

But when the Input array is in reverse condition, the algorithm takes the maximum time to sort the element i.e. the worst case (Big - O notation)

when the Input array is neither sorted nor in reverse order, then it takes average time ( $\Theta$  - notation) (Theta notation)

Q2 -

$$\sum_{i=1}^n 1 + 1 + \dots + k \text{ time}$$

$$\therefore 2^k \geq n$$

$$2^k = n$$

taking log both side

$$k \log 2 = \log n$$

$$k = \frac{\log n}{\log 2}$$

$$k = \log_2 n$$

$$O(\log n)$$

$$\left[ \log_b(x) = \frac{\log_a(x)}{\log_a(b)} \right]$$

Ques-

$$T(n) = \begin{cases} 3T(n-1) & n > 0 \\ 1 & n = 0 \end{cases}$$

$$T(n) = 3T(n-1) \text{ --- (1)}$$

$$\text{let } n = n-1$$

$$\text{Putting } n \text{ in eq (1)}$$

$$T(n-1) = 3T(n-2) \text{ --- (2)}$$

$$\text{Putting (2) in (1)}$$

$$T(n) = 3^2 T(n-2) \text{ --- (3)}$$

$$\text{let } n = n-2$$

$$\text{Putting } n \text{ in eq (1)}$$

$$T(n) = 3T(n-2) \text{ --- (4)}$$

$$\text{Putting (4) in (3)}$$

$$T(n) = 3^3 T(n-3)$$

$$T(n) = 3^k T(n-k)$$

$$\text{let } n-k = 0$$

$$n = k$$

$$T(n) = 3^n T(0)$$

$$= O(3^n)$$

Ques4 -

$$T(n) = \begin{cases} 2T(n-1) - 1 & n > 0 \\ 1 & n = 0 \end{cases}$$

$$T(n) = 2T(n-1) - 1 \text{ --- (1)}$$

Let  $n = n-1$  in eq (1)

$$T(n-1) = 2T(n-2) - 1 \text{ --- (2)}$$

put this value in eq (1)

$$T(n) = 2[2T(n-2) - 1] - 1$$

$$T(n) = 4T(n-2) - 2 - 1 \text{ --- (3)}$$

Let  $n = n-2$ 

$$T(n-2) = 2T(n-3) - 1 \text{ --- (4)}$$

put this value in eq (3)

$$T(n) = 4[2T(n-3) - 1] - 2 - 1$$

$$T(n) = 8T(n-3) - 4 - 2 - 1$$

$$T(n) = 2^k T(n-k) - 2^{k-1} - 2^{k-2} - \dots - 2^0$$

put  $n-k = 0$ 

$$n = k$$

$$T(n) = 2^n T(0) - 2^{n-1} - \dots - 2^0$$

$$= 2^n - [2^{n-1} + 2^{n-2} + \dots + 2^0]$$

$$\Rightarrow 2^n - 2^{n-1} \left(1 - \left(\frac{1}{2}\right)^n\right)^2$$

$$\Rightarrow 2^n [1 - (1 - (\frac{1}{2})^n)]$$

$$\Rightarrow 2^n [1 - 1 + (\frac{1}{2})^n]$$

$$\Rightarrow 2^n (\frac{1}{2})^n = 1$$

$$\Rightarrow O(1)$$

Ans

Ques 5 -

$$u^0 = 1, 2, 3, \dots$$

$$S = 1, 3, 6, 10, \dots, n \quad \text{--- (1)}$$

$$\text{also } S = 1, 3, 6, 10, \dots, n \quad \text{--- (2)}$$

Subst (1) - (2)

$$0 = 1 + 2 + 3 + \dots + n - T_n$$

$$T_n = 1 + 2 + 3 + \dots + k$$

for  $k$  iteration

$$1 + 2 + 3 + \dots + k \leq n$$

$$\frac{k(k+1)}{2} \leq n$$

$$\frac{k^2 + k}{2} \leq n$$

$$O(k^2) \leq n$$

$$k = O(\sqrt{n})$$

Ques 6 -

$$u^0 \leq n$$

$$u^0 \leq \sqrt{n}$$

$$u^0 = 1, 2, 3, \dots, \sqrt{n}$$

$$\sum_{u^0=1}^n 1 + 2 + 3 + \dots + \sqrt{n}$$

$$T(n) = \frac{\sqrt{n} \times (\sqrt{n} + 1)}{2}$$

$$T(n) = \frac{n + \sqrt{n}}{2}$$

$$= O(n)$$

Ans

Ques 7 -

Given  $K = K \neq 2$  $K = 1, 2, 4, 8, \dots, n$  $\log P, a = 1, a = 2$ 

$$= \frac{a(a^n - 1)}{a - 1}$$

$$= \frac{1(2^K - 1)}{1}$$

$$n = 2^K$$

$$\Rightarrow \log n = K$$

$$\begin{array}{ccccccc} a^0 & & f & & K & & \\ 1 & & \log n & & \log n * \log n & & \\ 2 & & \log n & & \vdots & & \\ 3 & & \vdots & & \vdots & & \\ \vdots & & \vdots & & \vdots & & \\ n & & \log n & & \log n * \log n & & \end{array}$$

$$\Rightarrow O(n * \log n * \log n)$$

$$O(n \log^2 n)$$

$$\text{Ques 8} \quad T(n) = T(n/3) + n^2$$

$$a = 1, b = 3 \quad f(n) = n^2$$

$$c = \log_3 1 = 0$$

$$\Rightarrow n^0 = 1 > [f(n) = n^2]$$

$$T(n) = O(n^2)$$

Ques 9-

$$\text{for } i^0 = 1 \Rightarrow j^0 = 1, 2, 3, 4 \dots n = n$$

$$\text{for } j^0 = 2 \Rightarrow j^0 = 1, 3, 5 \dots n = n/2$$

$$\vdots$$

$$\text{for } (i^0 = n) \Rightarrow j^0 = 1 \dots$$

$$\Rightarrow \sum_{j=n}^1 n + \frac{n}{2} + \frac{n}{3} + \frac{n}{4} \dots + 1$$

$$\Rightarrow \sum_{j=n}^1 n \left[ 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} \dots + \frac{1}{n} \right]$$

$$\Rightarrow \sum_{j=n}^1 n \log n$$

$$\Rightarrow O(n \log n) \quad \underline{\underline{\text{Ans}}}$$

Ques 10-

as given  $n^k \neq c^w$

relation b/w  $n^k$  &  $c^w$  is

$$n^k = O(c^w)$$

$$\text{as } n^k \leq a c^w$$

$$\forall n \geq n_0 \text{ \& some constant } n_0 > 0$$

$$\text{for } n_0 = 1$$

$$c = 2$$

$$\Rightarrow 1^k \leq 2^1$$

$$n_0 = 1 \text{ \& } c = 2$$

Ans