

Applied Linear Algebra

Standard Deviation

Devi Prasad M
Manipal School of Information Sciences
Manipal

September 2026

For any vector $\mathbf{x} \in \mathbb{R}^n$, the *mean* of its entries is

$$\text{avg}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n x_i$$

and the associated *de-meaned* vector is

$$\bar{\mathbf{x}} = \mathbf{x} - \text{avg}(\mathbf{x})\mathbf{1}_n$$

Notice $\text{avg}(\bar{\mathbf{x}}) = 0$, that is, the mean value of the entries of $\bar{\mathbf{x}}$ is zero.

Let $\mathbf{x} = (10, -2, 8, 23, 6, 75)$.

Then $\text{avg}(\mathbf{x}) = 20$ and the de-meaned vector is

$$\begin{aligned}\bar{\mathbf{x}} &= (10, -2, 8, 23, 6, 75) - (20, 20, 20, 20, 20, 20) \\ &= (10 - 20, -2 - 20, 8 - 20, 23 - 20, 6 - 20, 75 - 20) \\ &= (-10, -22, -12, 3, -14, 55)\end{aligned}$$

The mean of the entries of $\bar{\mathbf{x}}$ is zero.

What is $\bar{x}$ if all entries of $\mathbf{x}$ are equal?

Norm of a De-meaned Vector

Recall $\mathbf{x} = (10, -2, 8, 23, 6, 75)$ and $\text{avg}(\mathbf{x}) = 20$.

$$\|\mathbf{x}\| = \sqrt{100 + 4 + 64 + 529 + 36 + 5625} = \sqrt{6358} \approx 79.7371.$$

$$\begin{aligned}\bar{\mathbf{x}} &= (10, -2, 8, 23, 6, 75) - (20, 20, 20, 20, 20, 20) \\ &= (-10, -22, -12, 3, -14, 55)\end{aligned}$$

Norm of the de-meaned vector $\bar{\mathbf{x}}$ is

$$\begin{aligned}\|\bar{\mathbf{x}}\| &= \sqrt{(-10)^2 + (-22)^2 + (-12)^2 + 3^2 + (-14)^2 + 55^2} \\ &= \sqrt{10^2 + 22^2 + 12^2 + 3^2 + 14^2 + 55^2} \\ &= \sqrt{100 + 484 + 144 + 9 + 196 + 3025} \\ &= \sqrt{3958} \\ &\approx 62.9126\end{aligned}$$

The *standard deviation* of an n -vector $\mathbf{x}$ is defined as the RMS value of the de-meaned vector $\mathbf{x} - \text{avg}(\mathbf{x})\mathbf{1}_n$:

$$\begin{aligned}\text{std}(\mathbf{x}) &= \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \text{avg}(\mathbf{x}))^2} \\ &= \sqrt{\frac{(x_1 - \text{avg}(\mathbf{x}))^2 + \cdots + (x_n - \text{avg}(\mathbf{x}))^2}{n}} \\ &= \frac{\|\bar{\mathbf{x}}\|}{\sqrt{n}} \\ &= \frac{\|\mathbf{x} - ((\mathbf{1}_n^\top \mathbf{x})/n) \mathbf{1}_n\|}{\sqrt{n}}\end{aligned}$$

Consider our example

$$\mathbf{x} = (10, -2, 8, 23, 6, 75)$$

$$n = 6$$

$$\text{avg}(\mathbf{x}) = 20$$

$$\bar{\mathbf{x}} = (-10, -22, -12, 3, -14, 55)$$

$$\|\bar{\mathbf{x}}\| \approx 62.9126$$

Now we can compute the standard deviation of $\mathbf{x}$:

$$\begin{aligned}\text{std}(\mathbf{x}) &= \|\bar{\mathbf{x}}\|/\sqrt{n} \\ &= 62.9126/\sqrt{6} \\ &\approx 25.686\end{aligned}$$

- The standard deviation of a vector is **zero** only when all its entries are equal.
- The standard deviation of a vector is *small* when the entries of the vector are nearly the same.

The standard deviation of a vector $\mathbf{x}$ tells us the typical amount by which its entries deviate from their average value.

Consider the following 8-vector that has a mean of 0 and a standard deviation of 1:

$$\mathbf{x} = (-0.46, -0.51, -0.29, 0.18, 1.22, -0.79, -1.27, 1.92)$$

$$\sum_{i=1}^8 x_i = 0, \quad \text{avg}(\mathbf{x}) = 0$$

$$\sum_{i=1}^8 x_i^2 = 8, \quad \|\mathbf{x}\| = \sqrt{8}$$

$$\begin{aligned} \text{std}(\mathbf{x}) &= \|\mathbf{x}\|/\sqrt{8} \\ &= 1 \end{aligned}$$

Mean Absolute Deviation

Unit 1

The Mean Absolute Deviation (MAD) of a vector $\mathbf{x}$ is

$$\text{MAD}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n |x_i - \text{avg}(\mathbf{x})|$$

$$\begin{aligned}\text{MAD}(\mathbf{x}) &= \frac{0.46 + 0.51 + 0.29 + 0.18 + 1.22 + 0.79 + 1.27 + 1.92}{8} \\ &= \frac{6.64}{8} \\ &\approx 0.83\end{aligned}$$

Notice that the MAD is smaller than **Std**, and in this case, MAD looks like a better measure of the typical deviation compared to **Std**.

For most choices of $\mathbf{b}$, however, there is no n -vector $\mathbf{x}$ for which $\mathbf{Ax} = \mathbf{b}$. As a compromise, we seek an $\mathbf{x}$ for which $\mathbf{r} = \mathbf{Ax} - \mathbf{b}$, which we call the *residual* (for the equations $\mathbf{Ax} - \mathbf{b}$), is as small as possible.

We should choose $\mathbf{x}$ so as to minimize the norm of the residual, $\mathbf{Ax} - \mathbf{b}$.

Minimizing the norm of the residual and its square are the same, so we can just as well minimize

$$\|Ax - b\|^2 = \|r\|^2 = r_1^2 + \cdots + r_m^2,$$

the sum of squares of the residuals.