## GATE 2022 -AE 63

## EE23BTECH11057 - Shakunaveti Sai Sri Ram Varun

Question: Which one of the following is the closed form for the generating function of the sequence  $\{a\}_{n>0}$  defined below?

$$a_n = \begin{cases} n+1 & \text{, n is odd} \\ 1 & \text{otherwise} \end{cases}$$
 (1)

(A) 
$$\frac{x(1+x)^2}{(1-x^2)^2} + \frac{1}{1-x}$$
(B) 
$$\frac{x(3-x^2)}{(1-x^2)^2} + \frac{1}{1-x}$$
(C) 
$$\frac{2x}{(1-x^2)^2} + \frac{1}{1-x}$$
(D) 
$$\frac{x}{(1-x^2)^2} + \frac{1}{1-x}$$

(B) 
$$\frac{x(3-x^2)}{(1-x^2)^2} + \frac{1}{1-x}$$

(C) 
$$\frac{2x}{(1-x^2)^2} + \frac{1}{1-x}$$

(D) 
$$\frac{x}{(1-x^2)^2} + \frac{1}{1-x}$$

(GATE CS 2022 QUESTION 36)

$$X(z) = X_1(z) + X_2(z)$$
 (7)

$$X_1(z) = \frac{1}{1 - z^{-1}} \tag{8}$$

$$\implies x_1(n) = \frac{1}{1 - z^{-1}} \tag{9}$$

$$\implies a_n = x_1(n) + x_2(n) \tag{10}$$

To find inverse z-transform of  $X_2(z)$  we use contour integration technique:

$$x_2(n) = \frac{1}{2\pi j} \oint_C x_2(A; z^{-1}) z^{n-1} dz$$
 (11)

$$= \frac{1}{2\pi i} \oint_C \frac{z^n \left(z^2 + 1\right)}{\left(z^2 - 1\right)^2} dz \tag{12}$$

## **Solution:**

For the given sequence:

Parameter	Description	Value
X(z)	Generating function for a sequence $\{a_n\}$	?
$a_n$	$n^{th}$ term of the sequence	(n+1)u(n) (when odd)
		u(n) (when even)

TABLE I INPUT VALUES

$$X(z) = \sum_{k=-\infty}^{\infty} u(2k) z^{-2k} + \sum_{k=-\infty}^{\infty} ((2k+2) u(2k+1)) z^{-(2k+1)}$$

$$\implies X(z) = \left(1 + z^{-2} + z^{-4} + \dots\right) + \left(2z^{-1} + 4z^{-3} + 6z^{-5} + \dots\right)$$

$$\implies X(z) = \frac{1}{1 - z^{-2}} + \left(2z^{-1} + 4z^{-3} + 6z^{-5} \dots\right) (4)$$

$$\implies X(z) = \frac{1}{1 - z^{-2}} + 2z^{-1} \left(\frac{1}{1 - z^{-2}} + \frac{z^{-2}}{(1 - z^{-2})^2}\right)$$
(5)

$$\therefore X(z) = \frac{1}{1 - z^{-1}} + \frac{z^{-1} \left(1 + z^{-2}\right)}{\left(1 - z^{-2}\right)^2} \tag{6}$$

(6) is the closed form of generating function required in the question.

Hence, option (A) is correct.

We can observe that we have two poles at z = 1,-1. And poles are repeated twice, thus by applying residue theorem two times for poles 1 and -1:

$$x_{2}(n) = \frac{1}{(1)!} \lim_{z \to 1} \frac{d}{dz} \left( (z - 1)^{2} X_{2}(z) \right) + \frac{1}{(1)!} \lim_{z \to -1} \frac{d}{dz} \left( (z + 1)^{2} X_{2}(z) \right)$$

$$(13)$$

$$\implies x_{2}(n) = \lim_{z \to 1} \frac{d}{dz} \left( (z - 1)^{2} \frac{z^{n} \left( z^{2} + 1 \right)}{(z^{2} - 1)^{2}} \right) + \lim_{z \to -1} \frac{d}{dz} \left( (z + 1)^{2} \frac{z^{n} \left( z^{2} + 1 \right)}{(z^{2} - 1)^{2}} \right)$$

$$(14)$$

$$\implies x_{2}(n) = \lim_{z \to 1} \frac{(z + 1)^{2} \left( nz^{n-1} + (n + 2)z^{n+1} \right) - 2z^{n} \left( 1 + z^{2} \right) (z + 1)}{(z + 1)^{4}}$$

$$+ \lim_{z \to -1} \frac{(z - 1)^{2} \left( nz^{n-1} + (n + 2)z^{n+1} \right) - 2z^{n} \left( 1 + z^{2} \right) (z - 1)}{(z - 1)^{4}}$$

$$(15)$$

on simplification, we get

$$x_2(n) = \frac{n + n(-1)^{n-1}}{2} \tag{16}$$

$$\therefore a_n = u(n) + \frac{n + n(-1)^{n-1}}{2}u(n)$$
 (17)

Which is the sequence given in the Question.




Fig. 1. Terms of the sequence given