

# Streamlined Northern Rock Sole Analyses and Figures

Laura Vary

1/23/2022

## Contents

<b>Northern Rock Sole (NRS):</b>	<b>1</b>
Loading Data: . . . . .	1
Descriptive Information: . . . . .	1
Larval Generalized Additive Models: . . . . .	1

## Northern Rock Sole (NRS):

### Loading Data:

Northern Rock Sole: larval data only are included for this species. NRS spawn from December to March, live roughly 19 years, and transform to juveniles at standard lengths between 15 and 18 mm.

These data have been trimmed. The larval data are constrained to depths between 40 and 300 meters, to latitudes below 64.5 degrees north, and to days of year between 100 and 165. Larvae are linked to CTD-derived, *in situ* temperature and salinity measurements.

### Descriptive Information:

Table 1: Descriptive Metrics for NRS Larval Data

Lat Range	Lon Range	Day of Year Range	Bottom Depth Range
53.4-60.3	-178.2 to -158.6	101-159	41-298

The following two plots show *the day of year distribution for positive NRS larval catch* (left) and *the year distribution for positive NRS larval catch* (right).

### Larval Generalized Additive Models:

Now we'll move into the GAMs. The following code *is only necessary if the data were re-trimmed and new GAMs need to be run*. In this case, modify markdown document such that “{eval = TRUE}”. The other model figures are marked as “eval = FALSE” if they, as of the last model run, do not produce the best model results. **Make sure to save the new models as RDS objects.**




Figure 1: Northern Rock Sole Larvae

Northern rock sole larvae were best explained by the bivariate salinity-temperature model, in which the spatial and temporal distribution of larvae were modeled in association with a smooth containing *in situ* salinity-temperature data.

We begin with the base larval model:

```
lv.base<-readRDS("./GAM Models/nrs_larvae_base.rds")
summary(lv.base)

##
## Family: Tweedie(p=1.99)
## Link function: log
##
## Formula:
## (Cper10m2 + 1) ~ factor(year) + s(doy, k = 7) + s(lon, lat) +
##   s(bottom_depth, k = 5)
##
## Parametric coefficients:
##                               Estimate Std. Error t value Pr(>|t|)
## (Intercept)            3.9782    0.3107 12.805 < 2e-16 ***
## factor(year)1998     -0.5003    0.9961 -0.502 0.615591
## factor(year)1999     -1.4077    0.5851 -2.406 0.016266 *
## factor(year)2000     -1.4310    0.4798 -2.983 0.002909 **
## factor(year)2002      1.4793    0.3634  4.070 4.98e-05 ***
## factor(year)2003     -0.9398    0.3861 -2.434 0.015074 *
## factor(year)2005     -0.7234    0.3496 -2.069 0.038720 *
## factor(year)2006     -1.7510    0.3417 -5.125 3.43e-07 ***
## factor(year)2007     -1.2025    0.3790 -3.173 0.001546 **
## factor(year)2008     -3.1462    0.4977 -6.321 3.55e-10 ***
## factor(year)2009     -2.4943    0.3476 -7.177 1.20e-12 ***
## factor(year)2010     -1.9833    0.3382 -5.864 5.73e-09 ***
## factor(year)2011     -2.5878    0.4684 -5.524 3.99e-08 ***
```

```

## factor(year)2012 -0.9289      0.3313   -2.804 0.005125 ***
## factor(year)2013 -2.6212      0.4066   -6.446 1.61e-10 ***
## factor(year)2014 -1.1036      0.3328   -3.316 0.000939 ***
## factor(year)2015 -2.1946      0.4189   -5.239 1.88e-07 ***
## factor(year)2016 -1.8722      0.3339   -5.607 2.52e-08 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Approximate significance of smooth terms:
##          edf Ref.df    F p-value
## s(doy)      5.166 5.696 7.129 3.98e-06 ***
## s(lon,lat)  27.485 28.844 22.573 < 2e-16 ***
## s(bottom_depth) 3.663 3.925 11.699 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## R-sq.(adj) =  0.223 Deviance explained = 55.1%
## -REML = 4894.2 Scale est. = 1.743 n = 1358

```

AIC(lv.base)

```

## [1] 9755.342

##
## Family: Tweedie(p=1.99)
## Link function: log
##
## Formula:
## (Cper10m2 + 1) ~ factor(year) + s(doy, k = 7) + s(lon, lat) +
##   s(bottom_depth, k = 5) + s(temperature)
##
## Parametric coefficients:
##             Estimate Std. Error t value Pr(>|t|)
## (Intercept) 4.1517    0.2977 13.944 < 2e-16 ***
## factor(year)1998 -1.0782    0.5963 -1.808 0.07084 .
## factor(year)1999 -2.2057    0.3889 -5.671 1.75e-08 ***
## factor(year)2000 -1.5017    0.4649 -3.230 0.00127 **
## factor(year)2002  0.7160    0.3606  1.986 0.04729 *
## factor(year)2003 -1.6747    0.3807 -4.399 1.18e-05 ***
## factor(year)2005 -1.4158    0.3595 -3.939 8.63e-05 ***
## factor(year)2006 -1.8882    0.3157 -5.981 2.87e-09 ***
## factor(year)2007 -1.4764    0.3638 -4.059 5.23e-05 ***
## factor(year)2008 -1.8754    0.5010 -3.743 0.00019 ***
## factor(year)2009 -2.5539    0.3323 -7.684 3.01e-14 ***
## factor(year)2010 -1.6456    0.3309 -4.973 7.47e-07 ***
## factor(year)2011 -2.0361    0.4386 -4.642 3.79e-06 ***
## factor(year)2012 -0.6017    0.3315 -1.815 0.06970 .
## factor(year)2013 -1.6337    0.4065 -4.019 6.17e-05 ***
## factor(year)2014 -1.7847    0.3412 -5.230 1.97e-07 ***
## factor(year)2015 -2.8523    0.3943 -7.233 8.03e-13 ***
## factor(year)2016 -2.3765    0.3876 -6.131 1.15e-09 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

##
## Approximate significance of smooth terms:
##          edf Ref.df      F p-value
## s(doy)      1.000 1.001 0.035  0.852
## s(lon,lat) 27.506 28.848 24.626 <2e-16 ***
## s(bottom_depth) 3.700 3.939 15.342 <2e-16 ***
## s(temperature) 7.055 8.157 24.826 <2e-16 ***
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## R-sq.(adj) = 0.253 Deviance explained = 58.9%
## -REML = 4825.6 Scale est. = 1.6164 n = 1358

```

Then additive temperature and salinity, in individual additive terms. This is the second-best performing model.

```

lv.temp.sal<-readRDS("./GAM Models/nrs_larvae_addtempsal.rds")
summary(lv.temp.sal)

```

```

##
## Family: Tweedie(p=1.99)
## Link function: log
##
## Formula:
## (Cper10m2 + 1) ~ factor(year) + s(doy, k = 7) + s(lon, lat) +
##   s(bottom_depth, k = 5) + s(temperature) + s(salinity)
##
## Parametric coefficients:
##             Estimate Std. Error t value Pr(>|t|)
## (Intercept) 4.2268    0.3008 14.053 < 2e-16 ***
## factor(year)1998 -1.1341    0.5968 -1.900 0.057617 .
## factor(year)1999 -2.3507    0.3929 -5.983 2.83e-09 ***
## factor(year)2000 -1.5567    0.4687 -3.321 0.000921 ***
## factor(year)2002  0.5806    0.3664  1.585 0.113278
## factor(year)2003 -1.7345    0.3820 -4.540 6.13e-06 ***
## factor(year)2005 -1.4710    0.3629 -4.053 5.35e-05 ***
## factor(year)2006 -2.0232    0.3226 -6.271 4.86e-10 ***
## factor(year)2007 -1.6443    0.3677 -4.472 8.42e-06 ***
## factor(year)2008 -2.0644    0.5073 -4.069 5.00e-05 ***
## factor(year)2009 -2.7307    0.3376 -8.090 1.36e-15 ***
## factor(year)2010 -1.7789    0.3320 -5.358 9.97e-08 ***
## factor(year)2011 -2.1098    0.4381 -4.816 1.64e-06 ***
## factor(year)2012 -0.7948    0.3334 -2.384 0.017284 *
## factor(year)2013 -1.7046    0.4103 -4.155 3.47e-05 ***
## factor(year)2014 -1.7582    0.3440 -5.111 3.68e-07 ***
## factor(year)2015 -2.9554    0.3970 -7.444 1.77e-13 ***
## factor(year)2016 -2.3977    0.3908 -6.136 1.12e-09 ***
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Approximate significance of smooth terms:
##          edf Ref.df      F p-value
## s(doy)      1.000 1.000 0.208 0.64864

```

```

## s(lon,lat)      27.469 28.832 24.905 < 2e-16 ***
## s(bottom_depth) 3.641  3.916 14.505 < 2e-16 ***
## s(temperature)  6.956  8.079 18.627 < 2e-16 ***
## s(salinity)     4.855  6.078  3.748 0.00102 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## R-sq.(adj) =  0.278   Deviance explained = 59.4%
## -REML =    4822   Scale est. = 1.6021   n = 1358

```

```
AIC(lv.temp.sal)
```

```
## [1] 9595.43
```

And finally, the best performing model: the bivariate salinity-temperature additive term:

```
lv.2d<-readRDS("./GAM Models/nrs_larvae_2d.rds")
summary(lv.2d)
```

```

##
## Family: Tweedie(p=1.99)
## Link function: log
##
## Formula:
## (Cper10m2 + 1) ~ factor(year) + s(lon, lat) + s(doy, k = 7) +
##   s(bottom_depth) + s(salinity, temperature)
##
## Parametric coefficients:
##             Estimate Std. Error t value Pr(>|t|)
## (Intercept) 4.4280    0.3040 14.566 < 2e-16 ***
## factor(year)1998 -0.8756    0.5964 -1.468 0.142341
## factor(year)1999 -2.5886    0.3981 -6.502 1.13e-10 ***
## factor(year)2000 -1.6722    0.4648 -3.598 0.000333 ***
## factor(year)2002  0.3186    0.3761  0.847 0.397064
## factor(year)2003 -2.1518    0.3958 -5.437 6.48e-08 ***
## factor(year)2005 -1.6456    0.3679 -4.473 8.38e-06 ***
## factor(year)2006 -2.0281    0.3268 -6.206 7.31e-10 ***
## factor(year)2007 -2.0564    0.3754 -5.477 5.20e-08 ***
## factor(year)2008 -2.1762    0.5127 -4.245 2.35e-05 ***
## factor(year)2009 -2.7480    0.3388 -8.111 1.17e-15 ***
## factor(year)2010 -1.5555    0.3284 -4.737 2.41e-06 ***
## factor(year)2011 -2.5456    0.4398 -5.788 8.95e-09 ***
## factor(year)2012 -1.0979    0.3386 -3.242 0.001217 **
## factor(year)2013 -1.9252    0.4128 -4.664 3.43e-06 ***
## factor(year)2014 -2.2396    0.3509 -6.383 2.43e-10 ***
## factor(year)2015 -3.3242    0.3990 -8.332 < 2e-16 ***
## factor(year)2016 -2.8013    0.3922 -7.142 1.54e-12 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Approximate significance of smooth terms:
##          edf Ref.df      F p-value

```


```
## s(lon,lat)           27.300 28.781 19.444 <2e-16 ***
## s(doy)                1.001  1.001  1.871   0.172
## s(bottom_depth)       6.958  7.963  9.939 <2e-16 ***
## s(salinity,temperature) 24.122 27.598 12.146 <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## R-sq.(adj) =  0.279  Deviance explained = 62.1%
## -REML = 4796.5  Scale est. = 1.5229    n = 1358
```

```
AIC(lv.2d)
```


```
## [1] 9511.532
```


The following plot is the predicted NRS larval biogeography based on the best performing model, the bivariate salinity-temperature GAM. Observations (log transformed, n+1) are shown as well.

Predicted NRS Larval Biogeography, 2D Model




Predicted NRS Larval Biogeography, 2D Model





With this bivariate model, we can also calculate the predicted anomalous larval catch (more or less than expected) on a salinity-temperature plot. This figure shows that prediction, with observed larval catch ( $\log(n=1)$ ) overlaid.

```
## Warning in symbols(nrslarv.ctd$salinity[nrslarv.ctd$Cper10m2 > 0],  
## nrslarv.ctd$temperature[nrslarv.ctd$Cper10m2 > : "alpha" is not a graphical  
## parameter
```



To again share the improvements of the best performing models from the base models, we can look at the AIC division produces.

Table 2: Model Power through AIC Comparisons, Northern Rock Sole

	Best Divided By Base	Best Divided By Second Best
Larvae	0.9750076	0.9912565