



IIT ROORKEE



NPTEL ONLINE
CERTIFICATION COURSE

Confidence Interval Estimation: Single Population

Dr. A. Ramesh

Department of Management Studies
IIT ROORKEE



Goals

After completing this lecture, you should be able to:

- Distinguish between a point estimate and a confidence interval estimate
- Construct and interpret a confidence interval estimate for a single population mean using both the Z and t distributions
- Form and interpret a confidence interval estimate for a single population proportion
- Create confidence interval estimates for the variance of a normal population

Confidence Intervals


- Confidence Intervals for the Population Mean, μ
 - when Population Variance σ^2 is Known
 - when Population Variance σ^2 is Unknown
- Confidence Intervals for the Population Proportion, \hat{p} (large samples)
- Confidence interval estimates for the variance of a normal population

Definitions

- An **estimator** of a population parameter is
 - a random variable that depends on sample information . . .
 - whose value provides an approximation to this unknown parameter
- A specific value of that random variable is called an **estimate**

Point and Interval Estimates

- A **point estimate** is a single number,
- a **confidence interval** provides additional information about variability



Point Estimates

We can estimate a Population Parameter ...		with a Sample Statistic (a Point Estimate)
Mean	μ	\bar{x}
Proportion	P	\hat{p}

Unbiasedness

- A point estimator $\hat{\theta}$ is said to be an **unbiased estimator** of the parameter θ if the expected value, or mean, of the sampling distribution of $\hat{\theta}$ is θ ,


$$E(\hat{\theta}) = \theta$$

- Examples:
 - The sample mean \bar{x} is an unbiased estimator of μ
 - The sample variance s^2 is an unbiased estimator of σ^2
 - The sample proportion \hat{p} is an unbiased estimator of P

Unbiasedness

(continued)

- $\hat{\theta}_1$ is an unbiased estimator, $\hat{\theta}_2$ is biased:



Bias

- Let $\hat{\theta}$ be an estimator of θ
- The **bias** in $\hat{\theta}$ is defined as the difference between its mean and θ

$$\text{Bias}(\hat{\theta}) = E(\hat{\theta}) - \theta$$

- The bias of an unbiased estimator is 0

Most Efficient Estimator

- Suppose there are several unbiased estimators of θ
- The **most efficient estimator** or the **minimum variance unbiased estimator** of θ is the unbiased estimator with the **smallest variance**
- Let $\hat{\theta}_1$ and $\hat{\theta}_2$ be two unbiased estimators of θ , based on the same number of sample observations. Then,
 - $\hat{\theta}_1$ is said to be more efficient than $\hat{\theta}_2$ if $\text{Var}(\hat{\theta}_1) < \text{Var}(\hat{\theta}_2)$
 - The relative efficiency of $\hat{\theta}_1$ with respect to $\hat{\theta}_2$ is the ratio of their variances:

$$\text{Relative Efficiency} = \frac{\text{Var}(\hat{\theta}_2)}{\text{Var}(\hat{\theta}_1)}$$

Confidence Intervals

- How much uncertainty is associated with a point estimate of a population parameter?
- An **interval estimate** provides more information about a population characteristic than does a **point estimate**
- Such interval estimates are called confidence intervals


Confidence Interval Estimate

- An interval gives a **range** of values:
 - Takes into consideration variation in sample statistics from sample to sample
 - Based on observation from 1 sample
 - Gives information about closeness to unknown population parameters
 - Stated in terms of level of confidence
 - Can never be 100% confident

Confidence Interval and Confidence Level

- If $P(a < \theta < b) = 1 - \alpha$ then the interval from a to b is called a $100(1 - \alpha)\%$ confidence interval of θ .
- The quantity $(1 - \alpha)$ is called the confidence level of the interval (α between 0 and 1)
 - In repeated samples of the population, the true value of the parameter θ would be contained in $100(1 - \alpha)\%$ of intervals calculated this way.
 - The confidence interval calculated in this manner is written as $a < \theta < b$ with $100(1 - \alpha)\%$ confidence

Estimation Process



Confidence Level, $(1-\alpha)$

(continued)

- Suppose confidence level = 95%
- Also written $(1 - \alpha) = 0.95$
- A relative frequency interpretation:
 - From repeated samples, 95% of all the confidence intervals that can be constructed will contain the unknown true parameter
- A specific interval either will contain or will not contain the true parameter


General Formula

- The general formula for all confidence intervals is:

$$\text{Point Estimate} \pm (\text{Reliability Factor})(\text{Standard Error})$$

- The value of the reliability factor depends on the desired level of confidence

Confidence Intervals



Confidence Interval for μ (σ^2 Known)

- Assumptions
 - Population variance σ^2 is known
 - Population is normally distributed
 - If population is not normal, use large sample
- Confidence interval estimate:

$$\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} < \mu < \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

(where $z_{\alpha/2}$ is the normal distribution value for a probability of $\alpha/2$ in each tail)

Margin of Error

- The confidence interval,

$$\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} < \mu < \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

- Can also be written as $\bar{x} \pm ME$
where ME is called the **margin of error**

$$ME = z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

Reducing the Margin of Error


$$ME = z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

The margin of error can be reduced if

- the population standard deviation can be reduced ($\sigma \downarrow$)
- The sample size is increased ($n \uparrow$)
- The confidence level is decreased, $(1 - \alpha) \downarrow$

Finding the Reliability Factor, $z_{\alpha/2}$

- Consider a 95% confidence interval:



- Find $z_{.025} = \pm 1.96$ from the standard normal distribution table


Common Levels of Confidence

- Commonly used confidence levels are 90%, 95%, and 99%

Confidence Level	Confidence Coefficient, $1 - \alpha$	$Z_{\alpha/2}$ value
80%	.80	1.28
90%	.90	1.645
95%	.95	1.96
98%	.98	2.33
99%	.99	2.58
99.8%	.998	3.08
99.9%	.999	3.27

Intervals and Level of Confidence

Sampling Distribution of the Mean



Example

- A sample of 11 circuits from a large normal population has a mean resistance of 2.20 ohms. We know from past testing that the population standard deviation is 0.35 ohms.
- Determine a 95% confidence interval for the true mean resistance of the population.

Example

(continued)

- A sample of 11 circuits from a large normal population has a mean resistance of 2.20 ohms. We know from past testing that the population standard deviation is .35 ohms.

- Solution:**
$$\bar{x} \pm z \frac{\sigma}{\sqrt{n}}$$
$$= 2.20 \pm 1.96 (.35/\sqrt{11})$$
$$= 2.20 \pm .2068$$
$$1.9932 < \mu < 2.4068$$

Interpretation

- We are 95% confident that the true mean resistance is between 1.9932 and 2.4068 ohms
- Although the true mean may or may not be in this interval, 95% of intervals formed in this manner will contain the true mean

Confidence Intervals

