

***IN VITRO* CONSERVATION OF ENDANGERED ORCHID *Bulbophyllum auricomum***

**Lindl., THE ROYAL ORCHID OF MYANMAR**

**Myo Ma Ma Than<sup>1</sup>\***

Centre of Advanced Study, Department of Botany, University of Calcutta,

35, Ballygunge Circular Road, Kolkata-700 019, India

<sup>1</sup>Senior Researcher, Network Activities Group, Yangon, Myanmar

\*E-mail: myomamathan3@gmail.com

Telephone no.: 91-33-2461-5445

Fax: 91-33-2461-4849

## 24 Abstract

25 *In vitro* conservation method of *Bulbophyllum auricomum* Lindl. has been standardized/  
26 developed in the present study. Minimum shoot proliferation (7.68±0.71 mm in length) and at  
27 least 1 root/plantlet developed after 12 months when shoots germinated from protocorms were  
28 cultured incessantly on one-fourth strength MS nutrient medium with 60 gm l<sup>-1</sup> sucrose. In  
29 contrast, when the explants were maintained on full strength MS nutrient medium supplemented  
30 with 30 gm l<sup>-1</sup> sucrose, the cultures grew faster and filled the culture vessel (150 ml flask with 50  
31 ml medium) and difficult to maintain without subculture after 6 months. It was noted that all  
32 plantlets from different treatments were able to resume growth on MS basal medium after 3, 6 or  
33 12 months of continuous culture on the same medium in the same culture vessel. There was no  
34 significant difference in survival percentage of shoots (80-100 %) which were maintained on  
35 different strength of MS nutrient medium after 3, 6 and 12 months of conservation. While, one-  
36 fourth strength MS medium with high sucrose (60 gm l<sup>-1</sup>) showed a retardant effect on growth of  
37 *B. auricomum* plants.

38

39 **Key words:** *Bulbophyllum auricomum*, endangered species, *in vitro* conservation, nutrient  
40 medium, orchids

41

## 42 Introduction

43 Conservation of plant genetic resources is an essential component to sustain biodiversity for the  
44 agriculturally and horticulturally important species. It is estimated that genetic diversity of many  
45 plant species are being lost globally at a rate that is faster than at any previous time in history  
46 (IUCN 2004). This situation is even worse for orchid species partly because most of these have

Kommentar [P1]: I think "developed"

Formatiert: Hervorheben

Formatiert: Hervorheben

Kommentar [P2]: Growth?

Kommentar [P3]: After  
acclimatization?

47 habitat preference, pollinator dependence for the completion of life cycle (IUCN/SSC Orchid  
48 Specialist Group 1996) and symbiotic association with a class of fungi for seed germination  
49 under natural condition (Arditti 2008, Yam and Arditti 2009). The vast majority of orchids,  
50 especially those native to tropical regions, are currently endangered because of extensive  
51 disturbance of their natural habitat and indiscriminate harvesting of the naturally growing plants  
52 (Koopowitz et al. 1993, Than et al. 2009). Thus, *in situ* conservation became rather impossible  
53 due to the disappearance of large wild areas. *Ex situ* conservation under natural condition also  
54 limited due to labour costs for maintenance, trained personnel requirements, political and social  
55 issues, the risk of genetic wipe out as a result of natural disasters, etc. (Engelmann 1991). Hence,  
56 *in vitro* conservation of plant germplasm is becoming a complementary approach to conserve  
57 endemic and endangered plant species (Moges et al. 2003, Shibli et al. 2006). It is probably the  
58 only feasible way to maintain as a gene bank and has been applied widely for many plant species  
59 (Wannakraioj 1998, Sarkar et al. 2001, Minoo et al. 2006, Mweetwa et al. 2007, Rangsayatorn  
60 et al. 2009).

61 *In vitro* germplasm conservation can be performed using different techniques depending on the  
62 storage duration, i.e. long-term conservation (cryopreservation) and short- to medium-term  
63 conservation (to reduce the growth) (Engelmann 1991, 1998). Aseptic conservation of  
64 germplasm by controlling plant growth is a desirable method for developing country. It needs  
65 less space requirements and reduction of the expenses in labour costs for maintenance.  
66 Conservation can be executed by manipulating temperature (Withers 1982), osmotic stress  
67 (Schnapp and Preece 1986), mineral concentration (Withers 1982) or gas availability (Bridgen  
68 and Staby 1981). Synthetic seed production in several orchids via alginate encapsulation has also

69 been attempted as an alternate method for short- to mid-term storage (Saiprasad and Polisetty  
70 2003, Mohanraj et al. 2009, Gantait et al. 2012).

71 *Bulbophyllum auricomum*, a sympodial epiphytic orchid, having a geographical distribution  
72 ranging from Myanmar, Thailand, Sumatra to Java, blooms once a year (November-January) and  
73 is a profit-making orchid. This is 8-10 cm tall, ovoid-oblong pseudobulb carrying 1-2 leaves at  
74 the top and generally propagated through the division of pseudobulbs. However, the rate of  
75 multiplication is very slow as only one or two plants are produced per bulb per year (Myanma  
76 Encyclopedia 1972). Moreover, as in other orchids, the minute seeds are non-endospermic and  
77 require the association with appropriate fungi for seed germination and subsequent growth under  
78 natural condition. Due to its commercial importance and biological limitation to survive in  
79 natural environment, the species is now in the verge of extinction (Than et al. 2009, 2011, 2012).  
80 Consequently, *in vitro* conservation is of great interest for maintenance and storage of such  
81 endangered species. The objective of the present study was to develop method for *in vitro*  
82 conservation of endangered orchid *Bulbophyllum auricomum*, the royal orchid of Myanmar.

83

## 84 **Materials and methods**

### 85 **Plant Material**

86 *B. auricomum* plants were collected from their natural habitat of Yakhine Yoma mountain ranges  
87 in Myanmar and maintained in pots (Fig. 1a). Seeds of immature capsules (~ 3 months old) from  
88 a single donor plant were used in this study (Than et al. 2011). Seeds were aseptically cultured  
89 on semisolid MS medium (Murashige and Skoog 1962) (Fig. 1b). After two to three weeks of  
90 culture initiation, embryos swelled and turned into yellowish green spherules like protocorms  
91 (Fig. 1c). Shoots emerged from germinated protocorms (Fig. 1d) and shoot proliferation started

92 within six to seven weeks of *in vitro* culture. Axenic shoots of *B. auricomum* were maintained on  
93 MS (1962) nutrient medium by subculturing every two months.

#### 94 **Effect of sucrose concentrations and light regime on shoot retardation**

95 *B. auricomum* shoots derived from *in vitro* raised seedlings (~11.0 mm long) with  $19.73 \pm 2.43$   
96 mg fresh weight were used as source explant. To assess the slow growth of *B. auricomum* in  
97 culture, MS (1962) medium with 0, 20, 30, 40 and 60  $\text{gm l}^{-1}$  sucrose concentrations and light  
98 conditions were tested. Medium pH was adjusted to 5.7 and gelled with 0.75% w/v agar. Each  
99 flask contains 5 shoots and 15 replicates per treatment. Cultures were maintained at  $24 \pm 1^\circ \text{C}$   
100 under two different light regimes (16 h photoperiod or continuous illumination) with a photon  
101 flux density of  $80 \mu\text{mol m}^{-2} \text{s}^{-1}$  for 18 weeks. A better minimal growth condition was selected and  
102 employed next.

Kommentar [P4]: different

Kommentar [P5]: contained

Kommentar [P6]: flasks or shoots

Formatiert: Hervorheben

#### 103 **Effect of strength of MS basal salts**

104 *B. auricomum* shoots (~4.0 mm long) were used as explants for *in vitro* conservation. To  
105 standardize the suitable condition for maintaining the shoots under minimal growth condition,  
106 MS (1962) basal salts were used in full, half and one-fourth strength. The carbon source, sucrose,  
107 was compared at two different concentrations (30 and 60  $\text{gm l}^{-1}$ ). The 150 ml culture vessels  
108 containing 50 ml of medium were closed with aluminum foils to minimize the evaporation rate.  
109 Each flask contained 4 shoots and there were 20 replicates per treatment. Cultures were  
110 maintained at  $24 \pm 1^\circ \text{C}$  under 16 h photoperiod (artificial fluorescent light  $80 \mu\text{mol m}^{-2} \text{s}^{-1}$ ).  
111 Survival rate and viability were recorded after 3, 6 and 12 months of conservation. Survival rate  
112 was determined as the percentage of cultures showing growth of explants. To assess the recovery  
113 of storage plantlets, 20 randomly selected plantlets from different treatments were transferred to

Formatiert: Hervorheben

Kommentar [P7]: type?

114 MS medium with 30 gm l<sup>-1</sup> sucrose concentration after 3, 6 and 12 months of conservation and  
115 growth rate was recorded after 8 weeks of culture.

#### 116 **Hardening and transplantation**

117 Healthy *B. auricomum* plantlets with well-developed roots were taken out of the culture vessels  
118 and washed thoroughly under tap water to remove traces of adhering agar. These plantlets were  
119 transferred to the pots containing equal amount of charcoal and coconut husk. Pots were covered  
120 with polyethylene bags to maintain approximately 80-90 % humidity. Covers were withdrawn  
121 after 3-4 days and pots were maintained in a growth chamber at 25±1° C for 4 weeks. Seedling  
122 survival was recorded after 6 weeks of acclimatization.

#### 123 **Statistical analyses**

124 Tabulated results were analysed using one-way ANOVA (Sokal and Rohlf 1987) and standard  
125 errors (SE) of replicate data were calculated. Statistical difference between mean values was  
126 computed with algorithms of Duncan's Multiple Range Test using the Statistica Software v 5.0  
127 (StatSoft 1995). Growth Index (GI) was calculated as the final fresh weight of the explants  
128 divided by the initial fresh weight.

129

#### 130 **Results**

##### 131 **Effect of different sucrose concentrations and light regime on shoot retardation**

132 Shoots began to proliferate after four weeks of culture under different light regimes. Growth of  
133 culture regenerants is dependent on sucrose concentration in the culture medium. The minimal  
134 growth rate of shoot (10.53±0.58 mm) and root (1.73±0.33 mm) length was observed in the  
135 presence of 60 gm l<sup>-1</sup> sucrose concentration under 16 h photoperiod (Fig. 2 a,b). There were no  
136 significant differences in the number of leaf and root formation among different sucrose

Kommentar [P8]: was

137 concentrations under different light regimes (Table 1). Continuous illumination enhanced shoot  
138 multiplication and maximum number of shoots (20 shoots) was obtained in MS medium  
139 containing 30 gm l<sup>-1</sup> sucrose. Under 16 h photoperiod, 0, 30, 40 and 60 gm l<sup>-1</sup> sucrose  
140 concentrations inhibited multiple shoots formation and there was no significant difference noted  
141 in these concentrations. The minimum GI of shoot was retained when grown on MS medium  
142 supplemented with 60 gm l<sup>-1</sup> sucrose or without sucrose supplementation under 16 h photoperiod.

Kommentar [P9]: which ones?

#### 143 **Effect of MS nutrient medium strength and plant recovery**

Kommentar [P10]: What do you mean with recovery?

144 The effect of concentrations of sucrose and basal salts of MS nutrient medium on minimal  
145 growth of *B. auricomum* regenerants was studied. There was no significant difference in survival  
146 percentage of shoots (80-100 %) which were maintained on different culture medium after 3, 6  
147 and 12 months of conservation (Table 2). The rate of shoot growth was higher on full strength  
148 MS medium supplemented with 30 gm l<sup>-1</sup> sucrose concentration when compared to half or one-  
149 fourth strength MS medium. The maximum shoot length (23.60±2.74 mm) and shoot  
150 proliferation (~10 shoots) was observed on full strength MS nutrient medium supplemented with  
151 30 gm l<sup>-1</sup> sucrose concentration after 12 months of culture (Fig. 3a,b; Fig. 4a). These cultures  
152 grew faster and filled the culture vessel (150 ml flask containing 50 ml medium) after 6 months  
153 of culture if not subcultured. However, explants cultured on MS medium with sucrose 60 gm l<sup>-1</sup>  
154 resulted in retarded shoot growth. Minimum shoot length (7.68±0.71 mm) was noted when  
155 explants were cultured on one-fourth strength of MS nutrient medium with 60 gm l<sup>-1</sup> sucrose,  
156 after 12 months of culture without any subculture (Fig. 4b). Minimum shoot proliferation was  
157 obtained on one-fourth strength MS medium with 30 gm l<sup>-1</sup> or 60 gm l<sup>-1</sup> sucrose concentration,  
158 when the explants were cultured continuously for 12 months in the same culture vessels.

Kommentar [P11]: Not expected under this headline

Kommentar [P12]: It seems you like to have strong shoot growth. Did you explain what is the rate of shoot growth?

159 The optimum root induction (~ 7 roots/plant) was obtained when the shoots were grown on the  
160 medium containing full strength MS with 60 gm l<sup>-1</sup> sucrose after 6 months of culture (Fig. 3c). It  
161 was observed that prolonged period of culture inhibited new root induction and some roots  
162 turned brown when shoots were grown on the same medium for 12 months. The minimum root  
163 induction was noted on the medium containing one-fourth strength MS nutrient with 60 gm l<sup>-1</sup>  
164 sucrose (1 root/plant) after 3, 6 and 12 months of culture. The shoots grown on this culture  
165 medium supported normal growth and development of plantlets. In general, one-fourth strength  
166 MS medium with high sucrose (60 gm l<sup>-1</sup>) had a retardant effect on growth of *B. auricomum*  
167 plants.

168 Randomly selected plantlets from different treatments were transferred to normal MS medium  
169 after 3, 6 or 12 months in order to determine the growth and viability of cultures maintained in  
170 same culture vessel over long period without subculture. It was noted that all plantlets from  
171 different treatments were able to survive on MS basal medium without PGRs.

#### 172 **Hardening and transplantation of *B. auricomum* plants**

173 *B. auricomum* plants were maintained on MS basal medium devoid of PGRs (Fig. 4c) for 3  
174 months. Plantlets (4-5 cm) with well-developed roots were transplanted to the pot containing  
175 equal amount of charcoal and coconut husk for acclimatization. Charcoal was used as substrate  
176 together with coconut husk to avoid direct contact with the substrate, which retains excess  
177 moisture. Wood charcoal provides good drainage and adequate aeration to the roots, which is of  
178 primordial importance in the culture of orchids. These acclimatized plants survived and grew  
179 well but could not tolerate the high outdoor temperature during the summer season in Kolkata,  
180 India. The plantlets transplanted to the green house at Orchid Garden, Yangon, Myanmar with

**Kommentar [P13]:** From where do you know this is optimum?

**Kommentar [P14]:** Had they the same vigor?



181 80% success in establishment rate (Fig. 4d). The new shoot formation was initiated after 6  
182 weeks of acclimatization.

183

## 184 Discussion

185 The main objective of *in vitro* conservation of plant genetic resources is to increase the intervals  
186 between the subcultures under sterile condition maintaining their clonal fidelity (Moges et al.  
187 2003). Several types of plant materials such as bulblets (Kastner et al. 2001), nodal segments  
188 (Martin et al. 2007), protocorms (Rangsayatorn et al. 2009), PLBs (Kishi and Takagi 1997,  
189 Wannakraijong 1998), plantlets (Miedema 1982), seeds (Mweetwa et al. 2007), seedlings (Ming et  
190 al. 2000, Zhong et al. 2000), shoots (Agrawal et al. 1992, Minoo et al. 2006) and shoot tips (El-  
191 Gizawy and Ford-Lloyd 1987) were used as explants for *in vitro* conservation. Among them,  
192 shoot explants were most suitable for *in vitro* conservation due to high survival rate and high  
193 genetic stability (Reed et al. 1998). In the present study, axenic *B. auricomum* shoots were used  
194 as source of explants for *in vitro* conservation.

195 The growth rate of *in vitro* cultures could be retarded by using osmotic agents such as mannitol,  
196 sucrose, sorbitol, malic hydrazide, succinic acid and ancymidole (Moges et al. 2003). In the  
197 present study, sucrose is used as osmotic agent in the culture medium. Osmotic agents in the  
198 culture medium reduce mineral uptake by cells due to difference in osmotic pressures, thereby  
199 retarding plant growth (Thompson et al. 1986). Such stress condition inhibits shoot induction in  
200 cultured explants (Brown et al. 1979). The addition of osmoticum in culture medium has been  
201 proved to be efficient in reducing growth and increasing the storage life of many *in vitro* grown  
202 explants of different plant species (Wilson et al. 2000).

**Kommentar [P15]:** What is the standard interval. I guess for such slow growing cultures more than 4 weeks.

**Kommentar [P16]:** You used seed and therefore genetically heterogenic material and not clones.

**Kommentar [P17]:** Belongs to the introduction

**Kommentar [P18]:** introduction

203 Sucrose is widely used as carbon source in most of the tissue culture media (Hazarika 2003,  
 204 Arditti 2008). It functions as energy source and osmotic agent. Sucrose can also be used to retard  
 205 the plant growth *in vitro* (Moges et al. 2003). In the present study, shoot growth was retarded in  
 206 the presence of 60 gm l<sup>-1</sup> sucrose in the culture medium, which is consistent with the  
 207 observations of Van Waes and Debergh (1986) in Western European orchids, Rasmussen (1995)  
 208 in terrestrial orchids, and Wotavová-Novotná et al. (2007) in *Dactylorhiza* species. Shibli et al.  
 209 (1999) reported that the growth of bitter almond microshoots was significantly reduced in high  
 210 sucrose concentration. The similar findings have been reported in other plant species such as  
 211 *Solanum tuberosum* (Henshaw et al. 1980, Sarkar and Naik 1998), *Cymbidium* sp. (Homes et al.  
 212 1982), *Dendrobium caryseanthum* and *D. ochreatum* (Tandon and Sharma 1986), *Campsis*  
 213 *chinensis* (Paek and Kwang 1993) and *Anoectochilus formosanus* (Ket et al. 2004).  
 214 The reduction of nutrient concentration in the culture medium has been found to be beneficial for  
 215 storage of cultures (Kartha et al. 1981). The influence of low concentrations of nutrient elements  
 216 on growth limitation in *Lycopersicon esculentum* and *Dianthus caryophyllus* microplants has  
 217 been demonstrated by Schnapp and Preece (1986). They reported that microplant height and root  
 218 growth was significantly reduced when MS nutrient concentrations were decreased to one-fourth,  
 219 half- or three-fourth strength in the culture medium. Agrawal et al. (1992) also reported that  
 220 *Vanilla walkeriae* shoots were conserved for 7 months when maintained on half-strength MS  
 221 medium. In the present study, *B. auricomum* shoots were successfully maintained even on the  
 222 medium containing one-fourth strength MS nutrient. As observed in the present study, seedlings  
 223 of *Dendrobium candidum* (Ming et al. 2000) and *D. officinale* (Zhong et al. 2000) were  
 224 conserved *in vitro* for 12 months without subculture when maintained on one-fourth or half-

Kommentar [P19]: introduction

strength MS medium. In contrast, sweet potato plantlets grown on half-strength MS medium resulted in low survival (36 % - 48 %) after 4 months of conservation (Aguilar and Lopez 1993). The endemic orchid species, *Ipsea malabarica*, was maintained for 20 months on half-strength MS medium, without the addition of sucrose or PGRs (Martin and Pradeep 2003). In *Vanilla* sp., shoots were maintained for one year on full or half-strength MS nutrient with 15 gm l<sup>-1</sup> each of sucrose and mannitol (Minoo et al. 2006). In the present study, *B. auricomum* shoots could be successfully maintained for 12 months on the medium containing one-fourth strength MS nutrient with 60 gm l<sup>-1</sup> sucrose concentration. Thus, the reduction in concentrations of mineral nutrient and high sucrose concentration in the culture medium was sufficient to retard the *in vitro* growth in *B. auricomum*.

Availability of green shoots (for further micropropagation *vis-à-vis* regeneration) after protracted periods of slow-growth conservation is one of the most important features in any *in vitro* conservation programme (Roca et al. 1989). Too much reduction in microplant growth during conservation may be resulted in limited number of viable plantlets for further subculturing or regeneration (Sarkar et al. 2001). In the present investigation, shoots maintained on growth retardation medium (one-fourth strength MS medium with 60 gm l<sup>-1</sup> sucrose) survived after prolonged periods of storage and also fostered favourable plantlet to re-grow.

It may be concluded that a method for *in vitro* conservation of *B. auricomum* plants have been developed in the present study.

#### **Acknowledgements**

MMMT is thankful to the DBT-TWAS (Department of Biotechnology, Govt. of India and The World Academy of Sciences, Italy) for the award of Postdoctoral fellowship and Prof. Sumita

248 Jha, mentor, CAS, Department of Botany, University of Calcutta and Prof. Amita Pal, Bose  
249 Institute for their constant encouragement and supports.

250

251 **References**

252 Agrawal D.C., Morwal G.C., Mascarenhas A.F. (1992). *In vitro* propagation and slow growth  
253 storage of shoot cultures of *Vanilla walkeriae* Wight, an endangered orchid. *Lindleyana*, 7(2):  
254 95-99.

255 Aguilar M, Lopez M (1993) *In vitro* conservation of sweetpotato clone N-1437 with minimal  
256 growth rate. In: Reunion annual de la Sociedad del Programa Cooperativo Centro Americano de  
257 Cultivos y Animales (PCCMCA), Guatemala, 276-278 pp.

258 Arditti J. (2008). *Micropropagation of orchids*. Second Edition, John Wiley and Sons, New  
259 York, 1-88 pp.

260 Bridgen M.P., Staby G.L. (1981). Low pressure and low oxygen storage of *Nicotiana tabacum*  
261 and *Chrysanthemum Morifolium* tissue cultures. *Plant Science Letters*, 22: 177-186.

262 Brown D.C.W., Leung D.W.M., Thorpe T.A. (1979). Osmotic requirements for shoot formation  
263 in tobacco callus. *Physiologia Plantarum*, 46: 36-41.

264 El-Gizawy A.M., Ford-Lloyd B. (1987). An *in vitro* method for the conservation and storage of  
265 garlic (*Allium sativum*) germplasm. *Plant Cell, Tissue and Organ Culture*, 9: 147-150.

266 Engelmann F. (1991). *In vitro* conservation of tropical plant germplasm - a review. *Euphytica*,  
267 57: 227-243.

268 Engelmann F. (1998). *In vitro* conservation of horticultural genetic resources review of the state  
269 of the art. *ISHS: World Conference on Horticultural Research*, Rome, Italy, 17-20 June.

270 Gantait S., Bustam S., Sinniah U.R. (2012). Alginate-encapsulation, short-term storage and plant  
 271 regeneration from protocorm-like bodies of *Aranda* Wan Chark Kuan 'Blue' x *Vanda coerulea*  
 272 Griffth. ex. Lindl. (Orchidaceae). *Plant Growth Regulation*, 68(2): 303-311.  
 273 Hazarika B.N. (2003). Acclimatization of tissue-cultured plants. *Current Science*, 85(12): 25.  
 274 Henshaw G.G., O'Hara J.F., Westcott R.J. (1980) In: Ingram D.S., Helgelson J.P. (Eds) *Tissue*  
 275 *Culture Method for Plant Pathologists*. Blackwell Scientific Publications, UK, 71-76 pp.  
 276 Homes J., Dubus F., Bourdon J.L. (1982). Cold storage of plant tissue cultures. In: Fujiwara A.  
 277 (Eds). *Proceeding of 5<sup>th</sup> International Congress in Plant Tissue and Cell Culture*, Tokyo, 801-802  
 278 pp.  
 279 IUCN (2004). IUCN Red List of Threatened Species. [www.iucnredlist.org](http://www.iucnredlist.org).  
 280 IUCN/SSC Orchid Specialist Group (1996). *Orchids-Status Survey and Conservation Action*  
 281 *Plan*. Gland and Cambridge: IUCN, 3-4 pp.  
 282 Kartha K.K., Mroginski L.A., Pahl K., Leung N.L. (1981). Germplasm preservation of coffee  
 283 (*Coffea arabica* L.) by *in vitro* culture of shoot apical meristems. *Plant Science Letters*, 22: 301-  
 284 307.  
 285 Kastner U., Klahr A., Keller E.R.J., Kahane R. (2001). Formation of onion bulblets *in vitro* and  
 286 viability in medium-term storage. *Plant Cell Reports*, 20: 137-142.  
 287 Ket N.V., Hahn E.J., Park S.Y., Chakrabarty D., Paek K.Y. (2004). Micropropagation of an  
 288 endangered orchid *Anectochilus formosanus*. *Biologia Plantarum*, 48: 339-344.  
 289 Kishi F., Takagi K. (1997). Efficient method for the preservation and regeneration of orchid  
 290 protocorm-like bodies. *Scientia Horticulturae*, 68: 149-156.

291 Koopowitz H., Thornhill A., Anderson M. (1993). Species distribution profiles of the neotropical  
 292 orchids *Masdevallia* and *Dracula* (Pleurothallidinae, Orchidaceae); implications for  
 293 conservation. *Biodiversity and Conservation*, 2: 681–690.

294 Martin K.P., Pradeep A.K. (2003). Simple strategy for the *in vitro* conservation of *Ipsea*  
 295 *malabarica* an endemic and endangered orchid of the Western Ghats of Kerala, India. *Plant Cell,*  
 296 *Tissue and Organ Culture*, 74: 197–200.

297 Martin M.T., Pedranzani H.E., Sierra De Grado R. (2007). Behavior and preservation of an *in*  
 298 *vitro* collection of European aspen in Spain. *Biocell*, 31(1): 41-49.

299 Miedema P. (1982). A tissue culture technique for vegetative propagation and low temperature  
 300 preservation of *Beta vulgaris*. *Euphytica*, 31 : 635-643.

301 Ming X., Ye Q., Guo L., Zhong S.Y. (2000). *Dendrobium candidum* germplasm resources from  
 302 the low-temperature preservation. *Chinese Journal of Applied and Environmental Biology*, 6(4):  
 303 326-330.

304 Minoo D., Nirmal Babu K., Peter K.V. (2006). Conservation of *Vanilla* species, *in vitro*. *Scientia*  
 305 *Horticulturae*, 110: 175–180.

306 Moges A.D., Karam N.S., Shibli R.A. (2003). Slow growth *in vitro* preservation of African  
 307 violet (*Saintpaulia ionantha* Wendl.) shoot tips. *Advances in Horticultural Science*, 17: 1-8.

308 Mohanraj R., Ananthan R., Bai V.N. (2009). Production and storage of synthetic seeds in  
 309 *Coelogyne breviseapa* Lindl. *Asian Journal of Biotechnology*, 1: 124–128.

310 Murashige T., Skoog F. (1962). Revised medium for rapid growth and bioassays with tobacco  
 311 tissue cultures. *Physiologia Plantarum*, 15: 473-497.

312 Mweetwa A.M., Welbaum G.E., Tay D. (2007). Orchid seed storage for germplasm preservation.  
 313 *Acta Horticulturae*, 760: 2.

Formatiert: Deutsch (Deutschland)

314 Myanma Encyclopedia (1972). Tha Zin Pan. In: Myanma Encyclopedia, Yangon, Myanmar, 12:  
 315 465-466.

316 Paek K.Y., Kwang H.J. (1993). *In vitro* growth control of horticultural plants by manipulating  
 317 water potential of media with sucrose, manitol and agar. In: Quynh N.T., Uyen N.V. (Eds):  
 318 Adapted Propagation Techniques for Commercial Crops of the Tropics. IFS, Hochi Minh City,  
 319 126-136 pp.

320 Rangsayatorn N., Jina K., Pampasit S. (2009). Slow growth *in vitro* conservation of *Dendrobium*  
 321 *draconis* Rchb.f.: Effect of black Kwao Krua (*Mucuna collettii*, Lace). Acta Horticulturae, 829:  
 322 355-358.

323 Rasmussen H.N. (1995). Terrestrial orchids, from seed to mycotrophic plant. Cambridge  
 324 University Press, Cambridge, 184-196 pp.

325 Reed B.M., Paynter C.L., Denoma J., Chang Y. (1998). Techniques for medium- and long-term  
 326 storage of pear (*Pyrus* L.) genetic resources. Plant Genetic Resources Newsletter, 115: 1-5.

327 Roca W.M., Chavez R., Marin M.L., Arias D.I., Mafia G., Reyes R. (1989). *In vitro* methods of  
 328 germ-plasm conservation. Genome, 31: 813-817.

329 Saiprasad G.V.S., Polisetty R. (2003). Propagation of three orchid genera using encapsulated  
 330 protocorm-like bodies. In Vitro Cellular and Developmental Biology-Plant, 39: 42-48.

331 Sarkar D., Chakrabart S.K., Naik P.S. (2001). Slow-growth conservation of potato microplants:  
 332 efficacy of ancymidol for long-term storage *in vitro*. Euphytica, 117: 133-142.

333 Sarkar D., Naik P.S. (1998). Factors affecting minimal growth conservation of potato  
 334 micoroplants *in vitro*. Euphytica, 102: 275-280.

335 Schnapp R.S., Preece J.E. (1986). *In vitro* growth reduction of tomato and carnation microplants.  
 336 Plant Cell, Tissue and Organ Culture, 6: 3-8.

337 Shibli R.A., Shatnawi M.A., Ajlouni M.M., Jaradat A., Adham Y. (1999). Slow growth *in vitro*  
 338 conservation of bitter almond (*Amygdalus communis* L.). Advances in Horticultural Science, 13:  
 339 133-134.

340 Shibli R.A., Shatnawi M.A., Subaih W.S., Ajlouni M.M. (2006). *In vitro* conservation and  
 341 cryopreservation of plant genetic resources: A review. World Journal of Agricultural Sciences,  
 342 2(4): 372-382.

343 Sokal R., Rohlf F.J. (1987). Introduction to Biostatistics, Second Edition, Freeman W.H. and  
 344 Company, San Francisco, 132-183.

345 StatSoft INC. (1995). Statistica for Windows (Computer program). Statsoft INC., Tulsa,  
 346 Oklahoma, USA.

347 Tandon P., Sharma J. (1986). Regeneration of *Dendrobium* from cold preserved protocorms. In:  
 348 Somers D.A., Gengenbach B.G., Biesboer D.D., Hackett W.P., Green C.E. (Eds). Proceeding of  
 349 6<sup>th</sup> International Congress in Plant Tissue and Cell Culture, Minneapolis, USA, 425 pp.

350 Than M.M.M., Pal A., Jha S. (2009). *In vitro* flowering and propagation of *Bulbophyllum*  
 351 *auricomum* Lindl., the royal flower of Myanmar. Acta Horticulturae, 829: 105-111.

352 Than M.M.M., Pal A., Jha S. (2011). Chromosome number and modal karyotype in a  
 353 polysomatic endangered orchid, *Bulbophyllum auricomum* Lindl., the royal flower of Myanmar.  
 354 Plant Systematics and Evolution, 294: 167–175.

355 Than M.M.M., Pal A., Jha S. (2012) Plant regeneration from callus cultures in endangered orchid  
 356 *Bulbophyllum auricomum* lindl. Propagation of Ornamental Plants, 12(2): 102-108.

357 Thompson M.R., Douglas T.J., Obata-Sasamoto H., Thorpe T.A. (1986). Mannitol metabolism in  
 358 cultured plant cell. Physiologia Plantarum, 67: 365-369.



359 Van Waes J.M., Debergh P.C. (1986). *In vitro* germination of some Western European orchids.  
 360 *Physiologia Plantarum*, 67: 253-261.  
 361 Wannakraij S. (1998). *In vitro* preservation of *Vanda coerulea* Giff. Protocorm-like-bodies.  
 362 *Kasetsart Journal: Natural Science*, 32: 329-332.  
 363 Wilson S.B., Rajapakse N.C., Young R.E. (2000). Media composition and light affect storability  
 364 and post storage recovery of micropropagated hosta plantlets. *Journal of Horticultural Sciences*,  
 365 35: 1159-1162.  
 366 Withers L.A. (1982). *Institutes Working on the Tissue Culture for Genetic Conservation*, A  
 367 Revised List. IBPGR Secretariat, Rome, Italy, 104 pp.  
 368 Wotavová-Novotná K., Vejsadová H., Kindlmann P. (2007). Effects of sugars and growth  
 369 regulators on *in vitro* growth of *Dactylorhiza* species. *Biologia Plantarum*, 51(1): 198-200.  
 370 Yam T.W., Arditti J. (2009). History of orchid propagation: a mirror of the history of  
 371 biotechnology. *Plant Biotechnology Reports*, 3: 1–56.  
 372 Zhong S.Y., Pan R., Wang X., Ye Q., Guo L. (2000). *In vitro* conservation of *Dendrobium*  
 373 *officinale* at low temperature. *Chinese Journal of Applied and Environmental Biology*, 6(4): 326-  
 374 330.  
 375  
 376  
 377  
 378  
 379  
 380  
 381

## Tables

Table 1. Effect of different sucrose concentrations on *in vitro* shoot proliferation and root induction of *B. auricomum* under different light regimes.

Sucrose concentrations (gm l <sup>-1</sup> )	Mean no. of leaf/plant *	Mean no. of root/plant *	Mean no. of shoot/explant*	Growth Index *
<u>16 h photoperiod</u>				
0	1.27±0.17 <sup>a</sup>	4.53±0.17 <sup>a</sup>	2.67±0.24 <sup>a</sup>	5.0±0.70 <sup>a</sup>
20	1.33±0.24 <sup>a</sup>	6.27±1.09 <sup>a</sup>	4.67±0.35 <sup>ab</sup>	9.0±2.03 <sup>b</sup>
30	1.20±0.11 <sup>a</sup>	3.47±0.69 <sup>a</sup>	3.20±0.20 <sup>a</sup>	9.0±1.36 <sup>b</sup>
40	1.20±0.20 <sup>a</sup>	3.00±0.72 <sup>a</sup>	3.20±0.46 <sup>a</sup>	8.0±2.83 <sup>b</sup>
60	0.93±0.33 <sup>a</sup>	0.93±0.07 <sup>a</sup>	2.33±0.69 <sup>a</sup>	6.0±2.68 <sup>a</sup>
<u>Continuous light</u>				
0	1.53±0.59 <sup>a</sup>	3.80±2.01 <sup>a</sup>	6.20±0.87 <sup>b</sup>	9.0±1.45 <sup>b</sup>
20	1.07±0.26 <sup>a</sup>	4.07±0.35 <sup>a</sup>	3.27±0.17 <sup>a</sup>	9.0±0.40 <sup>b</sup>
30	1.67±0.17 <sup>a</sup>	5.27±0.25 <sup>a</sup>	20.27±1.35 <sup>c</sup>	28.0±3.67 <sup>c</sup>
40	1.40±0.05 <sup>a</sup>	4.93±1.79 <sup>a</sup>	4.80±0.87 <sup>ab</sup>	14.0±0.27 <sup>b</sup>
60	1.07±0.48 <sup>a</sup>	4.47±1.96 <sup>a</sup>	4.47±0.63 <sup>ab</sup>	14.0±0.53 <sup>b</sup>

\* The mean of 15 replicates (n=15) ± standard error (SE). Values followed by different letters within a column are significantly different according to Duncan's multiple range tests at  $P < 0.01$ .

Formatiert: Hervorheben

Table 2. Survival of *B. auricomum* shoots maintained on different culture media after 3, 6 and 12 months of conservation.

Basal medium	Sucrose concentrations (gm l <sup>-1</sup> )	% of survival		
		3 months	6 months	12 months
MS	30	100 <sup>a</sup>	100 <sup>a</sup>	90 <sup>a</sup>
1/2 strength MS	30	100 <sup>a</sup>	100 <sup>a</sup>	100 <sup>a</sup>
1/4 strength MS	30	100 <sup>a</sup>	95 <sup>a</sup>	95 <sup>a</sup>
MS	60	90 <sup>a</sup>	90 <sup>a</sup>	75 <sup>a</sup>
1/2 strength MS	60	90 <sup>a</sup>	85 <sup>a</sup>	85 <sup>a</sup>
1/4 strength MS	60	90 <sup>a</sup>	80 <sup>a</sup>	80 <sup>a</sup>

\* Mean of 20 replicates and values followed by same letters within a column are not significantly different according to Duncan's multiple range tests at P < 0.01.

**Kommentar [P20]:** these are qualitative data and therefore Duncans test is not possible.

419 **Legend of Figures**

420 Fig. 1 *In vitro* establishment of *B. auricomum* propagules. (a) Field grown donor plant, Bar 26  
421 mm; (b) isolated seeds *in vitro*, Bar 0.1 mm; (c) regenerated protocorms from germinated seeds,  
422 Bar 0.4 mm; (d) Emergence of shoots form protocorms, Bar 0.4 mm.

423

424 Fig. 2 Effect of different sucrose concentrations on length of (a) shoot and (b) root of *B.*  
425 *auricomum* under different light regimes after 18 weeks of *in vitro* culture.

426

427 Fig. 3 Effects of different salt strength of MS medium and sucrose concentrations on growth  
428 parameters of *B. auricomum* (a) shoot length; (b) shoots per explant and (c) number of roots after  
429 3, 6 and 12 months of *in vitro* culture without subculture.

430

431 Fig. 4 (a-b) Comparison of growth response of *B. auricomum* shoots maintained after 12 months  
432 of culture without any subculture. Shoots grown on (a) full strength MS nutrient medium  
433 supplemented with 30 gm l<sup>-1</sup> sucrose concentration, Bar 14 mm; and (b) one-fourth strength MS  
434 nutrient medium with 60 gm l<sup>-1</sup> sucrose, Bar 10 mm. (c-d) Hardening and transplantation of *B.*  
435 *auricomum*. (c) pseudobulb formation from the base of shoots, Bar 5.0 mm; (d) well-developed  
436 plantlets grown under natural condition, Bar 9.0 mm.

437

438

439

440

**Kommentar [P21]:** You cannot connect the results by lines, they are independent data. Please use columns.