


SDSS

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys. P_z : target-redshift probability conditional on being a quasar, under the quasar surface density Σ_Q .

Window: +/-2000 km/s. $\ln R$: ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

DECaLS

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys. P_z : target-redshift probability conditional on being a quasar, under the quasar surface density Σ_Q .

Window: ± 2000 km/s. $\ln R$: ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

ALLWISE

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys. P_z : target-redshift probability conditional on being a quasar, under the quasar surface density Σ_Q .

Window: +/-2000 km/s. $\ln R$: ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

PS1

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys. P_z : target-redshift probability conditional on being a quasar, under the quasar surface density Σ_Q .

Window: +/-2000 km/s. $\ln R$: ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

NSC

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys. P_z : target-redshift probability conditional on being a quasar, under the quasar surface density Σ_Q .

Window: ± 2000 km/s. $\ln R$: ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.


SkyMapper

The same five reserved quasars and five reserved field sources in every version



VHS

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys. P_z : target-redshift probability conditional on being a quasar, under the quasar surface density Σ_Q .

Window: ± 2000 km/s. $\ln R$: ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

DECaLS + ALLWISE

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys. P_z : target-redshift probability conditional on being a quasar, under the quasar surface density Σ_Q .

Window: ± 2000 km/s. $\ln R$: ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

SDSS + ALLWISE

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys. P_z : target-redshift probability conditional on being a quasar, under the quasar surface density Σ_Q .

Window: +/-2000 km/s. $\ln R$: ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

PS1 + ALLWISE

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys. P_z : target-redshift probability conditional on being a quasar, under the quasar surface density Σ_Q .

Window: +/-2000 km/s. $\ln R$: ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

ALLWISE + VHS

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys. P_z : target-redshift probability conditional on being a quasar, under the quasar surface density Σ_Q .

Window: ± 2000 km/s. $\ln R$: ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

All seven surveys

The same five reserved quasars and five reserved field sources in every version



Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys. P_z : target-redshift probability conditional on being a quasar, under the quasar surface density Σ_Q .

Window: +/-2000 km/s. $\ln R$: ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.