


# SDSS

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys.  $P_z$ : target-redshift probability conditional on being a quasar, under the quasar surface density  $\Sigma_Q$ .

Window: +/-2000 km/s.  $\ln R$ : ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

# DECaLS

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys.  $P_z$ : target-redshift probability conditional on being a quasar, under the quasar surface density  $\Sigma_Q$ .

Window:  $\pm 2000$  km/s.  $\ln R$ : ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

# ALLWISE

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys.  $P_z$ : target-redshift probability conditional on being a quasar, under the quasar surface density  $\Sigma_Q$ .

Window:  $\pm 2000$  km/s.  $\ln R$ : ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

# PS1

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys.  $P_z$ : target-redshift probability conditional on being a quasar, under the quasar surface density  $\Sigma_Q$ .

Window: +/-2000 km/s.  $\ln R$ : ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

# NSC

The same five reserved quasars and five reserved field sources in every version




Scores use ALL available bands of the selected surveys.  $P_z$ : target-redshift probability conditional on being a quasar, under the quasar surface density  $\Sigma_Q$ .

Window:  $\pm 2000$  km/s.  $\ln R$ : ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

# SkyMapper

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys.  $P_z$ : target-redshift probability conditional on being a quasar, under the quasar surface density  $\Sigma_Q$ .

Window:  $\pm 2000$  km/s.  $\ln R$ : ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

# VHS

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys.  $P_z$ : target-redshift probability conditional on being a quasar, under the quasar surface density  $\Sigma_Q$ .

Window: +/-2000 km/s.  $\ln R$ : ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

# DECaLS + ALLWISE

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys.  $P_z$ : target-redshift probability conditional on being a quasar, under the quasar surface density  $\Sigma_Q$ .

Window:  $\pm 2000$  km/s.  $\ln R$ : ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

# SDSS + ALLWISE

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys.  $P_z$ : target-redshift probability conditional on being a quasar, under the quasar surface density  $\Sigma_Q$ .

Window:  $\pm 2000$  km/s.  $\ln R$ : ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

# PS1 + ALLWISE

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys.  $P_z$ : target-redshift probability conditional on being a quasar, under the quasar surface density  $\Sigma_Q$ .

Window:  $\pm 2000$  km/s.  $\ln R$ : ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

# ALLWISE + VHS

The same five reserved quasars and five reserved field sources in every version




Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys.  $P_z$ : target-redshift probability conditional on being a quasar, under the quasar surface density  $\Sigma_Q$ .

Window: +/-2000 km/s.  $\ln R$ : ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.

# All seven surveys

The same five reserved quasars and five reserved field sources in every version



Axes: native-system luptitude colours (mag). Contours: two-colour marginals at 5%, 30%, 80% of peak; conditioned on the reference.

Scores use ALL available bands of the selected surveys.  $P_z$ : target-redshift probability conditional on being a quasar, under the quasar surface density  $\Sigma_Q$ .

Window:  $\pm 2000$  km/s.  $\ln R$ : ranking statistic with the reference-band priors and the unmodelled term. Field objects share one overlap field and have no class labels.