DMF DENTAL TRAINING CENTER: FINAL COACHING

Orthodontics

By: Dr. Mickhail Q. Francisco


1. Among the different facial types, which is associated with a paraboloid arch?
   a. Brachycephalic - wide face (Broad / square arch)
   b. Dolichocephalic - elongated face (tapering / narrow Arch)
   c. Mesocephalic
   d. Paracephalic
2. The undesirable side-effect most commonly associated with the use of a finger spring to tip the crown of a tooth is:
   a. Pain
   b. Gingival irritation
   c. Tendency for the tooth to intrude
   d. Severe mobility of the tooth
   e. Tendency for the root apex to move in the direction opposite from the crown
3. The "V" principle of growth is best illustrated by the:
   a. Nasal septum
   b. Mandibular ramus
   c. Mandibular symphysis
   d. Spheno-occipital synchondrosis
4. If the norm for the cephalometric angle SNA is 82°, a patient's reading of 90° for SNA most likely indicates:
   a. Maxillary protrusion
   b. The patient's ethnic background
   c. Protrusive maxillary incisors
   d. Dysplasia of the anterior cranial base
5. He is the father of modern orthodontics
   a. Norman Kingsley – pioneer of cleft palate therapy
   b. Edward Angle
   c. Martin Dewey - an orthodontist (subtype of agen classification) & malocclusion
   d. Pierre Fauchard - father of modern dentistry
   e. Joachim Lefoulon –
6. The person who coined the term "orthodontics"
   a. Norman Kingsley
   b. Edward Angle
   c. Martin Dewey
   d. Pierre Fauchard

e. Joachim Lefoulon


7. An orthodontist who pioneered the techniques in the cleft palate therapy
   a. Norman Kingsley
   b. Edward Angle
   c. Martin Dewey
   d. Pierre Fauchard
   e. Joachim Lefoulon
8. A technique to assess tooth size discrepancy that determines the ratio or proportion of the mesiodistal widths of the maxillary teeth to the mandibular teeth using a certain formula.
   a. MDA
   b. Bolton's analysis
   c. Howe's analysis
   d. Pont's index
9. Radiographs of a 5-year-old patient show that the permanent maxillary 1st molars are inclined mesially with associated resorption of the distal portions of the roots of primary second molars. This condition is known as:
   a. Ankylosis
   b. Ectopic eruption
   c. Premature eruption
   d. Internal resorption
   e. Intrafollicular resorption
10. The ideal eruption sequence of permanent teeth in the maxillary arch
    a. 61234578
    b. 61235478
    c. 61245378
    d. 61243578
11. Which of the following is not a space maintainer?
    a. Lingual arch
    b. Nance holding arch
    c. Class III restoration
    d. Stainless steel crown
    e. Palatal expansion appliance
12. On the first dental visit, the basic fear of a child under the age of 2 is related to:
    a. Fear of injection
    b. Anxiety over being separated from a parent
    c. Anxiety regarding the instruments used to perform dental treatment
    d. Not understanding the reason for dental treatment
13. The maxillary 2nd premolar erupts around what age?
    a. 11 years old
    b. 9 years old
    c. 13 years old
    d. 8 years old
14. A child who is reluctant to accept treatment, exhibit negative attitude, but not so pronounced is classified under Frankl?

a. I – definitely negative; crying/fearful
b. II – negative
c. III – positive; accepts the treatment/follows the dentist/xenjoy
d. IV – definitely positive; enjoys the treatment (love)

15. A major disadvantage of treatment using cervical headgear is:
    a. Impaction of maxillary canines
    b. Extrusion of maxillary incisors
    c. Extrusion of maxillary molars
    d. Potential deformity of the neck
    e. Psychologic trauma due to appearance



16. Crowding that occurs with mandibular incisors after age 18 is most often the result of:
    a. Orthodontic relapse
    b. Periodontal disease
    c. Impacted third molars
    d. Physiologic maturational change




17. Using a topical fluoride rinse before acid etch direct bonding of orthodontic brackets is contraindicated because fluoride:
    a. Decreases the solubility of enamel
    b. Increases the pH of the etching agent
    c. Causes copious amounts of saliva
    d. Directly reacts chemically with the bonding agent



18. Ellis classification of tooth trauma with pulpal exposure
    a. Class 1 – enamel only
    b. Class 2 – enamel & dentin w/out pulp exposure
    c. Class 3

19. The space maintainer of choice for early loss of primary mandibular canine is _____
    a. Lingual arch w/ spurs
    b. Band and loop
    c. Nance appliance
    d. Transpalatal arch



20. If a primary tooth has been extracted before succedaneous tooth has begun eruptive movement, its eruption will be _____.
    a. Hastened
    b. Delayed
    c. Not affected

21. Which of the following is a downside of using a properly fabricated lower lingual holding arch?
    a. Doesn't hold the leeway space
    b. May distalize the molars by 1 to 2mm if activated
    c. Food entrapment
    d. May cause interference with the eruption of premolars

22. The band and crib space maintainer is classified as
    a. Bilateral, fixed and functional
    b. Unilateral, fixed and functional
    c. Unilateral, fixed and nonfunctional

d. Unilateral, removable and nonfunctional

23. Frankfort horizontal is a reference plane constructed, by joining which of the following landmarks?
    a. Porion & Sella
    b. Porion & Nasion
    c. Porion & Orbitale
    d. Nasion & Sella
24. Following extraction of the mandibular primary second molar in a 3 ½ year old child, the treatment of choice is to
    a. Place a removable appliance to replace the extracted tooth
    b. Allow the permanent first molar to erupt and then maintain space for the premolar
    c. Place a fixed appliance to guide the eruption of the permanent first molar
    d. Place a removable appliance to replace the extracted tooth and prevent elongation of the opposing primary molar
25. In cephalometric radiographs, porion is found at the most _____ margin in relation to the EAM.
    a. Anterior
    b. Posterior
    c. Inferior
    d. Superior
26. According to the piezoelectric or bioelectric theory, certain ions are present during orthodontic tooth movements. Which ions are present in the bone receiving pressure?
    a. Cations
    b. Anions
    c. Both cations and anions
    d. None of the above




27. The crown of a maxillary central incisor is being tipped labially. During the process of tooth movement, the position of the apex is maintained. This is an example of _____
    a. Controlled tipping
    b. Uncontrolled tipping
    c. Labial torque
    d. Crown torque
28. Movement of canines when utilizing class III inter-arch elastics
    a. Mesial intrusion
    b. Distal intrusion
    c. Mesial extrusion
    d. Distal extrusion
29. It refers to the tendency of an object to rotate after a force is applied
    a. Torque
    b. Moment
    c. Tipping
    d. Rotation
30. The two most difficult orthodontic tooth movement to achieve are _____
    a. Intrusion and Extrusion

b. Extrusion and Tipping
c. Rotation and Intrusion
d. Intrusion and Translation

31. What appliance uses a band-free technique?
    a. Begg
    b. Ribbon arch
    c. Edgewise
    d. Universal
32. These are spaces found distal to the primary mandibular canines and mesial of the primary maxillary canines
    a. Diastema
    b. Developmental space
    c. Growth space
    d. Primate space
33. Among the following types of appliances, which is contraindicated for patients with open bite?
    a. Straight-pull headgear
    b. Cervical-pull headgear
    c. Tongue crib
    d. Palatal crib
34. Type of growth movement which refers to the movement of bone as whole unit due to the growth of adjacent bones
    a. Shifting
    b. Deposition
    c. Relocation
    d. Displacement
35. What is a second-order bend?
    a. A bend to provide correct angulation of a tooth in labiolingual direction (torque)
    b. A bend to position a tooth buccolingually
    c. A bend to rotate a tooth
    d. A bend to provide angulation of a tooth in mesiodistal direction (tip)
36. Arch shape & symmetry are best evaluated from the:
    a. Study cast
    b. Panoramic x-ray
    c. Frontal photograph
    d. Lateral cephalogram
    e. Intraoral photograph
37. A maxillary lateral incisor that is erupting is observed to be lingually locked between the central incisor and canine. Which of the following is the best treatment of choice?
    a. Refer to an orthodontist
    b. Observe
    c. Tongue blade
    d. Extract the deciduous canine
38. What is the process wherein the bone grows by the addition of bone tissue on one side of a bony cortex and taking away from the other side?
    a. Deposition
    b. Translocation

c. Drift
d. Resorption

39. The growth of the mandible primarily takes place at the
    a. Symphysis and alveolar bone
    b. Ramus and condyle
    c. Angle
    d. Teeth



40. In the cervical area of primary teeth, the enamel rods project _____ from the DEJ
    a. Superiorly
    b. Inferiorly
    c. Anteriorly
    d. Posteriorly

41. According to the principle of primary molar relationship. If there is a mesial step, the permanent first molars will have an initial relationship of _____ assuming that there is relatively equal growth between maxilla and mandible.
    a. Class I
    b. Class II
    c. Class III
    d. Cusp to cusp

42. It is the undesirable consequence for uprighting a mandibular first molar with a loop lingual arch
    a. Supraeruption of molar
    b. Mesial tipping of molar
    c. Flaring of incisors
    d. Mesial tilting of opposing molar

43. When the tooth is being translated, the center of rotation is _____
    a. Apex of the root
    b. Infinity
    c. Center of the root
    d. Fulcrum

44. Which of the following has the most fluoride content?
    a. Vegetable
    b. Meat
    c. Egg
    d. Salmon

45. Frankl behavioral rating scale of child that shows interest in the dental clinic and enjoys the treatment.
    a. Frankl I
    b. Frankl II
    c. Frankl III
    d. Frankl IV

46. 1ppm is equivalent to 1mg/_____
    a. 10ml
    b. 100ml
    c. 1000ml
    d. 10,000ml

47. The presence of multiple anterior crossbite in a 4-year-old child usually indicates which of the following developmental malocclusions?
    a. Skeletal class I malocclusion
    b. Skeletal class II malocclusion
    c. Skeletal class III malocclusion
    d. None. The child has normal development
48. It is the primary reason why we restore the proximal surfaces of deciduous teeth to its original MD dimension
    a. To prevent recurrent caries
    b. Maintenance of arch length
    c. Esthetics
    d. Prevent food impaction
49. Best restoration or treatment for primary teeth with dentinogesis imperfecta
    a. Crowns
    b. Laminates
    c. Partial crowns
    d. Any of the above
50. When comparing cementum to bone in their response to orthodontic forces, cementum resorbs:
    a. More readily
    b. Less readily
    c. Not at all
    d. Under lighter loads
    e. By undermining resorption
51. ANB of -2.5 degrees
    a. Class I
    b. Class II
    c. Class III
    d. Class IV
52. The ideal appliance to be used to correct a child with developing skeletal class II malocclusion.
    a. Cervical pull headgear
    b. Chin cup
    c. Facemask
    d. Edgewise appliance
53. It is the space maintainer of choice for a patient with bilateral and multiple tooth loss in the maxillary arch
    a. Nance
    b. RPD
    c. Band and loop
    d. LHA
54. A 9-year-old child with his parent comes into your clinic for routine dental examination. Upon examination, the permanent maxillary central and lateral incisors are already erupted. Beside the maxillary laterals are the primary canines. There is also a 1mm midline diastema. Which of the following is most probably true regarding the case of the patient?
    a. The patient has a thumb sucking habit
    b. There is a high chance of mesiodens present between 11 and 21
    c. The patient has an abnormal frenal attachment that causes the midline diastema

d. The child has normal development

55. All of the following are the primary considerations of an orthodontist during treatment except one. Which is it?
    a. Muscles
    b. Joints
    c. Teeth
    d. Gingiva
56. Which of the following is true regarding tongue thrusting? It is a/an:
    a. Retention of infantile swallowing
    b. Retention of adult swallowing
    c. Post-thumbsucking habit
    d. Indication of tongue anomaly
57. If a patient did not develop any tooth, this will primarily affect the growth of which of the following structures?
    a. Maxillary bone
    b. Mandibular bone
    c. Both maxillary and mandibular bone
    d. Alveolar bone
58. All are cephalometric landmarks, except one. Which is it?
    a. Sella
    b. Orbitale
    c. Pogonion
    d. Y-axis
59. A theory of craniofacial growth describing the importance of the nasal septum in the growth of the nasomaxillary complex.
    a. Moss' theory
    b. Scott's theory
    c. Sicher's theory
    d. Genetic theory
60. Which of the following is an indication of a distal shoe space maintainer?
    a. Extraction of first molars at age of 8
    b. Extraction of first molar at age of 12
    c. Early loss of primary 2nd molar before the eruption of permanent 1st molars
    d. All of the above
    e. None of the above

61. The least desirable method used in child management is
    a. Tell-show-do technique
    b. Gift before treatment
    c. Voice control
    d. Non-verbal communication
62. It is a space between the maxillary central incisors that is normal and usually present in a developing child
    a. Leeway space
    b. Diastema
    c. Primate space

d. Freeway space

63. It is a measure taken at the midline from a point midway between the central incisors to a tangent touching the distal surfaces of the second primary molars or the mesial surfaces of the first permanent molars.
    a. Arch perimeter
    b. Arch length
    c. Arch width
    d. Intercanine width
64. According to simon's classification of malocclusion, supraeruption of maxillary canine is an example of
    a. Attraction
    b. Abstraction
    c. Contraction
    d. Distraction
65. A lingual crossbite of maxillary 1st premolar is an example of _____
    a. Attraction
    b. Abstraction
    c. Contraction
    d. Distraction
66. During an orthodontic tooth movement, which cells are active in the pressure side?
    a. Osteoblast
    b. Osteoclast
    c. Undifferentiated mesenchymal cells
    d. Fibroblast
67. It is the other term for point A
    a. Subspinale
    b. Supramentale — point B
    c. Submentale
    d. Supraspinale
68. This is the primary anchorage component of the Hawley appliance
    a. Labial bow
    b. Clasp arms
    c. Interproximal clasp
    d. Acrylic base
69. Which may be a sign that a patient has a pseudoclass III and not a true skeletal class III malocclusion?
    a. The patient has a posterior crossbite
    b. Ability of the patient to retrude the mandible to correct the anterior crossbite
    c. Pain when retruding the mandible to correct the crossbite.
    d. Presence of an anterior crossbite
70. When treating pediatric patients with special conditions, which of the following is the best way to manage them?
    a. Ask the parent to bring the child regularly to the clinic so he/she will be accustomed to the environment
    b. Use general anesthetic drugs to put the patient to sleep while doing the procedure
    c. Use restraining devices to prevent the child from moving during treatment

d. Refer the child to a specialist

71. Which feature will you expect in a patient with a class II division II malocclusion?
    a. The MB cusp of the maxillary first molar coincides with the MB groove of the mandibular first molar.
    b. maxillary central incisors are inclined lingually
    c. The maxillary central incisors are proclined
    d. the mandibular plane is flat

72. hyalinization is present in which of the following types of orthodontic forces?
    a. Intermittent light force only
    b. Only continuous heavy force
    c. Any type of heavy forces
    d. Continuous light force only

73. During the treatment of a patient with posterior crossbite, how much expansion is produced per quarter turn of an expansion screw?
    a. 0.25mm
    b. 0.5mm
    c. 0.75mm
    d. 1.0mm

74. Which of the following is the correction definition of the center of resistance?
    a. The center of resistance is a point in which if a force is applied results to a tipping movement.
    b. The center of resistance is a point in which if a force is applied results to a rotational movement.
    c. The center of resistance is a point in which if a force is applied results to absence of movement.
    d. The center of resistance is a point in which if a force is applied results to a bodily movement.

75. The center of resistance of monorooted teeth is between 1/3 to ½ of root length apical to the alveolar crest. While for multirooted teeth the center of resistance is 1-2mm apical to the bifurcation.
    a. The first statement is true, the second is false
    b. The first statement is false, the second is true
    c. Both statements are true
    d. Both statements are false