

Ferrers graph

Ferrers graph

It is a graph to represent a partition by an array of dots.

Ferrers graph

It is a graph to represent a partition by an array of dots.
It has the following property.

Ferrers graph

It is a graph to represent a partition by an array of dots.

It has the following property.

- (i) There is one row for each part.
- (ii) The number of dots in any row is the size of that part.
- (iii) An upper row always contains at least as many dots as a lower row.
- (iv) The rows are aligned to the left.

Ferrers graph


It is a graph to represent a partition by an array of dots.

It has the following property.

- (i) There is one row for each part.
- (ii) The number of dots in any row is the size of that part.
- (iii) An upper row always contains at least as many dots as a lower row.
- (iv) The rows are aligned to the left.

Example:

Consider the partition 5 3 2 2.



The partition obtained by reading the Ferrers graph by column is called *conjugate partition*.

The partition obtained by reading the Ferrers graph by column is called *conjugate partition*.

The conjugate partition of 5 3 2 2 is 4 4 2 1 1.

The partition obtained by reading the Ferrers graph by column is called *conjugate partition*.

The conjugate partition of 5 3 2 2 is 4 4 2 1 1.

A partition whose Ferrers graph reads the same by rows and by columns is called **self-conjugate**.

Example: 5 4 2 2 1, 4 3 2 1.

• • • • •

• • • •

• •

• •

•

Problems

Problems

Q1. Show that the number of partitions of n is equal to number of partitions of $2n$ with exactly n parts.

Problems

Q1. Show that the number of partitions of n is equal to number of partitions of $2n$ with exactly n parts.

Soln: Consider a Ferrers graph of a partition of n . Add a column of n dots on the left of the graph.

Problems

Q1. Show that the number of partitions of n is equal to number of partitions of $2n$ with exactly n parts.

Soln: Consider a Ferrers graph of a partition of n . Add a column of n dots on the left of the graph. Then the graph represents a partition of $2n$ with exactly n parts.


Problems

Q1. Show that the number of partitions of n is equal to number of partitions of $2n$ with exactly n parts.

Soln: Consider a Ferrers graph of a partition of n . Add a column of n dots on the left of the graph. Then the graph represents a partition of $2n$ with exactly n parts.



Partition of $n=7$



Partition of 14 with 7 parts.



Partition of $n=7$



Partition of 14 with 7 parts.

Problems

Q1. Show that the number of partitions of n is equal to number of partitions of $2n$ with exactly n parts.

Soln: Consider a Ferrers graph of a partition of n . Add a column of n dots on the left of the graph. Then the graph represents a partition of $2n$ with exactly n parts.



Partition of $n=7$



Partition of 14 with 7 parts.



Partition of $n=7$



Partition of 14 with 7 parts.

Consider a partition of $2n$ with exactly n parts. Then the left most column contains n dots.

Problems

Q1. Show that the number of partitions of n is equal to number of partitions of $2n$ with exactly n parts.

Soln: Consider a Ferrers graph of a partition of n . Add a column of n dots on the left of the graph. Then the graph represents a partition of $2n$ with exactly n parts.



Partition of $n=7$



Partition of 14 with 7 parts.



Partition of $n=7$



Partition of 14 with 7 parts.

Consider a partition of $2n$ with exactly n parts. Then the left most column contains n dots. Eliminating the first column, results in a partition of n .

Problems

Q1. Show that the number of partitions of n is equal to number of partitions of $2n$ with exactly n parts.

Soln: Consider a Ferrers graph of a partition of n . Add a column of n dots on the left of the graph. Then the graph represents a partition of $2n$ with exactly n parts.



Partition of $n=7$



Partition of 14 with 7 parts.



Partition of $n=7$



Partition of 14 with 7 parts.

Consider a partition of $2n$ with exactly n parts. Then the left most column contains n dots. Eliminating the first column, results in a partition of n . Thus, for every partition of n dots there corresponds a partition of $2n$ with exactly n parts and vice versa.

Problems

Q1. Show that the number of partitions of n is equal to number of partitions of $2n$ with exactly n parts.

Soln: Consider a Ferrers graph of a partition of n . Add a column of n dots on the left of the graph. Then the graph represents a partition of $2n$ with exactly n parts.



Partition of $n=7$



Partition of 14 with 7 parts.



Partition of $n=7$



Partition of 14 with 7 parts.

Consider a partition of $2n$ with exactly n parts. Then the left most column contains n dots. Eliminating the first column, results in a partition of n . Thus, for every partition of n dots there corresponds a partition of $2n$ with exactly n parts and vice versa. Hence, the number of partitions of n is equal to number of partitions of $2n$ with exactly n parts.

Q2. Show that number of partitions of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Q2. Show that number of partitions of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Soln: Consider the Ferrers graph representation of a partition of n with no part greater than k .

Q2. Show that number of partitions of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Soln: Consider the Ferrers graph representation of a partition of n with no part greater than k .

The number of dots in each row is less than or equal to k .

Q2. Show that number of partitions of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Soln: Consider the Ferrers graph representation of a partition of n with no part greater than k .

The number of dots in each row is less than or equal to k .

Read the partition column-wise.

Q2. Show that number of partitions of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Soln: Consider the Ferrers graph representation of a partition of n with no part greater than k .

The number of dots in each row is less than or equal to k .

Read the partition column-wise. Then the number of parts is $\leq k$.

Q2. Show that number of partitions of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Soln: Consider the Ferrers graph representation of a partition of n with no part greater than k .

The number of dots in each row is less than or equal to k .

Read the partition column-wise. Then the number of parts is $\leq k$.

Thus, for a partition of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Q2. Show that number of partitions of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Soln: Consider the Ferrers graph representation of a partition of n with no part greater than k .

The number of dots in each row is less than or equal to k .

Read the partition column-wise. Then the number of parts is $\leq k$.

Thus, for a partition of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Conversely, consider a partition with atmost k parts, then the number of rows is $\leq k$ in the corresponding Ferrers graph.

Q2. Show that number of partitions of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Soln: Consider the Ferrers graph representation of a partition of n with no part greater than k .

The number of dots in each row is less than or equal to k .

Read the partition column-wise. Then the number of parts is $\leq k$.

Thus, for a partition of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Conversely, consider a partition with atmost k parts, then the number of rows is $\leq k$ in the corresponding Ferrers graph.

Hence, in the conjugate partition, size of any column is $\leq k$.

Q2. Show that number of partitions of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Soln: Consider the Ferrers graph representation of a partition of n with no part greater than k .

The number of dots in each row is less than or equal to k .

Read the partition column-wise. Then the number of parts is $\leq k$.

Thus, for a partition of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Conversely, consider a partition with atmost k parts, then the number of rows is $\leq k$ in the corresponding Ferrers graph.

Hence, in the conjugate partition, size of any column is $\leq k$.

Thus, for every partition with atmost k parts, there corresponds a partition with no part greater than k .

Q2. Show that number of partitions of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Soln: Consider the Ferrers graph representation of a partition of n with no part greater than k .

The number of dots in each row is less than or equal to k .

Read the partition column-wise. Then the number of parts is $\leq k$.

Thus, for a partition of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Conversely, consider a partition with atmost k parts, then the number of rows is $\leq k$ in the corresponding Ferrers graph.

Hence, in the conjugate partition, size of any column is $\leq k$.

Thus, for every partition with atmost k parts, there corresponds a partition with no part greater than k .

Hence, number of partitions of an integer n with no part greater than k is equal to the number of partition of n with atmost k parts.

Q3. Show that the number of partitions of n in which no part is smaller than 3 is equal to the number of partitions of $n + 3$ in which the three largest parts are consecutive integers.

Soln: Consider the Ferrers diagram of a partition of n with no part smaller than 3. Then the last row contains at least 3 dots, and therefore in the conjugate of this diagram the first three rows have the same length. Now, add 2 dots to the first row, and 1 dot to the second row. The resulting diagram has $n + 3$ dots and the first three rows have consecutive lengths. Thus, it represents a partition of $n + 3$ in which the three largest parts are consecutive integers. Both the operations of conjugations and adding 2 dots to the first row and 1 dot to the second are invertible, and therefore there is a bijection between the partitions of the two kinds, showing that they are equinumerous.

Q4. Obtain a generating function for the partition of n with exactly k parts.

Q4. Obtain a generating function for the partition of n with exactly k parts.

Solution:

Q4. Obtain a generating function for the partition of n with exactly k parts.

Solution: Enumerator for partitions of n with exactly k parts =
(Enumerator with atmost k parts) $-$ (Enumerator with atmost $(k - 1)$ parts).

Q4. Obtain a generating function for the partition of n with exactly k parts.

Solution: Enumerator for partitions of n with exactly k parts =
(Enumerator with atmost k parts) $-$ (Enumerator with atmost $(k - 1)$ parts).
 $=$ (Enumerator with no part greater than k) $-$ (Enumerator with no part greater than $(k - 1)$)

Q4. Obtain a generating function for the partition of n with exactly k parts.

Solution: Enumerator for partitions of n with exactly k parts =
(Enumerator with atmost k parts) – (Enumerator with atmost $(k - 1)$ parts).

= (Enumerator with no part greater than k) – (Enumerator with no part greater than $(k - 1)$)

$$= \{(1-x)^{-1}(1-x^2)^{-1} \dots (1-x^k)^{-1}\} - \{(1-x)^{-1}(1-x^2)^{-1} \dots (1-x^{k-1})^{-1}\}$$

Q4. Obtain a generating function for the partition of n with exactly k parts.

Solution: Enumerator for partitions of n with exactly k parts =
(Enumerator with at most k parts) – (Enumerator with at most $(k - 1)$ parts).

= (Enumerator with no part greater than k) – (Enumerator with no part greater than $(k - 1)$)

$$= \{(1-x)^{-1}(1-x^2)^{-1} \dots (1-x^k)^{-1}\} - \{(1-x)^{-1}(1-x^2)^{-1} \dots (1-x^{k-1})^{-1}\}$$

$$= \{(1-x)^{-1}(1-x^2)^{-1} \dots (1-x^{k-1})^{-1}\}((1-x^k)^{-1} - 1)$$

Q4. Obtain a generating function for the partition of n with exactly k parts.

Solution: Enumerator for partitions of n with exactly k parts =
(Enumerator with at most k parts) – (Enumerator with at most $(k - 1)$ parts).

= (Enumerator with no part greater than k) – (Enumerator with no part greater than $(k - 1)$)

$$= \{(1-x)^{-1}(1-x^2)^{-1} \dots (1-x^k)^{-1}\} - \{(1-x)^{-1}(1-x^2)^{-1} \dots (1-x^{k-1})^{-1}\}$$

$$= \{(1-x)^{-1}(1-x^2)^{-1} \dots (1-x^{k-1})^{-1}\}((1-x^k)^{-1} - 1)$$

$$= (1-x)^{-1}(1-x^2)^{-1} \dots (1-x^{k-1})^{-1} x^k (1-x^k)^{-1}$$

Q4. Obtain a generating function for the partition of n with exactly k parts.

Solution: Enumerator for partitions of n with exactly k parts =
(Enumerator with at most k parts) – (Enumerator with at most $(k - 1)$ parts).

= (Enumerator with no part greater than k) – (Enumerator with no part greater than $(k - 1)$)

$$= \{(1-x)^{-1}(1-x^2)^{-1} \dots (1-x^k)^{-1}\} - \{(1-x)^{-1}(1-x^2)^{-1} \dots (1-x^{k-1})^{-1}\}$$

$$= \{(1-x)^{-1}(1-x^2)^{-1} \dots (1-x^{k-1})^{-1}\}((1-x^k)^{-1} - 1)$$

$$= (1-x)^{-1}(1-x^2)^{-1} \dots (1-x^{k-1})^{-1} x^k (1-x^k)^{-1}$$

Coefficient of x^n in $(1-x)^{-1}(1-x^2)^{-1} \dots (1-x^{k-1})^{-1} x^k (1-x^k)^{-1}$ is the number of partitions of n with exactly k parts.

Q5. Prove that the number of partition of n with exactly k parts is equal to the number of partition of $n - k$ with no part greater than k .

Q5. Prove that the number of partition of n with exactly k parts is equal to the number of partition of $n - k$ with no part greater than k .

Solution:

Q5. Prove that the number of partition of n with exactly k parts is equal to the number of partition of $n - k$ with no part greater than k .

Solution:

The number of partition of $n - k$ with no part greater than k is coefficient of x^{n-k} in $(1 - x)^{-1}(1 - x^2)^{-1} \dots (1 - x^{k-1})^{-1}(1 - x^k)^{-1}$.

Q5. Prove that the number of partition of n with exactly k parts is equal to the number of partition of $n - k$ with no part greater than k .

Solution:

The number of partition of $n - k$ with no part greater than k is coefficient of x^{n-k} in $(1 - x)^{-1}(1 - x^2)^{-1} \dots (1 - x^{k-1})^{-1}(1 - x^k)^{-1}$.

The number of partition of n with exactly k parts is equal to coefficient of x^n in $x^k(1 - x)^{-1}(1 - x^2)^{-1} \dots (1 - x^{k-1})^{-1}(1 - x^k)^{-1}$

Q5. Prove that the number of partition of n with exactly k parts is equal to the number of partition of $n - k$ with no part greater than k .

Solution:

The number of partition of $n - k$ with no part greater than k is coefficient of x^{n-k} in $(1 - x)^{-1}(1 - x^2)^{-1} \dots (1 - x^{k-1})^{-1}(1 - x^k)^{-1}$.

The number of partition of n with exactly k parts is equal to coefficient of x^n in $x^k(1 - x)^{-1}(1 - x^2)^{-1} \dots (1 - x^{k-1})^{-1}(1 - x^k)^{-1}$

which is equal to the coefficient of x^{n-k} in $(1 - x)^{-1}(1 - x^2)^{-1} \dots (1 - x^{k-1})^{-1}(1 - x^k)^{-1}$.

Q5. Prove that the number of partition of n with exactly k parts is equal to the number of partition of $n - k$ with no part greater than k .

Solution:

The number of partition of $n - k$ with no part greater than k is coefficient of x^{n-k} in $(1 - x)^{-1}(1 - x^2)^{-1} \dots (1 - x^{k-1})^{-1}(1 - x^k)^{-1}$.

The number of partition of n with exactly k parts is equal to coefficient of x^n in $x^k(1 - x)^{-1}(1 - x^2)^{-1} \dots (1 - x^{k-1})^{-1}(1 - x^k)^{-1}$

which is equal to the coefficient of x^{n-k} in $(1 - x)^{-1}(1 - x^2)^{-1} \dots (1 - x^{k-1})^{-1}(1 - x^k)^{-1}$.

Which is equal to the number of partition of $n - k$ with no part greater than k .

Q5. Prove that the number of partition of n with exactly k parts is equal to the number of partition of $n - k$ with no part greater than k .

Solution:

The number of partition of $n - k$ with no part greater than k is coefficient of x^{n-k} in $(1 - x)^{-1}(1 - x^2)^{-1} \dots (1 - x^{k-1})^{-1}(1 - x^k)^{-1}$.

The number of partition of n with exactly k parts is equal to coefficient of x^n in $x^k(1 - x)^{-1}(1 - x^2)^{-1} \dots (1 - x^{k-1})^{-1}(1 - x^k)^{-1}$

which is equal to the coefficient of x^{n-k} in $(1 - x)^{-1}(1 - x^2)^{-1} \dots (1 - x^{k-1})^{-1}(1 - x^k)^{-1}$.

Which is equal to the number of partition of $n - k$ with no part greater than k .

Hence, the number of partition of n with exactly k parts is equal to the number of partition of $n - k$ with no part greater than k .