

COMBINATIONS AND PERMUTATIONS

COMBINATIONS AND PERMUTATIONS

COMBINATIONS AND PERMUTATIONS

Reference books;

1. C L Liu, Elements of Discrete Mathematics.
2. E S Page and L B Wilson, An introduction to Computational Combinatorics.
3. Alan Takkar, Applied Combinatorics.

COMBINATIONS AND PERMUTATIONS

COMBINATIONS AND PERMUTATIONS

An r -combination of n elements: is a selection of r of the objects where the order of the objects in the selection is immaterial.

COMBINATIONS AND PERMUTATIONS

An r -combination of n elements: is a selection of r of the objects where the order of the objects in the selection is immaterial.

An r - permutation of n elements is an ordered selection of r of the objects.

COMBINATIONS AND PERMUTATIONS

An r -combination of n elements: is a selection of r of the objects where the order of the objects in the selection is immaterial.

An r - permutation of n elements is an ordered selection of r of the objects.

- **1. Permutation with no repetition:** The number of r -permutations of n objects,
 $P(n, r) = {}^n P_r$

COMBINATIONS AND PERMUTATIONS

An r -combination of n elements: is a selection of r of the objects where the order of the objects in the selection is immaterial.

An r - permutation of n elements is an ordered selection of r of the objects.

- **1. Permutation with no repetition:** The number of r -permutations of n objects, $P(n, r) = {}^n P_r$
- **2. Permutation with unlimited repetition:** The no of r - permutations with unlimited repetition n^r .

COMBINATIONS AND PERMUTATIONS

An r -combination of n elements: is a selection of r of the objects where the order of the objects in the selection is immaterial.

An r - permutation of n elements is an ordered selection of r of the objects.

- **1. Permutation with no repetition:** The number of r -permutations of n objects, $P(n, r) = {}^n P_r$

- **2. Permutation with unlimited repetition:** The no of r - permutations with unlimited repetition n^r .

- **3. Permutation with restricted repetition:** If there n objects of which m_1 are of the first kind, m_2 are of the second kind, ..., m_k are of the k^{th} kind, so that $\sum_{i=1}^k m_i = n$. The number of permutations of all the objects in this case is

COMBINATIONS AND PERMUTATIONS

An r -combination of n elements: is a selection of r of the objects where the order of the objects in the selection is immaterial.

An r - permutation of n elements is an ordered selection of r of the objects.

- **1. Permutation with no repetition:** The number of r -permutations of n objects, $P(n, r) = {}^n P_r$

- **2. Permutation with unlimited repetition:** The no of r - permutations with unlimited repetition n^r .

- **3. Permutation with restricted repetition:** If there n objects of which m_1 are of the first kind, m_2 are of the second kind, ..., m_k are of the k^{th} kind, so that $\sum_{i=1}^k m_i = n$. The number of permutations of all the objects in this case is
$$\frac{n!}{m_1! m_2! \dots m_k!}$$

COMBINATIONS AND PERMUTATIONS

An r -combination of n elements: is a selection of r of the objects where the order of the objects in the selection is immaterial.

An r - permutation of n elements is an ordered selection of r of the objects.

- **1. Permutation with no repetition:** The number of r -permutations of n objects, $P(n, r) = {}^n P_r$

- **2. Permutation with unlimited repetition:** The no of r - permutations with unlimited repetition n^r .

- **3. Permutation with restricted repetition:** If there n objects of which m_1 are of the first kind, m_2 are of the second kind, ..., m_k are of the k^{th} kind, so that $\sum_{i=1}^k m_i = n$. The number of permutations of all the objects in this case is
$$\frac{n!}{m_1! m_2! \dots m_k!}$$

- **4. Combinations without repetition:** The number of r -combinations of n objects without repetition is $C(n, r) = {}^n C_r = \frac{n!}{r!(n-r)!}$

COMBINATIONS AND PERMUTATIONS

An r -combination of n elements: is a selection of r of the objects where the order of the objects in the selection is immaterial.

An r - permutation of n elements is an ordered selection of r of the objects.

- **1. Permutation with no repetition:** The number of r -permutations of n objects, $P(n, r) = {}^n P_r$

- **2. Permutation with unlimited repetition:** The no of r - permutations with unlimited repetition n^r .

- **3. Permutation with restricted repetition:** If there n objects of which m_1 are of the first kind, m_2 are of the second kind, ..., m_k are of the k^{th} kind, so that $\sum_{i=1}^k m_i = n$. The number of permutations of all the objects in this case is
$$\frac{n!}{m_1! m_2! \dots m_k!}$$

- **4. Combinations without repetition:** The number of r -combinations of n objects without repetition is $C(n, r) = {}^n C_r = \frac{n!}{r!(n-r)!}$

- **5. Combinations with repetition:** The number of r -combinations of n objects with repetition is $C(n+r-1, r) = {}^{n+r-1} C_r$

Distribution

Distributing r different objects to n distinct cells:

Distribution

Distributing r different objects to n distinct cells:

- 1. Such that each cell has at most one object: nP_r

Distribution

Distributing r different objects to n distinct cells:

- 1. Such that each cell has at most one object: nP_r
- 2. If we allow each cell to hold any number of objects: n^r

Distribution

Distributing r different objects to n distinct cells:

- 1. Such that each cell has at most one object: nP_r
- 2. If we allow each cell to hold any number of objects: n^r
- 3. When the r objects to be distributed are not all different i.e.

m_1 are of first kind

m_2 of the second kind,

\vdots

m_k of them of the k^{th} kind s.t $r = \sum_{i=1}^k m_i$.

Suppose that each of the n distinct cells may hold at most one object

($n \geq r$) is

Distribution

Distributing r different objects to n distinct cells:

- 1. Such that each cell has at most one object: nP_r
- 2. If we allow each cell to hold any number of objects: n^r
- 3. When the r objects to be distributed are not all different i.e.

m_1 are of first kind

m_2 of the second kind,

\vdots

m_k of them of the k^{th} kind s.t $r = \sum_{i=1}^k m_i$.

Suppose that each of the n distinct cells may hold at most one object

($n \geq r$) is $\frac{n!}{m_1!m_2!\dots m_k!}$

Distribution

Distributing r different objects to n distinct cells:

- 1. Such that each cell has at most one object: nP_r
- 2. If we allow each cell to hold any number of objects: n^r
- 3. When the r objects to be distributed are not all different i.e.

m_1 are of first kind

m_2 of the second kind,

\vdots

m_k of them of the k^{th} kind s.t $r = \sum_{i=1}^k m_i$.

Suppose that each of the n distinct cells may hold at most one object

($n \geq r$) is $\frac{n!}{m_1!m_2!\dots m_k!}$

Distributing r identical objects to n distinct cells:

Distribution

Distributing r different objects to n distinct cells:

- 1. Such that each cell has at most one object: nP_r
- 2. If we allow each cell to hold any number of objects: n^r
- 3. When the r objects to be distributed are not all different i.e.

m_1 are of first kind

m_2 of the second kind,

\vdots

m_k of them of the k^{th} kind s.t $r = \sum_{i=1}^k m_i$.

Suppose that each of the n distinct cells may hold at most one object

($n \geq r$) is $\frac{n!}{m_1!m_2!\dots m_k!}$

Distributing r identical objects to n distinct cells:

- 4. Such that each cell has at most one object: nC_r

Distribution

Distributing r different objects to n distinct cells:

- 1. Such that each cell has at most one object: nP_r
- 2. If we allow each cell to hold any number of objects: n^r
- 3. When the r objects to be distributed are not all different i.e.

m_1 are of first kind

m_2 of the second kind,

\vdots

m_k of them of the k^{th} kind s.t $r = \sum_{i=1}^k m_i$.

Suppose that each of the n distinct cells may hold at most one object

($n \geq r$) is $\frac{n!}{m_1!m_2!\dots m_k!}$

Distributing r identical objects to n distinct cells:

- 4. Such that each cell has at most one object: nC_r
- 5. If we allow each cell to hold any number of objects: ${}^{n+r-1}C_r$

Problems

Q1. Find the no of ways in which 3 exams can be scheduled in a 5 day period s.t

- (i) No two exams are scheduled on the same day?
- (ii) There are no restrictions on number of exams conducted on a day?

Problems

Q1. Find the no of ways in which 3 exams can be scheduled in a 5 day period s.t

- (i) No two exams are scheduled on the same day?
- (ii) There are no restrictions on number of exams conducted on a day?

Q2. Find the no of permutations of the word INSTITUTION?

- (i) How many of them begin with I and end with N?
- (ii) How many permutations are with 3 Ts not together? How many of these begin with I?

Problems

Q1. Find the no of ways in which 3 exams can be scheduled in a 5 day period s.t

- (i) No two exams are scheduled on the same day?
- (ii) There are no restrictions on number of exams conducted on a day?

Q2. Find the no of permutations of the word INSTITUTION?

- (i) How many of them begin with I and end with N?
- (ii) How many permutations are with 3 *T*s not together? How many of these begin with *I*?

Q3. In how many ways 3 integers can be selected from $3n$ consecutive integers such that the sum is a multiple of 3?


Q4. If 5 men A,B,C,D,E intend to speak at a meeting, (i) in how many orders can they do so without B speaking before A?
(ii) how many orders are there in which A speaks immediately before B?

Q4. If 5 men A,B,C,D,E intend to speak at a meeting, (i) in how many orders can they do so without B speaking before A?


(ii) how many orders are there in which A speaks immediately before B?

(i) Soution:

Case 1: A talks First



Case 2: A Talks second




Q4. If 5 men A,B,C,D,E intend to speak at a meeting, (i) in how many orders can they do so without B speaking before A?


(ii) how many orders are there in which A speaks immediately before B?

(i) Soution:


Case 1: A talks First




Case 2: A Talks second



Case 3: A talks third



Case 4: A talks fourth




Q4. If 5 men A,B,C,D,E intend to speak at a meeting, (i) in how many orders can they do so without B speaking before A?


(ii) how many orders are there in which A speaks immediately before B?

(i) Soution:


Case 1: A talks First




Case 2: A Talks second



Case 3: A talks third



Case 4: A talks fourth




Answer is $4! + 3 \times 3! + 3 \times 2 \times 2! + 3!$

Q5. A new national flag has to be designed with 6 vertical strips in yellow, green, blue and red. In how many ways can this be done s.t no two adjacent strips have same color?

Q5. A new national flag has to be designed with 6 vertical strips in yellow, green, blue and red. In how many ways can this be done s.t no two adjacent strips have same color?


Q5. A new national flag has to be designed with 6 vertical strips in yellow, green, blue and red. In how many ways can this be done s.t no two adjacent strips have same color?



The first strip can be colored using any of the four colors.

Second strip can be colored using the remaining 3 colors leaving the one which is used to color first strip

Q5. A new national flag has to be designed with 6 vertical strips in yellow, green, blue and red. In how many ways can this be done s.t no two adjacent strips have same color?




The first strip can be colored using any of the four colors.

Second strip can be colored using the remaining 3 colors leaving the one which is used to color first strip


Answer is 4×3^5 .

Q6. In how many ways can 2 squares be selected one by one from 8×8 chess board such that they are not in the same row and same columns?

Q6. In how many ways can 2 squares be selected one by one from 8×8 chess board such that they are not in the same row and same columns?




Q6. In how many ways can 2 squares be selected one by one from 8×8 chess board such that they are not in the same row and same columns?



For the first square we have 64 choices.


Q6. In how many ways can 2 squares be selected one by one from 8×8 chess board such that they are not in the same row and same columns?



For the first square we have 64 choices.

And for the second square we have 7×7 Choices.

Q6. In how many ways can 2 squares be selected one by one from 8×8 chess board such that they are not in the same row and same columns?



For the first square we have 64 choices.

And for the second square we have 7×7 Choices.

Therefore Ans is 64×49 .

Q7. Find the sum of all 4 digit numbers that can be obtained using the digits 1,2,3,4 once in each?

Q7. Find the sum of all 4 digit numbers that can be obtained using the digits 1,2,3,4 once in each?

Q7. Find the sum of all 4 digit numbers that can be obtained using the digits 1,2,3,4 once in each?

Solution: Total number of 4 digit numbers using the digits 1, 2, 3, 4 are

Q7. Find the sum of all 4 digit numbers that can be obtained using the digits 1,2,3,4 once in each?

Solution: Total number of 4 digit numbers using the digits 1, 2, 3, 4 are $4!$

Q7. Find the sum of all 4 digit numbers that can be obtained using the digits 1,2,3,4 once in each?

Solution: Total number of 4 digit numbers using the digits 1, 2, 3, 4 are $4!$

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1
-----	-----	-----	-----

Q7. Find the sum of all 4 digit numbers that can be obtained using the digits 1,2,3,4 once in each?

Solution: Total number of 4 digit numbers using the digits 1, 2, 3, 4 are $4!$

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1
-----	-----	-----	-----

Each digit occupies each place 6 times.

Q7. Find the sum of all 4 digit numbers that can be obtained using the digits 1,2,3,4 once in each?

Solution: Total number of 4 digit numbers using the digits 1, 2, 3, 4 are $4!$

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1
-----	-----	-----	-----

Each digit occupies each place 6 times.

The answer is

Q7. Find the sum of all 4 digit numbers that can be obtained using the digits 1,2,3,4 once in each?

Solution: Total number of 4 digit numbers using the digits 1, 2, 3, 4 are 4!

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1
-----	-----	-----	-----

Each digit occupies each place 6 times.

The answer is

$$6 \times \{(1+2+3+4)1000 + (1+2+3+4)100 + (1+2+3+4)10 + (1+2+3+4)\} \\ = 66660.$$

Q7. Find the sum of all 4 digit numbers that can be obtained using the digits 1,2,3,4 once in each?

Solution: Total number of 4 digit numbers using the digits 1, 2, 3, 4 are $4!$

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1
-----	-----	-----	-----

Each digit occupies each place 6 times.

The answer is

$$6 \times \{(1+2+3+4)1000 + (1+2+3+4)100 + (1+2+3+4)10 + (1+2+3+4)\} \\ = 66660.$$

What is the sum if repetition of the 4 digits allowed?

Q8. How many positive integers less than **one million** can be formed using (i) 7's, 8's, 9's only?

(ii) Using 0's, 8's, 9's only?

(i) Solution:

Q8. How many positive integers less than **one million** can be formed using (i) 7's, 8's, 9's only?

(ii) Using 0's, 8's, 9's only?

(i) Solution:

$< 10,00,000$

Q8. How many positive integers less than **one million** can be formed using (i) 7's, 8's, 9's only?

(ii) Using 0's, 8's, 9's only?


(i) Solution:

$< 10,00,000$

6 digit numbers



5 digit numbers



1 digit number



Q8. How many positive integers less than **one million** can be formed using (i) 7's, 8's, 9's only?

(ii) Using 0's, 8's, 9's only?


(i) Solution:

$< 10,00,000$

6 digit numbers



5 digit numbers



1 digit number



Answer is $3^6 + 3^5 + 3^4 + 3^3 + 3^2 + 3 = 1092$.

Q9. 6 distinct symbols are transmitted through a communication channel. A total of 12 blanks are to be inserted between the symbols with at least two blank spaces between every pair of symbols. In how many ways can we arrange symbols and blanks?

Solution:

Q9. 6 distinct symbols are transmitted through a communication channel. A total of 12 blanks are to be inserted between the symbols with at least two blank spaces between every pair of symbols. In how many ways can we arrange symbols and blanks?

Solution:

1 2 3 4 5 6
1 _ _ 2 _ _ 3 _ _ 4 _ _ 5 _ _ 6

Q9. 6 distinct symbols are transmitted through a communication channel. A total of 12 blanks are to be inserted between the symbols with at least two blank spaces between every pair of symbols. In how many ways can we arrange symbols and blanks?

Solution:

1 2 3 4 5 6
1 _ _ 2 _ _ 3 _ _ 4 _ _ 5 _ _ 6

Now we have 2 identical blanks left which is to be inserted in the 5 gaps between the 6 distinct symbols with allowing repetition.

Q9. 6 distinct symbols are transmitted through a communication channel. A total of 12 blanks are to be inserted between the symbols with at least two blank spaces between every pair of symbols. In how many ways can we arrange symbols and blanks?

Solution:

1 2 3 4 5 6
1 _ _ 2 _ _ 3 _ _ 4 _ _ 5 _ _ 6

Now we have 2 identical blanks left which is to be inserted in the 5 gaps between the 6 distinct symbols with allowing repetition.

Which is same as selecting 2 out of 5 gaps with repetition allowed

Q9. 6 distinct symbols are transmitted through a communication channel. A total of 12 blanks are to be inserted between the symbols with at least two blank spaces between every pair of symbols. In how many ways can we arrange symbols and blanks?

Solution:

1 2 3 4 5 6
 1 _ _ 2 _ _ 3 _ _ 4 _ _ 5 _ _ 6

Now we have 2 identical blanks left which is to be inserted in the 5 gaps between the 6 distinct symbols with allowing repetition.

Which is same as selecting 2 out of 5 gaps with repetition allowed

Distributing 2 identical blanks (objects) to 5 distinct gaps (cells), such that each cell can hold any number of objects (with allowing repetition of gaps). Therefore Answer is ${}^{5+2-1}C_2 = 15$.

Q10. Three identical dice are rolled. How many outcomes can be recorded?

Solution:

Q10. Three identical dice are rolled. How many outcomes can be recorded?

Solution:

Out of 6 outcomes choose 3 numbers with allowing repetition.

Q10. Three identical dice are rolled. How many outcomes can be recorded?

Solution:

Out of 6 outcomes choose 3 numbers with allowing repetition.

Q10. Three identical dice are rolled. How many outcomes can be recorded?

Solution:

Out of 6 outcomes choose 3 numbers with allowing repetition.

Distribution of 3 identical dice (objects) to 6 distinct outcomes (cells), such that each cell can hold any number of objects (with allowing repetition of outcomes).

Q10. Three identical dice are rolled. How many outcomes can be recorded?

Solution:

Out of 6 outcomes choose 3 numbers with allowing repetition.

Distribution of 3 identical dice (objects) to 6 distinct outcomes (cells), such that each cell can hold any number of objects (with allowing repetition of outcomes).

Answer is ${}^{6+3-1}C_3 = {}^8C_3 = 56$.

Q10. Three identical dice are rolled. How many outcomes can be recorded?

Solution:

Out of 6 outcomes choose 3 numbers with allowing repetition.

Distribution of 3 identical dice (objects) to 6 distinct outcomes (cells), such that each cell can hold any number of objects (with allowing repetition of outcomes).

Answer is ${}^{6+3-1}C_3 = {}^8C_3 = 56$.

111 222 333 444 555 666

Q10. Three identical dice are rolled. How many outcomes can be recorded?

Solution:

Out of 6 outcomes choose 3 numbers with allowing repetition.

Distribution of 3 identical dice (objects) to 6 distinct outcomes (cells), such that each cell can hold any number of objects (with allowing repetition of outcomes).

Answer is ${}^{6+3-1}C_3 = {}^8C_3 = 56$.

111 222 333 444 555 666

112 113 114 115 116

Q10. Three identical dice are rolled. How many outcomes can be recorded?

Solution:

Out of 6 outcomes choose 3 numbers with allowing repetition.

Distribution of 3 identical dice (objects) to 6 distinct outcomes (cells), such that each cell can hold any number of objects (with allowing repetition of outcomes).

Answer is ${}^{6+3-1}C_3 = {}^8C_3 = 56$.

111 222 333 444 555 666

112 113 114 115 116

221 223 224 225 226

Q10. Three identical dice are rolled. How many outcomes can be recorded?

Solution:

Out of 6 outcomes choose 3 numbers with allowing repetition.

Distribution of 3 identical dice (objects) to 6 distinct outcomes (cells), such that each cell can hold any number of objects (with allowing repetition of outcomes).

Answer is ${}^{6+3-1}C_3 = {}^8C_3 = 56$.

111 222 333 444 555 666

112 113 114 115 116

221 223 224 225 226

...

⋮

661 662 663 664 665

Q10. Three identical dice are rolled. How many outcomes can be recorded?

Solution:

Out of 6 outcomes choose 3 numbers with allowing repetition.

Distribution of 3 identical dice (objects) to 6 distinct outcomes (cells), such that each cell can hold any number of objects (with allowing repetition of outcomes).

Answer is ${}^{6+3-1}C_3 = {}^8C_3 = 56$.

111 222 333 444 555 666

112 113 114 115 116

221 223 224 225 226

...

⋮

661 662 663 664 665

123 124 125 126 134 135 136 145 146 156

234 235 236 245 246 256 345 346 356 456.

Q11. In how many ways can an examiner assign 30 Marks to 8 questions such that no question receives less than 2 marks?

Solution: Initially we give 2 marks each to all the 8 questions.

Q11. In how many ways can an examiner assign 30 Marks to 8 questions such that no question receives less than 2 marks?

Solution: Initially we give 2 marks each to all the 8 questions.

Now the remaining 14 Marks to be given to 8 questions with allowing repetition.

Q11. In how many ways can an examiner assign 30 Marks to 8 questions such that no question receives less than 2 marks?

Solution: Initially we give 2 marks each to all the 8 questions.

Now the remaining 14 Marks to be given to 8 questions with allowing repetition.


Each mark can be given to any of the 8 questions allowing repetition of questions.

Q11. In how many ways can an examiner assign 30 Marks to 8 questions such that no question receives less than 2 marks?

Solution: Initially we give 2 marks each to all the 8 questions.

Now the remaining 14 Marks to be given to 8 questions with allowing repetition.

Each mark can be given to any of the 8 questions allowing repetition of questions.




Q11. In how many ways can an examiner assign 30 Marks to 8 questions such that no question receives less than 2 marks?

Solution: Initially we give 2 marks each to all the 8 questions.

Now the remaining 14 Marks to be given to 8 questions with allowing repetition.

Each mark can be given to any of the 8 questions allowing repetition of questions.




Distribution of 14 identical marks(objects) to 8 distinct questions (cells) such that each cell can hold any number of objects (with allowing repetition of questions).

Q11. In how many ways can an examiner assign 30 Marks to 8 questions such that no question receives less than 2 marks?

Solution: Initially we give 2 marks each to all the 8 questions.

Now the remaining 14 Marks to be given to 8 questions with allowing repetition.

Each mark can be given to any of the 8 questions allowing repetition of questions.




Distribution of 14 identical marks(objects) to 8 distinct questions (cells) such that each cell can hold any number of objects (with allowing repetition of questions).


Answer is ${}^{8+14-1}C_{14} = {}^{21}C_{14}$

Q12. In how many ways can two adjacent squares can be selected from an 8×8 chess board?

Q12. In how many ways can two adjacent squares can be selected from an 8×8 chess board?



Q12. In how many ways can two adjacent squares can be selected from an 8×8 chess board?



Answer is $4 \times 2 + 24 \times 3 + 36 \times 4$

Q13. Among all 7 digits numbers, how many of them contain exactly three 9s?

Q13. Among all 7 digits numbers, how many of them contain exactly three 9s?

Solution:



Diagram illustrating the first case: numbers starting with 9. A horizontal line represents the 7-digit number. The first digit is labeled '9' above it. Below the line, there are seven upward-pointing arrows. The first arrow is positioned under the '9'. Below the first arrow is the text "Begin with 9". Below the remaining six arrows is the expression $(6C2) * 9 * 9 * 9 * 9$.




Diagram illustrating the second case: numbers starting with 1, 2, ..., 8. A horizontal line represents the 7-digit number. Below the line, there are seven upward-pointing arrows. Below the first arrow is the text "Begin with 1, 2, ..., 8". Below the remaining six arrows is the expression $8 * (6C3) * 9 * 9 * 9$.


Q13. Among all 7 digits numbers, how many of them contain exactly three 9s?

Solution:



Begin with 9

$$(6C2) * 9 * 9 * 9 * 9$$



Begin with 1,2,...,8

$$8 * (6C3) * 9 * 9 * 9$$

Answer is ${}^6C_2 \times 9^4 + 8 \times {}^6C_3 \times 9^3$

Q13. Among all 7 digits numbers, how many of them contain exactly three 9s?

Solution:

9

↑ ↑ ↑ ↑ ↑ ↑ ↑

Begin with 9

$(6C2) \cdot 9 \cdot 9 \cdot 9 \cdot 9$

↑ ↑ ↑ ↑ ↑ ↑ ↑

Begin with 1,2,...,8


$8 \cdot (6C3) \cdot 9 \cdot 9 \cdot 9$

Answer is ${}^6C_2 \times 9^4 + 8 \times {}^6C_3 \times 9^3$

Q14. The number of ways to choose 3 days out of 7 days (with repetition) is _____

Q13. Among all 7 digits numbers, how many of them contain exactly three 9s?

Solution:



Answer is ${}^6C_2 \times 9^4 + 8 \times {}^6C_3 \times 9^3$

Q14. The number of ways to choose 3 days out of 7 days (with repetition) is _____

$$7+3-1C_3 = {}^9C_3.$$


Q15. How many points of intersection are formed by n lines drawn in a plane if no two are parallel and no three concurrent? Into how many regions is the plane divided?

Q15. How many points of intersection are formed by n lines drawn in a plane if no two are parallel and no three concurrent? Into how many regions is the plane divided?

Solution:

Q15. How many points of intersection are formed by n lines drawn in a plane if no two are parallel and no three concurrent? Into how many regions is the plane divided?

Solution:




Two lines 1 intersecting point

3 lines $(1+2)$ intersecting points

4 lines $(1+2+3)$ intersecting points

Q15. How many points of intersection are formed by n lines drawn in a plane if no two are parallel and no three concurrent? Into how many regions is the plane divided?

Solution:







Two lines 1 intersecting point

3 lines $(1+2)$ intersecting points

4 lines $(1+2+3)$ intersecting points





So n lines $1+2+\dots+(n-1)$ intersecting points.

Number of intersecting points is nC_2 .



1 line 2 regions
 2 line 4 regions
 3 line 7 regions
 4 line 11 regions

=intersecting points of 2 lines +1
 =intersecting points of 3 lines +1
 =intersecting points of 4 lines +1
 =intersecting points of 5 lines +1



1 line 2 regions
2 line 4 regions
3 line 7 regions
4 line 11 regions

=intersecting points of 2 lines +1
=intersecting points of 3 lines +1
=intersecting points of 4 lines +1
=intersecting points of 5 lines +1

So, number of regions created by n lines is number of intersecting points created by $n+1$ lines + 1 = ${}^{n+1}C_2 + 1 = \frac{(n+1)n}{2} + 1$ regions.