

Derangements

Derangements

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Derangements

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Q2. Show that the proportion of the Permutation of $\{1, 2, \dots, n\}$ which contains no consecutive pair $(i, i + 1)$ for any i is approximately $\frac{n + 1}{ne}$ as n becomes large.

PARTITIONS AND COMPOSITIONS

PARTITIONS AND COMPOSITIONS

Any positive integer n can be represented as a sum of one or more positive integers (a_i) ,

PARTITIONS AND COMPOSITIONS

Any positive integer n can be represented as a sum of one or more positive integers (a_i) ,

i.e., $n = a_1 + a_2 + \dots + a_m$

PARTITIONS AND COMPOSITIONS

Any positive integer n can be represented as a sum of one or more positive integers (a_i) ,

$$\text{i.e., } n = a_1 + a_2 + \dots + a_m$$

Divisions of a positive integer n are of two types depending on whether the ordering of the parts a_1, a_2, \dots, a_m is regarded as important or not.

PARTITIONS AND COMPOSITIONS

Any positive integer n can be represented as a sum of one or more positive integers (a_i) ,

$$\text{i.e., } n = a_1 + a_2 + \dots + a_m$$

Divisions of a positive integer n are of two types depending on whether the ordering of the parts a_1, a_2, \dots, a_m is regarded as important or not.

Ordered divisions of a positive integer n is called **compositions** while unordered divisions is called as **partitions**.

Consider, for example, the partitions and compositions of the integer 5.

Consider, for example, the partitions and compositions of the integer 5.

There are *seven* **unrestricted partitions**, namely

Consider, for example, the partitions and compositions of the integer 5.

There are *seven* **unrestricted partitions**, namely
5,

Consider, for example, the partitions and compositions of the integer 5.

There are *seven* **unrestricted partitions**, namely
5, 4+1,

Consider, for example, the partitions and compositions of the integer 5.

There are *seven* **unrestricted partitions**, namely
5, 4+1, 3+2,

Consider, for example, the partitions and compositions of the integer 5.

There are *seven* **unrestricted partitions**, namely

5, 4+1, 3+2, 3+1+1, 2+2+1, 2+1+1+1, 1+1+1+1+1.

Consider, for example, the partitions and compositions of the integer 5.

There are *seven* **unrestricted partitions**, namely

5, 4+1, 3+2, 3+1+1, 2+2+1, 2+1+1+1, 1+1+1+1+1.

And two of these, namely 4+1 and 3+2 , have exactly two parts.

Consider, for example, the partitions and compositions of the integer 5.

There are *seven* **unrestricted partitions**, namely

5, 4+1, 3+2, 3+1+1, 2+2+1, 2+1+1+1, 1+1+1+1+1.

And two of these, namely 4+1 and 3+2, have exactly two parts.

There are *sixteen* **unrestricted compositions** of $n=5$.

Consider, for example, the partitions and compositions of the integer 5.

There are *seven* **unrestricted partitions**, namely

5, 4+1, 3+2, 3+1+1, 2+2+1, 2+1+1+1, 1+1+1+1+1.

And two of these, namely 4+1 and 3+2, have exactly two parts.

There are *sixteen* **unrestricted compositions** of $n=5$. They are,

Consider, for example, the partitions and compositions of the integer 5.

There are *seven* **unrestricted partitions**, namely

5, 4+1, 3+2, 3+1+1, 2+2+1, 2+1+1+1, 1+1+1+1+1.

And two of these, namely 4+1 and 3+2, have exactly two parts.

There are *sixteen* **unrestricted compositions** of $n=5$. They are,

5, 4+1, 1+4, 3+2, 2+3, 3+1+1, 1+3+1, 1+1+3, 2+2+1, 2+1+2, 1+2+2, 2+1+1+1, 1+2+1+1, 1+1+2+1, 1+1+1+2, 1+1+1+1+1 and four of these have exactly two parts.

Consider, for example, the partitions and compositions of the integer 5.

There are *seven* **unrestricted partitions**, namely

5, 4+1, 3+2, 3+1+1, 2+2+1, 2+1+1+1, 1+1+1+1+1.

And two of these, namely 4+1 and 3+2, have exactly two parts.

There are *sixteen* **unrestricted compositions** of $n=5$. They are,

5, 4+1, 1+4, 3+2, 2+3, 3+1+1, 1+3+1, 1+1+3, 2+2+1, 2+1+2, 1+2+2, 2+1+1+1, 1+2+1+1, 1+1+2+1, 1+1+1+2, 1+1+1+1+1 and four of these have exactly two parts.

When we write partitions or compositions we will omit the + signs, thus 2+1+1+1 will be written 2111, or 21^3 , and for partitions the largest parts will be written first.

Enumerators for Compositions

Enumerators for Compositions

Consider a positive integer n and n ones in a row.

Enumerators for Compositions

Consider a positive integer n and n ones in a row.
There are $(n - 1)$ spaces between the n ones.

Enumerators for Compositions

Consider a positive integer n and n ones in a row.

There are $(n - 1)$ spaces between the n ones.


If there is **no restriction** on the number of parts, we may or may not put a marker in any of the $(n - 1)$ spaces between the ones in order to form groups.

Enumerators for Compositions

Consider a positive integer n and n ones in a row.

There are $(n - 1)$ spaces between the n ones.

If there is **no restriction** on the number of parts, we may or may not put a marker in any of the $(n - 1)$ spaces between the ones in order to form groups.




Enumerators for Compositions

Consider a positive integer n and n ones in a row.

There are $(n - 1)$ spaces between the n ones.

If there is **no restriction** on the number of parts, we may or may not put a marker in any of the $(n - 1)$ spaces between the ones in order to form groups.




This can be done in 2^{n-1} ways.

Enumerators for Compositions

Consider a positive integer n and n ones in a row.

There are $(n - 1)$ spaces between the n ones.

If there is **no restriction** on the number of parts, we may or may not put a marker in any of the $(n - 1)$ spaces between the ones in order to form groups.



This can be done in 2^{n-1} ways.

The number of compositions of n with no restriction on the number of parts is 2^{n-1} .

For $n = 5$ we have $2^4 = 16$ unrestricted compositions.

For $n = 5$ we have $2^4 = 16$ unrestricted compositions.

If we restrict the compositions to have **exactly m parts**,

For $n = 5$ we have $2^4 = 16$ unrestricted compositions.

If we restrict the compositions to have **exactly m parts**, just $(m - 1)$ markers are needed to form m groups and the number of ways of placing $(m - 1)$ markers in the $(n - 1)$ spaces between the ones is ${}^{n-1}C_{m-1}$.

For $n = 5$ we have $2^4 = 16$ unrestricted compositions.

If we restrict the compositions to have **exactly m parts**, just $(m - 1)$ markers are needed to form m groups and the number of ways of placing $(m - 1)$ markers in the $(n - 1)$ spaces between the ones is ${}^{n-1}C_{m-1}$.

The number of compositions of n with exactly m parts is ${}^{n-1}C_{m-1}$.

For $n = 5$ we have $2^4 = 16$ unrestricted compositions.

If we restrict the compositions to have **exactly m parts**, just $(m - 1)$ markers are needed to form m groups and the number of ways of placing $(m - 1)$ markers in the $(n - 1)$ spaces between the ones is ${}^{n-1}C_{m-1}$.

The number of compositions of n with exactly m parts is ${}^{n-1}C_{m-1}$.

For $n = 5$ with $m = 2$ we have ${}^{n-1}C_{m-1} = {}^4C_1$, compositions.

For $n = 5$ we have $2^4 = 16$ unrestricted compositions.

If we restrict the compositions to have **exactly m parts**, just $(m - 1)$ markers are needed to form m groups and the number of ways of placing $(m - 1)$ markers in the $(n - 1)$ spaces between the ones is ${}^{n-1}C_{m-1}$.

The number of compositions of n with exactly m parts is ${}^{n-1}C_{m-1}$.

For $n = 5$ with $m = 2$ we have ${}^{n-1}C_{m-1} = {}^4C_1$, compositions.
They are 14, 41, 23, 32.

For $n = 5$ we have $2^4 = 16$ unrestricted compositions.

If we restrict the compositions to have **exactly m parts**, just $(m - 1)$ markers are needed to form m groups and the number of ways of placing $(m - 1)$ markers in the $(n - 1)$ spaces between the ones is ${}^{n-1}C_{m-1}$.

The number of compositions of n with exactly m parts is ${}^{n-1}C_{m-1}$.

For $n = 5$ with $m = 2$ we have ${}^{n-1}C_{m-1} = {}^4C_1$, compositions. They are 14, 41, 23, 32.

For $n = 6$ and $m = 3$ we have ${}^5C_2 = 10$ compositions. They are

For $n = 5$ we have $2^4 = 16$ unrestricted compositions.

If we restrict the compositions to have **exactly m parts**, just $(m - 1)$ markers are needed to form m groups and the number of ways of placing $(m - 1)$ markers in the $(n - 1)$ spaces between the ones is ${}^{n-1}C_{m-1}$.

The number of compositions of n with exactly m parts is ${}^{n-1}C_{m-1}$.

For $n = 5$ with $m = 2$ we have ${}^{n-1}C_{m-1} = {}^4C_1$, compositions. They are 14, 41, 23, 32.

For $n = 6$ and $m = 3$ we have ${}^5C_2 = 10$ compositions. They are 123, 132, 213, 231, 312, 321, 222, 411, 114, 141.

Q1. How many compositions of n with m parts are there when zero as a part are allowed.

Q1. How many compositions of n with m parts are there when zero as a part are allowed.

Ans: Consider a compositions of n with m parts when zero as a part are allowed.

Q1. How many compositions of n with m parts are there when zero as a part are allowed.

Ans: Consider a compositions of n with m parts when zero as a part are allowed.

Add one to each part,

Q1. How many compositions of n with m parts are there when zero as a part are allowed.

Ans: Consider a compositions of n with m parts when zero as a part are allowed.

Add one to each part, then it will represent a composition of $(n + m)$ with exactly m parts.

Q1. How many compositions of n with m parts are there when zero as a part are allowed.

Ans: Consider a compositions of n with m parts when zero as a part are allowed.

Add one to each part, then it will represent a composition of $(n + m)$ with exactly m parts.

Therefore, number of compositions $= {}^{n+m-1}C_{m-1}$.

Q1. How many compositions of n with m parts are there when zero as a part are allowed.

Ans: Consider a compositions of n with m parts when zero as a part are allowed.

Add one to each part, then it will represent a composition of $(n + m)$ with exactly m parts.

Therefore, number of compositions = $^{n+m-1}C_{m-1}$.

Example: $n=5, m=2$.

Q1. How many compositions of n with m parts are there when zero as a part are allowed.

Ans: Consider a compositions of n with m parts when zero as a part are allowed.

Add one to each part, then it will represent a composition of $(n + m)$ with exactly m parts.

Therefore, number of compositions = ${}^{n+m-1}C_{m-1}$.

Example: $n=5, m=2$.

Then $(1, 4), (4, 1), (2, 3), (3, 2), (0, 5), (5, 0)$ are compositions of 5 with 2 parts when zero parts allowed.

Q1. How many compositions of n with m parts are there when zero as a part are allowed.

Ans: Consider a compositions of n with m parts when zero as a part are allowed.

Add one to each part, then it will represent a composition of $(n + m)$ with exactly m parts.

Therefore, number of compositions = $^{n+m-1}C_{m-1}$.

Example: $n=5, m=2$.

Then $(1, 4), (4, 1), (2, 3), (3, 2), (0, 5), (5, 0)$ are compositions of 5 with 2 parts when zero parts allowed.

Add one to each part, then we get $(2, 5), (5, 2), (3, 4), (4, 3), (1, 6), (6, 1)$. This is a composition of $n + m = 7$ with exactly $m = 2$ parts.

Q1. How many compositions of n with m parts are there when zero as a part are allowed.

Ans: Consider a compositions of n with m parts when zero as a part are allowed.

Add one to each part, then it will represent a composition of $(n + m)$ with exactly m parts.

Therefore, number of compositions = $^{n+m-1}C_{m-1}$.

Example: $n=5, m=2$.

Then $(1, 4), (4, 1), (2, 3), (3, 2), (0, 5), (5, 0)$ are compositions of 5 with 2 parts when zero parts allowed.

Add one to each part, then we get $(2, 5), (5, 2), (3, 4), (4, 3), (1, 6), (6, 1)$. This is a composition of $n + m = 7$ with exactly $m = 2$ parts.

Such compositions are equal to $^{n+m-1}C_{m-1} = {}^6C_1 = 6$.

Q2. In how many ways can an examiner assign 30 Marks to 8 questions such that no question receives less than 2 marks?

Q2. In how many ways can an examiner assign 30 Marks to 8 questions such that no question receives less than 2 marks?

Solution: Assign 2 marks to each of 8 questions. Remaining 14 marks to 8 questions allowing 0 mark is same as

Q2. In how many ways can an examiner assign 30 Marks to 8 questions such that no question receives less than 2 marks?

Solution: Assign 2 marks to each of 8 questions. Remaining 14 marks to 8 questions allowing 0 mark is same as

"composition of 14 (n) with 8 (m) parts allowing zero part" is

$${}^{n+m-1}C_{m-1} = {}^{14+8-1}C_{8-1} = {}^{21}C_7.$$

Q2. In how many ways can an examiner assign 30 Marks to 8 questions such that no question receives less than 2 marks?

Solution: Assign 2 marks to each of 8 questions. Remaining 14 marks to 8 questions allowing 0 mark is same as

"composition of 14 (n) with 8 (m) parts allowing zero part" is

$${}^{n+m-1}C_{m-1} = {}^{14+8-1}C_{8-1} = {}^{21}C_7.$$

OR

Assign 1 mark to 8 questions.

Q2. In how many ways can an examiner assign 30 Marks to 8 questions such that no question receives less than 2 marks?

Solution: Assign 2 marks to each of 8 questions. Remaining 14 marks to 8 questions allowing 0 mark is same as

"composition of 14 (n) with 8 (m) parts allowing zero part" is

$${}^{n+m-1}C_{m-1} = {}^{14+8-1}C_{8-1} = {}^{21}C_7.$$

OR

Assign 1 mark to 8 questions. Remaining 22 marks to 8 questions such that each question gets at least 1 marks is same as

Q2. In how many ways can an examiner assign 30 Marks to 8 questions such that no question receives less than 2 marks?

Solution: Assign 2 marks to each of 8 questions. Remaining 14 marks to 8 questions allowing 0 mark is same as

"composition of 14 (n) with 8 (m) parts allowing zero part" is

$${}^{n+m-1}C_{m-1} = {}^{14+8-1}C_{8-1} = {}^{21}C_7.$$

OR

Assign 1 mark to 8 questions. Remaining 22 marks to 8 questions such that each question gets at least 1 marks is same as

"composition of 22 (n) with 8 (m) parts with no zero parts" is

Q2. In how many ways can an examiner assign 30 Marks to 8 questions such that no question receives less than 2 marks?

Solution: Assign 2 marks to each of 8 questions. Remaining 14 marks to 8 questions allowing 0 mark is same as

"composition of 14 (n) with 8 (m) parts allowing zero part" is

$${}^{n+m-1}C_{m-1} = {}^{14+8-1}C_{8-1} = {}^{21}C_7.$$

OR

Assign 1 mark to 8 questions. Remaining 22 marks to 8 questions such that each question gets at least 1 marks is same as

"composition of 22 (n) with 8 (m) parts with no zero parts" is

$${}^{n-1}C_{m-1} = {}^{22-1}C_{8-1} = {}^{21}C_7.$$

Enumerator for compositions Using Generating function

Enumerator for compositions Using Generating function

Let $C_m(x)$ be the enumerator for compositions of n with exactly m parts, where $C_m(x) = \sum_n C_{mn} x^n$ and C_{mn} , the coefficient of x^n in this series, is the number of compositions of n into exactly m parts.

Enumerator for compositions Using Generating function

Let $C_m(x)$ be the enumerator for compositions of n with exactly m parts, where $C_m(x) = \sum_n C_{mn}x^n$ and C_{mn} , the coefficient of x^n in this series, is the number of compositions of n into exactly m parts. Each part of any composition can be one, two, three or any greater number.

Enumerator for compositions Using Generating function

Let $C_m(x)$ be the enumerator for compositions of n with exactly m parts, where $C_m(x) = \sum_n C_{mn} x^n$ and C_{mn} , the coefficient of x^n in this series, is the number of compositions of n into exactly m parts.

Each part of any composition can be one, two, three or any greater number.

The factor in the enumerator must contain each of these powers of x .

Enumerator for compositions Using Generating function

Let $C_m(x)$ be the enumerator for **compositions of n with exactly m parts**, where $C_m(x) = \sum_n C_{mn}x^n$ and C_{mn} , the coefficient of x^n in this series, is the number of compositions of n into exactly m parts.

Each part of any composition can be one, two, three or any greater number.

The factor in the enumerator must contain each of these powers of x .

The factor for any part is $x + x^2 + x^3 + \dots + x^k + \dots =$

Enumerator for compositions Using Generating function

Let $C_m(x)$ be the enumerator for **compositions of n with exactly m parts**, where $C_m(x) = \sum_n C_{mn}x^n$ and C_{mn} , the coefficient of x^n in this series, is the number of compositions of n into exactly m parts.

Each part of any composition can be one, two, three or any greater number.

The factor in the enumerator must contain each of these powers of x .

The factor for any part is $x + x^2 + x^3 + \dots + x^k + \dots = x(1 - x)^{-1}$.

Enumerator for compositions Using Generating function

Let $C_m(x)$ be the enumerator for **compositions of n with exactly m parts**, where $C_m(x) = \sum_n C_{mn}x^n$ and C_{mn} , the coefficient of x^n in this series, is the number of compositions of n into exactly m parts.

Each part of any composition can be one, two, three or any greater number.

The factor in the enumerator must contain each of these powers of x .

The factor for any part is $x + x^2 + x^3 + \dots + x^k + \dots = x(1 - x)^{-1}$.

Since there are exactly m parts, the GF is the product of m such factors.

Enumerator for compositions Using Generating function

Let $C_m(x)$ be the enumerator for **compositions of n with exactly m parts**, where $C_m(x) = \sum_n C_{mn}x^n$ and C_{mn} , the coefficient of x^n in this series, is the number of compositions of n into exactly m parts.

Each part of any composition can be one, two, three or any greater number.

The factor in the enumerator must contain each of these powers of x .

The factor for any part is $x + x^2 + x^3 + \dots + x^k + \dots = x(1 - x)^{-1}$.

Since there are exactly m parts, the GF is the product of m such factors.

$$C_m(x) = (x + x^2 + \dots)^m =$$

Enumerator for compositions Using Generating function

Let $C_m(x)$ be the enumerator for **compositions of n with exactly m parts**, where $C_m(x) = \sum_n C_{mn}x^n$ and C_{mn} , the coefficient of x^n in this series, is the number of compositions of n into exactly m parts.

Each part of any composition can be one, two, three or any greater number.

The factor in the enumerator must contain each of these powers of x .

The factor for any part is $x + x^2 + x^3 + \dots + x^k + \dots = x(1 - x)^{-1}$.

Since there are exactly m parts, the GF is the product of m such factors.

$$C_m(x) = (x + x^2 + \dots)^m = x^m(1 - x)^{-m}$$

Enumerator for compositions Using Generating function

Let $C_m(x)$ be the enumerator for compositions of n with exactly m parts, where $C_m(x) = \sum_n C_{mn}x^n$ and C_{mn} , the coefficient of x^n in this series, is the number of compositions of n into exactly m parts.

Each part of any composition can be one, two, three or any greater number.

The factor in the enumerator must contain each of these powers of x .

The factor for any part is $x + x^2 + x^3 + \dots + x^k + \dots = x(1 - x)^{-1}$.

Since there are exactly m parts, the GF is the product of m such factors.

$$C_m(x) = (x + x^2 + \dots)^m = x^m(1 - x)^{-m}$$

$$C_m(x) = x^m(1 + {}^m C_1 x + {}^{m+1} C_2 x^2 + \dots).$$

Enumerator for compositions Using Generating function

Let $C_m(x)$ be the enumerator for **compositions of n with exactly m parts**, where $C_m(x) = \sum_n C_{mn}x^n$ and C_{mn} , the coefficient of x^n in this series, is the number of compositions of n into exactly m parts.

Each part of any composition can be one, two, three or any greater number.

The factor in the enumerator must contain each of these powers of x .

The factor for any part is $x + x^2 + x^3 + \dots + x^k + \dots = x(1 - x)^{-1}$.

Since there are exactly m parts, the GF is the product of m such factors.

$$C_m(x) = (x + x^2 + \dots)^m = x^m(1 - x)^{-m}$$

$$C_m(x) = x^m(1 + {}^m C_1 x + {}^{m+1} C_2 x^2 + \dots).$$

The coefficient of x^n in this enumerator $C_m(x)$ is the coefficient of x^{n-m} in $(1 + {}^m C_1 x + {}^{m+1} C_2 x^2 + \dots)$ which is equal to ${}^{n-1} C_{n-m} = {}^{n-1} C_{m-1}$.

Enumerator for compositions Using Generating function

Let $C_m(x)$ be the enumerator for **compositions of n with exactly m parts**, where $C_m(x) = \sum_n C_{mn}x^n$ and C_{mn} , the coefficient of x^n in this series, is the number of compositions of n into exactly m parts.

Each part of any composition can be one, two, three or any greater number.

The factor in the enumerator must contain each of these powers of x .

The factor for any part is $x + x^2 + x^3 + \dots + x^k + \dots = x(1 - x)^{-1}$.

Since there are exactly m parts, the GF is the product of m such factors.

$$C_m(x) = (x + x^2 + \dots)^m = x^m(1 - x)^{-m}$$

$$C_m(x) = x^m(1 + {}^m C_1 x + {}^{m+1} C_2 x^2 + \dots).$$

The coefficient of x^n in this enumerator $C_m(x)$ is the coefficient of x^{n-m} in $(1 + {}^m C_1 x + {}^{m+1} C_2 x^2 + \dots)$ which is equal to ${}^{n-1} C_{n-m} = {}^{n-1} C_{m-1}$.

So number of compositions of n with exactly m parts is equal to ${}^{n-1} C_{m-1}$.

The enumerating GF for **composition with no restriction on the number of parts** $C(x)$ can be obtained from $C_m(x)$.

The enumerating GF for **composition with no restriction on the number of parts** $C(x)$ can be obtained from $C_m(x)$.

$$C(x) = \sum_{m=1}^{\infty} C_m(x) =$$

The enumerating GF for **composition with no restriction on the number of parts** $C(x)$ can be obtained from $C_m(x)$.

$$C(x) = \sum_{m=1}^{\infty} C_m(x) = \sum_{m=1}^{\infty} x^m (1-x)^{-m} = \sum_{m=1}^{\infty} \left(\frac{x}{1-x} \right)^m.$$

The enumerating GF for **composition with no restriction on the number of parts** $C(x)$ can be obtained from $C_m(x)$.

$$C(x) = \sum_{m=1}^{\infty} C_m(x) = \sum_{m=1}^{\infty} x^m (1-x)^{-m} = \sum_{m=1}^{\infty} \left(\frac{x}{1-x}\right)^m.$$

Put $\left(\frac{x}{1-x}\right) = t$.

The enumerating GF for **composition with no restriction on the number of parts** $C(x)$ can be obtained from $C_m(x)$.

$$C(x) = \sum_{m=1}^{\infty} C_m(x) = \sum_{m=1}^{\infty} x^m (1-x)^{-m} = \sum_{m=1}^{\infty} \left(\frac{x}{1-x}\right)^m.$$

Put $\left(\frac{x}{1-x}\right) = t$.

$$\text{So, } C(x) = \sum_{m=1}^{\infty} t^m = t + t^2 + t^3 + \dots = t(1 + t + t^2 + \dots) = \left(\frac{t}{1-t}\right)$$

The enumerating GF for **composition with no restriction on the number of parts** $C(x)$ can be obtained from $C_m(x)$.

$$C(x) = \sum_{m=1}^{\infty} C_m(x) = \sum_{m=1}^{\infty} x^m (1-x)^{-m} = \sum_{m=1}^{\infty} \left(\frac{x}{1-x}\right)^m.$$

Put $\left(\frac{x}{1-x}\right) = t$.

$$\text{So, } C(x) = \sum_{m=1}^{\infty} t^m = t + t^2 + t^3 + \dots = t(1 + t + t^2 + \dots) = \left(\frac{t}{1-t}\right)$$

$$C(x) =$$

The enumerating GF for **composition with no restriction on the number of parts** $C(x)$ can be obtained from $C_m(x)$.

$$C(x) = \sum_{m=1}^{\infty} C_m(x) = \sum_{m=1}^{\infty} x^m (1-x)^{-m} = \sum_{m=1}^{\infty} \left(\frac{x}{1-x}\right)^m.$$

Put $\left(\frac{x}{1-x}\right) = t$.

$$\text{So, } C(x) = \sum_{m=1}^{\infty} t^m = t + t^2 + t^3 + \dots = t(1 + t + t^2 + \dots) = \left(\frac{t}{1-t}\right)$$

$$C(x) = x(1-2x)^{-1}$$

The enumerating GF for **composition with no restriction on the number of parts** $C(x)$ can be obtained from $C_m(x)$.

$$C(x) = \sum_{m=1}^{\infty} C_m(x) = \sum_{m=1}^{\infty} x^m (1-x)^{-m} = \sum_{m=1}^{\infty} \left(\frac{x}{1-x}\right)^m.$$

$$\text{Put } \left(\frac{x}{1-x}\right) = t.$$

$$\text{So, } C(x) = \sum_{m=1}^{\infty} t^m = t + t^2 + t^3 + \dots = t(1 + t + t^2 + \dots) = \left(\frac{t}{1-t}\right)$$

$$C(x) = x(1-2x)^{-1}$$

$$C(x) = x(1 + 2x + (2x)^2 + \dots) = \sum_{n=1}^{\infty} 2^{n-1} x^n.$$

The enumerating GF for **composition with no restriction on the number of parts** $C(x)$ can be obtained from $C_m(x)$.

$$C(x) = \sum_{m=1}^{\infty} C_m(x) = \sum_{m=1}^{\infty} x^m (1-x)^{-m} = \sum_{m=1}^{\infty} \left(\frac{x}{1-x}\right)^m.$$

Put $\left(\frac{x}{1-x}\right) = t$.

So, $C(x) = \sum_{m=1}^{\infty} t^m = t + t^2 + t^3 + \dots = t(1 + t + t^2 + \dots) = \left(\frac{t}{1-t}\right)$

$$C(x) = x(1-2x)^{-1}$$

$$C(x) = x(1 + 2x + (2x)^2 + \dots) = \sum_{n=1}^{\infty} 2^{n-1} x^n.$$

Number of compositions with no restriction on the number of parts is the coefficient of x^n in $\sum_{n=1}^{\infty} 2^{n-1} x^n$ which is equal to 2^{n-1} .

Generating function for unrestricted partitions

Generating function for unrestricted partitions

The polynomial $1 + x + x^2 + \dots + x^k + \dots + x^n$ is concerned with one's in the partition.

Generating function for unrestricted partitions

The polynomial $1 + x + x^2 + \dots + x^k + \dots + x^n$ is concerned with one's in the partition.

The polynomial $1 + x^2 + (x^2)^2 + \dots x^{2k} + \dots$ is concerned with two's in the partition and in particular, the coefficient of $x^{2k} = (x^2)^k$ represents the case of just k number of 2's in the partition.

Generating function for unrestricted partitions

The polynomial $1 + x + x^2 + \dots + x^k + \dots + x^n$ is concerned with one's in the partition.

The polynomial $1 + x^2 + (x^2)^2 + \dots x^{2k} + \dots$ is concerned with two's in the partition and in particular, the coefficient of $x^{2k} = (x^2)^k$ represents the case of just k number of 2's in the partition.

So the GF for partition should contain one factor for 1's, one for 2's and so on.

Generating function for unrestricted partitions

The polynomial $1 + x + x^2 + \dots + x^k + \dots + x^n$ is concerned with one's in the partition.

The polynomial $1 + x^2 + (x^2)^2 + \dots x^{2k} + \dots$ is concerned with two's in the partition and in particular, the coefficient of $x^{2k} = (x^2)^k$ represents the case of just k number of 2's in the partition.

So the GF for partition should contain one factor for 1's, one for 2's and so on.

$$GF = (1 + x + x^2 + \dots)(1 + x^2 + x^4 + \dots)(1 + x^3 + x^6 + \dots) \dots$$

Generating function for unrestricted partitions

The polynomial $1 + x + x^2 + \dots + x^k + \dots + x^n$ is concerned with one's in the partition.

The polynomial $1 + x^2 + (x^2)^2 + \dots x^{2k} + \dots$ is concerned with two's in the partition and in particular, the coefficient of $x^{2k} = (x^2)^k$ represents the case of just k number of 2's in the partition.

So the GF for partition should contain one factor for 1's, one for 2's and so on.

$$\text{GF} = (1 + x + x^2 + \dots)(1 + x^2 + x^4 + \dots)(1 + x^3 + x^6 + \dots) \dots$$

$$= (1 - x)^{-1}(1 - x^2)^{-1}(1 - x^3)^{-1} \dots$$

Problems

Problems

Q1. Prove that number of partitions of n in which no integer occurs more than twice as a part is equal to the number of partitions of n into parts not divisible by 3.

Problems

Q1. Prove that number of partitions of n in which no integer occurs more than twice as a part is equal to the number of partitions of n into parts not divisible by 3.

Solution: Enumerator for partitions of n into parts not divisible by 3 is

Problems

Q1. Prove that number of partitions of n in which no integer occurs more than twice as a part is equal to the number of partitions of n into parts not divisible by 3.

Solution: Enumerator for partitions of n into parts not divisible by 3 is

$$G_0(x) = (1 - x)^{-1}(1 - x^2)^{-1}(1 - x^4)^{-1} \dots$$

Problems

Q1. Prove that number of partitions of n in which no integer occurs more than twice as a part is equal to the number of partitions of n into parts not divisible by 3.

Solution: Enumerator for partitions of n into parts not divisible by 3 is

$$G_0(x) = (1 - x)^{-1}(1 - x^2)^{-1}(1 - x^4)^{-1} \dots$$

Enumerator for partitions of n in which no integer occurs more than twice as a part is equal to

Problems

Q1. Prove that number of partitions of n in which no integer occurs more than twice as a part is equal to the number of partitions of n into parts not divisible by 3.

Solution: Enumerator for partitions of n into parts not divisible by 3 is

$$G_0(x) = (1 - x)^{-1}(1 - x^2)^{-1}(1 - x^4)^{-1} \dots$$

Enumerator for partitions of n in which no integer occurs more than twice as a part is equal to

$$G_1(x) = (1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$$

Problems

Q1. Prove that number of partitions of n in which no integer occurs more than twice as a part is equal to the number of partitions of n into parts not divisible by 3.

Solution: Enumerator for partitions of n into parts not divisible by 3 is

$$G_0(x) = (1 - x)^{-1}(1 - x^2)^{-1}(1 - x^4)^{-1} \dots$$

Enumerator for partitions of n in which no integer occurs more than twice as a part is equal to

$$G_1(x) = (1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$$

$$G_1(x) = \frac{(1 - x)(1 + x + x^2)}{(1 - x)} \cdot \frac{(1 - x^2)(1 + x^2 + x^4)}{(1 - x^2)} \cdot \frac{(1 - x^3)(1 + x^3 + x^6)}{(1 - x^3)} \dots$$

Problems

Q1. Prove that number of partitions of n in which no integer occurs more than twice as a part is equal to the number of partitions of n into parts not divisible by 3.

Solution: Enumerator for partitions of n into parts not divisible by 3 is

$$G_0(x) = (1 - x)^{-1}(1 - x^2)^{-1}(1 - x^4)^{-1} \dots$$

Enumerator for partitions of n in which no integer occurs more than twice as a part is equal to

$$G_1(x) = (1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$$

$$\begin{aligned} G_1(x) &= \frac{(1 - x)(1 + x + x^2)}{(1 - x)} \cdot \frac{(1 - x^2)(1 + x^2 + x^4)}{(1 - x^2)} \cdot \frac{(1 - x^3)(1 + x^3 + x^6)}{(1 - x^3)} \dots \\ &= \frac{(1 - x^3)(1 - x^6)(1 - x^9)(1 - x^{12})(1 - x^{15})(1 - x^{18}) \dots}{(1 - x)(1 - x^2)(1 - x^3)(1 - x^4)(1 - x^5)(1 - x^6) \dots} \end{aligned}$$

Problems

Q1. Prove that number of partitions of n in which no integer occurs more than twice as a part is equal to the number of partitions of n into parts not divisible by 3.

Solution: Enumerator for partitions of n into parts not divisible by 3 is

$$G_0(x) = (1 - x)^{-1}(1 - x^2)^{-1}(1 - x^4)^{-1} \dots$$

Enumerator for partitions of n in which no integer occurs more than twice as a part is equal to

$$G_1(x) = (1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$$

$$G_1(x) = \frac{(1 - x)(1 + x + x^2)}{(1 - x)} \cdot \frac{(1 - x^2)(1 + x^2 + x^4)}{(1 - x^2)} \cdot \frac{(1 - x^3)(1 + x^3 + x^6)}{(1 - x^3)} \dots$$

$$= \frac{(1 - x^3)(1 - x^6)(1 - x^9)(1 - x^{12})(1 - x^{15})(1 - x^{18}) \dots}{(1 - x)(1 - x^2)(1 - x^3)(1 - x^4)(1 - x^5)(1 - x^6) \dots}$$

$$= (1 - x)^{-1}(1 - x^2)^{-1}(1 - x^4)^{-1} \dots = G_0(x).$$

Let $n = 7$

Let $n = 7$

Partition of 7 such that no part is divisible by 3 are

Let $n = 7$

Partition of 7 such that no part is divisible by 3 are
7, 511, 52, 2221, 421, 4111, 211111, 1111111, 22111.

Let $n = 7$

Partition of 7 such that no part is divisible by 3 are

7, 511, 52, 2221, 421, 4111, 211111, 1111111, 22111.

Partitions of $n = 7$ in which no integer occurs more than twice are

Let $n = 7$

Partition of 7 such that no part is divisible by 3 are

7, 511, 52, 2221, 421, 4111, 211111, 1111111, 22111.

Partitions of $n = 7$ in which no integer occurs more than twice are

7, 511, 322, 421, 331, 52, 43, 61, 3211.

Q2. Show that the number of partitions of n in which every part is odd is equal to the number of partitions of n with unequal (distinct) parts.

Q2. Show that the number of partitions of n in which every part is odd is equal to the number of partitions of n with unequal (distinct) parts.

Solution:

Q2. Show that the number of partitions of n in which every part is odd is equal to the number of partitions of n with unequal (distinct) parts.

Solution: GF for the number of partitions of n in which every part is odd is equal to

Q2. Show that the number of partitions of n in which every part is odd is equal to the number of partitions of n with unequal (distinct) parts.

Solution: GF for the number of partitions of n in which every part is odd is equal to

$$G_0(x) = (1 - x)^{-1}(1 - x^3)^{-1}(1 - x^5)^{-1}(1 - x^7)^{-1} \dots$$

Q2. Show that the number of partitions of n in which every part is odd is equal to the number of partitions of n with unequal (distinct) parts.

Solution: GF for the number of partitions of n in which every part is odd is equal to

$$G_0(x) = (1 - x)^{-1}(1 - x^3)^{-1}(1 - x^5)^{-1}(1 - x^7)^{-1} \dots$$

GF for the number of partitions of n with unequal (distinct) parts is equal to

Q2. Show that the number of partitions of n in which every part is odd is equal to the number of partitions of n with unequal (distinct) parts.

Solution: GF for the number of partitions of n in which every part is odd is equal to

$$G_0(x) = (1 - x)^{-1}(1 - x^3)^{-1}(1 - x^5)^{-1}(1 - x^7)^{-1} \dots$$

GF for the number of partitions of n with unequal (distinct) parts is equal to

$$G_1(x) = (1 + x)(1 + x^2)(1 + x^3) \dots$$

Q2. Show that the number of partitions of n in which every part is odd is equal to the number of partitions of n with unequal (distinct) parts.

Solution: GF for the number of partitions of n in which every part is odd is equal to

$$G_0(x) = (1 - x)^{-1}(1 - x^3)^{-1}(1 - x^5)^{-1}(1 - x^7)^{-1} \dots$$

GF for the number of partitions of n with unequal (distinct) parts is equal to

$$G_1(x) = (1 + x)(1 + x^2)(1 + x^3) \dots$$

(All distinct means each integer can occur 0 or 1 time only)

Q2. Show that the number of partitions of n in which every part is odd is equal to the number of partitions of n with unequal (distinct) parts.

Solution: GF for the number of partitions of n in which every part is odd is equal to

$$G_0(x) = (1 - x)^{-1}(1 - x^3)^{-1}(1 - x^5)^{-1}(1 - x^7)^{-1} \dots$$

GF for the number of partitions of n with unequal (distinct) parts is equal to

$$G_1(x) = (1 + x)(1 + x^2)(1 + x^3) \dots$$

(All distinct means each integer can occur 0 or 1 time only)

Consider, $G_1(x) = (1 + x)(1 + x^2)(1 + x^3) \dots$

Q2. Show that the number of partitions of n in which every part is odd is equal to the number of partitions of n with unequal (distinct) parts.

Solution: GF for the number of partitions of n in which every part is odd is equal to

$$G_0(x) = (1 - x)^{-1}(1 - x^3)^{-1}(1 - x^5)^{-1}(1 - x^7)^{-1} \dots$$

GF for the number of partitions of n with unequal (distinct) parts is equal to

$$G_1(x) = (1 + x)(1 + x^2)(1 + x^3) \dots$$

(All distinct means each integer can occur 0 or 1 time only)

Consider, $G_1(x) = (1 + x)(1 + x^2)(1 + x^3) \dots$

$$= \frac{(1+x)(1-x)}{(1-x)} \cdot \frac{(1+x^2)(1-x^2)}{(1-x^2)} \cdot \frac{(1+x^3)(1-x^3)}{(1-x^3)} \dots$$

Q2. Show that the number of partitions of n in which every part is odd is equal to the number of partitions of n with unequal (distinct) parts.

Solution: GF for the number of partitions of n in which every part is odd is equal to

$$G_0(x) = (1 - x)^{-1}(1 - x^3)^{-1}(1 - x^5)^{-1}(1 - x^7)^{-1} \dots$$

GF for the number of partitions of n with unequal (distinct) parts is equal to

$$G_1(x) = (1 + x)(1 + x^2)(1 + x^3) \dots$$

(All distinct means each integer can occur 0 or 1 time only)

Consider, $G_1(x) = (1 + x)(1 + x^2)(1 + x^3) \dots$

$$\begin{aligned} &= \frac{(1+x)(1-x)}{(1-x)} \cdot \frac{(1+x^2)(1-x^2)}{(1-x^2)} \cdot \frac{(1+x^3)(1-x^3)}{(1-x^3)} \dots \\ &= \frac{(1-x^2)}{(1-x)} \frac{(1-x^4)}{(1-x^2)} \frac{(1-x^6)}{(1-x^3)} \frac{(1-x^8)}{(1-x^4)} \dots \end{aligned}$$

Q2. Show that the number of partitions of n in which every part is odd is equal to the number of partitions of n with unequal (distinct) parts.

Solution: GF for the number of partitions of n in which every part is odd is equal to

$$G_0(x) = (1 - x)^{-1}(1 - x^3)^{-1}(1 - x^5)^{-1}(1 - x^7)^{-1} \dots$$

GF for the number of partitions of n with unequal (distinct) parts is equal to

$$G_1(x) = (1 + x)(1 + x^2)(1 + x^3) \dots$$

(All distinct means each integer can occur 0 or 1 time only)

Consider, $G_1(x) = (1 + x)(1 + x^2)(1 + x^3) \dots$

$$= \frac{(1+x)(1-x)}{(1-x)} \cdot \frac{(1+x^2)(1-x^2)}{(1-x^2)} \cdot \frac{(1+x^3)(1-x^3)}{(1-x^3)} \dots$$

$$= \frac{(1-x^2)}{(1-x)} \frac{(1-x^4)}{(1-x^2)} \frac{(1-x^6)}{(1-x^3)} \frac{(1-x^8)}{(1-x^4)} \dots$$

$$= (1 - x)^{-1}(1 - x^3)^{-1}(1 - x^5)^{-1} \dots$$

$$= G_0(x).$$

Let $n = 7$

Let $n = 7$

Partition of 7 such that every part is odd are

Let $n = 7$

Partition of 7 such that every part is odd are
511, 331, 31111, 1111111, 7.

Let $n = 7$

Partition of 7 such that every part is odd are
511, 331, 31111, 1111111, 7.

Partitions of $n = 7$ with distinct parts are

Let $n = 7$

Partition of 7 such that every part is odd are
511, 331, 31111, 1111111, 7.

Partitions of $n = 7$ with distinct parts are
421, 52, 43, 61, 7.

Ferrors graph

Ferrers graph

It is a graph to represent a partition by an array of dots.

Ferrors graph

It is a graph to represent a partition by an array of dots.
It has the following property.

Ferrers graph

It is a graph to represent a partition by an array of dots.

It has the following property.

- (i) There is one row for each part.
- (ii) The number of dots in any row is the size of that part.
- (iii) An upper row always contains at least as many dots as a lower row.
- (iv) The rows are aligned to the left.

Ferrers graph

It is a graph to represent a partition by an array of dots.

It has the following property.

- (i) There is one row for each part.
- (ii) The number of dots in any row is the size of that part.
- (iii) An upper row always contains at least as many dots as a lower row.
- (iv) The rows are aligned to the left.

Example:

Consider the partition 5 3 2 2.

Ferrers graph

It is a graph to represent a partition by an array of dots.

It has the following property.

- (i) There is one row for each part.
- (ii) The number of dots in any row is the size of that part.
- (iii) An upper row always contains at least as many dots as a lower row.
- (iv) The rows are aligned to the left.

Example:

Consider the partition 5 3 2 2.

• • • • •

Ferrers graph


It is a graph to represent a partition by an array of dots.

It has the following property.

- (i) There is one row for each part.
- (ii) The number of dots in any row is the size of that part.
- (iii) An upper row always contains at least as many dots as a lower row.
- (iv) The rows are aligned to the left.

Example:

Consider the partition 5 3 2 2.



The partition obtained by reading the Ferrer's graph by column is called *conjugate partition*.

The partition obtained by reading the Ferrer's graph by column is called *conjugate partition*.

The conjugate partition of 5 3 2 2 is 4 4 2 1 1.

The partition obtained by reading the Ferrer's graph by column is called *conjugate partition*.

The conjugate partition of 5 3 2 2 is 4 4 2 1 1.

A partition whose Ferrer's graph reads the same by rows and by columns is called **self-conjugate**.

The partition obtained by reading the Ferrer's graph by column is called *conjugate partition*.

The conjugate partition of 5 3 2 2 is 4 4 2 1 1.

A partition whose Ferrer's graph reads the same by rows and by columns is called **self-conjugate**.

Example: 5 4 2 2 1, 4 3 2 1.

The partition obtained by reading the Ferrer's graph by column is called *conjugate partition*.

The conjugate partition of 5 3 2 2 is 4 4 2 1 1.

A partition whose Ferrer's graph reads the same by rows and by columns is called **self-conjugate**.

Example: 5 4 2 2 1, 4 3 2 1.

• • • • •

The partition obtained by reading the Ferrer's graph by column is called *conjugate partition*.

The conjugate partition of 5 3 2 2 is 4 4 2 1 1.

A partition whose Ferrer's graph reads the same by rows and by columns is called **self-conjugate**.

Example: 5 4 2 2 1, 4 3 2 1.

• • • • •

• • • •

• •

• •

•