

THE PRINCIPLE OF INCLUSION AND EXCLUSION

THE PRINCIPLE OF INCLUSION AND EXCLUSION

Consider N objects and the two properties a and b ; then the number of objects without both these properties is given by

THE PRINCIPLE OF INCLUSION AND EXCLUSION

Consider N objects and the two properties a and b ; then the number of objects without both these properties is given by

$N(a'b') = N - N(a) - N(b) + N(ab)$, where $N(a)$ is number of objects with property a and $N(b)$ is those with property b .

THE PRINCIPLE OF INCLUSION AND EXCLUSION

Consider N objects and the two properties a and b ; then the number of objects without both these properties is given by

$N(a'b') = N - N(a) - N(b) + N(ab)$, where $N(a)$ is number of objects with property a and $N(b)$ is those with property b .

Theorem 1.1

Consider the N objects, of which $N(a_1)$ having the property a_1 , $N(a_2)$ the property a_2, \dots , $N(a_r)$ the property a_r , $N(a_1 a_2)$ having both the properties a_1 and a_2 , ..., $N(a_1 a_2 \dots a_r)$ having all the properties a_1, a_2, \dots, a_r , then the number $N(a'_1 a'_2 \dots a'_r)$ with none of these properties is given by

$$\begin{aligned} N(a'_1 a'_2 \dots a'_r) &= \\ N - N(a_1) - N(a_2) - \dots - N(a_r) &+ N(a_1 a_2) + N(a_1 a_3) + \dots + (-1)^r N(a_1 \dots a_r) \\ &= N - \sum N(a_i) + \sum_{i < j} N(a_i a_j) - \sum N(a_i a_j a_k) + \dots + (-1)^r N(a_1 a_2 \dots a_r). \end{aligned}$$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$N = 6300, N(a_1) =$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$

$N(a_1 a_2) =$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180,$$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) =$$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

$$N(a'_1 a'_2 a'_3) =$$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

$$N(a'_1 a'_2 a'_3) = 6300 - 2100 - 1260 - 900 + 420 + 300 + 180 - 60 = 2880.$$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

$$N(a'_1 a'_2 a'_3) = 6300 - 2100 - 1260 - 900 + 420 + 300 + 180 - 60 = 2880.$$

Q2. How many arrangements of the digits 0, 1, 2, ..., 9 are there in which the first digit is greater than 1 and the last digit is less than 8?

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

$$N(a'_1 a'_2 a'_3) = 6300 - 2100 - 1260 - 900 + 420 + 300 + 180 - 60 = 2880.$$

Q2. How many arrangements of the digits 0, 1, 2, ..., 9 are there in which the first digit is greater than 1 and the last digit is less than 8?

Ans:

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

$$N(a'_1 a'_2 a'_3) = 6300 - 2100 - 1260 - 900 + 420 + 300 + 180 - 60 = 2880.$$

Q2. How many arrangements of the digits 0, 1, 2, ..., 9 are there in which the first digit is greater than 1 and the last digit is less than 8?

Ans:

Let F: Set of all arrangements with 0 or 1 in the first digit.

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

$$N(a'_1 a'_2 a'_3) = 6300 - 2100 - 1260 - 900 + 420 + 300 + 180 - 60 = 2880.$$

Q2. How many arrangements of the digits 0, 1, 2, ..., 9 are there in which the first digit is greater than 1 and the last digit is less than 8?

Ans:

Let F: Set of all arrangements with 0 or 1 in the first digit.

L: Set of all arrangements with 8 or 9 in the last digit.

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

$$N(a'_1 a'_2 a'_3) = 6300 - 2100 - 1260 - 900 + 420 + 300 + 180 - 60 = 2880.$$

Q2. How many arrangements of the digits 0, 1, 2, ..., 9 are there in which the first digit is greater than 1 and the last digit is less than 8?

Ans:

Let F: Set of all arrangements with 0 or 1 in the first digit.

L: Set of all arrangements with 8 or 9 in the last digit.

$$N = 10!,$$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

$$N(a'_1 a'_2 a'_3) = 6300 - 2100 - 1260 - 900 + 420 + 300 + 180 - 60 = 2880.$$

Q2. How many arrangements of the digits 0, 1, 2, ..., 9 are there in which the first digit is greater than 1 and the last digit is less than 8?

Ans:

Let F: Set of all arrangements with 0 or 1 in the first digit.

L: Set of all arrangements with 8 or 9 in the last digit.

$$N = 10!, N(F) =$$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

$$N(a'_1 a'_2 a'_3) = 6300 - 2100 - 1260 - 900 + 420 + 300 + 180 - 60 = 2880.$$

Q2. How many arrangements of the digits 0, 1, 2, ..., 9 are there in which the first digit is greater than 1 and the last digit is less than 8?

Ans:

Let F: Set of all arrangements with 0 or 1 in the first digit.

L: Set of all arrangements with 8 or 9 in the last digit.

$$N = 10!, N(F) = 2 \times 9!, N(L) =$$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

$$N(a'_1 a'_2 a'_3) = 6300 - 2100 - 1260 - 900 + 420 + 300 + 180 - 60 = 2880.$$

Q2. How many arrangements of the digits 0, 1, 2, ..., 9 are there in which the first digit is greater than 1 and the last digit is less than 8?

Ans:

Let F: Set of all arrangements with 0 or 1 in the first digit.

L: Set of all arrangements with 8 or 9 in the last digit.

$$N = 10!, N(F) = 2 \times 9!, N(L) = 2 \times 9!,$$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

$$N(a'_1 a'_2 a'_3) = 6300 - 2100 - 1260 - 900 + 420 + 300 + 180 - 60 = 2880.$$

Q2. How many arrangements of the digits 0, 1, 2, ..., 9 are there in which the first digit is greater than 1 and the last digit is less than 8?

Ans:

Let F: Set of all arrangements with 0 or 1 in the first digit.

L: Set of all arrangements with 8 or 9 in the last digit.

$$N = 10!, N(F) = 2 \times 9!, N(L) = 2 \times 9!, N(FL) =$$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

$$N(a'_1 a'_2 a'_3) = 6300 - 2100 - 1260 - 900 + 420 + 300 + 180 - 60 = 2880.$$

Q2. How many arrangements of the digits 0, 1, 2, ..., 9 are there in which the first digit is greater than 1 and the last digit is less than 8?

Ans:

Let F: Set of all arrangements with 0 or 1 in the first digit.

L: Set of all arrangements with 8 or 9 in the last digit.

$$N = 10!, N(F) = 2 \times 9!, N(L) = 2 \times 9!, N(FL) = 2 \times 2 \times 8!.$$

Problems

Q1. How many integers between 1 and 6300 are not divisible by 3, 5 or 7?

Ans: Let a_1, a_2, a_3 be the properties of divisibility by 3, 5 and 7 respectively.

$$N = 6300, N(a_1) = 2100, N(a_2) = 1260, N(a_3) = 900,$$

$$N(a_1 a_2) = 420, N(a_1 a_3) = 300, N(a_2 a_3) = 180, N(a_1 a_2 a_3) = 60.$$

$$N(a'_1 a'_2 a'_3) = 6300 - 2100 - 1260 - 900 + 420 + 300 + 180 - 60 = 2880.$$

Q2. How many arrangements of the digits 0, 1, 2, ..., 9 are there in which the first digit is greater than 1 and the last digit is less than 8?

Ans:

Let F : Set of all arrangements with 0 or 1 in the first digit.

L : Set of all arrangements with 8 or 9 in the last digit.

$$N = 10!, N(F) = 2 \times 9!, N(L) = 2 \times 9!, N(FL) = 2 \times 2 \times 8!.$$

$$N(F'L') = 10! - (2 \times 9!) - (2 \times 9!) + 2 \times 2 \times 8!$$

Q3. How many positive integers ≤ 70 are relatively prime to 70?

Q3. How many positive integers ≤ 70 are relatively prime to 70?

Relatively prime to 70 means the numbers that have no common divisors with 70.

Q3. How many positive integers ≤ 70 are relatively prime to 70?

Relatively prime to 70 means the numbers that have no common divisors with 70.

The prime divisors of 70 are 2,5,7.

We want to count the number of integers ≤ 70 that do not have 2 or 5 or 7 as divisors.

Q3. How many positive integers ≤ 70 are relatively prime to 70?

Relatively prime to 70 means the numbers that have no common divisors with 70.

The prime divisors of 70 are 2,5,7.

We want to count the number of integers ≤ 70 that do not have 2 or 5 or 7 as divisors.

Let a_1 be the set of integers that are divisible by 2.

Q3. How many positive integers ≤ 70 are relatively prime to 70?

Relatively prime to 70 means the numbers that have no common divisors with 70.

The prime divisors of 70 are 2,5,7.

We want to count the number of integers ≤ 70 that do not have 2 or 5 or 7 as divisors.

Let a_1 be the set of integers that are divisible by 2.

Let a_2 be the set of integers that are divisible by 5.

Q3. How many positive integers ≤ 70 are relatively prime to 70?

Relatively prime to 70 means the numbers that have no common divisors with 70.

The prime divisors of 70 are 2,5,7.

We want to count the number of integers ≤ 70 that do not have 2 or 5 or 7 as divisors.

Let a_1 be the set of integers that are divisible by 2.

Let a_2 be the set of integers that are divisible by 5.

Let a_3 be the set of integers that are divisible by 7.

Q3. How many positive integers ≤ 70 are relatively prime to 70?

Relatively prime to 70 means the numbers that have no common divisors with 70.

The prime divisors of 70 are 2,5,7.

We want to count the number of integers ≤ 70 that do not have 2 or 5 or 7 as divisors.

Let a_1 be the set of integers that are divisible by 2.

Let a_2 be the set of integers that are divisible by 5.

Let a_3 be the set of integers that are divisible by 7.

$$N(a_1) =$$

Q3. How many positive integers ≤ 70 are relatively prime to 70?

Relatively prime to 70 means the numbers that have no common divisors with 70.

The prime divisors of 70 are 2,5,7.

We want to count the number of integers ≤ 70 that do not have 2 or 5 or 7 as divisors.

Let a_1 be the set of integers that are divisible by 2.

Let a_2 be the set of integers that are divisible by 5.

Let a_3 be the set of integers that are divisible by 7.

$$N(a_1) = 35, \quad N(a_2) = 14, \quad N(a_3) = 10,$$

Q3. How many positive integers ≤ 70 are relatively prime to 70?

Relatively prime to 70 means the numbers that have no common divisors with 70.

The prime divisors of 70 are 2,5,7.

We want to count the number of integers ≤ 70 that do not have 2 or 5 or 7 as divisors.

Let a_1 be the set of integers that are divisible by 2.

Let a_2 be the set of integers that are divisible by 5.

Let a_3 be the set of integers that are divisible by 7.

$$N(a_1) = 35, \quad N(a_2) = 14, \quad N(a_3) = 10, \quad N(a_1 a_2) =$$

Q3. How many positive integers ≤ 70 are relatively prime to 70?

Relatively prime to 70 means the numbers that have no common divisors with 70.

The prime divisors of 70 are 2,5,7.

We want to count the number of integers ≤ 70 that do not have 2 or 5 or 7 as divisors.

Let a_1 be the set of integers that are divisible by 2.

Let a_2 be the set of integers that are divisible by 5.

Let a_3 be the set of integers that are divisible by 7.

$$N(a_1) = 35, \quad N(a_2) = 14, \quad N(a_3) = 10, \quad N(a_1 a_2) = 7,$$

$$N(a_1 a_3) = 5, \quad N(a_2 a_3) = 2, \quad N(a_1 a_2 a_3) =$$

Q3. How many positive integers ≤ 70 are relatively prime to 70?

Relatively prime to 70 means the numbers that have no common divisors with 70.

The prime divisors of 70 are 2,5,7.

We want to count the number of integers ≤ 70 that do not have 2 or 5 or 7 as divisors.

Let a_1 be the set of integers that are divisible by 2.

Let a_2 be the set of integers that are divisible by 5.

Let a_3 be the set of integers that are divisible by 7.

$$N(a_1) = 35, \quad N(a_2) = 14, \quad N(a_3) = 10, \quad N(a_1 a_2) = 7,$$

$$N(a_1 a_3) = 5, \quad N(a_2 a_3) = 2, \quad N(a_1 a_2 a_3) = 1.$$

Q3. How many positive integers ≤ 70 are relatively prime to 70?

Relatively prime to 70 means the numbers that have no common divisors with 70.

The prime divisors of 70 are 2, 5, 7.

We want to count the number of integers ≤ 70 that do not have 2 or 5 or 7 as divisors.

Let a_1 be the set of integers that are divisible by 2.

Let a_2 be the set of integers that are divisible by 5.

Let a_3 be the set of integers that are divisible by 7.

$$N(a_1) = 35, \quad N(a_2) = 14, \quad N(a_3) = 10, \quad N(a_1 a_2) = 7,$$

$$N(a_1 a_3) = 5, \quad N(a_2 a_3) = 2, \quad N(a_1 a_2 a_3) = 1.$$

$$N(a'_1 a'_2 a'_3) = 70 - (35 + 14 + 10) + (7 + 5 + 2) - 1 = 24.$$

Derangements

Derangements

It is a permutation of objects that leaves no object in its position.

Derangements

It is a permutation of objects that leaves no object in its position.

Example: $(2,3,1)$ and $(3,1,2)$ are the derangements of $1,2,3$.

Derangements

It is a permutation of objects that leaves no object in its position.

Example: $(2,3,1)$ and $(3,1,2)$ are the derangements of $1,2,3$.

For $n = 4$ the total number of derangements are——

Derangements

It is a permutation of objects that leaves no object in its position.

Example: $(2,3,1)$ and $(3,1,2)$ are the derangements of $1,2,3$.

For $n = 4$ the total number of derangements are—— 9.

Derangements

It is a permutation of objects that leaves no object in its position.

Example: (2,3,1) and (3,1,2) are the derangements of 1,2,3.

For $n = 4$ the total number of derangements are—— 9.

i.e., total number of possibilities such that i^{th} digit not in i^{th} position.

Derangements

It is a permutation of objects that leaves no object in its position.

Example: (2,3,1) and (3,1,2) are the derangements of 1,2,3.

For $n = 4$ the total number of derangements are—— 9.

i.e., total number of possibilities such that i^{th} digit not in i^{th} position.

And they are the permutations

Derangements

It is a permutation of objects that leaves no object in its position.

Example: (2,3,1) and (3,1,2) are the derangements of 1,2,3.

For $n = 4$ the total number of derangements are—— 9.

i.e., total number of possibilities such that i^{th} digit not in i^{th} position.

And they are the permutations

2143, 2341, 2413, 3142, 3412, 3421, 4123, 4312, 4321.

Derangements

It is a permutation of objects that leaves no object in its position.

Example: (2,3,1) and (3,1,2) are the derangements of 1,2,3.

For $n = 4$ the total number of derangements are—— 9.

i.e., total number of possibilities such that i^{th} digit not in i^{th} position.

And they are the permutations

2143, 2341, 2413, 3142, 3412, 3421, 4123, 4312, 4321.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

Derangements

It is a permutation of objects that leaves no object in its position.

Example: (2,3,1) and (3,1,2) are the derangements of 1,2,3.

For $n = 4$ the total number of derangements are—— 9.

i.e., total number of possibilities such that i^{th} digit not in i^{th} position.

And they are the permutations

2143, 2341, 2413, 3142, 3412, 3421, 4123, 4312, 4321.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

$$N = 4!, \quad N(a_1) = N(a_2) = N(a_3) = N(a_4) =$$

Derangements

It is a permutation of objects that leaves no object in its position.

Example: (2,3,1) and (3,1,2) are the derangements of 1,2,3.

For $n = 4$ the total number of derangements are—— 9.

i.e., total number of possibilities such that i^{th} digit not in i^{th} position.

And they are the permutations

2143, 2341, 2413, 3142, 3412, 3421, 4123, 4312, 4321.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

$$N = 4!, \quad N(a_1) = N(a_2) = N(a_3) = N(a_4) = 3! = 6,$$

Derangements

It is a permutation of objects that leaves no object in its position.

Example: (2,3,1) and (3,1,2) are the derangements of 1,2,3.

For $n = 4$ the total number of derangements are—— 9.

i.e., total number of possibilities such that i^{th} digit not in i^{th} position.

And they are the permutations

2143, 2341, 2413, 3142, 3412, 3421, 4123, 4312, 4321.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

$$N = 4!, \quad N(a_1) = N(a_2) = N(a_3) = N(a_4) = 3! = 6,$$

$$N(a_1 a_2) = N(a_1 a_3) = N(a_1 a_4) = N(a_2 a_3) = N(a_2 a_4) = N(a_3 a_4) =$$

Derangements

It is a permutation of objects that leaves no object in its position.

Example: (2,3,1) and (3,1,2) are the derangements of 1,2,3.

For $n = 4$ the total number of derangements are—— 9.

i.e., total number of possibilities such that i^{th} digit not in i^{th} position.

And they are the permutations

2143, 2341, 2413, 3142, 3412, 3421, 4123, 4312, 4321.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

$$N = 4!, \quad N(a_1) = N(a_2) = N(a_3) = N(a_4) = 3! = 6,$$

$$N(a_1 a_2) = N(a_1 a_3) = N(a_1 a_4) = N(a_2 a_3) = N(a_2 a_4) = N(a_3 a_4) = 2!.$$

Derangements

It is a permutation of objects that leaves no object in its position.

Example: (2,3,1) and (3,1,2) are the derangements of 1,2,3.

For $n = 4$ the total number of derangements are—— 9.

i.e., total number of possibilities such that i^{th} digit not in i^{th} position.

And they are the permutations

2143, 2341, 2413, 3142, 3412, 3421, 4123, 4312, 4321.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

$$N = 4!, \quad N(a_1) = N(a_2) = N(a_3) = N(a_4) = 3! = 6,$$

$$N(a_1 a_2) = N(a_1 a_3) = N(a_1 a_4) = N(a_2 a_3) = N(a_2 a_4) = N(a_3 a_4) = 2!.$$

$$N(a_1 a_2 a_3) = N(a_1 a_3 a_4) = N(a_2 a_3 a_4) = 1, \quad N(a_1 a_2 a_3 a_4) =$$

Derangements

It is a permutation of objects that leaves no object in its position.

Example: (2,3,1) and (3,1,2) are the derangements of 1,2,3.

For $n = 4$ the total number of derangements are—— 9.

i.e., total number of possibilities such that i^{th} digit not in i^{th} position.

And they are the permutations

2143, 2341, 2413, 3142, 3412, 3421, 4123, 4312, 4321.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

$$N = 4!, \quad N(a_1) = N(a_2) = N(a_3) = N(a_4) = 3! = 6,$$

$$N(a_1 a_2) = N(a_1 a_3) = N(a_1 a_4) = N(a_2 a_3) = N(a_2 a_4) = N(a_3 a_4) = 2!.$$

$$N(a_1 a_2 a_3) = N(a_1 a_3 a_4) = N(a_2 a_3 a_4) = 1, \quad N(a_1 a_2 a_3 a_4) = 1.$$

Derangements

It is a permutation of objects that leaves no object in its position.

Example: (2,3,1) and (3,1,2) are the derangements of 1,2,3.

For $n = 4$ the total number of derangements are—— 9.

i.e., total number of possibilities such that i^{th} digit not in i^{th} position.

And they are the permutations

2143, 2341, 2413, 3142, 3412, 3421, 4123, 4312, 4321.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

$$N = 4!, \quad N(a_1) = N(a_2) = N(a_3) = N(a_4) = 3! = 6,$$

$$N(a_1 a_2) = N(a_1 a_3) = N(a_1 a_4) = N(a_2 a_3) = N(a_2 a_4) = N(a_3 a_4) = 2!.$$

$$N(a_1 a_2 a_3) = N(a_1 a_3 a_4) = N(a_2 a_3 a_4) = 1, \quad N(a_1 a_2 a_3 a_4) = 1.$$

$$\text{So, } N(a'_1 a'_2 a'_3 a'_4) = 4! - 4 \times 6 + {}^4C_2 \times 2 - {}^4C_3 \times 1 + 1 = 9.$$

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Ans: Let a_k be the property that the element k is in the k^{th} position, $1 \leq k \leq n$.

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Ans: Let a_k be the property that the element k is in the k^{th} position, $1 \leq k \leq n$.

The required number is $N(a'_1 a'_2 \dots a'_n)$.

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Ans: Let a_k be the property that the element k is in the k^{th} position, $1 \leq k \leq n$.

The required number is $N(a'_1 a'_2 \dots a'_n)$.

Total permutation is $N = n!$.

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Ans: Let a_k be the property that the element k is in the k^{th} position, $1 \leq k \leq n$.

The required number is $N(a'_1 a'_2 \dots a'_n)$.

Total permutation is $N = n!$.

$N(a_i) =$

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Ans: Let a_k be the property that the element k is in the k^{th} position, $1 \leq k \leq n$.

The required number is $N(a'_1 a'_2 \dots a'_n)$.

Total permutation is $N = n!$.

$N(a_i) = (n - 1)!$,

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Ans: Let a_k be the property that the element k is in the k^{th} position, $1 \leq k \leq n$.

The required number is $N(a'_1 a'_2 \dots a'_n)$.

Total permutation is $N = n!$.

$N(a_i) = (n-1)!$, $N(a_i a_j) =$

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Ans: Let a_k be the property that the element k is in the k^{th} position, $1 \leq k \leq n$.

The required number is $N(a'_1 a'_2 \dots a'_n)$.

Total permutation is $N = n!$.

$N(a_i) = (n-1)!$, $N(a_i a_j) = (n-2)!$,

Q1. How many permutations of the n distinct elements $(1,2,3,\dots,n)$ are there in which the element k is not in the k^{th} position?

Ans: Let a_k be the property that the element k is in the k^{th} position, $1 \leq k \leq n$.

The required number is $N(a'_1 a'_2 \dots a'_n)$.

Total permutation is $N = n!$.

$N(a_i) = (n-1)!$, $N(a_i a_j) = (n-2)!$, $N(a_i a_j a_k) = (n-3)!$,

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Ans: Let a_k be the property that the element k is in the k^{th} position, $1 \leq k \leq n$.

The required number is $N(a'_1 a'_2 \dots a'_n)$.

Total permutation is $N = n!$.

$N(a_i) = (n-1)!$, $N(a_i a_j) = (n-2)!$, $N(a_i a_j a_k) = (n-3)!$,

So on $N(a_1 a_2 \dots a_n) = 1$.

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Ans: Let a_k be the property that the element k is in the k^{th} position, $1 \leq k \leq n$.

The required number is $N(a'_1 a'_2 \dots a'_n)$.

Total permutation is $N = n!$.

$N(a_i) = (n-1)!$, $N(a_i a_j) = (n-2)!$, $N(a_i a_j a_k) = (n-3)!$,

So on $N(a_1 a_2 \dots a_n) = 1$.

Then the principle of inclusion and exclusion gives

$$N(a'_1 a'_2 \dots a'_n) = N - \sum_{i=1}^n N(a_i) + \sum_{i < j} N(a_i a_j) - \sum N(a_i a_j a_k) \dots + (-1)^n N(a_1 \dots a_n).$$

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Ans: Let a_k be the property that the element k is in the k^{th} position, $1 \leq k \leq n$.

The required number is $N(a'_1 a'_2 \dots a'_n)$.

Total permutation is $N = n!$.

$N(a_i) = (n-1)!$, $N(a_i a_j) = (n-2)!$, $N(a_i a_j a_k) = (n-3)!$,

So on $N(a_1 a_2 \dots a_n) = 1$.

Then the principle of inclusion and exclusion gives

$$\begin{aligned} N(a'_1 a'_2 \dots a'_n) &= N - \sum_{i=1}^n N(a_i) + \sum_{i < j} N(a_i a_j) - \sum N(a_i a_j a_k) \dots + (-1)^n N(a_1 \dots a_n). \\ &= n! - \end{aligned}$$

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Ans: Let a_k be the property that the element k is in the k^{th} position, $1 \leq k \leq n$.

The required number is $N(a'_1 a'_2 \dots a'_n)$.

Total permutation is $N = n!$.

$N(a_i) = (n-1)!$, $N(a_i a_j) = (n-2)!$, $N(a_i a_j a_k) = (n-3)!$,

So on $N(a_1 a_2 \dots a_n) = 1$.

Then the principle of inclusion and exclusion gives

$$N(a'_1 a'_2 \dots a'_n) = N - \sum_{i=1}^n N(a_i) + \sum_{i < j} N(a_i a_j) - \sum N(a_i a_j a_k) \dots + (-1)^n N(a_1 \dots a_n).$$

$$= n! - n(n-1)! + {}^n C_2 (n-2)! - {}^n C_3 (n-3)! + \dots + (-1)^i {}^n C_i (n-i)! + \dots + (-1)^n$$

Q1. How many permutations of the n distinct elements $(1, 2, 3, \dots, n)$ are there in which the element k is not in the k^{th} position?

Ans: Let a_k be the property that the element k is in the k^{th} position, $1 \leq k \leq n$.

The required number is $N(a'_1 a'_2 \dots a'_n)$.

Total permutation is $N = n!$.

$N(a_i) = (n-1)!$, $N(a_i a_j) = (n-2)!$, $N(a_i a_j a_k) = (n-3)!$,

So on $N(a_1 a_2 \dots a_n) = 1$.

Then the principle of inclusion and exclusion gives

$$N(a'_1 a'_2 \dots a'_n) = N - \sum_{i=1}^n N(a_i) + \sum_{i < j} N(a_i a_j) - \sum N(a_i a_j a_k) \dots + (-1)^n N(a_1 \dots a_n).$$

$$= n! - n(n-1)! + {}^n C_2 (n-2)! - {}^n C_3 (n-3)! + \dots + (-1)^i {}^n C_i (n-i)! + \dots + (-1)^n$$

$$N(a'_1 a'_2 \dots a'_n) = n! \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \dots + \frac{(-1)^n}{n!} \right).$$

We note that $N(a'_1 a'_2 \dots, a'_n) = n!(1 - \frac{1}{1!} + \frac{1}{2!} - \dots + \frac{(-1)^n}{n!}) \cong \frac{n!}{e}$, if n large.

We note that $N(a'_1 a'_2 \dots, a'_n) = n!(1 - \frac{1}{1!} + \frac{1}{2!} - \dots + \frac{(-1)^n}{n!}) \cong \frac{n!}{e}$, if n large.

$$(e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots)$$

We note that $N(a'_1 a'_2 \dots, a'_n) = n!(1 - \frac{1}{1!} + \frac{1}{2!} - \dots + \frac{(-1)^n}{n!}) \cong \frac{n!}{e}$, if n large.

$$(e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots)$$

For $n = 6$ the total number of derangements are

We note that $N(a'_1 a'_2 \dots, a'_n) = n!(1 - \frac{1}{1!} + \frac{1}{2!} - \dots + \frac{(-1)^n}{n!}) \cong \frac{n!}{e}$, if n large.

$$(e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots)$$

For $n = 6$ the total number of derangements are
 $6!(\frac{1}{2} - \frac{1}{3!} + \dots + \frac{1}{6!}) = 265$.

Q2. Show that the proportion of the Permutatition of $\{1, 2, \dots, n\}$ which contains no consecutive pair $(i, i + 1)$ for any i is approximately $\frac{n+1}{ne}$ as n becomes large.

Q2. Show that the proportion of the Permutatition of $\{1, 2, \dots, n\}$ which contains no consecutive pair $(i, i + 1)$ for any i is approximately $\frac{n+1}{ne}$ as n becomes large.

Solution: Let a_i be the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq n - 1$.

Q2. Show that the proportion of the Permutatition of $\{1, 2, \dots, n\}$ which contains no consecutive pair $(i, i + 1)$ for any i is approximately $\frac{n+1}{ne}$ as n becomes large.

Solution: Let a_i be the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq n - 1$.

$N = n!$, $N(a_i) =$

Q2. Show that the proportion of the Permutatition of $\{1, 2, \dots, n\}$ which contains no consecutive pair $(i, i + 1)$ for any i is approximately $\frac{n+1}{ne}$ as n becomes large.

Solution: Let a_i be the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq n - 1$.

$$N = n!, \quad N(a_i) = (n - 1)!.$$

Because, for a_1 i.e., $(1, 2)$ occurs consecutively, we consider $(1, 2)$ as one number. Therefore, we have $(n - 1)$ numbers.

Q2. Show that the proportion of the Permutatition of $\{1, 2, \dots, n\}$ which contains no consecutive pair $(i, i + 1)$ for any i is approximately $\frac{n+1}{ne}$ as n becomes large.

Solution: Let a_i be the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq n - 1$.

$$N = n!, \quad N(a_i) = (n - 1)!.$$

Because, for a_1 i.e., $(1, 2)$ occurs consecutively, we consider $(1, 2)$ as one number. Therefore, we have $(n - 1)$ numbers.

$$N(a_i a_j) =$$

Q2. Show that the proportion of the Permutatition of $\{1, 2, \dots, n\}$ which contains no consecutive pair $(i, i + 1)$ for any i is approximately $\frac{n+1}{ne}$ as n becomes large.

Solution: Let a_i be the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq n - 1$.

$$N = n!, \quad N(a_i) = (n - 1)!.$$

Because, for a_1 i.e., $(1, 2)$ occurs consecutively, we consider $(1, 2)$ as one number. Therefore, we have $(n - 1)$ numbers.

$$N(a_i a_j) = (n - 2)!$$

Q2. Show that the proportion of the Permutatition of $\{1, 2, \dots, n\}$ which contains no consecutive pair $(i, i + 1)$ for any i is approximately $\frac{n+1}{ne}$ as n becomes large.

Solution: Let a_i be the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq n - 1$.

$$N = n!, \quad N(a_i) = (n - 1)!.$$

Because, for a_1 i.e., $(1, 2)$ occurs consecutively, we consider $(1, 2)$ as one number. Therefore, we have $(n - 1)$ numbers.

$$N(a_i a_j) = (n - 2)!$$

$$N(a_i a_j a_k) = (n - 3)!$$

Q2. Show that the proportion of the Permutatition of $\{1, 2, \dots, n\}$ which contains no consecutive pair $(i, i + 1)$ for any i is approximately $\frac{n+1}{ne}$ as n becomes large.

Solution: Let a_i be the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq n - 1$.

$$N = n!, \quad N(a_i) = (n - 1)!.$$

Because, for a_1 i.e., $(1, 2)$ occurs consecutively, we consider $(1, 2)$ as one number. Therefore, we have $(n - 1)$ numbers.

$$N(a_i a_j) = (n - 2)!$$

$$N(a_i a_j a_k) = (n - 3)!$$

\vdots

$$N(a_1 a_2 \dots a_{n-1}) = 1.$$

Q2. Show that the proportion of the Permutatition of $\{1, 2, \dots, n\}$ which contains no consecutive pair $(i, i + 1)$ for any i is approximately $\frac{n+1}{ne}$ as n becomes large.

Solution: Let a_i be the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq n - 1$.

$$N = n!, \quad N(a_i) = (n - 1)!.$$

Because, for a_1 i.e., $(1, 2)$ occurs consecutively, we consider $(1, 2)$ as one number. Therefore, we have $(n - 1)$ numbers.

$$N(a_i a_j) = (n - 2)!$$

$$N(a_i a_j a_k) = (n - 3)!$$

\vdots

$$N(a_1 a_2 \dots a_{n-1}) = 1.$$

$$N(a'_1 a'_2 \dots) = N - \sum_{i=1}^{n-1} N(a_i) + \sum_{i < j} N(a_i a_j) - \sum N(a_i a_j a_k) - \dots$$

Q2. Show that the proportion of the Permutatition of $\{1, 2, \dots, n\}$ which contains no consecutive pair $(i, i + 1)$ for any i is approximately $\frac{n+1}{ne}$ as n becomes large.

Solution: Let a_i be the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq n - 1$.

$$N = n!, \quad N(a_i) = (n - 1)!.$$

Because, for a_1 i.e., $(1, 2)$ occurs consecutively, we consider $(1, 2)$ as one number. Therefore, we have $(n - 1)$ numbers.

$$N(a_i a_j) = (n - 2)!$$

$$N(a_i a_j a_k) = (n - 3)!$$

\vdots

$$N(a_1 a_2 \dots a_{n-1}) = 1.$$

$$N(a'_1 a'_2 \dots) = N - \sum_{i=1}^{n-1} N(a_i) + \sum_{i < j} N(a_i a_j) - \sum N(a_i a_j a_k) - \dots$$

$$= n! - {}^{n-1}C_1(n-1)! + {}^{n-1}C_2(n-2)! - {}^{n-1}C_3(n-3)! + \dots$$

$$N(a'_1 a'_2 \dots) = n! - {}^{n-1}C_1(n-1)! + {}^{n-1}C_2(n-2)! - {}^{n-1}C_3(n-3)! + \dots$$

$$\begin{aligned}
 N(a'_1 a'_2 \dots) &= n! - {}^{n-1}C_1(n-1)! + {}^{n-1}C_2(n-2)! - {}^{n-1}C_3(n-3)! + \dots \\
 &= (n-1)! \left\{ n - (n-1) + \frac{n-2}{2!} - \frac{n-3}{3!} + \dots \right\}
 \end{aligned}$$

$$\begin{aligned}
N(a'_1 a'_2 \dots) &= n! - {}^{n-1}C_1(n-1)! + {}^{n-1}C_2(n-2)! - {}^{n-1}C_3(n-3)! + \dots \\
&= (n-1)! \left\{ n - (n-1) + \frac{n-2}{2!} - \frac{n-3}{3!} + \dots \right\} \\
&= (n-1)! \left\{ \left(1 - \frac{2}{2!} + \frac{3}{3!} - \frac{4}{4!} + \dots\right) + n \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots\right) \right\}
\end{aligned}$$

$$\begin{aligned}
N(a'_1 a'_2 \dots) &= n! - {}^{n-1}C_1(n-1)! + {}^{n-1}C_2(n-2)! - {}^{n-1}C_3(n-3)! + \dots \\
&= (n-1)! \left\{ n - (n-1) + \frac{n-2}{2!} - \frac{n-3}{3!} + \dots \right\} \\
&= (n-1)! \left\{ \left(1 - \frac{2}{2!} + \frac{3}{3!} - \frac{4}{4!} + \dots\right) + n \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots\right) \right\} \\
&= (n-1)! \left\{ \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots\right) + n \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots\right) \right\}
\end{aligned}$$

$$\begin{aligned}
N(a'_1 a'_2 \dots) &= n! - {}^{n-1}C_1(n-1)! + {}^{n-1}C_2(n-2)! - {}^{n-1}C_3(n-3)! + \dots \\
&= (n-1)! \left\{ n - (n-1) + \frac{n-2}{2!} - \frac{n-3}{3!} + \dots \right\} \\
&= (n-1)! \left\{ \left(1 - \frac{2}{2!} + \frac{3}{3!} - \frac{4}{4!} + \dots\right) + n \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots\right) \right\} \\
&= (n-1)! \left\{ \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots\right) + n \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots\right) \right\} \\
&= (n-1)! \left(\frac{1}{e} + \frac{n}{e} \right)
\end{aligned}$$

$$\begin{aligned}
N(a'_1 a'_2 \dots) &= n! - {}^{n-1}C_1(n-1)! + {}^{n-1}C_2(n-2)! - {}^{n-1}C_3(n-3)! + \dots \\
&= (n-1)! \left\{ n - (n-1) + \frac{n-2}{2!} - \frac{n-3}{3!} + \dots \right\} \\
&= (n-1)! \left\{ \left(1 - \frac{2}{2!} + \frac{3}{3!} - \frac{4}{4!} + \dots\right) + n \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots\right) \right\} \\
&= (n-1)! \left\{ \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots\right) + n \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots\right) \right\} \\
&= (n-1)! \left(\frac{1}{e} + \frac{n}{e} \right) = (n-1)! \left(\frac{n+1}{e} \right).
\end{aligned}$$

$$\begin{aligned}
N(a'_1 a'_2 \dots) &= n! - {}^{n-1}C_1(n-1)! + {}^{n-1}C_2(n-2)! - {}^{n-1}C_3(n-3)! + \dots \\
&= (n-1)! \left\{ n - (n-1) + \frac{n-2}{2!} - \frac{n-3}{3!} + \dots \right\} \\
&= (n-1)! \left\{ \left(1 - \frac{2}{2!} + \frac{3}{3!} - \frac{4}{4!} + \dots\right) + n \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots\right) \right\} \\
&= (n-1)! \left\{ \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots\right) + n \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots\right) \right\} \\
&= (n-1)! \left(\frac{1}{e} + \frac{n}{e} \right) = (n-1)! \left(\frac{n+1}{e} \right).
\end{aligned}$$

Proportion of permutation is $\frac{(n-1)! \left(\frac{n+1}{e} \right)}{n!} = \frac{n+1}{ne}.$

Consider $n = 4$.

Consider $n = 4$. Here, a_i is the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq 3$.

Consider $n = 4$. Here, a_i is the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq 3$.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

Consider $n = 4$. Here, a_i is the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq 3$.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

Here, a_1 is the property that 12 is together.

Consider $n = 4$. Here, a_i is the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq 3$.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

Here, a_1 is the property that 12 is together.

$N(a_1) = 6$ which are 1234, 1243, 3124, 3412, 4123, 4312.

Consider $n = 4$. Here, a_i is the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq 3$.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

Here, a_1 is the property that 12 is together.

$N(a_1) = 6$ which are 1234, 1243, 3124, 3412, 4123, 4312.

Then, $a_1 a_2$ is the property that 12 is together and 23 is together which means 123 is together.

Consider $n = 4$. Here, a_i is the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq 3$.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

Here, a_1 is the property that 12 is together.

$N(a_1) = 6$ which are 1234, 1243, 3124, 3412, 4123, 4312.

Then, $a_1 a_2$ is the property that 12 is together and 23 is together which means 123 is together.

$N(a_1 a_2) = 2$, which are 1234, 4123.

Consider $n = 4$. Here, a_i is the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq 3$.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

Here, a_1 is the property that 12 is together.

$N(a_1) = 6$ which are 1234, 1243, 3124, 3412, 4123, 4312.

Then, $a_1 a_2$ is the property that 12 is together and 23 is together which means 123 is together.

$N(a_1 a_2) = 2$, which are 1234, 4123.

$a_1 a_3$ is the property that 12 is together and 34 is together.

Consider $n = 4$. Here, a_i is the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq 3$.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

Here, a_1 is the property that 12 is together.

$N(a_1) = 6$ which are 1234, 1243, 3124, 3412, 4123, 4312.

Then, $a_1 a_2$ is the property that 12 is together and 23 is together which means 123 is together.

$N(a_1 a_2) = 2$, which are 1234, 4123.

$a_1 a_3$ is the property that 12 is together and 34 is together.

$N(a_1 a_3) = 2$ which are 1234, 3412.

Consider $n = 4$. Here, a_i is the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq 3$.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

Here, a_1 is the property that 12 is together.

$N(a_1) = 6$ which are 1234, 1243, 3124, 3412, 4123, 4312.

Then, $a_1 a_2$ is the property that 12 is together and 23 is together which means 123 is together.

$N(a_1 a_2) = 2$, which are 1234, 4123.

$a_1 a_3$ is the property that 12 is together and 34 is together.

$N(a_1 a_3) = 2$ which are 1234, 3412.

$N(a_1 a_2 a_3) = 1$.

Consider $n = 4$. Here, a_i is the property that $(i, i + 1)$ occurs consecutively, $1 \leq i \leq 3$.

1 2 3 4	2 1 3 4	3 1 2 4	4 1 2 3
1 2 4 3	2 1 4 3	3 1 4 2	4 1 3 2
1 3 2 4	2 3 1 4	3 2 1 4	4 2 1 3
1 3 4 2	2 3 4 1	3 2 4 1	4 2 3 1
1 4 2 3	2 4 1 3	3 4 1 2	4 3 1 2
1 4 3 2	2 4 3 1	3 4 2 1	4 3 2 1

Here, a_1 is the property that 12 is together.

$N(a_1) = 6$ which are 1234, 1243, 3124, 3412, 4123, 4312.

Then, $a_1 a_2$ is the property that 12 is together and 23 is together which means 123 is together.

$N(a_1 a_2) = 2$, which are 1234, 4123.

$a_1 a_3$ is the property that 12 is together and 34 is together.

$N(a_1 a_3) = 2$ which are 1234, 3412.

$N(a_1 a_2 a_3) = 1$.

$$N(a'_1 a'_2 a'_3 a'_4) = 4! - {}^3C_1(3!) + {}^3C_2(2!) - 1 = 11.$$