


## INTRODUCTION

Phytoplankton cause algal blooms, some of which are harmful - HABs



Massive amounts of fish kills due to past events of HABs in Singapore




**Dinoflagellate blooms**  
 2009- 200,000 farmed fishes  
 2014- 500 tonnes of fish from 53 farms  
 2015 - 600 tonnes of fish in 77 farms/ loss at \$1.3 million/farmer

- HAB mitigation & control:** Early & precise characterization of bloom causing organisms.
- Challenge:** Tedious conventional methods requiring expertise and slow analysis.
- Solution:** Third Generation Sequencing with MinION™ - relatively lower costs, precise & rapid analyses.


## OBJECTIVES

### 1. Design and optimize:


- Methodology for MinION sequencing of eukaryotic microalgal rDNA unit (18S - 28S)
- Bioinformatics pipeline for error correction and analyses of sequence




### 2. Build a reference database containing these rDNA error corrected sequences




## SEQUENCED SAMPLES



## EXPERIMENTAL METHODS



## RESULTS



### Single species sequences analyses

Table 1: Similarity of canu (MinION) sequences compared to PR2 references (containing only 18S sequences) based on the pairwise distance matrix using MEGA version 7.2

	<i>Coscinodiscus</i> sp.	<i>Cyclotella</i> sp.	<i>Scrippsiella</i> sp.	<i>Karlodinium veneficum</i>	<i>Karenia mikimotoi</i>	<i>Tetraselmis</i> sp.
Accuracy of consensus compared to PR2	99.50%	98.95%	99.70%	99.50%	99.00%	99.30%

Table 2: Similarity of canu (MinION) sequences compared to NCBI-WGS references (full length rDNA sequences) based on the pairwise distance matrix using MEGA version 7.2

	<i>Picochlorum</i> sp SENEW3	<i>Ostreococcus tauri</i>	<i>Phaeodactylum tricornutum</i>
Accuracy of consensus compared to PR2	99.55%	98.80%	99.20%
Accuracy of consensus compared to WGS	99.50%	99.20%	99.20%

Table 3: Reduction in error rates and increased Q-scores of MinION


Single species sequences	Sequencing error rate	Q-score
Raw reads	12 - 22%	6.6 - 9
Error corrected reads	≤ 0.7%	21.5

$$Q \text{ (Phred score)} = -10 \log_{10} P$$

Average accuracy of the reads = 99.3%


### Microscopy Vs MinION™

Factor	Microscope	MinION™
Analysis time	~5-48hrs. per sample	~<2 hrs. for 96 samples/MinION
Identification level	Phylum, genus and sometimes species level.	Phylum, genus and species level identification
Detection limit	2 cells/mL	<10 cells/mL
Accuracy	Dependent on taxonomic expertise	Taxonomist not required due to the availability of a bioinformatics pipeline




Phylogenetic tree constructed using the Neighbor Joining method of canu consensus sequences clustered with PR2 and WGS reference sequences

### Environmental samples analyses



### 'LASer' :Long Amplicon Sequence Repository- error corrected rDNA sequences



## REFERENCES

- Piles of dead fish at Pasir Ris beach. (2015, March 1). Retrieved July 20, 2017, from <http://www.todayonline.com/singapore/piles-dead-fish-pasir-ris-beach>.
- De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M., & Van Broeckhoven, C. (2018). NanoPack: visualizing and processing long-read sequencing data. *Bioinformatics*, 34(15), 2666-2669. doi:10.1093/bioinformatics/bty149
- Koren, S., Walenz, B.P., Berlin, K., Miller, J.R., Phillippy, A.M. *Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation*. Genome Research. (2017)

## ACKNOWLEDGEMENT

This research is supported by the National Research Foundation, Prime Minister's Office, Singapore under its Marine Science Research and Development Programme Award No. MSRD-P13. I would like to thank Edwin Tat Seng, IT Senior Engineer at NTU for all the help with installation of the required software for data analysis and for all the associated troubleshooting procedures..