

Many Treatments, Some Instruments: Sharp Identification and Specification Testing in General Selection Models

Atom Vayalinkal

University of Toronto

February 21, 2025

University of Toronto

This paper: Instrumental Variables (IV) with Multiple Treatments

Setting:

- Finitely-supported treatment: $\mathcal{D}$ is finite, ordered or unordered

This paper: Instrumental Variables (IV) with Multiple Treatments

Setting:

- Finitely-supported treatment: $\mathcal{D}$ is finite, ordered or unordered
- Finitely-supported instrument: $\mathcal{Z}$ is finite, common in empirical work

This paper: Instrumental Variables (IV) with Multiple Treatments

Setting:

- Finitely-supported treatment: $\mathcal{D}$ is finite, ordered or unordered
- Finitely-supported instrument: $\mathcal{Z}$ is finite, common in empirical work
- Y may be continuous: $\mathcal{Y}$ can be any closed subset of $\mathbb{R}^n$
- Randomly assigned instruments, not just mean independent

This paper: Instrumental Variables (IV) with Multiple Treatments

Setting:

- Finitely-supported treatment: $\mathcal{D}$ is finite, ordered or unordered
- Finitely-supported instrument: $\mathcal{Z}$ is finite, common in empirical work
- Y may be continuous: $\mathcal{Y}$ can be any closed subset of $\mathbb{R}^n$
- Randomly assigned instruments, not just mean independent

Goal: Unified (partial) identification approach that allows flexible choice of target parameter and treatment selection (or choice) model

This paper: Instrumental Variables (IV) with Multiple Treatments

Setting:

- Finitely-supported treatment: $\mathcal{D}$ is finite, ordered or unordered
- Finitely-supported instrument: $\mathcal{Z}$ is finite, common in empirical work
- Y may be continuous: $\mathcal{Y}$ can be any closed subset of $\mathbb{R}^n$
- Randomly assigned instruments, not just mean independent

Goal: Unified (partial) identification approach that allows flexible choice of target parameter and treatment selection (or choice) model

Scope: Models that impose assumptions on selection, but not outcome

Model

Observables (Y, D, Z) , unobservables $(\mathbf{V}, \varepsilon_Y)$, functions g_D and g_Y :

$$\text{Treatment Selection Equation: } D = g_D(Z, \mathbf{V}) , \quad (1)$$

$$\text{Outcome Equation: } Y = g_Y(D, \mathbf{V}, \varepsilon_Y) , \quad (2)$$

$$\text{Independence Condition: } Z, \mathbf{V}, \varepsilon_Y \text{ are mutually independent.} \quad (3)$$

Model

Observables (Y, D, Z) , unobservables $(\mathbf{V}, \varepsilon_Y)$, functions g_D and g_Y :

$$\text{Treatment Selection Equation: } D = g_D(Z, \mathbf{V}) , \quad (1)$$

$$\text{Outcome Equation: } Y = g_Y(D, \mathbf{V}, \varepsilon_Y) , \quad (2)$$

$$\text{Independence Condition: } Z, \mathbf{V}, \varepsilon_Y \text{ are mutually independent.} \quad (3)$$

So far, **formally equivalent** to usual potential outcomes IV model, define potential outcomes and treatments:

$$Y_d := g_Y(d, \mathbf{V}, \varepsilon_Y) , \quad (4)$$

$$D_z := g_D(z, \mathbf{V}) . \quad (5)$$

Model

Observables (Y, D, Z) , unobservables $(\mathbf{V}, \varepsilon_Y)$, functions g_D and g_Y :

$$\text{Treatment Selection Equation: } D = g_D(Z, \mathbf{V}) , \quad (1)$$

$$\text{Outcome Equation: } Y = g_Y(D, \mathbf{V}, \varepsilon_Y) , \quad (2)$$

$$\text{Independence Condition: } Z, \mathbf{V}, \varepsilon_Y \text{ are mutually independent.} \quad (3)$$

So far, **formally equivalent** to usual potential outcomes IV model, define potential outcomes and treatments:

$$Y_d := g_Y(d, \mathbf{V}, \varepsilon_Y) , \quad (4)$$

$$D_z := g_D(z, \mathbf{V}) . \quad (5)$$

This paper:

Models that impose assumptions on selection, but not outcome

- **Allow:** assumptions on treatment selection equation,
i.e. restrictions on g_D and/or on $\mathbf{V}$
- **Avoid:** assumptions on outcome equation

Selection Models and Response-Types

Define an individual's *response-type*, denoted T , as the random vector

$$T := (D_z : z \in \mathcal{Z})$$

Selection Models and Response-Types

Define an individual's *response-type*, denoted T , as the random vector

$$T := (D_z : z \in \mathcal{Z})$$

→ Conditional on T , D is randomly assigned (determined by Z)

Selection Models and Response-Types

Define an individual's *response-type*, denoted $\mathbf{T}$, as the random vector

$$\mathbf{T}(\mathbf{V}) = \mathbf{T} := (D_z : z \in \mathcal{Z}) = (\mathbf{g}_D(z, \mathbf{V}) : z \in \mathcal{Z})$$

- Conditional on $\mathbf{T}$, D is randomly assigned (determined by $\mathcal{Z}$)
- Selection models restrict the set of admissible distributions for $\mathbf{T}$

Selection Models and Response-Types

Define an individual's **response-type**, denoted $\mathbf{T}$, as the random vector

$$\mathbf{T}(\mathbf{V}) = \mathbf{T} := (D_z : z \in \mathcal{Z}) = (g_D(z, \mathbf{V}) : z \in \mathcal{Z})$$

- Conditional on $\mathbf{T}$, D is randomly assigned (determined by $\mathcal{Z}$)
- Selection models restrict the set of admissible distributions for $\mathbf{T}$
- Distribution of $\mathbf{T}$ summarized by **response-type probabilities**:

$$\boldsymbol{\pi} := \left(\mathbb{P}(\mathbf{T} = \boldsymbol{\tau}) : \boldsymbol{\tau} \in \mathcal{D}^{\mathcal{Z}} \right)$$

Selection Models and Response-Types

Define an individual's **response-type**, denoted T , as the random vector

$$T(V) = T := (D_z : z \in \mathcal{Z}) = (g_D(z, V) : z \in \mathcal{Z})$$

- Conditional on T , D is randomly assigned (determined by Z)
- Selection models restrict the set of admissible distributions for T
- Distribution of T summarized by **response-type probabilities**:

$$\pi := \left(\mathbb{P}(T = \tau) : \tau \in \mathcal{D}^{\mathcal{Z}} \right)$$

Heckman and Pinto (2018): **discrete IV identification problem is finite mixture problem** involving response-type conditional distributions

- **This paper**: **attack mixture problem directly**, develop new results on (partial) identification of components in such mixture models

Overview of Results

For selection models, paper provides **tractable** characterizations of

1. their empirical content and (sharp) testable implications,

Overview of Results

For selection models, paper provides **tractable** characterizations of

1. their empirical content and (sharp) testable implications,
2. the identified set for a large class of target parameters, including mean counterfactuals such as response-type probabilities, mean outcomes, and mean treatment effects,

Overview of Results

For selection models, paper provides **tractable** characterizations of

1. their empirical content and (sharp) testable implications,
2. the identified set for a large class of target parameters, including mean counterfactuals such as response-type probabilities, mean outcomes, and mean treatment effects,
3. the identified set for (joint) potential outcome distributions.

Overview of Results

For selection models, paper provides **tractable** characterizations of

1. their empirical content and (sharp) testable implications,
2. the identified set for a large class of target parameters, including mean counterfactuals such as response-type probabilities, mean outcomes, and mean treatment effects,
3. the identified set for (joint) potential outcome distributions.

→ Results are **sharp under random assignment** of Z

Overview of Results

For selection models, paper provides **tractable** characterizations of

1. their empirical content and (sharp) testable implications,
2. the identified set for a large class of target parameters, including mean counterfactuals such as response-type probabilities, mean outcomes, and mean treatment effects,
3. the identified set for (joint) potential outcome distributions.

→ Results are **sharp under random assignment** of Z

The proposed approach can be used to derive:

- **Analytic bounds** on target parameters → not “black box”

Overview of Results

For selection models, paper provides **tractable** characterizations of

1. their empirical content and (sharp) testable implications,
2. the identified set for a large class of target parameters, including mean counterfactuals such as response-type probabilities, mean outcomes, and mean treatment effects,
3. the identified set for (joint) potential outcome distributions.

→ Results are **sharp under random assignment** of Z

The proposed approach can be used to derive:

- **Analytic bounds** on target parameters → not “black box”
- Inclusion tests for inference on target parameters
→ can be inverted to construct confidence regions

Overview of Results

For selection models, paper provides **tractable** characterizations of

1. their empirical content and (sharp) testable implications,
2. the identified set for a large class of target parameters, including mean counterfactuals such as response-type probabilities, mean outcomes, and mean treatment effects,
3. the identified set for (joint) potential outcome distributions.

→ Results are **sharp under random assignment** of Z

The proposed approach can be used to derive:

- **Analytic bounds** on target parameters → not “black box”
- Inclusion tests for inference on target parameters
→ can be inverted to construct confidence regions
- **Specification tests** for treatment selection models

Overview of Results

For selection models, paper provides **tractable** characterizations of

1. their empirical content and (sharp) testable implications,
2. the identified set for a large class of target parameters, including mean counterfactuals such as response-type probabilities, mean outcomes, and mean treatment effects,
3. the identified set for (joint) potential outcome distributions.

→ Results are **sharp under random assignment** of Z

The proposed approach can be used to derive:

- **Analytic bounds** on target parameters → not “black box”
- Inclusion tests for inference on target parameters
→ can be inverted to construct confidence regions
- **Specification tests** for treatment selection models

For “large” problems where analytic methods may be impractical, I propose computationally feasible algorithm.

Example: Moving to Opportunity (MTO) Experiment

Let D denote neighbourhood choice. Here, individuals choose from three different types of neighbourhoods: “high” poverty (H), “medium” poverty (M), and “low” poverty (L).

Example: Moving to Opportunity (MTO) Experiment

Let D denote neighbourhood choice. Here, individuals choose from three different types of neighbourhoods: “high” poverty (H), “medium” poverty (M), and “low” poverty (L).

Let $Z = (Z_M, Z_L)$ be the randomly assigned instrument. Individuals living in high poverty neighbourhoods were randomly offered rent vouchers:

$(Z_M, Z_L) = (0, 0)$: No vouchers

$(Z_M, Z_L) = (0, 1)$: Vouchers for L neighbourhoods

$(Z_M, Z_L) = (1, 1)$: Vouchers for M and L neighbourhoods

Example: Moving to Opportunity (MTO) Experiment

Let D denote neighbourhood choice. Here, individuals choose from three different types of neighbourhoods: “high” poverty (H), “medium” poverty (M), and “low” poverty (L).

Let $Z = (Z_M, Z_L)$ be the randomly assigned instrument. Individuals living in high poverty neighbourhoods were randomly offered rent vouchers:

$(Z_M, Z_L) = (0, 0)$: No vouchers

$(Z_M, Z_L) = (0, 1)$: Vouchers for L neighbourhoods

$(Z_M, Z_L) = (1, 1)$: Vouchers for M and L neighbourhoods

→ Therefore, $D \in \{H, M, L\}$ and $Z \in \{(0, 0), (0, 1), (1, 1)\}$.

→ Response-type: $T = (D_{(0,0)}, D_{(0,1)}, D_{(1,1)})$.

Example: Moving to Opportunity (MTO) Experiment

Suppose we impose “Unordered Partial Monotonicity” (Mountjoy, 2022):

- Z_M incentivizes $D = M$ and Z_L incentivizes $D = L$
- Changes in Z don't effect choices among **irrelevant** alternatives (e.g., $Z : (0, 0) \rightarrow (0, 1)$ cannot cause $D : H \rightarrow M$)

Example: Moving to Opportunity (MTO) Experiment

Suppose we impose “Unordered Partial Monotonicity” (Mountjoy, 2022):

- Z_M incentivizes $D = M$ and Z_L incentivizes $D = L$
- Changes in Z don't effect choices among **irrelevant** alternatives (e.g., $Z : (0, 0) \rightarrow (0, 1)$ **cannot cause** $D : H \rightarrow M$)

Shifting $Z : (1, 1) \rightarrow (0, 1)$ causes two types of moves:

- Individuals with $T \in \{(H, L, M), (M, L, M), (L, L, M)\}$: $M \rightarrow L$
- Individuals with $T = (H, H, M)$: $M \rightarrow H$

Example: Moving to Opportunity (MTO) Experiment

Suppose we impose “Unordered Partial Monotonicity” (Mountjoy, 2022):

- Z_M incentivizes $D = M$ and Z_L incentivizes $D = L$
- Changes in Z don't effect choices among irrelevant alternatives (e.g., $Z : (0, 0) \rightarrow (0, 1)$ cannot cause $D : H \rightarrow M$)

Shifting $Z : (1, 1) \rightarrow (0, 1)$ causes two types of moves:

- Individuals with $T \in \{(H, L, M), (M, L, M), (L, L, M)\}$: $M \rightarrow L$
- Individuals with $T = (H, H, M)$: $M \rightarrow H$

The following weighted average can be (point) identified

$$\omega_{M \rightarrow L} E \left[Y_L - Y_M \mid D_{(0,1)} = L, D_{(1,1)} = M \right] \\ + \omega_{M \rightarrow H} E \left[Y_H - Y_M \mid D_{(0,1)} = H, D_{(1,1)} = M \right] ,$$

but the two component effects are only partially identified.

Other Examples of Potential Applications

- Multi-dimensional instrument(s) or treatment(s)
 - **Multi-dimensional treatment:** disentangling different dimensions
 - Mediation: want to separate direct, indirect, and total effects
 - More generally: settings with complex incentive schemes, or complicated relationship between Z and D
- Ordered Treatments: e.g., Impact of Incarceration, Schooling etc.
- Nonparametric Discrete Choice Models
 - **MTO, Head Start**, College Choice...
- Other settings with multi-dimensional heterogeneity
 - Survey/RCT Non-Response Bias (e.g., Dutz et al., 2022)
- Exclusion Restriction Violations and/or Sample Selection Problems
 - See, e.g., Kroft, Mourifié, and Vayalinkal (2024)
- Sensitivity analysis for point-identified models
 - In the spirit of Noack (2021)

Related IV Literature

Point Identification: Angrist and Imbens (1995)... Heckman and Vytlacil (1999, 2001, 2007a,b)... Kirkebøen et al. (2016); [Heckman and Pinto \(2018\)](#); Lee and Salanié (2018); Mountjoy (2022)...

Partial Identification: Manski (1989, 1990); Balke and Pearl (1997)... [Beresteanu et al. \(2012\)](#)... Mogstad et al. (2018); Torgovitsky (2019); Kitagawa (2021); Kamat et al. (2023); Marx (2024); [Bai et al. \(2024\)](#)...

- **This paper:** sharp under full IV independence for any selection model; allows multivalued treatments and continuous outcomes; tractable, but not “black box”
- **This paper cannot:** incorporate general assumptions on outcomes, or continuous instruments

Specification Testing : Kitagawa (2015); Mourifié and Wan (2017); Kédagni and Mourifié (2020); Sun (2023)...

- **This paper:** sharp test for general selection models with discrete IV

Preliminaries: Setup

P denotes distribution of **observed** data (Y, D, Z) .

Preliminaries: Setup

P denotes distribution of **observed** data (Y, D, Z) .

Q denotes **unknown** (“parent”) distribution of

$$((Y_d : d \in \mathcal{D}), (D_z : z \in \mathcal{Z}), Z) = ((Y_d : d \in \mathcal{D}), \mathbf{T}, Z).$$

Preliminaries: Setup

P denotes distribution of **observed** data (Y, D, Z) .

Q denotes **unknown** (“parent”) distribution of

$$((Y_d : d \in \mathcal{D}), (D_z : z \in \mathcal{Z}), Z) = ((Y_d : d \in \mathcal{D}), \mathbf{T}, Z).$$

$\mathcal{M}$ denotes set of parent dists Q **consistent with model**.

Preliminaries: Setup

P denotes distribution of **observed** data (Y, D, Z) .

Q denotes **unknown** (“parent”) distribution of

$$((Y_d : d \in \mathcal{D}), (D_z : z \in \mathcal{Z}), Z) = ((Y_d : d \in \mathcal{D}), \mathbf{T}, Z).$$

$\mathcal{M}$ denotes set of parent dists Q **consistent with model**.

For any $Q \in \mathcal{M}$, define P_Q as dist. of (Y, D, Z) **generated by Q** .

Preliminaries: Setup

P denotes distribution of **observed** data (Y, D, Z) .

Q denotes **unknown** (“parent”) distribution of

$$((Y_d : d \in \mathcal{D}), (D_z : z \in \mathcal{Z}), Z) = ((Y_d : d \in \mathcal{D}), \mathbf{T}, Z).$$

$\mathcal{M}$ denotes set of parent dists Q **consistent with model**.

For any $Q \in \mathcal{M}$, define P_Q as dist. of (Y, D, Z) **generated by Q** .

Define $\mathcal{Q}$ as **identified set for Q** , i.e. set of Q **consistent w/ model & data**:

$$\mathcal{Q} := \{Q \in \mathcal{M} \mid P_Q = P\}$$

Preliminaries: Setup

P denotes distribution of **observed** data (Y, D, Z) .

Q denotes **unknown** (“parent”) distribution of

$$((Y_d : d \in \mathcal{D}), (D_z : z \in \mathcal{Z}), Z) = ((Y_d : d \in \mathcal{D}), \mathbf{T}, Z).$$

$\mathcal{M}$ denotes set of parent dists Q **consistent with model**.

For any $Q \in \mathcal{M}$, define P_Q as dist. of (Y, D, Z) **generated by Q** .

Define $\mathcal{Q}$ as **identified set for Q** , i.e. set of Q **consistent w/ model & data**:

$$\mathcal{Q} := \{Q \in \mathcal{M} \mid P_Q = P\}$$

For any $Q \in \mathcal{M}$, denote by π_Q **the dist. of $\mathbf{T}$ under Q** , i.e.

$$\pi_Q := \left(Q(\mathbf{T} = \tau) : \tau \in \mathcal{D}^{\mathcal{Z}} \right)$$

Preliminaries: Treatment Selection Model

Definition.

A **treatment selection model** is a 3-tuple $\mathbb{S} = (\mathcal{V}_{\mathbb{S}}, \mathcal{F}_{\mathbb{S}}, \mathcal{G}_{\mathbb{S}})$, where

$\mathcal{V}_{\mathbb{S}}$ is a set ,

$\mathcal{F}_{\mathbb{S}}$ is a subset of $\{F \mid F \text{ is a distribution with } \text{supp}(F) \subseteq \mathcal{V}_{\mathbb{S}}\}$,

and $\mathcal{G}_{\mathbb{S}}$ is a subset of the set of functions $\{\mathcal{Z} \times \mathcal{V}_{\mathbb{S}} \rightarrow \mathcal{D}\}$.

- $\mathcal{V}_{\mathbb{S}}$ represents restrictions on $\text{supp}(V)$,
- $\mathcal{F}_{\mathbb{S}}$ represents restrictions on the distribution of V ,
- $\mathcal{G}_{\mathbb{S}}$ represents shape or functional-form restrictions on g_D .

Preliminaries: Treatment Selection Model

Definition.

A **treatment selection model** is a 3-tuple $\mathbb{S} = (\mathcal{V}_{\mathbb{S}}, \mathcal{F}_{\mathbb{S}}, \mathcal{G}_{\mathbb{S}})$, where

$\mathcal{V}_{\mathbb{S}}$ is a set ,

$\mathcal{F}_{\mathbb{S}}$ is a subset of $\{F \mid F \text{ is a distribution with } \text{supp}(F) \subseteq \mathcal{V}_{\mathbb{S}}\}$,

and $\mathcal{G}_{\mathbb{S}}$ is a subset of the set of functions $\{\mathcal{Z} \times \mathcal{V}_{\mathbb{S}} \rightarrow \mathcal{D}\}$.

$Q \in \mathcal{M}$ **satisfies** $\mathbb{S}$ iff there exists $(Y, D, Z, \mathbf{V}, \varepsilon_Y, g_D, g_Y)$ such that:

- (i) $\text{supp}(\mathbf{V}) \subseteq \mathcal{V}_{\mathbb{S}}$, $\text{Law}(\mathbf{V}) \in \mathcal{F}_{\mathbb{S}}$ and $g_D \in \mathcal{G}_{\mathbb{S}}$
- (ii) they satisfy model (1)-(3) and generate Q

→ $\mathcal{V}_{\mathbb{S}}$ represents restrictions on $\text{supp}(\mathbf{V})$,

→ $\mathcal{F}_{\mathbb{S}}$ represents restrictions on the distribution of $\mathbf{V}$,

→ $\mathcal{G}_{\mathbb{S}}$ represents shape or functional-form restrictions on g_D .

Preliminaries: Treatment Selection Model

Definition.

A **treatment selection model** is a 3-tuple $\mathbb{S} = (\mathcal{V}_{\mathbb{S}}, \mathcal{F}_{\mathbb{S}}, \mathcal{G}_{\mathbb{S}})$, where

$\mathcal{V}_{\mathbb{S}}$ is a set ,

$\mathcal{F}_{\mathbb{S}}$ is a subset of $\{F \mid F \text{ is a distribution with } \text{supp}(F) \subseteq \mathcal{V}_{\mathbb{S}}\}$,

and $\mathcal{G}_{\mathbb{S}}$ is a subset of the set of functions $\{\mathcal{Z} \times \mathcal{V}_{\mathbb{S}} \rightarrow \mathcal{D}\}$.

$Q \in \mathcal{M}$ **satisfies** $\mathbb{S}$ iff there exists $(Y, D, Z, \mathbf{V}, \epsilon_Y, g_D, g_Y)$ such that:

- (i) $\text{supp}(\mathbf{V}) \subseteq \mathcal{V}_{\mathbb{S}}$, $\text{Law}(\mathbf{V}) \in \mathcal{F}_{\mathbb{S}}$ and $g_D \in \mathcal{G}_{\mathbb{S}}$
- (ii) they satisfy model (1)-(3) and generate Q

→ $\mathcal{V}_{\mathbb{S}}$ represents restrictions on $\text{supp}(\mathbf{V})$,

→ $\mathcal{F}_{\mathbb{S}}$ represents restrictions on the distribution of $\mathbf{V}$,

→ $\mathcal{G}_{\mathbb{S}}$ represents shape or functional-form restrictions on g_D .

Identification Power of Selection Models

Denote by $\mathcal{Q}_{\mathbb{S}}$ the identified set for parent distributions under $\mathbb{S}$, i.e.

$$\mathcal{Q}_{\mathbb{S}} := \{Q \in \mathcal{Q} \mid Q \text{ satisfies } \mathbb{S}\} .$$

Identification Power of Selection Models

Denote by $\mathcal{Q}_{\mathbb{S}}$ the identified set for parent distributions under $\mathbb{S}$, i.e.

$$\mathcal{Q}_{\mathbb{S}} := \{Q \in \mathcal{Q} \mid Q \text{ satisfies } \mathbb{S}\} .$$

Define $\mathcal{F}_T(\mathbb{S})$ as set of *a priori* restrictions on distribution of T , i.e.

$$\begin{aligned} \mathcal{F}_T(\mathbb{S}) &:= \left\{ \left(\mathbb{P}_F \left((g_D(z, V) : z \in \mathcal{Z}) = \tau \right) : \tau \in \mathcal{D}^{\mathcal{Z}} \right) \mid g_D \in \mathcal{G}_{\mathbb{S}}, F \in \mathcal{F}_{\mathbb{S}} \right\} \\ &= \{ \pi_Q \mid Q \in \mathcal{M} \text{ and } Q \text{ satisfies } \mathbb{S} \} . \end{aligned}$$

Identification Power of Selection Models

Denote by $\mathcal{Q}_{\mathbb{S}}$ the identified set for parent distributions under $\mathbb{S}$, i.e.

$$\mathcal{Q}_{\mathbb{S}} := \{Q \in \mathcal{Q} \mid Q \text{ satisfies } \mathbb{S}\} .$$

Define $\mathcal{F}_T(\mathbb{S})$ as set of *a priori* restrictions on distribution of T , i.e.

$$\begin{aligned} \mathcal{F}_T(\mathbb{S}) &:= \left\{ \left(\mathbb{P}_F \left((g_D(z, \mathbf{V}) : z \in \mathcal{Z}) = \tau \right) : \tau \in \mathcal{D}^{\mathcal{Z}} \right) \mid g_D \in \mathcal{G}_{\mathbb{S}}, F \in \mathcal{F}_{\mathbb{S}} \right\} \\ &= \{ \pi_Q \mid Q \in \mathcal{M} \text{ and } Q \text{ satisfies } \mathbb{S} \} . \end{aligned}$$

$\rightarrow \mathcal{F}_T(\mathbb{S})$ is a feature of $\mathbb{S}$, does not depend on data P

Identification Power of Selection Models

Denote by $\mathcal{Q}_{\mathbb{S}}$ the identified set for parent distributions under $\mathbb{S}$, i.e.

$$\mathcal{Q}_{\mathbb{S}} := \{Q \in \mathcal{Q} \mid Q \text{ satisfies } \mathbb{S}\} .$$

Define $\mathcal{F}_T(\mathbb{S})$ as set of *a priori* restrictions on distribution of T , i.e.

$$\begin{aligned} \mathcal{F}_T(\mathbb{S}) &:= \left\{ \left(\mathbb{P}_F \left((g_D(z, V) : z \in \mathcal{Z}) = \tau \right) : \tau \in \mathcal{D}^{\mathcal{Z}} \right) \mid g_D \in \mathcal{G}_{\mathbb{S}}, F \in \mathcal{F}_{\mathbb{S}} \right\} \\ &= \{ \pi_Q \mid Q \in \mathcal{M} \text{ and } Q \text{ satisfies } \mathbb{S} \} . \end{aligned}$$

→ $\mathcal{F}_T(\mathbb{S})$ is a feature of $\mathbb{S}$, does not depend on data P

Theorem 1.

For any treatment selection model $\mathbb{S}$, $\mathcal{Q}_{\mathbb{S}} = \{Q \in \mathcal{Q} \mid \pi_Q \in \mathcal{F}_T(\mathbb{S})\}$.

→ $\mathbb{S}$ has identifying power on features of Q **only** through dist. of T

Parameters of Interest

Today: For simplicity:

- assume **scalar continuous** Y , supported on an interval $\mathcal{Y} \subseteq \mathbb{R}$
- focus on models that simply **eliminate** some response-types
→ i.e. models that restrict

$$\mathcal{T} := \text{supp}(T) = \{\tau_1, \dots, \tau_{N_T}\}$$

(basic example: “**monotonicity**” = “no defiers”)

Parameters of Interest

Today: For simplicity:

- assume **scalar continuous** Y , supported on an interval $\mathcal{Y} \subseteq \mathbb{R}$
- focus on models that simply **eliminate** some response-types
 - i.e. models that restrict

$$\mathcal{T} := \text{supp}(\mathbf{T}) = \{\tau_1, \dots, \tau_{N_T}\}$$

(basic example: “**monotonicity**” = “**no defiers**”)

- focus on (weighted averages of) parameters of the form

$$E[Y_d - Y_{d'} \mid \mathbf{T} = \boldsymbol{\tau}],$$

→ examples include **ATE**, **ATT**, **ATUT**, (generalized) **LATE**, **PRTE** etc.

IV Mixture Model: Basic Idea

For any $d \in \mathcal{D}$ and each $z \in \mathcal{Z}$,

We observe Y for those assigned $Z = z$ and took treatment d

We want to learn about Y_d for those of type $T = \tau$, for some $\tau \in \mathcal{T}$

IV Mixture Model: Basic Idea

For any $d \in \mathcal{D}$ and each $z \in \mathcal{Z}$,

We observe Y for those assigned $Z = z$ and took treatment d

We want to learn about Y_d for those of type $T = \tau$, for some $\tau \in \mathcal{T}$

Key: Individuals who take treatment $D = d$ when $Z = z$ consists of the mixture of individuals of types $T = \tau$ such that $\tau(z) = d$.

IV Mixture Model: Basic Idea

For any $d \in \mathcal{D}$ and each $z \in \mathcal{Z}$,

We observe Y for those assigned $Z = z$ and took treatment d

We want to learn about Y_d for those of type $T = \tau$, for some $\tau \in \mathcal{T}$

Key: Individuals who take treatment $D = d$ when $Z = z$ consists of the mixture of individuals of types $T = \tau$ such that $\tau(z) = d$.

→ Observed outcome distributions are mixture distributions:

$$\begin{aligned} \sum_{\tau \in \mathcal{T}} \mathbb{1} \{ \tau(z) = d \} \mathbb{P}(T = \tau) \mathbb{P}(Y_d \leq y \mid T = \tau) \\ = \mathbb{P}(D = d \mid Z = z) \mathbb{P}(Y \leq y \mid D = d, Z = z), \end{aligned}$$

IV Mixture Model: Basic Idea

For any $d \in \mathcal{D}$ and each $z \in \mathcal{Z}$,

We observe Y for those assigned $Z = z$ and took treatment d

We want to learn about Y_d for those of type $T = \tau$, for some $\tau \in \mathcal{T}$

Key: Individuals who take treatment $D = d$ when $Z = z$ consists of the mixture of individuals of types $T = \tau$ such that $\tau(z) = d$.

→ Observed outcome distributions are mixture distributions:

$$\begin{aligned} \sum_{\tau \in \mathcal{T}} \underbrace{\mathbb{1}\{\tau(z) = d\}}_{\text{known}} \underbrace{\mathbb{P}(T = \tau) \mathbb{P}(Y_d \leq y \mid T = \tau)}_{\text{unobserved}} \\ = \underbrace{\mathbb{P}(D = d \mid Z = z) \mathbb{P}(Y \leq y \mid D = d, Z = z)}_{\text{observed}}, \end{aligned}$$

IV Mixture Model: Basic Idea

For any $d \in \mathcal{D}$ and each $z \in \mathcal{Z}$,

We observe Y for those assigned $Z = z$ and took treatment d

We want to learn about Y_d for those of type $T = \tau$, for some $\tau \in \mathcal{T}$

Key: Individuals who take treatment $D = d$ when $Z = z$ consists of the mixture of individuals of types $T = \tau$ such that $\tau(z) = d$.

→ Observed outcome distributions are mixture distributions:

$$\begin{aligned} \sum_{\tau \in \mathcal{T}} \underbrace{\mathbb{1}\{\tau(z) = d\}}_{\text{known, feature of } \mathbb{S}} \underbrace{Q(T = \tau) Q(Y_d \leq y \mid T = \tau)}_{\text{unobserved, feature of } Q} \\ = \underbrace{P(D = d \mid Z = z) P(Y \leq y \mid D = d, Z = z)}_{\text{observed, feature of } P}, \end{aligned}$$

IV Mixture Model: Basic Idea

For any $d \in \mathcal{D}$ and each $z \in \mathcal{Z}$,

We observe Y for those assigned $Z = z$ and took treatment d

We want to learn about Y_d for those of type $T = \tau$, for some $\tau \in \mathcal{T}$

Key: Individuals who take treatment $D = d$ when $Z = z$ consists of the mixture of individuals of types $T = \tau$ such that $\tau(z) = d$.

→ Observed outcome distributions are mixture distributions:

$$\begin{aligned} \sum_{\tau \in \mathcal{T}} \underbrace{\mathbb{1}\{\tau(z) = d\}}_{\text{known, feature of } \mathbb{S}} \underbrace{Q(T = \tau) Q(Y_d \leq y \mid T = \tau)}_{\text{unobserved, feature of } Q} \\ = \underbrace{P(D = d \mid Z = z) P(Y \leq y \mid D = d, Z = z)}_{\text{observed, feature of } P}, \end{aligned}$$

→ Allowing Z to vary, get linear system with known coefficients

“Trimming” Bounds of Horowitz and Manski (1995)

Suppose all we observe is the following mixture distribution, with CDF F ,

$$F(y) = \gamma F_1(y) + (1 - \gamma) F_2(y) ,$$

where $\gamma \in (0, 1)$ is known, but F_1 and F_2 are unknown CDFs. Let $Y_1 \sim F_1$.


“Trimming” Bounds of Horowitz and Manski (1995)

Suppose all we observe is the following mixture distribution, with CDF F ,

$$F(y) = \gamma F_1(y) + (1 - \gamma) F_2(y) ,$$

where $\gamma \in (0, 1)$ is known, but F_1 and F_2 are unknown CDFs. Let $Y_1 \sim F_1$.

If $\gamma = 0.1$, sharp lower bound for $E[Y_1]$ is constructed as follows:



(a) Lower bound is area of shaded region

Figure 1: Sharp lower bound for $E[Y_1]$, with $\gamma = 0.1$


“Trimming” Bounds of Horowitz and Manski (1995)

Suppose all we observe is the following mixture distribution, with CDF F ,


$$F(y) = \gamma F_1(y) + (1 - \gamma) F_2(y),$$

where $\gamma \in (0, 1)$ is known, but F_1 and F_2 are unknown CDFs. Let $Y_1 \sim F_1$.

If $\gamma = 0.1$, sharp lower bound for $E[Y_1]$ is constructed as follows:



(a) Lower bound is area of shaded region



(b) Density of F_1

Figure 1: Sharp lower bound for $E[Y_1]$, with $\gamma = 0.1$


“Trimming” Bounds of Horowitz and Manski (1995)

Suppose all we observe is the following mixture distribution, with CDF F ,

$$F(y) = \gamma F_1(y) + (1 - \gamma) F_2(y),$$


where $\gamma \in (0, 1)$ is known, but F_1 and F_2 are unknown CDFs. Let $Y_1 \sim F_1$.

If $\gamma = 0.1$, sharp **upper** bound for $E[Y_1]$ is constructed as follows:



(a) Upper bound is area of shaded region

Density Implied by Upper Bound




(b) Density of F_1

Figure 1: Sharp upper bound for $E[Y_1]$, with $\gamma = 0.1$


“Trimming” Bounds of Horowitz and Manski (1995)

If instead we have $\gamma = 0.9$, then lower bound is



(a) Lower bound is area of shaded region

Density Implied by Lower Bound



(b) Density of F_1

Figure 2: Sharp lower bound for $E[Y_1]$, with $\gamma = 0.9$

“Trimming” Bounds of Horowitz and Manski (1995)

and upper bound is




Figure 2: Sharp upper bound for $E[Y_1]$, with $\gamma = 0.9$

Horowitz and Manski (1995) Bounds: Takeaway

The observed mixture distribution, written in terms of densities, is

$$f(y) = \gamma f_1(y) + (1 - \gamma) f_2(y) ,$$

where f_1 is the density of F_1 , and f_2 is the density of F_2 .

Horowitz and Manski (1995) Bounds: Takeaway

The observed mixture distribution, written in terms of densities, is

$$f(y) = \gamma f_1(y) + (1 - \gamma) f_2(y) ,$$

where f_1 is the density of F_1 , and f_2 is the density of F_2 .

This implies two restrictions on the function $\gamma f_1(\cdot)$:

→ point-wise restriction: $0 \leq \gamma f_1(y) \leq f(y)$, for all y

Horowitz and Manski (1995) Bounds: Takeaway

The observed mixture distribution, written in terms of densities, is

$$f(y) = \gamma f_1(y) + (1 - \gamma) f_2(y) ,$$

where f_1 is the density of F_1 , and f_2 is the density of F_2 .

This implies two restrictions on the function $\gamma f_1(\cdot)$:

→ point-wise restriction: $0 \leq \gamma f_1(y) \leq f(y)$, for all y

→ mass restriction: $\int_{-\infty}^{\infty} f_1(y) d\mu(y) = 1 \implies \int_{-\infty}^{\infty} \gamma f_1(y) d\mu(y) = \gamma$

Horowitz and Manski (1995) Bounds: Takeaway

The observed mixture distribution, written in terms of densities, is

$$f(y) = \gamma f_1(y) + (1 - \gamma) f_2(y) ,$$

where f_1 is the density of F_1 , and f_2 is the density of F_2 .

This implies two restrictions on the function $\gamma f_1(\cdot)$:

→ point-wise restriction: $0 \leq \gamma f_1(y) \leq f(y)$, for all y

→ mass restriction: $\int_{-\infty}^{\infty} f_1(y) d\mu(y) = 1 \implies \int_{-\infty}^{\infty} \gamma f_1(y) d\mu(y) = \gamma$

Note that $E[Y_1] := \frac{1}{\gamma} \int_{-\infty}^{\infty} y \gamma f_1(y) d\mu(y)$

Horowitz and Manski (1995) Bounds: Takeaway

The observed mixture distribution, written in terms of densities, is

$$f(y) = \gamma f_1(y) + (1 - \gamma) f_2(y),$$

where f_1 is the density of F_1 , and f_2 is the density of F_2 .

This implies two restrictions on the function $\gamma f_1(\cdot)$:

→ point-wise restriction: $0 \leq \gamma f_1(y) \leq f(y)$, for all y

→ mass restriction: $\int_{-\infty}^{\infty} f_1(y) d\mu(y) = 1 \implies \int_{-\infty}^{\infty} \gamma f_1(y) d\mu(y) = \gamma$

Note that $E[Y_1] := \frac{1}{\gamma} \int_{-\infty}^{\infty} y \gamma f_1(y) d\mu(y)$

→ optimizing the right hand side over all functions $\gamma f_1(\cdot)$ that satisfy the two conditions above results in the Horowitz-Manski bounds

Example 1: Head Start Impact Study

Let D denote preschool choice. Families choose between no preschool (n), competing preschools (c), and Head Start (h) for their child.

Example 1: Head Start Impact Study

Let D denote preschool choice. Families choose between no preschool (n), competing preschools (c), and Head Start (h) for their child.

The Head Start Impact Study randomized applicants into two groups:

$Z = 0$: Not offered admission to Head Start

$Z = 1$: Offered admission to Head Start.

Example 1: Head Start Impact Study

Let D denote preschool choice. Families choose between no preschool (n), competing preschools (c), and Head Start (h) for their child.

The Head Start Impact Study randomized applicants into two groups:

$Z = 0$: Not offered admission to Head Start

$Z = 1$: Offered admission to Head Start.

→ Therefore, $D \in \{n, c, h\}$ and $Z \in \{0, 1\}$.

→ Response-type: $T = (D_0, D_1)$.

Example 1 (Head Start): Partial Identification

Assume that the following assumption holds for everyone

- either $D_1 = D_0$ or $\mathbb{1}\{D_1 = h\} > \mathbb{1}\{D_0 = h\}$

Example 1 (Head Start): Partial Identification

Assume that the following assumption holds for everyone

- either $D_1 = D_0$ or $\mathbb{1}\{D_1 = h\} > \mathbb{1}\{D_0 = h\}$

Shifting $Z : 0 \rightarrow 1$ causes two types of enrolment changes:

- Individuals with $T = (n, h)$: $n \rightarrow h$
- Individuals with $T = (c, h)$: $c \rightarrow h$

Example 1 (Head Start): Partial Identification

Assume that the following assumption holds for everyone

- either $D_1 = D_0$ or $\mathbb{1}\{D_1 = h\} > \mathbb{1}\{D_0 = h\}$

Shifting $Z : 0 \rightarrow 1$ causes two types of enrolment changes:

- Individuals with $T = (n, h)$: $n \rightarrow h$
- Individuals with $T = (c, h)$: $c \rightarrow h$

The following weighted average can be (point) identified

$$\omega_{n \rightarrow h} E[Y_h - Y_n \mid D_0 = n, D_1 = h] \\ + \omega_{c \rightarrow h} E[Y_h - Y_c \mid D_0 = c, D_1 = h] ,$$

but the two component effects are only partially identified.

Example 1 (Head Start): “ h ”-response-type

Consider the effect $\mathbb{E}[Y_h - Y_c \mid \mathbf{T} = (ch)]$

Example 1 (Head Start): “ h ”-response-type

Consider the effect $\mathbb{E}[Y_h - Y_c \mid \mathbf{T} = (ch)]$

$\rightarrow \mathbb{P}[\mathbf{T} = (ch)]$ and $\mathbb{E}[Y_c \mid \mathbf{T} = (ch)]$ are point-identified

Example 1 (Head Start): “ h ”-response-type

Consider the effect $\mathbb{E}[Y_h - Y_c \mid \mathbf{T} = (ch)]$

→ $\mathbb{P}[\mathbf{T} = (ch)]$ and $\mathbb{E}[Y_c \mid \mathbf{T} = (ch)]$ are point-identified

→ Therefore, it suffices to find the bounds for

$$\mathbb{P}[\mathbf{T} = (ch)] \mathbb{E}[Y_h \mid \mathbf{T} = (ch)]$$

Example 1 (Head Start): “ h ”-response-type

Consider the effect $\mathbb{E}[Y_h - Y_c \mid \mathbf{T} = (ch)]$

→ $\mathbb{P}[\mathbf{T} = (ch)]$ and $\mathbb{E}[Y_c \mid \mathbf{T} = (ch)]$ are point-identified

→ Therefore, it suffices to find the bounds for

$$\begin{aligned}\mathbb{P}[\mathbf{T} = (ch)] \mathbb{E}[Y_h \mid \mathbf{T} = (ch)] &= \mathbb{E}[Y_h \mathbb{1}\{\mathbf{T} = (ch)\}] \\ &= \int_{\mathcal{Y}} \mathbb{P}[\mathbf{T} = (ch)] f_{Y_h|\mathbf{T}=(ch)}(y) d\mu(y)\end{aligned}$$

Example 1 (Head Start): “ h ”-response-type

Consider the effect $\mathbb{E}[Y_h - Y_c \mid \mathbf{T} = (ch)]$

→ $\mathbb{P}[\mathbf{T} = (ch)]$ and $\mathbb{E}[Y_c \mid \mathbf{T} = (ch)]$ are point-identified

→ Therefore, it suffices to find the bounds for

$$\begin{aligned}\mathbb{P}[\mathbf{T} = (ch)] \mathbb{E}[Y_h \mid \mathbf{T} = (ch)] &= \mathbb{E}[Y_h \mathbf{1}\{\mathbf{T} = (ch)\}] \\ &= \int_{\mathcal{Y}} \mathbb{P}[\mathbf{T} = (ch)] f_{Y_h \mid \mathbf{T}=(ch)}(y) d\mu(y)\end{aligned}$$

There are two types such that $D_0 \neq h$ and $D_1 = h$:

$$\{(ch), (nh)\}$$

Example 1 (Head Start): Pointwise Restriction

Define

$$\begin{aligned}\xi_{Y_h|T=(ch)}(y) &:= \mathbb{P}[T = (ch)] f_{Y_h|T=(ch)}(y) , \\ \text{and } \xi_{Y_h|T=(nh)}(y) &:= \mathbb{P}[T = (nh)] f_{Y_h|T=(nh)}(y) .\end{aligned}$$

Example 1 (Head Start): Pointwise Restriction

Define

$$\begin{aligned}\xi_{Y_h|T=(ch)}(y) &:= \mathbb{P}[T = (ch)] f_{Y_h|T=(ch)}(y) , \\ \text{and } \xi_{Y_h|T=(nh)}(y) &:= \mathbb{P}[T = (nh)] f_{Y_h|T=(nh)}(y) .\end{aligned}$$

Now, note that

$$P(D = h|Z = 1) f_{Y|D=h,Z=1}(y)$$

Example 1 (Head Start): Pointwise Restriction

Define

$$\begin{aligned}\xi_{Y_h|T=(ch)}(y) &:= \mathbb{P}[T = (ch)] f_{Y_h|T=(ch)}(y) , \\ \text{and } \xi_{Y_h|T=(nh)}(y) &:= \mathbb{P}[T = (nh)] f_{Y_h|T=(nh)}(y) .\end{aligned}$$

Now, note that

$$\begin{aligned}&P(D = h|Z = 1) f_{Y|D=h,Z=1}(y) - P(D = h|Z = 0) f_{Y|D=h,Z=0}(y) \\ &= \xi_{Y_h|T=(ch)}(y) + \xi_{Y_h|T=(nh)}(y)\end{aligned}$$

Example 1 (Head Start): Pointwise Restriction

Define

$$\begin{aligned}\xi_{Y_h|T=(ch)}(y) &:= \mathbb{P}[T = (ch)] f_{Y_h|T=(ch)}(y) , \\ \text{and } \xi_{Y_h|T=(nh)}(y) &:= \mathbb{P}[T = (nh)] f_{Y_h|T=(nh)}(y) .\end{aligned}$$

Now, note that

$$\begin{aligned}&P(D = h|Z = 1) f_{Y|D=h,Z=1}(y) - P(D = h|Z = 0) f_{Y|D=h,Z=0}(y) \\ &= \xi_{Y_h|T=(ch)}(y) + \xi_{Y_h|T=(nh)}(y)\end{aligned}$$

Example 1 (Head Start): Pointwise Restriction

Define

$$\begin{aligned}\xi_{Y_h|T=(ch)}(y) &:= \mathbb{P}[T=(ch)] f_{Y_h|T=(ch)}(y) , \\ \text{and } \xi_{Y_h|T=(nh)}(y) &:= \mathbb{P}[T=(nh)] f_{Y_h|T=(nh)}(y) .\end{aligned}$$

Now, note that

$$\begin{aligned}&P(D=h|Z=1) f_{Y|D=h,Z=1}(y) - P(D=h|Z=0) f_{Y|D=h,Z=0}(y) \\ &= \xi_{Y_h|T=(ch)}(y) + \xi_{Y_h|T=(nh)}(y)\end{aligned}$$

Defining $\xi_{Y_h|T \in [(\cdot h)]} := \xi_{Y_h|T=(ch)}(y) + \xi_{Y_h|T=(nh)}(y)$

Example 1 (Head Start): Pointwise Restriction

Define

$$\xi_{Y_h|T=(ch)}(y) := \mathbb{P}[T = (ch)] f_{Y_h|T=(ch)}(y) ,$$

and $\xi_{Y_h|T=(nh)}(y) := \mathbb{P}[T = (nh)] f_{Y_h|T=(nh)}(y) .$


Now, note that

$$\begin{aligned} & P(D = h|Z = 1) f_{Y|D=h,Z=1}(y) - P(D = h|Z = 0) f_{Y|D=h,Z=0}(y) \\ &= \xi_{Y_h|T=(ch)}(y) + \xi_{Y_h|T=(nh)}(y) \end{aligned}$$


Defining $\xi_{Y_h|T \in [(\cdot h)]} := \xi_{Y_h|T=(ch)}(y) + \xi_{Y_h|T=(nh)}(y)$, we can “trim” the identified function $\xi_{Y_h|T \in [(\cdot h)]}$ to get sharp bounds on $E[Y_h|T = (ch)]$.

Example 1 (Head Start): Sharp Lower Bound

Suppose $\mathbb{P}[T = (ch)] = 0.2$.



(a) Dark blue region has area 0.2.



(b) Density $\frac{\xi_{Y_h|T=(ch)}}{P[T=(ch)]} = f_{Y_h|T=(ch)}$ implied by sharp lower bound

Figure 3: (Simulated) Construction of sharp lower bound for $E[Y_h|T = (ch)]$

Example 2 (MTO): “ M ”-response-type

Recall: $D \in \{H, M, L\}$ and $Z = (Z_M, Z_L) \in \{(0, 0), (0, 1), (1, 1)\}$.

Response-type: $\mathbf{T} = (D_{(0,0)} D_{(0,1)} D_{(1,1)})$.

Consider the effect $\mathbb{E}[Y_H - Y_M \mid \mathbf{T} = (HHM)]$

Example 2 (MTO): “M”-response-type

Recall: $D \in \{H, M, L\}$ and $Z = (Z_M, Z_L) \in \{(0, 0), (0, 1), (1, 1)\}$.

Response-type: $\mathbf{T} = (D_{(0,0)} D_{(0,1)} D_{(1,1)})$.

Consider the effect $\mathbb{E}[Y_H - Y_M \mid \mathbf{T} = (HHM)]$

$\rightarrow \mathbb{P}[\mathbf{T} = (HHM)]$ and $\mathbb{E}[Y_H \mid \mathbf{T} = (HHM)]$ are point-identified

Example 2 (MTO): “M”-response-type

Recall: $D \in \{H, M, L\}$ and $Z = (Z_M, Z_L) \in \{(0, 0), (0, 1), (1, 1)\}$.

Response-type: $\mathbf{T} = (D_{(0,0)} D_{(0,1)} D_{(1,1)})$.

Consider the effect $\mathbb{E}[Y_H - Y_M \mid \mathbf{T} = (HHM)]$

→ $\mathbb{P}[\mathbf{T} = (HHM)]$ and $\mathbb{E}[Y_H \mid \mathbf{T} = (HHM)]$ are point-identified

→ Therefore, it suffices to find the bounds for

$$\begin{aligned}\mathbb{P}[\mathbf{T} = (HHM)] \mathbb{E}[Y_M \mid \mathbf{T} = (HHM)] &= \int_{\mathcal{Y}} \mathbb{P}[\mathbf{T} = (HHM)] f_{Y_M \mid \mathbf{T} = (HHM)}(y) d\mu(y) \\ &=: \int_{\mathcal{Y}} \xi_{Y_M \mid \mathbf{T} = (HHM)}(y) d\mu(y)\end{aligned}$$

Example 2 (MTO): “M”-response-type

Recall: $D \in \{H, M, L\}$ and $Z = (Z_M, Z_L) \in \{(0, 0), (0, 1), (1, 1)\}$.

Response-type: $\mathbf{T} = (D_{(0,0)} D_{(0,1)} D_{(1,1)})$.

Consider the effect $\mathbb{E}[Y_H - Y_M \mid \mathbf{T} = (HHM)]$

→ $\mathbb{P}[\mathbf{T} = (HHM)]$ and $\mathbb{E}[Y_H \mid \mathbf{T} = (HHM)]$ are point-identified

→ Therefore, it suffices to find the bounds for

$$\begin{aligned}\mathbb{P}[\mathbf{T} = (HHM)] \mathbb{E}[Y_M \mid \mathbf{T} = (HHM)] &= \int_{\mathcal{Y}} \mathbb{P}[\mathbf{T} = (HHM)] f_{Y_M \mid \mathbf{T} = (HHM)}(y) d\mu(y) \\ &=: \int_{\mathcal{Y}} \xi_{Y_M \mid \mathbf{T} = (HHM)}(y) d\mu(y)\end{aligned}$$

There are three types such that $D_{(0,0)} \neq M$, $D_{(0,1)} \neq M$ and $D_{(1,1)} = M$:

$$\{(HHM), (HLM), (LLM)\}$$

Example 2 (MTO): Pointwise Restrictions

In this case, the aggregate density

$$\xi_{Y_M|T \in [\cdot M]}(y) := \xi_{Y_M|T=(HHM)}(y) + \xi_{Y_M|T=(HLM)}(y) + \xi_{Y_M|T=(LLM)}(y)$$

is not point-identified, but it can be bounded.

Example 2 (MTO): Pointwise Restrictions

In this case, the aggregate density

$$\xi_{Y_M|T \in [(\cdot M)]}(y) := \xi_{Y_M|T=(HHM)}(y) + \xi_{Y_M|T=(HLM)}(y) + \xi_{Y_M|T=(LLM)}(y)$$

is not point-identified, but it can be bounded.

Here,

$$\begin{aligned} & P[D = M | Z = (1, 1)] f_{Y|D=M, Z=(1,1)}(y) \\ &= \xi_{Y_M|T \in [(\cdot M)]}(y) + \xi_{Y_M|T=(MMM)}(y) + \xi_{Y_M|T=(MLM)}(y) . \end{aligned}$$

Example 2 (MTO): Pointwise Restrictions

In this case, the aggregate density

$$\xi_{Y_M|T \in [(\cdot M)]}(y) := \xi_{Y_M|T=(HHM)}(y) + \xi_{Y_M|T=(HLM)}(y) + \xi_{Y_M|T=(LLM)}(y)$$

is not point-identified, but it can be bounded.

Here,

$$\begin{aligned} & P[D = M|Z = (1, 1)] f_{Y|D=M, Z=(1,1)}(y) \\ &= \xi_{Y_M|T \in [(\cdot M)]}(y) + \xi_{Y_M|T=(MMM)}(y) + \xi_{Y_M|T=(MLM)}(y) . \end{aligned}$$

First, we have that

$$\begin{aligned} & P[D = M|Z = (1, 1)] f_{Y|D=M, Z=(1,1)} - P[D = M|Z = (0, 1)] f_{Y|D=M, Z=(0,1)} \\ &= \xi_{Y_M|T \in [(\cdot M)]} + \xi_{Y_M|T=(MLM)} \end{aligned}$$

Example 2 (MTO): Pointwise Restrictions

In this case, the aggregate density

$$\xi_{Y_M|T \in [(\cdot M)]}(y) := \xi_{Y_M|T=(HHM)}(y) + \xi_{Y_M|T=(HLM)}(y) + \xi_{Y_M|T=(LLM)}(y)$$

is not point-identified, but it can be bounded.

Here,

$$\begin{aligned} & P[D = M|Z = (1, 1)] f_{Y|D=M, Z=(1,1)}(y) \\ &= \xi_{Y_M|T \in [(\cdot M)]}(y) + \xi_{Y_M|T=(MMM)}(y) + \xi_{Y_M|T=(MLM)}(y) . \end{aligned}$$

First, we have that

$$\begin{aligned} & P[D = M|Z = (1, 1)] f_{Y|D=M, Z=(1,1)} - P[D = M|Z = (0, 1)] f_{Y|D=M, Z=(0,1)} \\ &= \xi_{Y_M|T \in [(\cdot M)]} + \xi_{Y_M|T=(MLM)} \\ &\geq \xi_{Y_M|T \in [(\cdot M)]} \end{aligned}$$

Example 2 (MTO): Pointwise Restrictions

In this case, the aggregate density

$$\xi_{Y_M|T \in [(\cdot M)]}(y) := \xi_{Y_M|T=(HHM)}(y) + \xi_{Y_M|T=(HLM)}(y) + \xi_{Y_M|T=(LLM)}(y)$$

is not point-identified, but it can be bounded.

Here,

$$\begin{aligned} & P[D = M|Z = (1, 1)] f_{Y|D=M, Z=(1,1)}(y) \\ &= \xi_{Y_M|T \in [(\cdot M)]}(y) + \xi_{Y_M|T=(MMM)}(y) + \xi_{Y_M|T=(MLM)}(y) . \end{aligned}$$

First, we have that

$$\begin{aligned} & P[D = M|Z = (1, 1)] f_{Y|D=M, Z=(1,1)} - P[D = M|Z = (0, 1)] f_{Y|D=M, Z=(0,1)} \\ &= \xi_{Y_M|T \in [(\cdot M)]} + \xi_{Y_M|T=(MLM)} \\ &\geq \xi_{Y_M|T \in [(\cdot M)]} \end{aligned}$$

Define $\bar{f}^*(y)$ to be this bound.

Example 2 (MTO): Pointwise Restrictions

In this case, the aggregate density

$$\xi_{Y_M|T \in [(\cdot M)]}(y) := \xi_{Y_M|T=(HHM)}(y) + \xi_{Y_M|T=(HLM)}(y) + \xi_{Y_M|T=(LLM)}(y)$$

is not point-identified, but it can be bounded.

Here,

$$\begin{aligned} & P[D = M|Z = (1, 1)] f_{Y|D=M, Z=(1,1)}(y) \\ &= \xi_{Y_M|T \in [(\cdot M)]}(y) + \xi_{Y_M|T=(MMM)}(y) + \xi_{Y_M|T=(MLM)}(y) . \end{aligned}$$

Next, we have that

$$\begin{aligned} & P[D = M|Z = (1, 1)] f_{Y|D=M, Z=(1,1)} - P[D = M|Z = (0, 0)] f_{Y|D=M, Z=(0,0)} \\ &= \xi_{Y_M|T \in [(\cdot M)]} - \xi_{Y_M|T=(MLL)} \end{aligned}$$

Example 2 (MTO): Pointwise Restrictions

In this case, the aggregate density

$$\xi_{Y_M|T \in [(\cdot M)]}(y) := \xi_{Y_M|T=(HHM)}(y) + \xi_{Y_M|T=(HLM)}(y) + \xi_{Y_M|T=(LLM)}(y)$$

is not point-identified, but it can be bounded.

Here,

$$\begin{aligned} & P[D = M|Z = (1, 1)] f_{Y|D=M, Z=(1,1)}(y) \\ &= \xi_{Y_M|T \in [(\cdot M)]}(y) + \xi_{Y_M|T=(MMM)}(y) + \xi_{Y_M|T=(MLM)}(y) . \end{aligned}$$

Next, we have that

$$\begin{aligned} & P[D = M|Z = (1, 1)] f_{Y|D=M, Z=(1,1)} - P[D = M|Z = (0, 0)] f_{Y|D=M, Z=(0,0)} \\ &= \xi_{Y_M|T \in [(\cdot M)]} - \xi_{Y_M|T=(MLL)} \\ &\leq \xi_{Y_M|T \in [(\cdot M)]} \end{aligned}$$

Example 2 (MTO): Pointwise Restrictions

In this case, the aggregate density

$$\xi_{Y_M|T \in [(\cdot M)]}(y) := \xi_{Y_M|T=(HHM)}(y) + \xi_{Y_M|T=(HLM)}(y) + \xi_{Y_M|T=(LLM)}(y)$$

is not point-identified, but it can be bounded.

Here,

$$\begin{aligned} & P[D = M|Z = (1, 1)] f_{Y|D=M, Z=(1,1)}(y) \\ &= \xi_{Y_M|T \in [(\cdot M)]}(y) + \xi_{Y_M|T=(MMM)}(y) + \xi_{Y_M|T=(MLM)}(y) . \end{aligned}$$

Next, we have that

$$\begin{aligned} & P[D = M|Z = (1, 1)] f_{Y|D=M, Z=(1,1)} - P[D = M|Z = (0, 0)] f_{Y|D=M, Z=(0,0)} \\ &= \xi_{Y_M|T \in [(\cdot M)]} - \xi_{Y_M|T=(MLL)} \\ &\leq \xi_{Y_M|T \in [(\cdot M)]} \end{aligned}$$

Define $\underline{f}^*(y)$ to be the pointwise maximum of 0 and this bound.

Example 2 (MTO): Simple Bounds

So, we have the pointwise bounds $\underline{f}^*(y) \leq \xi_{Y_M|T \in [\cdot, M]}(y) \leq \overline{f}^*(y)$:

Example 2 (MTO): Simple Bounds

So, we have the pointwise bounds $\underline{f}^*(y) \leq \xi_{Y_M|T \in [\cdot, M]}(y) \leq \overline{f}^*(y)$:

→ valid bound on $\mathbb{P}[T = (HHM)] \mathbb{E}[Y_M | T = (HHM)]$ can be obtained by trimming $\overline{f}^*(y)$

Example 2 (MTO): Simple Bounds

So, we have the pointwise bounds $\underline{f}^*(y) \leq \xi_{Y_M|T \in [\cdot, M]}(y) \leq \overline{f}^*(y)$:

- valid bound on $\mathbb{P}[T = (HHM)] \mathbb{E}[Y_M | T = (HHM)]$ can be obtained by trimming $\overline{f}^*(y)$
- however, this bound is not generally sharp

Example 2 (MTO): Simple Bounds

So, we have the pointwise bounds $\underline{f}^*(y) \leq \mathbb{E}[Y_M | \mathbf{T} \in [(\cdot, M)]](y) \leq \overline{f}^*(y)$:

- valid bound on $\mathbb{P}[\mathbf{T} = (HHM)] \mathbb{E}[Y_M | \mathbf{T} = (HHM)]$ can be obtained by trimming $\overline{f}^*(y)$
- however, this bound is not generally sharp


This is because the resulting bound may imply a value of

$$\mathbb{P}[\mathbf{T} \in [(\cdot, M)]] := \mathbb{P}[\mathbf{T} = (HHM)] + \mathbb{P}[\mathbf{T} = (HLM)] + \mathbb{P}[\mathbf{T} = (LLM)]$$

that may not be feasible.

Example 2 (MTO): Non-Sharpness of Simple Bounds

Suppose the sharp upper bound on $\mathbb{P}[T \in [(\cdot \cdot M)]]$ is 0.36, and $\mathbb{P}[T = (HHM)] = 0.28$.




Pointwise bounds on $\xi_{Y_M|T \in [(\cdot \cdot M)]}$

Figure 4: (Simulated) Construction of lower bounds for $E[Y_M|T = (HHM)]$

Example 2 (MTO): Non-Sharpness of Simple Bounds

Suppose the sharp upper bound on $\mathbb{P}[T \in [(\cdot \cdot M)]]$ is 0.36, and $\mathbb{P}[T = (HHM)] = 0.28$.

Construction of Simple Lower Bound




dark blue area = 0.28

Figure 4: (Simulated) Construction of lower bounds for $E[Y_M | T = (HHM)]$

Example 2 (MTO): Non-Sharpness of Simple Bounds


Suppose the sharp upper bound on $\mathbb{P}[T \in [(\cdot \cdot M)]]$ is 0.36, and $\mathbb{P}[T = (HHM)] = 0.28$.

Construction of Simple Lower Bound



dark blue area = 0.28

Density $f_{Y_M|T=(HHM)}$ Implied by Simple Lower Bound




This density implies the simple lower bound of 3.289

Figure 4: (Simulated) Construction of lower bounds for $E[Y_M|T = (HHM)]$

Example 2 (MTO): Non-Sharpness of Simple Bounds

Suppose the sharp upper bound on $\mathbb{P}[T \in [(\cdot \cdot M)]]$ is 0.36, and $\mathbb{P}[T = (HHM)] = 0.28$.


Construction of Simple Lower Bound



dark blue area = 0.28

dark blue area + light blue area = 0.423

Density $f_{Y_M|T=(HHM)}$ Implied by Simple Lower Bound



This density implies the simple lower bound of 3.289

Figure 4: (Simulated) Construction of lower bounds for $E[Y_M|T = (HHM)]$

Example 2 (MTO): Non-Sharpness of Simple Bounds

Suppose the sharp upper bound on $\mathbb{P}[T \in [(\cdot \cdot M)]]$ is 0.36, and $\mathbb{P}[T = (HHM)] = 0.28$.

Construction of Sharp Lower Bound





Figure 4: (Simulated) Construction of lower bounds for $E[Y_M | T = (HHM)]$

Example 2 (MTO): Non-Sharpness of Simple Bounds

Suppose the sharp upper bound on $\mathbb{P}[T \in [(\cdot \cdot M)]]$ is 0.36, and $\mathbb{P}[T = (HHM)] = 0.28$.


Construction of Sharp Lower Bound



dark blue area = 0.28

dark blue area + light blue area = 0.36

Density $f_{Y_M|T=(HHM)}$ Implied by Sharp Lower Bound



This density implies the sharp lower bound of 3.665

Figure 4: (Simulated) Construction of lower bounds for $E[Y_M|T = (HHM)]$

Example 3: Multi-dimensional Treatment

Consider a setting with a **main treatment**, $M \in \{0, 1\}$, and **supplementary treatment**, $S \in \{0, 1\}$, and define $D = (M, S)$.

Example 3: Multi-dimensional Treatment

Consider a setting with a **main treatment**, $M \in \{0, 1\}$, and **supplementary treatment**, $S \in \{0, 1\}$, and define $D = (M, S)$.

Some individuals are randomly offered incentives to participate in treatment. Let $Z = (Z_M, Z_S)$ where

- $Z_M \in \{0, 1\}$ represents an **incentive to participate in M**
- $Z_S \in \{0, 1\}$ represents an **additional incentive to participate in S**

Example 3: Multi-dimensional Treatment

Consider a setting with a **main treatment**, $M \in \{0, 1\}$, and **supplementary treatment**, $S \in \{0, 1\}$, and define $D = (M, S)$.

Some individuals are randomly offered incentives to participate in treatment. Let $Z = (Z_M, Z_S)$ where

- $Z_M \in \{0, 1\}$ represents an **incentive to participate in M**
- $Z_S \in \{0, 1\}$ represents an **additional incentive to participate in S**

Suppose individuals are

- only offered supplementary incentive if offered main incentive
- only allowed supplementary treatment if they take main treatment

Example 3: Multi-dimensional Treatment

Consider a setting with a **main treatment**, $M \in \{0, 1\}$, and **supplementary treatment**, $S \in \{0, 1\}$, and define $D = (M, S)$.

Some individuals are randomly offered incentives to participate in treatment. Let $Z = (Z_M, Z_S)$ where

- $Z_M \in \{0, 1\}$ represents an **incentive to participate in M**
- $Z_S \in \{0, 1\}$ represents an **additional incentive to participate in S**

Suppose individuals are

- only offered supplementary incentive if offered main incentive
- only allowed supplementary treatment if they take main treatment

→ Therefore, $D \in \{(0, 0), (1, 0), (1, 1)\}$ and $Z \in \{(0, 0), (1, 0), (1, 1)\}$.

→ Response-type: $T = (D_{(0,0)}, D_{(1,0)}, D_{(1,1)})$.

Multi-dimensional Treatment Example

Assume that the following holds for all individuals:

1. $Z_M : 0 \rightarrow 1$ (weakly) increases $\mathbb{1}\{D \in \{(1, 0), (1, 1)\}\}$ and $\mathbb{1}\{S = 1\}$,
2. $Z_S : 0 \rightarrow 1$ (weakly) increases $\mathbb{1}\{D = (1, 1)\}$ and $\mathbb{1}\{M = 1\}$.

Multi-dimensional Treatment Example

Assume that the following holds for all individuals:

1. $Z_M : 0 \rightarrow 1$ (weakly) increases $\mathbb{1}\{D \in \{(1, 0), (1, 1)\}\}$ and $\mathbb{1}\{S = 1\}$,
2. $Z_S : 0 \rightarrow 1$ (weakly) increases $\mathbb{1}\{D = (1, 1)\}$ and $\mathbb{1}\{M = 1\}$.

Many potential parameters of interest...

Multi-dimensional Treatment Example

Assume that the following holds for all individuals:

1. $Z_M : 0 \rightarrow 1$ (weakly) increases $\mathbb{1}\{D \in \{(1, 0), (1, 1)\}\}$ and $\mathbb{1}\{S = 1\}$,
2. $Z_S : 0 \rightarrow 1$ (weakly) increases $\mathbb{1}\{D = (1, 1)\}$ and $\mathbb{1}\{M = 1\}$.

Many potential parameters of interest...

For example, consider the decomposition

$$\begin{aligned} & \underbrace{\mathbb{E} [Y_{(1,1)} - Y_{(0,0)} \mid \mathbf{T} = ((0, 0), (1, 0), (1, 1))]}_{\text{Total Effect}} \\ &= \underbrace{\mathbb{E} [Y_{(1,1)} - Y_{(1,0)} \mid \mathbf{T} = ((0, 0), (1, 0), (1, 1))]}_{\text{Supplementary Effect}} \\ &+ \underbrace{\mathbb{E} [Y_{(1,0)} - Y_{(0,0)} \mid \mathbf{T} = ((0, 0), (1, 0), (1, 1))]}_{\text{Primary Effect}} \end{aligned}$$

Multi-dimensional Treatment Example

Assume that the following holds for all individuals:

1. $Z_M : 0 \rightarrow 1$ (weakly) increases $\mathbb{1}\{D \in \{(1, 0), (1, 1)\}\}$ and $\mathbb{1}\{S = 1\}$,
2. $Z_S : 0 \rightarrow 1$ (weakly) increases $\mathbb{1}\{D = (1, 1)\}$ and $\mathbb{1}\{M = 1\}$.

Many potential parameters of interest...

For example, consider the decomposition

$$\begin{aligned} & \underbrace{\mathbb{E} [Y_{(1,1)} - Y_{(0,0)} \mid \mathbf{T} = ((0, 0), (1, 0), (1, 1))]}_{\text{Total Effect}} \\ &= \underbrace{\mathbb{E} [Y_{(1,1)} - Y_{(1,0)} \mid \mathbf{T} = ((0, 0), (1, 0), (1, 1))]}_{\text{Supplementary Effect}} \\ &+ \underbrace{\mathbb{E} [Y_{(1,0)} - Y_{(0,0)} \mid \mathbf{T} = ((0, 0), (1, 0), (1, 1))]}_{\text{Primary Effect}} \end{aligned}$$

→ **All components are only partially identified!**

Multi-dim. Treatment example: Singleton “(1, 0)”-response-type

Let's focus on the mean outcome $\mathbb{E} \left[Y_{(1,0)} \mid \mathbf{T} = (00, 10, 11) \right]$.

Multi-dim. Treatment example: Singleton “(1, 0)”-response-type

Let's focus on the mean outcome $\mathbb{E} [Y_{(1,0)} \mid \mathbf{T} = (00, 10, 11)]$.

→ there are no other response-types such that

$$D_{(0,0)} \neq (1, 0), D_{(1,1)} \neq (1, 0) \text{ and } D_{(1,0)} = (1, 0)$$

Multi-dim. Treatment example: Singleton “(1, 0)”-response-type

Let's focus on the mean outcome $\mathbb{E} [Y_{(1,0)} \mid \mathbf{T} = (00, 10, 11)]$.

→ there are no other response-types such that

$$D_{(0,0)} \neq (1, 0), D_{(1,1)} \neq (1, 0) \text{ and } D_{(1,0)} = (1, 0)$$

→ $\xi_{Y_{(1,0)}|\mathbf{T}=(00,10,11)}$ is not point-identified, but can be bounded

Multi-dim. Treatment example: Singleton “(1, 0)”-response-type

Let's focus on the mean outcome $\mathbb{E} [Y_{(1,0)} \mid \mathbf{T} = (00, 10, 11)]$.

→ there are no other response-types such that

$$D_{(0,0)} \neq (1, 0), D_{(1,1)} \neq (1, 0) \text{ and } D_{(1,0)} = (1, 0)$$

→ $\xi_{Y_{(1,0)}|\mathbf{T}=(00,10,11)}$ is not point-identified, but can be bounded

Define $\mathbf{pf}_{d|Z=z}(y) := P(D = d | Z = z) f_{Y|D=d, Z=z}(y)$.

Multi-dim. Treatment example: Singleton “(1, 0)”-response-type

Let's focus on the mean outcome $\mathbb{E} [Y_{(1,0)} \mid \mathbf{T} = (00, 10, 11)]$.

→ there are no other response-types such that

$$D_{(0,0)} \neq (1, 0), D_{(1,1)} \neq (1, 0) \text{ and } D_{(1,0)} = (1, 0)$$

→ $\xi_{Y_{(1,0)}|\mathbf{T}=(00,10,11)}$ is not point-identified, but can be bounded

Define $\mathbf{pf}_{d|Z=z}(y) := P(D = d | Z = z) f_{Y|D=d, Z=z}(y)$.

Proceeding as before, we have the bounds

$$\begin{aligned} \underline{f}^*(y) &:= \max \left\{ 0, \mathbf{pf}_{(1,0)|Z=(1,0)}(y) - \mathbf{pf}_{(1,0)|Z=(1,1)}(y) - \mathbf{pf}_{(1,0)|Z=(0,0)}(y) \right\} \\ &\leq \xi_{Y_{(1,0)}|\mathbf{T}=(00,10,11)}(y) \leq \\ &\quad \mathbf{pf}_{(1,0)|Z=(1,0)}(y) - \mathbf{pf}_{(1,0)|Z=(1,1)}(y) =: \bar{f}^*(y) , \end{aligned}$$

Multi-dim. Treatment example: Trimming Failure

We have $\underline{f}^*(y) \leq \xi_{Y_{(1,0)}|\mathbf{T}=(00,10,11)}(y) \leq \overline{f}^*(y)$

For simplicity, assume that $\mathbb{P}[\mathbf{T} = (00, 10, 11)]$ is identified, and define

$$\bar{p} := \mathbb{P}[\mathbf{T} = (00, 10, 11)] - \int_{\mathcal{Y}} \underline{f}^*(y) \, d\mu(y) .$$

Multi-dim. Treatment example: Trimming Failure

We have $\underline{f}^*(y) \leq \xi_{Y_{(1,0)}|T=(00,10,11)}(y) \leq \overline{f}^*(y)$

For simplicity, assume that $\mathbb{P}[T = (00, 10, 11)]$ is identified, and define

$$\bar{p} := \mathbb{P}[T = (00, 10, 11)] - \int_{\mathcal{Y}} \underline{f}^*(y) \, d\mu(y) .$$

Simple bounds can be constructed as follows:

- (i) Trim $\overline{f}^* - \underline{f}^*$ up to mass $\bar{p}$
- (ii) Add this to the density implied by lower bound, $\underline{f}^*(y)$

Multi-dim. Treatment example: Trimming Failure

We have $\underline{f}^*(y) \leq \xi_{Y_{(1,0)}|T=(00,10,11)}(y) \leq \overline{f}^*(y)$

For simplicity, assume that $\mathbb{P}[T = (00, 10, 11)]$ is identified, and define

$$\bar{p} := \mathbb{P}[T = (00, 10, 11)] - \int_{\mathcal{Y}} \underline{f}^*(y) \, d\mu(y) .$$


Simple bounds can be constructed as follows:

- (i) Trim $\overline{f}^* - \underline{f}^*$ up to mass $\bar{p}$
- (ii) Add this to the density implied by lower bound, $\underline{f}^*(y)$

→ The resulting bounds are valid, but **still not generally sharp**.

Multi-dim. Treatment example: Tighter Bounds due to Pointwise Dependence

Suppose $\mathbb{P}[T = (00, 10, 11)] = 0.25$.



Pointwise bounds on $\xi_{Y_M|T=(00,10,11)}$

Figure 5: Construction of lower bounds for $E[Y_{(1,0)}|T = (00, 10, 11)]$

Multi-dim. Treatment example: Tighter Bounds due to Pointwise Dependence

Suppose $\mathbb{P}[T = (00, 10, 11)] = 0.25$.

Construction of Simple Lower Bound





Figure 5: Construction of lower bounds for $E[Y_{(1,0)} | T = (00, 10, 11)]$

Multi-dim. Treatment example: Tighter Bounds due to Pointwise Dependence


Suppose $\mathbb{P}[T = (00, 10, 11)] = 0.25$.

Construction of Simple Lower Bound



dark blue area = 0.25

Density $f_{Y_{(1,0)}|T=(00,10,11)}$ Implied by
Simple Lower Bound




This density implies the simple lower bound
of 4.865

Figure 5: Construction of lower bounds for $E[Y_{(1,0)}|T = (00, 10, 11)]$

Multi-dim. Treatment example: Tighter Bounds due to Pointwise Dependence


Suppose $\mathbb{P}[T = (00, 10, 11)] = 0.25$.

Construction of Sharp Lower Bound



dark blue area = 0.25

Density $f_{Y_{(1,0)}|T=(00,10,11)}$ Implied by
Sharp Lower Bound



This density implies the sharp lower bound
of 5.106

Figure 5: Construction of lower bounds for $E[Y_{(1,0)}|T = (00, 10, 11)]$

IV Mixture Model: Notation

Enumerate instrument values as $\mathcal{Z} := \{1, \dots, N_Z\}$. For any random variables X_1, X_2 , let $f_{X_1|X_2}$ be the density of $X_1|X_2$.

For $d \in \mathcal{D}$ and $y \in \mathcal{Y}$, define $\mathbf{pf}_d$ as the **observed** vector of weighted densities

$$\mathbf{pf}_d(y) := \begin{bmatrix} P(D = d|Z = 1) f_{Y|D=d,Z=1}(y) \\ \vdots \\ P(D = d|Z = N_Z) f_{Y|D=d,Z=N_Z}(y) \end{bmatrix}$$

IV Mixture Model: Notation

Enumerate instrument values as $\mathcal{Z} := \{1, \dots, N_Z\}$. For any random variables X_1, X_2 , let $f_{X_1|X_2}$ be the density of $X_1|X_2$.

For $d \in \mathcal{D}$ and $y \in \mathcal{Y}$, define $\mathbf{pf}_d$ as the **observed** vector of weighted densities

$$\mathbf{pf}_d(y) := \begin{bmatrix} P(D = d|Z = 1) f_{Y|D=d,Z=1}(y) \\ \vdots \\ P(D = d|Z = N_Z) f_{Y|D=d,Z=N_Z}(y) \end{bmatrix}$$

Define $\boldsymbol{\pi}$ as the **unobserved** vector of response-type probabilities,

$$\boldsymbol{\pi} := \begin{bmatrix} \mathbb{P}(\mathbf{T} = \boldsymbol{\tau}_1) \\ \vdots \\ \mathbb{P}(\mathbf{T} = \boldsymbol{\tau}_{N_T}) \end{bmatrix}$$

IV Mixture Model: Notation

Enumerate instrument values as $\mathcal{Z} := \{1, \dots, N_Z\}$. For any random variables X_1, X_2 , let $f_{X_1|X_2}$ be the density of $X_1|X_2$.

For $d \in \mathcal{D}$ and $y \in \mathcal{Y}$, define $\mathbf{pf}_d$ as the **observed** vector of weighted densities

$$\mathbf{pf}_d(y) := \begin{bmatrix} P(D = d|Z = 1) f_{Y|D=d,Z=1}(y) \\ \vdots \\ P(D = d|Z = N_Z) f_{Y|D=d,Z=N_Z}(y) \end{bmatrix}$$

Define $\boldsymbol{\pi}$ as the **unobserved** vector of response-type probabilities,

$$\boldsymbol{\pi} := \begin{bmatrix} \mathbb{P}(\mathbf{T} = \boldsymbol{\tau}_1) \\ \vdots \\ \mathbb{P}(\mathbf{T} = \boldsymbol{\tau}_{N_T}) \end{bmatrix}$$

and define $\boldsymbol{\xi}_d$ as the **unobserved** vector of weighted response-type densities

$$\boldsymbol{\xi}_d(y) := \begin{bmatrix} \mathbb{P}(\mathbf{T} = \boldsymbol{\tau}_1) f_{Y_d|\mathbf{T}=\boldsymbol{\tau}_1}(y) \\ \vdots \\ \mathbb{P}(\mathbf{T} = \boldsymbol{\tau}_{N_T}) f_{Y_d|\mathbf{T}=\boldsymbol{\tau}_{N_T}}(y) \end{bmatrix}$$

IV Mixture Model: Notation

Finally, define the d -response-matrix as

$$\mathbf{R}_d := \begin{bmatrix} \mathbb{1} \{ \tau_1 (1) = d \} & \dots & \mathbb{1} \{ \tau_{N_T} (1) = d \} \\ \vdots & \ddots & \vdots \\ \mathbb{1} \{ \tau_0 (N_Z) = d \} & \dots & \mathbb{1} \{ \tau_{N_T} (N_Z) = d \} \end{bmatrix},$$

which is the **known** binary matrix that indicates **when each response-type takes treatment d** (as in Heckman and Pinto, 2018).

IV Mixture Model: Notation

Finally, define the d -response-matrix as

$$\mathbf{R}_d := \begin{bmatrix} \mathbb{1}\{\tau_1(1) = d\} & \dots & \mathbb{1}\{\tau_{N_T}(1) = d\} \\ \vdots & \ddots & \vdots \\ \mathbb{1}\{\tau_0(N_Z) = d\} & \dots & \mathbb{1}\{\tau_{N_T}(N_Z) = d\} \end{bmatrix},$$

which is the **known** binary matrix that indicates **when each response-type takes treatment d** (as in Heckman and Pinto, 2018).

Basic Example ($Z \in \{0, 1\}$, $D \in \{0, 1\}$, no restrictions on selection) : In this case, the response types are always-takers (AT), compliers (C), defiers (D), never-takers (NT) , leading to

$$\mathbf{R}_0 = \begin{matrix} & \begin{matrix} \text{(AT)} & \text{(C)} & \text{(D)} & \text{(NT)} \end{matrix} \\ \begin{bmatrix} 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{bmatrix} & \begin{matrix} (Z = 0) \\ (Z = 1) \end{matrix} \end{matrix} \quad \text{and} \quad \mathbf{R}_1 = \begin{matrix} & \begin{matrix} \text{(AT)} & \text{(C)} & \text{(D)} & \text{(NT)} \end{matrix} \\ \begin{bmatrix} 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \end{bmatrix} & \begin{matrix} (Z = 0) \\ (Z = 1) \end{matrix} \end{matrix}$$

IV Mixture Model

Using this notation, we have the following system for $d \in \mathcal{D}$ and $y \in \mathcal{Y}$

$$\mathbf{R}_d \boldsymbol{\xi}_d(y) = \mathbf{p} \mathbf{f}_d(y) \quad \text{and} \quad \boldsymbol{\xi}_d(y) \geq 0 \quad (\star)$$

When $(\star)$ is underdetermined, $\boldsymbol{\xi}_d(y)$ is generally not point-identified.

IV Mixture Model

Using this notation, we have the following system for $d \in \mathcal{D}$ and $y \in \mathcal{Y}$

$$\mathbf{R}_d \boldsymbol{\xi}_d(y) = \mathbf{p} \mathbf{f}_d(y) \quad \text{and} \quad \boldsymbol{\xi}_d(y) \geq 0 \quad (\star)$$

When $(\star)$ is underdetermined, $\boldsymbol{\xi}_d(y)$ is generally not point-identified.

Proposition 1 (due to Heckman and Pinto (2018))

For any $\mathbf{c} \in \mathbb{R}^{|\mathcal{T}|}$,

$\mathbf{c} \in \text{row-span}(\mathbf{R}_d) \implies \mathbf{c}^\top \boldsymbol{\xi}_d$ is point-identified ,

$\mathbf{c} \in \text{row-span} \left(\begin{bmatrix} \mathbf{R}_1 \\ \vdots \\ \mathbf{R}_{N_D} \end{bmatrix} \right) \implies \mathbf{c}^\top \boldsymbol{\pi}$ is point-identified .

→ Equivalent to Theorem T-2 in Heckman and Pinto (2018), where it is stated in terms of the Moore-Penrose generalized inverse of $\mathbf{R}_d$

Informal Outline of Approach

Step 0: Selection model determines set of possible response-types, $\mathcal{T}$

Informal Outline of Approach

Step 0: Selection model determines set of possible response-types, $\mathcal{T}$

Step 1: For each $d \in \mathcal{D}$, $\xi_d(y)$ must be a solution to $(\star)$

Informal Outline of Approach

Step 0: Selection model determines set of possible response-types, $\mathcal{T}$

Step 1: For each $d \in \mathcal{D}$, $\xi_d(y)$ must be a solution to $(\star)$ which imposes

1a. restrictions on π , i.e. distribution of T

1b. restrictions on the conditional distribution of $Y_d|T$

Informal Outline of Approach

Step 0: Selection model determines set of possible response-types, $\mathcal{T}$

Step 1: For each $d \in \mathcal{D}$, $\xi_d(y)$ must be a solution to $(\star)$ which imposes

1a. restrictions on π , i.e. distribution of T

1b. restrictions on the conditional distribution of $Y_d|T$

Step 2: Get identified set for π by combining restrictions from Step 1 over treatments $d \in \mathcal{D}$

Informal Outline of Approach

Step 0: Selection model determines set of possible response-types, $\mathcal{T}$

Step 1: For each $d \in \mathcal{D}$, $\xi_d(y)$ must be a solution to $(\star)$ which imposes

1a. restrictions on π , i.e. distribution of T

1b. restrictions on the conditional distribution of $Y_d|T$

Step 2: Get identified set for π by combining restrictions from Step 1 over treatments $d \in \mathcal{D}$

- Optional: incorporate additional assumptions on treatment selection
- Reject selection model iff this set is empty (sharp specification test)

Informal Outline of Approach

Step 0: Selection model determines set of possible response-types, $\mathcal{T}$

Step 1: For each $d \in \mathcal{D}$, $\xi_d(y)$ must be a solution to $(\star)$ which imposes

1a. restrictions on π , i.e. distribution of T

1b. restrictions on the conditional distribution of $Y_d|T$

Step 2: Get identified set for π by combining restrictions from Step 1 over treatments $d \in \mathcal{D}$

- Optional: incorporate additional assumptions on treatment selection
- Reject selection model iff this set is empty (sharp specification test)

Step 3: Define parameter of interest (e.g., ATE)

- Possible values for this parameter are those consistent with restrictions from Steps 1 and 2

$\xi_d(y)$ must be a solution to $(\star)$

Fix $d \in \mathcal{D}$. For $y \in \mathcal{Y}$, the set of solutions to $(\star)$ define a convex polytope

$\xi_d(y)$ must be a solution to $(\star)$

Fix $d \in \mathcal{D}$. For $y \in \mathcal{Y}$, the set of solutions to $(\star)$ define a convex polytope

$$\mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y)) := \left\{ \mathbf{x} \in \mathbb{R}_{\geq 0}^{|\mathcal{T}|} \mid \mathbf{R}_d \mathbf{x} = \mathbf{p}\mathbf{f}_d(y) \right\}.$$


→ By $(\star)$, we require that $\xi_d(y) \in \mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y))$ for each $y \in \mathcal{Y}$.

$\xi_d(y)$ must be a solution to $(\star)$


Fix $d \in \mathcal{D}$. For $y \in \mathcal{Y}$, the set of solutions to $(\star)$ define a convex polytope

$$\mathcal{B}_d(\mathbf{pf}_d(y)) := \left\{ \mathbf{x} \in \mathbb{R}_{\geq 0}^{|\mathcal{T}|} \mid \mathbf{R}_d \mathbf{x} = \mathbf{pf}_d(y) \right\}.$$


→ By $(\star)$, we require that $\xi_d(y) \in \mathcal{B}_d(\mathbf{pf}_d(y))$ for each $y \in \mathcal{Y}$.



(a) $\tilde{\mathcal{B}}_d(\mathbf{pf}_d(0))$



(b) $\tilde{\mathcal{B}}_d(\mathbf{pf}_d(1.5))$



(c) $\tilde{\mathcal{B}}_d(\mathbf{pf}_d(2))$

Figure 6: Hypothetical $\tilde{\mathcal{B}}_d(\mathbf{pf}_d(\cdot))$ in a case with 2 “degrees of freedom”

$\xi_d(y)$ must be a solution to $(\star)$




Figure 7: Hypothetical graph of the correspondence $y \mapsto \tilde{\mathcal{B}}_d(\mathbf{pf}_d(y))$

$\xi_d(y)$ must be a solution to $(\star)$




Figure 7: Hypothetical graph of the correspondence $y \mapsto \tilde{\mathcal{B}}_d(\mathbf{p}\mathbf{f}_d(y))$

$\xi_d(y)$ must be a solution to $(\star)$




Figure 7: Hypothetical graph of the correspondence $y \mapsto \tilde{\mathcal{B}}_d(\mathbf{pf}_d(y))$

$\xi_d(y)$ must be a solution to $(\star)$




Figure 7: Hypothetical graph of the correspondence $y \mapsto \tilde{\mathcal{B}}_d(\mathbf{pf}_d(y))$

$\xi_d(y)$ must be a solution to $(\star)$

→ If $\mathbf{x}$ is a vector-valued function such that $\mathbf{x}(y) \in \mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y))$ for all $y \in \mathcal{Y}$, then $\mathbf{x}$ is a “candidate” for $\xi_d(y)$.




Figure 7: Hypothetical graph of the correspondence $y \mapsto \tilde{\mathcal{B}}_d(\mathbf{p}\mathbf{f}_d(y))$

Candidates for π


If $\mathbf{x}$ is a candidate for $\xi_d(y)$, then $\int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y)$ is a candidate for π , the vector of response-type probabilities. Precisely,

$$\mathbf{x} = \xi_d \implies \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) = \pi$$


Candidates for π

If $\mathbf{x}$ is a candidate for $\xi_d(y)$, then $\int_{\mathcal{Y}} \mathbf{x}(y) d\mu(y)$ is a candidate for π , the vector of response-type probabilities. Precisely,

$$\mathbf{x} = \xi_d \implies \int_{\mathcal{Y}} \mathbf{x}(y) d\mu(y) = \pi$$



(a) $\mathbf{x}(y) \in \mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y)) \forall y \in \mathcal{Y}$



(b) $\int_{\mathcal{Y}} \mathbf{x}(y) d\mu(y)$

Candidates for π

If $\mathbf{x}$ is a candidate for $\xi_d(y)$, then $\int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y)$ is a candidate for π , the vector of response-type probabilities. Precisely,

$$\mathbf{x} = \xi_d \implies \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) = \pi$$

Define $\Pi_{\mathcal{T}|d}$ as the set of such candidates for π :

$$\begin{aligned} \Pi_{\mathcal{T}|d} &:= \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}_{\geq 0}^{\mathcal{T}} \\ \mathbf{R}_d \mathbf{x}(y) = \mathbf{p} \mathbf{f}_d(y), \, \forall y \in \mathcal{Y} \end{array} \right\} \\ &= \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{\mathcal{T}} \\ \mathbf{x}(y) \in \mathcal{B}_d(\mathbf{p} \mathbf{f}_d(y)), \, \forall y \in \mathcal{Y} \end{array} \right\}. \end{aligned}$$

Candidates for π

If $\mathbf{x}$ is a candidate for $\xi_d(y)$, then $\int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y)$ is a candidate for π , the vector of response-type probabilities. Precisely,

$$\mathbf{x} = \xi_d \implies \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) = \pi$$

Define $\Pi_{\mathcal{T}|d}$ as the set of such candidates for π :


$$\begin{aligned} \Pi_{\mathcal{T}|d} &:= \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}_{\geq 0}^{\mathcal{T}} \\ \mathbf{R}_d \mathbf{x}(y) = \mathbf{p} \mathbf{f}_d(y), \, \forall y \in \mathcal{Y} \end{array} \right\} \\ &= \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{\mathcal{T}} \\ \mathbf{x}(y) \in \mathcal{B}_d(\mathbf{p} \mathbf{f}_d(y)), \, \forall y \in \mathcal{Y} \end{array} \right\}. \end{aligned}$$

→ $\Pi_{\mathcal{T}|d}$ is a convex polytope, and its boundary can be obtained by integrating along the boundary of $\mathcal{B}_d(\mathbf{p} \mathbf{f}_d(y))$


Candidates for π

Define $\Pi_{\mathcal{T}|d}$ as the set of such candidates for π :

$$\Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{\mathcal{T}} \\ \mathbf{x}(y) \in \mathcal{B}_d(\mathbf{pf}_d(y)), \forall y \in \mathcal{Y} \end{array} \right\}.$$



(a) $\mathbf{x}(y) \in \mathcal{B}_d(\mathbf{pf}_d(y)) \forall y \in \mathcal{Y}$




(b) $\int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y)$


Candidates for π

Define $\Pi_{\mathcal{T}|d}$ as the set of such candidates for π :

$$\Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{\mathcal{T}} \\ \mathbf{x}(y) \in \mathcal{B}_d(\mathbf{pf}_d(y)), \forall y \in \mathcal{Y} \end{array} \right\}.$$



(a) $\mathbf{x}(y) \in \mathcal{B}_d(\mathbf{pf}_d(y)) \forall y \in \mathcal{Y}$




(b) $\int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y)$


Candidates for π

Define $\Pi_{\mathcal{T}|d}$ as the set of such candidates for π :

$$\Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{\mathcal{T}} \\ \mathbf{x}(y) \in \mathcal{B}_d(\mathbf{pf}_d(y)), \forall y \in \mathcal{Y} \end{array} \right\}.$$



(a) $\mathbf{x}(y) \in \mathcal{B}_d(\mathbf{pf}_d(y)) \forall y \in \mathcal{Y}$




(b) $\int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y)$


Candidates for π

Define $\Pi_{\mathcal{T}|d}$ as the set of such candidates for π :

$$\Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{\mathcal{T}} \\ \mathbf{x}(y) \in \mathcal{B}_d(\mathbf{pf}_d(y)), \forall y \in \mathcal{Y} \end{array} \right\}.$$



(a) $\mathbf{x}(y) \in \mathcal{B}_d(\mathbf{pf}_d(y)) \forall y \in \mathcal{Y}$




(b) $\int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y)$

Candidates for π

Define $\Pi_{\mathcal{T}|d}$ as the set of such candidates for π :

$$\Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{\mathcal{T}} \\ \mathbf{x}(y) \in \mathcal{B}_d(\mathbf{pf}_d(y)), \forall y \in \mathcal{Y} \end{array} \right\}.$$



Candidates for π

Define $\Pi_{\mathcal{T}|d}$ as the set of such candidates for π :

$$\Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{\mathcal{T}} \\ \mathbf{x}(y) \in \mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y)), \forall y \in \mathcal{Y} \end{array} \right\}.$$

Candidates for π

Define $\Pi_{\mathcal{T}|d}$ as the set of such candidates for π :

$$\Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{\mathcal{T}} \\ \mathbf{x}(y) \in \mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y)), \forall y \in \mathcal{Y} \end{array} \right\}.$$

Comments

- We must have $\pi \in \Pi_{\mathcal{T}|d}$ for every $d \in \mathcal{D}$.

Candidates for π

Define $\Pi_{\mathcal{T}|d}$ as the set of such candidates for π :

$$\Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{\mathcal{T}} \\ \mathbf{x}(y) \in \mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y)), \forall y \in \mathcal{Y} \end{array} \right\}.$$

Comments

- We must have $\pi \in \Pi_{\mathcal{T}|d}$ for every $d \in \mathcal{D}$.
- $\Pi_{\mathcal{T}|d} = \emptyset$ iff $\mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y)) = \emptyset$ for some $y \in \mathcal{Y}$.

Candidates for π

Define $\Pi_{\mathcal{T}|d}$ as the set of such candidates for π :

$$\Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{\mathcal{T}} \\ \mathbf{x}(y) \in \mathcal{B}_d(\mathbf{pf}_d(y)), \forall y \in \mathcal{Y} \end{array} \right\}.$$

Comments

- We must have $\pi \in \Pi_{\mathcal{T}|d}$ for every $d \in \mathcal{D}$.
- $\Pi_{\mathcal{T}|d} = \emptyset$ iff $\mathcal{B}_d(\mathbf{pf}_d(y)) = \emptyset$ for some $y \in \mathcal{Y}$.
- $\Pi_{\mathcal{T}|d}$ is the *Minkowski integral* of the map $y \mapsto \mathcal{B}_d(\mathbf{pf}_d(y))$
 - Continuous analogue to Minkowski sum (Billera and Sturmfels, 1992)

Candidates for π

Define $\Pi_{\mathcal{T}|d}$ as the set of such candidates for π :

$$\Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) \, d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{\mathcal{T}} \\ \mathbf{x}(y) \in \mathcal{B}_d(\mathbf{pf}_d(y)), \forall y \in \mathcal{Y} \end{array} \right\}.$$

Comments

- We must have $\pi \in \Pi_{\mathcal{T}|d}$ for every $d \in \mathcal{D}$.
- $\Pi_{\mathcal{T}|d} = \emptyset$ iff $\mathcal{B}_d(\mathbf{pf}_d(y)) = \emptyset$ for some $y \in \mathcal{Y}$.
- $\Pi_{\mathcal{T}|d}$ is the *Minkowski integral* of the map $y \mapsto \mathcal{B}_d(\mathbf{pf}_d(y))$
 - Continuous analogue to Minkowski sum (Billera and Sturmfels, 1992)
 - Close connection to *Aumann expectation* from theory of random sets (Artstein and Vitale, 1975): both are adaptations of Aumann (1965)'s integral for set-valued measures

Candidates for π : Support Function

Every non-empty closed convex set $C \subseteq \mathbb{R}^n$ is uniquely determined by its **support function** $h_C : \mathbb{R}^n \rightarrow \mathbb{R}$, defined as

$$h_C(\mathbf{c}) := \sup_{\mathbf{x} \in C} \mathbf{c}^\top \mathbf{x} .$$

Define the shorthand $\mathbf{h}_d(\mathbf{c}; \mathbf{p}\mathbf{f}_d(y)) := h_{\mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y))}(\mathbf{c})$.

Candidates for π : Support Function

Every non-empty closed convex set $C \subseteq \mathbb{R}^n$ is uniquely determined by its **support function** $h_C : \mathbb{R}^n \rightarrow \mathbb{R}$, defined as

$$h_C(\mathbf{c}) := \sup_{\mathbf{x} \in C} \mathbf{c}^\top \mathbf{x} .$$

Define the shorthand $h_d(\mathbf{c}; \mathbf{p}\mathbf{f}_d(y)) := h_{\mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y))}(\mathbf{c})$.

Lemma 1

Suppose $\mathcal{T} \subseteq \mathcal{D}^{\mathcal{Z}}$ is such that $\mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y)) \neq \emptyset$ for all $y \in \mathcal{Y}$. Then $\Pi_{\mathcal{T}|d}$ is a non-empty convex set such that, for all $\mathbf{c} \in \mathbb{R}^{\mathcal{T}}$

$$h_{\Pi_{\mathcal{T}|d}}(\mathbf{c}) = \int_{\mathcal{Y}} h_d(\mathbf{c}; \mathbf{p}\mathbf{f}_d(y)) \, d\mu(y) .$$

Further, $\Pi_{\mathcal{T}|d}$ is a convex polytope.

Candidates for π : Support Function

Corollary 1

Suppose $\mathcal{T} \subseteq \mathcal{D}^{\mathcal{Z}}$ is such that $\mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y)) \neq \emptyset$ for all $y \in \mathcal{Y}$.

Given matrix $\mathbf{A} = \begin{bmatrix} \mathbf{a}_1^\top \\ \vdots \\ \mathbf{a}_n^\top \end{bmatrix} \in \mathbb{R}^{n \times \mathcal{T}}$, such that, for some $\mathbf{b} : \mathcal{Y} \rightarrow \mathbb{R}^n$,

$$\mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y)) = \{\mathbf{x} \in \mathbb{R}^{\mathcal{T}} \mid \mathbf{A}\mathbf{x} \leq \mathbf{b}(y)\} \quad \forall y \in \mathcal{Y},$$

$\Pi_{\mathcal{T}|d}$ can be represented as

$$\Pi_{\mathcal{T}|d} = \left\{ \mathbf{x} \in \mathbb{R}^{\mathcal{T}} \mid \mathbf{A}\mathbf{x} \leq \begin{bmatrix} \int_{\mathcal{Y}} h_d(\mathbf{a}_1; \mathbf{p}\mathbf{f}_d(y)) d\mu(y) \\ \vdots \\ \int_{\mathcal{Y}} h_d(\mathbf{a}_n; \mathbf{p}\mathbf{f}_d(y)) d\mu(y) \end{bmatrix} \right\}.$$

Identified Set for Response-Type Probabilities

Given selection model $\mathbb{S}$, denote by $\mathcal{T}(\mathbb{S})$ the set of response-types not *a priori* eliminated by $\mathbb{S}$, i.e

$$\mathcal{T}(\mathbb{S}) := \left\{ \tau \in \mathcal{D}^Z \mid \exists x \in \mathcal{F}_T(\mathbb{S}), x_{(\tau)} > 0 \right\} .$$

Identified Set for Response-Type Probabilities

Given selection model $\mathbb{S}$, denote by $\mathcal{T}(\mathbb{S})$ the set of response-types *not a priori* eliminated by $\mathbb{S}$, i.e

$$\mathcal{T}(\mathbb{S}) := \left\{ \tau \in \mathcal{D}^Z \mid \exists x \in \mathcal{F}_T(\mathbb{S}), x_{(\tau)} > 0 \right\} .$$

Denote by $\Pi_{\mathbb{S}}$ the *identified set for π* under $\mathbb{S}$, i.e. $\Pi_{\mathbb{S}} := \{ \pi_Q \mid Q \in \mathcal{Q}_{\mathbb{S}} \}$

Theorem 2 (Sharp Testable Implications of Selection Models)

For any treatment selection model $\mathbb{S}$,

$$\Pi_{\mathbb{S}} = \left(\bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}(\mathbb{S})|d} \right) \cap \mathcal{F}_T(\mathbb{S}) .$$

In particular, $\mathcal{Q}_{\mathbb{S}} \neq \emptyset$ if and only if $\left(\bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}(\mathbb{S})|d} \right) \cap \mathcal{F}_T(\mathbb{S}) \neq \emptyset$.

Identified Set for Response-Type Probabilities

Given selection model $\mathbb{S}$, denote by $\mathcal{T}(\mathbb{S})$ the set of response-types *not a priori* eliminated by $\mathbb{S}$, i.e

$$\mathcal{T}(\mathbb{S}) := \left\{ \tau \in \mathcal{D}^Z \mid \exists x \in \mathcal{F}_T(\mathbb{S}), x_{(\tau)} > 0 \right\} .$$

Denote by $\Pi_{\mathbb{S}}$ the *identified set for π* under $\mathbb{S}$, i.e. $\Pi_{\mathbb{S}} := \{ \pi_Q \mid Q \in \mathcal{Q}_{\mathbb{S}} \}$

Theorem 2 (Sharp Testable Implications of Selection Models)

For any treatment selection model $\mathbb{S}$,

$$\Pi_{\mathbb{S}} = \left(\bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}(\mathbb{S})|d} \right) \cap \mathcal{F}_T(\mathbb{S}) .$$

In particular, $\mathcal{Q}_{\mathbb{S}} \neq \emptyset$ if and only if $\left(\bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}(\mathbb{S})|d} \right) \cap \mathcal{F}_T(\mathbb{S}) \neq \emptyset$.

Identified Set for Response-Type Probabilities

Given selection model $\mathbb{S}$, denote by $\mathcal{T}(\mathbb{S})$ the set of response-types *not a priori* eliminated by $\mathbb{S}$, i.e

$$\mathcal{T}(\mathbb{S}) := \left\{ \tau \in \mathcal{D}^Z \mid \exists x \in \mathcal{F}_T(\mathbb{S}), x_{(\tau)} > 0 \right\} .$$

Denote by $\Pi_{\mathbb{S}}$ the *identified set for π* under $\mathbb{S}$, i.e. $\Pi_{\mathbb{S}} := \{ \pi_Q \mid Q \in \mathcal{Q}_{\mathbb{S}} \}$

Theorem 2 (Sharp Testable Implications of Selection Models)

For any treatment selection model $\mathbb{S}$,

$$\Pi_{\mathbb{S}} = \left(\bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}(\mathbb{S})|d} \right) \cap \mathcal{F}_T(\mathbb{S}) .$$

In particular, $\mathcal{Q}_{\mathbb{S}} \neq \emptyset$ if and only if $\left(\bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}(\mathbb{S})|d} \right) \cap \mathcal{F}_T(\mathbb{S}) \neq \emptyset$.

Identified Set for Response-Type Probabilities

Given selection model $\mathbb{S}$, denote by $\mathcal{T}(\mathbb{S})$ the set of response-types *not a priori* eliminated by $\mathbb{S}$, i.e

$$\mathcal{T}(\mathbb{S}) := \left\{ \tau \in \mathcal{D}^Z \mid \exists x \in \mathcal{F}_T(\mathbb{S}), x_{(\tau)} > 0 \right\} .$$

Denote by $\Pi_{\mathbb{S}}$ the *identified set for π* under $\mathbb{S}$, i.e. $\Pi_{\mathbb{S}} := \{ \pi_Q \mid Q \in \mathcal{Q}_{\mathbb{S}} \}$


Theorem 2 (Sharp Testable Implications of Selection Models)

For any treatment selection model $\mathbb{S}$,

$$\Pi_{\mathbb{S}} = \left(\bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}(\mathbb{S})|d} \right) \cap \mathcal{F}_T(\mathbb{S}) .$$


In particular, $\mathcal{Q}_{\mathbb{S}} \neq \emptyset$ if and only if $\left(\bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}(\mathbb{S})|d} \right) \cap \mathcal{F}_T(\mathbb{S}) \neq \emptyset$.

(Stylized) Example with 3 Treatments




(a) Hypothetical $\Pi_{\mathcal{T}|1}$

(Stylized) Example with 3 Treatments




(a) Hypothetical $\Pi_{\mathcal{T}|1}, \Pi_{\mathcal{T}|2}$

(Stylized) Example with 3 Treatments




(a) Hypothetical $\Pi_{\mathcal{T}|1}, \Pi_{\mathcal{T}|2}, \Pi_{\mathcal{T}|3}$

(Stylized) Example with 3 Treatments



(a) Hypothetical $\Pi_{\mathcal{T}} := \bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}|d}$

(Stylized) Example with 3 Treatments



(a) Hypothetical $\Pi_{\mathcal{T}} := \bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}|d}$

Comments

Theorem 2 suggests joint test of $\mathbb{S}$ and basic model

- Assume potential outcomes exist, impose no further restrictions in Theorem 2 $\implies$ sharp test for random assignment of Z

Comments

Theorem 2 suggests joint test of $\mathbb{S}$ and basic model

- Assume potential outcomes exist, impose no further restrictions in Theorem 2 $\implies$ sharp test for random assignment of Z

Suppose $\mathcal{F}_T(\mathbb{S})$ is a convex polytope.

Comments

Theorem 2 suggests joint test of $\mathbb{S}$ and basic model

- Assume potential outcomes exist, impose no further restrictions in Theorem 2 $\implies$ sharp test for random assignment of Z

Suppose $\mathcal{F}_{\mathcal{T}}(\mathbb{S})$ is a convex polytope. Then,

- Theorem 2 requires (i) $\Pi_{\mathcal{T}|d} \neq \emptyset$ and (ii) $\mathcal{F}_{\mathcal{T}}(\mathbb{S}) \cap \left(\bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}|d}\right) \neq \emptyset$
 - (i) is automatically satisfied in some cases; e.g., $\mathcal{T} = \mathcal{D}^{|Z|}$
 - (ii) is equivalent to testing feasibility of a linear program (LP)

Comments

Theorem 2 suggests joint test of $\mathbb{S}$ and basic model

- Assume potential outcomes exist, impose no further restrictions in Theorem 2 $\implies$ sharp test for random assignment of Z

Suppose $\mathcal{F}_{\mathcal{T}}(\mathbb{S})$ is a convex polytope. Then,

- Theorem 2 requires (i) $\Pi_{\mathcal{T}|d} \neq \emptyset$ and (ii) $\mathcal{F}_{\mathcal{T}}(\mathbb{S}) \cap \left(\bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}|d}\right) \neq \emptyset$
 - (i) is automatically satisfied in some cases; e.g., $\mathcal{T} = \mathcal{D}^{|Z|}$
 - (ii) is equivalent to testing feasibility of a linear program (LP)
- $\Pi_{\mathcal{T}|d} \neq \emptyset \iff \mathcal{B}_d(\mathbf{pf}_d(y)) \neq \emptyset, \forall y \iff \mathbf{pf}_d(y) \in \text{cone}(\mathbf{R}_d), \forall y$
 - $\text{cone}(\mathbf{R}_d)$ is the polyhedral cone generated by columns of $\mathbf{R}_d$

Comments

Theorem 2 suggests joint test of $\mathbb{S}$ and basic model

- Assume potential outcomes exist, impose no further restrictions in Theorem 2 $\implies$ sharp test for random assignment of Z

Suppose $\mathcal{F}_T(\mathbb{S})$ is a convex polytope. Then,

- Theorem 2 requires (i) $\Pi_{\mathcal{T}|d} \neq \emptyset$ and (ii) $\mathcal{F}_T(\mathbb{S}) \cap \left(\bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}|d}\right) \neq \emptyset$
 - (i) is automatically satisfied in some cases; e.g., $\mathcal{T} = \mathcal{D}^{|Z|}$
 - (ii) is equivalent to testing feasibility of a linear program (LP)
- $\Pi_{\mathcal{T}|d} \neq \emptyset \iff \mathcal{B}_d(\mathbf{pf}_d(y)) \neq \emptyset, \forall y \iff \mathbf{pf}_d(y) \in \text{cone}(\mathbf{R}_d), \forall y$
 - $\text{cone}(\mathbf{R}_d)$ is the polyhedral cone generated by columns of $\mathbf{R}_d$
- Can find matrix $\mathbf{H}$ so that, $\forall y, \mathbf{pf}_d(y) \in \text{cone}(\mathbf{R}_d) \iff \mathbf{H}\mathbf{pf}_d(y) \geq 0$

Comments

Theorem 2 suggests **joint test of $\mathbb{S}$ and basic model**

- Assume potential outcomes exist, impose no further restrictions in Theorem 2 $\implies$ **sharp test for random assignment of Z**

Suppose $\mathcal{F}_T(\mathbb{S})$ is a **convex polytope**. Then,

- Theorem 2 requires (i) $\Pi_{\mathcal{T}|d} \neq \emptyset$ and (ii) $\mathcal{F}_T(\mathbb{S}) \cap \left(\bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}|d}\right) \neq \emptyset$
 - (i) is automatically satisfied in some cases; e.g., $\mathcal{T} = \mathcal{D}^{|Z|}$
 - (ii) is equivalent to testing feasibility of a linear program (LP)
- $\Pi_{\mathcal{T}|d} \neq \emptyset \iff \mathcal{B}_d(\mathbf{pf}_d(y)) \neq \emptyset, \forall y \iff \mathbf{pf}_d(y) \in \text{cone}(\mathbf{R}_d), \forall y$
 - $\text{cone}(\mathbf{R}_d)$ is the polyhedral cone generated by columns of $\mathbf{R}_d$
- Can find matrix $\mathbf{H}$ so that, $\forall y, \mathbf{pf}_d(y) \in \text{cone}(\mathbf{R}_d) \iff \mathbf{H}\mathbf{pf}_d(y) \geq 0$
 - $\mathbf{H}$ can be found by applying a facet-enumeration algorithm to $\mathbf{R}_d$
 - Then, can jointly test (i-ii) as **conditional moment inequalities**

Comments

Theorem 2 suggests **joint test of $\mathbb{S}$ and basic model**

- Assume potential outcomes exist, impose no further restrictions in Theorem 2 $\implies$ **sharp test for random assignment of Z**

Suppose $\mathcal{F}_T(\mathbb{S})$ is a **convex polytope**. Then,

- Theorem 2 requires (i) $\Pi_{\mathcal{T}|d} \neq \emptyset$ and (ii) $\mathcal{F}_T(\mathbb{S}) \cap \left(\bigcap_{d \in \mathcal{D}} \Pi_{\mathcal{T}|d}\right) \neq \emptyset$
 - (i) is automatically satisfied in some cases; e.g., $\mathcal{T} = \mathcal{D}^{|Z|}$
 - (ii) is equivalent to testing feasibility of a linear program (LP)
- $\Pi_{\mathcal{T}|d} \neq \emptyset \iff \mathcal{B}_d(\mathbf{pf}_d(y)) \neq \emptyset, \forall y \iff \mathbf{pf}_d(y) \in \text{cone}(\mathbf{R}_d), \forall y$
 - $\text{cone}(\mathbf{R}_d)$ is the polyhedral cone generated by columns of $\mathbf{R}_d$
- Can find matrix $\mathbf{H}$ so that, $\forall y, \mathbf{pf}_d(y) \in \text{cone}(\mathbf{R}_d) \iff \mathbf{H}\mathbf{pf}_d(y) \geq 0$
 - $\mathbf{H}$ can be found by applying a facet-enumeration algorithm to $\mathbf{R}_d$
 - Then, can jointly test (i-ii) as **conditional moment inequalities**
- Or, use Fang and Seo (2021) to test (i) and Fang et al. (2023) to test (ii)

What about outcomes?

What about outcomes?

$$\text{Recall } \Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{|\mathbf{T}|} \\ \mathbf{R}_d \mathbf{x} = \mathbf{p}\mathbf{f}_d(y) \quad \forall y \in \mathcal{Y} \end{array} \right\}$$

What about outcomes?

$$\text{Recall } \Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{|\mathbf{T}|} \\ \mathbf{R}_d \mathbf{x} = \mathbf{p}\mathbf{f}_d(y) \quad \forall y \in \mathcal{Y} \end{array} \right\}$$

Can define a similar set of “candidates”:

$$\Theta_{\mathcal{T}|d} := \left\{ \left(\int_{\mathcal{Y}} y \mathbf{x}(y) d\mu(y), \int_{\mathcal{Y}} \mathbf{x}(y) d\mu(y) \right) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{|\mathbf{T}|} \\ \mathbf{R}_d \mathbf{x} = \mathbf{p}\mathbf{f}_d(y) \quad \forall y \in \mathcal{Y} \end{array} \right\}$$

What about outcomes?

$$\text{Recall } \Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{|\mathbf{T}|} \\ \mathbf{R}_d \mathbf{x} = \mathbf{p} \mathbf{f}_d(y) \quad \forall y \in \mathcal{Y} \end{array} \right\}$$

Can define a similar set of “candidates”:

$$\Theta_{\mathcal{T}|d} := \left\{ \left(\int_{\mathcal{Y}} y \mathbf{x}(y) d\mu(y), \int_{\mathcal{Y}} \mathbf{x}(y) d\mu(y) \right) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{|\mathbf{T}|} \\ \mathbf{R}_d \mathbf{x} = \mathbf{p} \mathbf{f}_d(y) \quad \forall y \in \mathcal{Y} \end{array} \right\}$$

$$\rightarrow \text{candidates for } (\boldsymbol{\theta}_d, \boldsymbol{\pi}), \text{ where } \boldsymbol{\theta}_d := \begin{bmatrix} \mathbb{P}(\mathbf{T} = \tau_1) \mathbb{E}[Y_d \mid \mathbf{T} = \tau_1] \\ \vdots \\ \mathbb{P}(\mathbf{T} = \tau_{N_T}) \mathbb{E}[Y_d \mid \mathbf{T} = \tau_{N_T}] \end{bmatrix}$$

What about outcomes?

$$\text{Recall } \Pi_{\mathcal{T}|d} := \left\{ \int_{\mathcal{Y}} \mathbf{x}(y) d\mu(y) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{|\mathbf{T}|} \\ \mathbf{R}_d \mathbf{x} = \mathbf{p}\mathbf{f}_d(y) \quad \forall y \in \mathcal{Y} \end{array} \right\}$$

Can define a similar set of “candidates”:

$$\Theta_{\mathcal{T}|d} := \left\{ \left(\int_{\mathcal{Y}} y \mathbf{x}(y) d\mu(y), \int_{\mathcal{Y}} \mathbf{x}(y) d\mu(y) \right) \mid \begin{array}{l} \text{measurable } \mathbf{x} : \mathcal{Y} \rightarrow \mathbb{R}^{|\mathbf{T}|} \\ \mathbf{R}_d \mathbf{x} = \mathbf{p}\mathbf{f}_d(y) \quad \forall y \in \mathcal{Y} \end{array} \right\}$$

$$\rightarrow \text{candidates for } (\boldsymbol{\theta}_d, \boldsymbol{\pi}), \text{ where } \boldsymbol{\theta}_d := \begin{bmatrix} \mathbb{P}(\mathbf{T} = \tau_1) \mathbb{E}[Y_d \mid \mathbf{T} = \tau_1] \\ \vdots \\ \mathbb{P}(\mathbf{T} = \tau_{N_T}) \mathbb{E}[Y_d \mid \mathbf{T} = \tau_{N_T}] \end{bmatrix}$$

Lemma 2

Suppose $\mathcal{B}_d(\mathbf{p}\mathbf{f}_d(y)) \neq \emptyset, \forall y \in \mathcal{Y}$. Then, for all $(\mathbf{c}_1, \mathbf{c}_2) \in \mathbb{R}^{\mathcal{T}} \times \mathbb{R}^{\mathcal{T}}$,

$$h_{\Theta_{\mathcal{T}|d}}(\mathbf{c}_1, \mathbf{c}_2) = \int_{\mathcal{Y}} h_d(y \mathbf{c}_1 + \mathbf{c}_2; \mathbf{p}\mathbf{f}_d(y)) d\mu(y).$$

Conditional Mean Potential Outcomes

Target Parameter: $\mathbb{E} [Y_d \mid \boldsymbol{T} \in \mathcal{T}_\dagger]$, for some subset $\mathcal{T}_\dagger \subseteq \mathcal{T}$.

Conditional Mean Potential Outcomes

Target Parameter: $\mathbb{E} [Y_d \mid \mathbf{T} \in \mathcal{T}_\dagger]$, for some subset $\mathcal{T}_\dagger \subseteq \mathcal{T}$.

Define $\mathbf{c}_\dagger := (\mathbb{1} \{ \boldsymbol{\tau} \in \mathcal{T}_\dagger \} : \boldsymbol{\tau} \in \mathcal{T}(\mathbb{S}))$ and note that

$$\mathbb{E} [Y_d \mid \mathbf{T} \in \mathcal{T}_\dagger] = \frac{\sum_{\boldsymbol{\tau} \in \mathcal{T}_\dagger} \mathbb{P}(\mathbf{T} = \boldsymbol{\tau}) \mathbb{E} [Y_d \mid \mathbf{T} = \boldsymbol{\tau}]}{\sum_{\boldsymbol{\tau} \in \mathcal{T}_\dagger} \mathbb{P}(\mathbf{T} = \boldsymbol{\tau})} = \frac{\mathbf{c}_\dagger^\top \boldsymbol{\theta}_d}{\mathbf{c}_\dagger^\top \boldsymbol{\pi}}$$

Conditional Mean Potential Outcomes

Target Parameter: $\mathbb{E} [Y_d \mid \mathbf{T} \in \mathcal{T}_\dagger]$, for some subset $\mathcal{T}_\dagger \subseteq \mathcal{T}$.

Define $\mathbf{c}_\dagger := (\mathbb{1} \{ \tau \in \mathcal{T}_\dagger \} : \tau \in \mathcal{T}(\mathbb{S}))$ and note that

$$\mathbb{E} [Y_d \mid \mathbf{T} \in \mathcal{T}_\dagger] = \frac{\sum_{\tau \in \mathcal{T}_\dagger} \mathbb{P}(\mathbf{T} = \tau) \mathbb{E} [Y_d \mid \mathbf{T} = \tau]}{\sum_{\tau \in \mathcal{T}_\dagger} \mathbb{P}(\mathbf{T} = \tau)} = \frac{\mathbf{c}_\dagger^\top \boldsymbol{\theta}_d}{\mathbf{c}_\dagger^\top \boldsymbol{\pi}}$$

→ The identified set for $\mathbb{E} [\kappa(Y_d) \mid \mathbf{T} \in \mathcal{T}_\dagger]$ under $\mathbb{S}$ is given by

$$\left\{ \frac{\mathbf{c}_\dagger^\top \boldsymbol{\theta}_d}{\mathbf{c}_\dagger^\top \boldsymbol{\pi}} \mid \begin{array}{l} (\boldsymbol{\theta}_d, \boldsymbol{\pi}) \in \Theta_{\mathcal{T}(\mathbb{S})|d} , \\ \boldsymbol{\pi} \in \Pi_{\mathbb{S}} \end{array} \right\}$$

Conditional Mean Potential Outcomes

Target Parameter: $\mathbb{E} [Y_d \mid \mathbf{T} \in \mathcal{T}_\dagger]$, for some subset $\mathcal{T}_\dagger \subseteq \mathcal{T}$.

Define $\mathbf{c}_\dagger := (\mathbb{1} \{ \tau \in \mathcal{T}_\dagger \} : \tau \in \mathcal{T}(\mathbb{S}))$ and note that

$$\mathbb{E} [Y_d \mid \mathbf{T} \in \mathcal{T}_\dagger] = \frac{\sum_{\tau \in \mathcal{T}_\dagger} \mathbb{P}(\mathbf{T} = \tau) \mathbb{E} [Y_d \mid \mathbf{T} = \tau]}{\sum_{\tau \in \mathcal{T}_\dagger} \mathbb{P}(\mathbf{T} = \tau)} = \frac{\mathbf{c}_\dagger^\top \boldsymbol{\theta}_d}{\mathbf{c}_\dagger^\top \boldsymbol{\pi}}$$

→ The identified set for $\mathbb{E} [\kappa(Y_d) \mid \mathbf{T} \in \mathcal{T}_\dagger]$ under $\mathbb{S}$ is given by

$$\left\{ \frac{\mathbf{c}_\dagger^\top \boldsymbol{\theta}_d}{\mathbf{c}_\dagger^\top \boldsymbol{\pi}} \mid \begin{array}{l} (\boldsymbol{\theta}_d, \boldsymbol{\pi}) \in \Theta_{\mathcal{T}(\mathbb{S})|d} , \\ \boldsymbol{\pi} \in \Pi_{\mathbb{S}} \end{array} \right\} = \left\{ \gamma \in \mathbb{R} \mid \begin{array}{l} (\boldsymbol{\theta}_d, \boldsymbol{\pi}) \in \Theta_{\mathcal{T}(\mathbb{S})|d} , \\ \boldsymbol{\pi} \in \Pi_{\mathbb{S}}, \mathbf{c}_\dagger^\top \boldsymbol{\theta}_d = \gamma \mathbf{c}_\dagger^\top \boldsymbol{\pi} \end{array} \right\}$$

Conditional Mean Potential Outcomes

Target Parameter: $\mathbb{E} [Y_d \mid \mathbf{T} \in \mathcal{T}_\dagger]$, for some subset $\mathcal{T}_\dagger \subseteq \mathcal{T}$.

Define $\mathbf{c}_\dagger := (\mathbb{1} \{ \tau \in \mathcal{T}_\dagger \} : \tau \in \mathcal{T}(\mathbb{S}))$ and note that

$$\mathbb{E} [Y_d \mid \mathbf{T} \in \mathcal{T}_\dagger] = \frac{\sum_{\tau \in \mathcal{T}_\dagger} \mathbb{P}(\mathbf{T} = \tau) \mathbb{E} [Y_d \mid \mathbf{T} = \tau]}{\sum_{\tau \in \mathcal{T}_\dagger} \mathbb{P}(\mathbf{T} = \tau)} = \frac{\mathbf{c}_\dagger^\top \boldsymbol{\theta}_d}{\mathbf{c}_\dagger^\top \boldsymbol{\pi}}$$

→ The **identified set** for $\mathbb{E} [\kappa(Y_d) \mid \mathbf{T} \in \mathcal{T}_\dagger]$ under $\mathbb{S}$ is given by

$$\left\{ \frac{\mathbf{c}_\dagger^\top \boldsymbol{\theta}_d}{\mathbf{c}_\dagger^\top \boldsymbol{\pi}} \mid \begin{array}{l} (\boldsymbol{\theta}_d, \boldsymbol{\pi}) \in \Theta_{\mathcal{T}(\mathbb{S})|d}, \\ \boldsymbol{\pi} \in \Pi_{\mathbb{S}} \end{array} \right\} = \left\{ \gamma \in \mathbb{R} \mid \begin{array}{l} (\boldsymbol{\theta}_d, \boldsymbol{\pi}) \in \Theta_{\mathcal{T}(\mathbb{S})|d}, \\ \boldsymbol{\pi} \in \Pi_{\mathbb{S}}, \mathbf{c}_\dagger^\top \boldsymbol{\theta}_d = \gamma \mathbf{c}_\dagger^\top \boldsymbol{\pi} \end{array} \right\}$$

→ for each γ , check if intersection of three **closed convex sets** is empty

Conditional Mean Potential Outcomes

Target Parameter: $\mathbb{E} [Y_d \mid \mathbf{T} \in \mathcal{T}_\dagger]$, for some subset $\mathcal{T}_\dagger \subseteq \mathcal{T}$.

Define $\mathbf{c}_\dagger := (\mathbb{1} \{ \tau \in \mathcal{T}_\dagger \} : \tau \in \mathcal{T}(\mathbb{S}))$ and note that

$$\mathbb{E} [Y_d \mid \mathbf{T} \in \mathcal{T}_\dagger] = \frac{\sum_{\tau \in \mathcal{T}_\dagger} \mathbb{P}(\mathbf{T} = \tau) \mathbb{E} [Y_d \mid \mathbf{T} = \tau]}{\sum_{\tau \in \mathcal{T}_\dagger} \mathbb{P}(\mathbf{T} = \tau)} = \frac{\mathbf{c}_\dagger^\top \boldsymbol{\theta}_d}{\mathbf{c}_\dagger^\top \boldsymbol{\pi}}$$

→ The **identified set** for $\mathbb{E} [\kappa(Y_d) \mid \mathbf{T} \in \mathcal{T}_\dagger]$ under $\mathbb{S}$ is given by

$$\left\{ \frac{\mathbf{c}_\dagger^\top \boldsymbol{\theta}_d}{\mathbf{c}_\dagger^\top \boldsymbol{\pi}} \mid \begin{array}{l} (\boldsymbol{\theta}_d, \boldsymbol{\pi}) \in \Theta_{\mathcal{T}(\mathbb{S})|d}, \\ \boldsymbol{\pi} \in \Pi_{\mathbb{S}} \end{array} \right\} = \left\{ \gamma \in \mathbb{R} \mid \begin{array}{l} (\boldsymbol{\theta}_d, \boldsymbol{\pi}) \in \Theta_{\mathcal{T}(\mathbb{S})|d}, \\ \boldsymbol{\pi} \in \Pi_{\mathbb{S}}, \mathbf{c}_\dagger^\top \boldsymbol{\theta}_d = \gamma \mathbf{c}_\dagger^\top \boldsymbol{\pi} \end{array} \right\}$$

→ for each γ , check if intersection of three **closed convex sets** is empty

→ (under some conditions) can use **hyperplane separation theorem**

Mean Outcomes: Inclusion Test

Theorem 3

Suppose $\mathcal{Q}_S \neq \emptyset$ and $\mathcal{F}_T(S)$ is closed and convex.

Mean Outcomes: Inclusion Test

Theorem 3

Suppose $\mathcal{Q}_{\mathbb{S}} \neq \emptyset$ and $\mathcal{F}_{\mathcal{T}}(\mathbb{S})$ is closed and convex.

Let $\mathcal{T}_{\dagger} \subseteq \mathcal{T}(\mathbb{S})$ and $d \in \mathcal{D}$ be given, such that

$$\mathcal{T}_{\dagger} \subseteq \{\tau \in \mathcal{T}(\mathbb{S}) \mid \tau^{-1}(d) \neq \emptyset\} .$$

Mean Outcomes: Inclusion Test

Theorem 3

Suppose $\mathcal{Q}_{\mathbb{S}} \neq \emptyset$ and $\mathcal{F}_{\mathcal{T}}(\mathbb{S})$ is closed and convex.

Let $\mathcal{T}_{\dagger} \subseteq \mathcal{T}(\mathbb{S})$ and $d \in \mathcal{D}$ be given, such that

$$\mathcal{T}_{\dagger} \subseteq \{\boldsymbol{\tau} \in \mathcal{T}(\mathbb{S}) \mid \boldsymbol{\tau}^{-1}(d) \neq \emptyset\} .$$

Then, $\{\mathbb{E}_Q[Y_d \mid \boldsymbol{T} \in \mathcal{T}_{\dagger}] \mid Q \in \mathcal{Q}_{\mathbb{S}}\}$ is equivalent to

$$\left\{ \gamma \in \mathbb{R} \mid \inf_{\boldsymbol{u}, \boldsymbol{v} \in \mathbb{R}^{\mathcal{T}}} \left(\int_{\mathcal{Y}} \boldsymbol{h}_{d(y)}((y - \gamma)(\boldsymbol{c}_{\dagger} \odot \boldsymbol{v}) + \boldsymbol{u}) d\mu(y) + \boldsymbol{h}_{\Pi_{\mathbb{S}}}(-\boldsymbol{u}) \right) \geq 0 \right\}$$

where $\odot$ denotes the coordinate-wise product of two vectors (Hadamard product).

Mean Outcomes: Inclusion Test

Theorem 3

Suppose $\mathcal{Q}_{\mathbb{S}} \neq \emptyset$ and $\mathcal{F}_{\mathcal{T}}(\mathbb{S})$ is closed and convex.

Let $\mathcal{T}_{\dagger} \subseteq \mathcal{T}(\mathbb{S})$ and $d \in \mathcal{D}$ be given, such that

$$\mathcal{T}_{\dagger} \subseteq \{\boldsymbol{\tau} \in \mathcal{T}(\mathbb{S}) \mid \boldsymbol{\tau}^{-1}(d) \neq \emptyset\} .$$

Then, $\{\mathbb{E}_Q[Y_d \mid \boldsymbol{T} \in \mathcal{T}_{\dagger}] \mid Q \in \mathcal{Q}_{\mathbb{S}}\}$ is equivalent to

$$\left\{ \gamma \in \mathbb{R} \mid \inf_{\boldsymbol{u}, \boldsymbol{v} \in \mathbb{R}^{\mathcal{T}}} \left(\int_{\mathcal{Y}} \boldsymbol{h}_{d(y)}((y - \gamma)(\boldsymbol{c}_{\dagger} \odot \boldsymbol{v}) + \boldsymbol{u}) d\mu(y) + \boldsymbol{h}_{\Pi_{\mathbb{S}}}(-\boldsymbol{u}) \right) \geq 0 \right\}$$

where $\odot$ denotes the coordinate-wise product of two vectors (Hadamard product).

→ Need to solve one unconstrained convex program per γ

Theorem 4 (Sharp Bounds on T -Conditional Y_d -Means)

Suppose $\mathbb{S}$, d , and $\mathcal{T}_\dagger$ are as in Theorem 3. Define

$$\underline{p}_\dagger := \inf_{\mathbf{x} \in \Pi_{\mathbb{S}}} \mathbf{c}_\dagger^\top \mathbf{x} \quad \text{and} \quad \bar{p}_\dagger := \sup_{\mathbf{x} \in \Pi_{\mathbb{S}}} \mathbf{c}_\dagger^\top \mathbf{x} .$$

Theorem 4 (Sharp Bounds on T -Conditional Y_d -Means)

Suppose $\mathbb{S}$, d , and $\mathcal{T}_\dagger$ are as in Theorem 3. Define

$$\underline{p}_\dagger := \inf_{\mathbf{x} \in \Pi_{\mathbb{S}}} \mathbf{c}_\dagger^\top \mathbf{x} \text{ and } \bar{p}_\dagger := \sup_{\mathbf{x} \in \Pi_{\mathbb{S}}} \mathbf{c}_\dagger^\top \mathbf{x} .$$

For each $p \in [\underline{p}_\dagger, \bar{p}_\dagger]$, $\mathbf{v} \in \mathbb{R}^{\mathcal{T}(\mathbb{S})}$, and $\lambda \in \mathbb{R}$, define

$$L_d(p; \mathbf{v}, \lambda) := - \int_{\mathcal{Y}} \mathbf{h}_{d(y)}((-y + \lambda) \mathbf{c}_\dagger + \mathbf{v}) \, d\mu(y) - \mathbf{h}_{\Pi_{\mathbb{S}}}(-\mathbf{v}) + \lambda p ,$$

$$U_d(p; \mathbf{v}, \lambda) := \int_{\mathcal{Y}} \mathbf{h}_{d(y)}((y - \lambda) \mathbf{c}_\dagger + \mathbf{v}) \, d\mu(y) + \mathbf{h}_{\Pi_{\mathbb{S}}}(-\mathbf{v}) + \lambda p .$$

Theorem 4 (Sharp Bounds on T -Conditional Y_d -Means)

Suppose $\mathbb{S}$, d , and $\mathcal{T}_\dagger$ are as in Theorem 3. Define

$$\underline{p}_\dagger := \inf_{\mathbf{x} \in \Pi_{\mathbb{S}}} \mathbf{c}_\dagger^\top \mathbf{x} \quad \text{and} \quad \bar{p}_\dagger := \sup_{\mathbf{x} \in \Pi_{\mathbb{S}}} \mathbf{c}_\dagger^\top \mathbf{x} .$$

For each $p \in [\underline{p}_\dagger, \bar{p}_\dagger]$, $\mathbf{v} \in \mathbb{R}^{\mathcal{T}(\mathbb{S})}$, and $\lambda \in \mathbb{R}$, define

$$L_d(p; \mathbf{v}, \lambda) := - \int_{\mathcal{Y}} \mathbf{h}_{d(y)}((-y + \lambda) \mathbf{c}_\dagger + \mathbf{v}) \, d\mu(y) - \mathbf{h}_{\Pi_{\mathbb{S}}}(-\mathbf{v}) + \lambda p ,$$

$$U_d(p; \mathbf{v}, \lambda) := \int_{\mathcal{Y}} \mathbf{h}_{d(y)}((y - \lambda) \mathbf{c}_\dagger + \mathbf{v}) \, d\mu(y) + \mathbf{h}_{\Pi_{\mathbb{S}}}(-\mathbf{v}) + \lambda p .$$

Suppose $\underline{p}_\dagger > 0$.

Theorem 4 (Sharp Bounds on T -Conditional Y_d -Means)

Suppose $\mathbb{S}$, d , and $\mathcal{T}_\dagger$ are as in Theorem 3. Define

$$p_{-\dagger} := \inf_{\mathbf{x} \in \Pi_{\mathbb{S}}} \mathbf{c}_{\dagger}^\top \mathbf{x} \quad \text{and} \quad \bar{p}_{\dagger} := \sup_{\mathbf{x} \in \Pi_{\mathbb{S}}} \mathbf{c}_{\dagger}^\top \mathbf{x}.$$

For each $p \in [p_{-\dagger}, \bar{p}_{\dagger}]$, $\mathbf{v} \in \mathbb{R}^{\mathcal{T}(\mathbb{S})}$, and $\lambda \in \mathbb{R}$, define

$$L_d(p; \mathbf{v}, \lambda) := - \int_{\mathcal{Y}} \mathbf{h}_{d(y)}((-y + \lambda) \mathbf{c}_{\dagger} + \mathbf{v}) \, d\mu(y) - \mathbf{h}_{\Pi_{\mathbb{S}}}(-\mathbf{v}) + \lambda p,$$

$$U_d(p; \mathbf{v}, \lambda) := \int_{\mathcal{Y}} \mathbf{h}_{d(y)}((y - \lambda) \mathbf{c}_{\dagger} + \mathbf{v}) \, d\mu(y) + \mathbf{h}_{\Pi_{\mathbb{S}}}(-\mathbf{v}) + \lambda p.$$

Suppose $p_{-\dagger} > 0$. Then, the following bounds are sharp under $\mathbb{S}$:

$$\min_{p_{-\dagger} \leq p \leq \bar{p}_{\dagger}} \frac{L_d(p)}{p} \leq \mathbb{E}_Q[Y_d \mid \mathbf{T} \in \mathcal{T}_{\dagger}] \leq \max_{p_{-\dagger} \leq p \leq \bar{p}_{\dagger}} \frac{U_d(p)}{p},$$

where $L_d(p) := \sup_{\lambda, \mathbf{v}} \{L_d(p; \mathbf{v}, \lambda)\}$ and $U_d(p) := \inf_{\lambda, \mathbf{v}} \{U_d(p; \mathbf{v}, \lambda)\}$.

Comments

→ For each bound, solve **sequence of unconstrained convex programs**

Comments

- For each bound, solve sequence of unconstrained convex programs
- Easily extends to treatment effect case.

Comments

- For each bound, solve **sequence of unconstrained convex programs**
- **Easily extends to treatment effect case.** For mean $Y_d - Y_{d'}$, can replace $U_d(p; \mathbf{v}, \lambda)$ with $U_{(d,d')}(p; \mathbf{v}_d, \mathbf{v}_{d'}, \lambda)$, defined as

$$\int_{\mathcal{Y}} \mathbf{h}_{d(y)}(y \mathbf{c}_{\dagger} + \mathbf{v}_d) \, d\mu(y) + \int_{\mathcal{Y}} \mathbf{h}_{d'(y)}(-y \mathbf{c}_{\dagger} + \mathbf{v}_{d'}) \, d\mu(y) + \mathbf{h}_{\Pi_S}(-\mathbf{v}_d - \mathbf{v}_{d'} - \lambda \mathbf{c}_{\dagger}) + \lambda p.$$

Comments

→ For each bound, solve **sequence of unconstrained convex programs**

→ **Easily extends to treatment effect case.** For mean $Y_d - Y_{d'}$, can replace $U_d(p; \mathbf{v}, \lambda)$ with $U_{(d,d')}(p; \mathbf{v}_d, \mathbf{v}_{d'}, \lambda)$, defined as

$$\int_{\mathcal{Y}} \mathbf{h}_{d(y)}(y \mathbf{c}_{\dagger} + \mathbf{v}_d) d\mu(y) + \int_{\mathcal{Y}} \mathbf{h}_{d'(y)}(-y \mathbf{c}_{\dagger} + \mathbf{v}_{d'}) d\mu(y) + \mathbf{h}_{\Pi_{\mathbb{S}}}(-\mathbf{v}_d - \mathbf{v}_{d'} - \lambda \mathbf{c}_{\dagger}) + \lambda p.$$

- If $\mathcal{F}_T(\mathbb{S})$ is a **convex polytope**, there exists $\{\mathbf{c}_{k|d}\}_{k=1}^n$ such that

$$\Pi_{\mathbb{S}} = \left\{ \mathbf{x} \in \mathbb{R} \mid \mathbf{c}_{k|d}^{\top} \mathbf{x} \leq \mathbf{h}_{\Pi_{\mathbb{S}}}(\mathbf{c}_{k|d}) \forall k = 1, \dots, n \right\}.$$

Comments

→ For each bound, solve **sequence of unconstrained convex programs**

→ **Easily extends to treatment effect case.** For mean $Y_d - Y_{d'}$, can replace $U_d(p; \mathbf{v}, \lambda)$ with $U_{(d,d')}(p; \mathbf{v}_d, \mathbf{v}_{d'}, \lambda)$, defined as

$$\int_{\mathcal{Y}} \mathbf{h}_{d(y)}(y \mathbf{c}_{\dagger} + \mathbf{v}_d) d\mu(y) + \int_{\mathcal{Y}} \mathbf{h}_{d'(y)}(-y \mathbf{c}_{\dagger} + \mathbf{v}_{d'}) d\mu(y) + \mathbf{h}_{\Pi_{\mathbb{S}}}(-\mathbf{v}_d - \mathbf{v}_{d'} - \lambda \mathbf{c}_{\dagger}) + \lambda p.$$

- If $\mathcal{F}_T(\mathbb{S})$ is a **convex polytope**, there exists $\{\mathbf{c}_{k|d}\}_{k=1}^n$ such that

$$\Pi_{\mathbb{S}} = \left\{ \mathbf{x} \in \mathbb{R} \mid \mathbf{c}_{k|d}^{\top} \mathbf{x} \leq \mathbf{h}_{\Pi_{\mathbb{S}}}(\mathbf{c}_{k|d}) \forall k = 1, \dots, n \right\}.$$

In such cases, $U_d(p)$ has the following **“multiplier”** representation

$$U_d(p) = \inf_{\substack{\lambda \in \mathbb{R}, \\ \gamma \in \mathbb{R}_{\geq 0}^n}} \left\{ \int_{\mathcal{Y}} \mathbf{h}_{d(y)} \left((\kappa(y) - \lambda) \mathbf{c}_{\dagger} - \sum_{k=1}^n \gamma_{(k)} \mathbf{c}_{k|d} \right) d\mu(y) + \sum_{k=1}^n \gamma_{(k)} \mathbf{h}_{\Pi_{\mathbb{S}}}(\mathbf{c}_{k|d}) + \lambda p \right\}$$

Comments

→ For each bound, solve **sequence of unconstrained convex programs**

→ **Easily extends to treatment effect case.** For mean $Y_d - Y_{d'}$, can replace $U_d(p; \mathbf{v}, \lambda)$ with $U_{(d,d')}(p; \mathbf{v}_d, \mathbf{v}_{d'}, \lambda)$, defined as

$$\int_{\mathcal{Y}} \mathbf{h}_{d(y)}(y \mathbf{c}_{\dagger} + \mathbf{v}_d) d\mu(y) + \int_{\mathcal{Y}} \mathbf{h}_{d'(y)}(-y \mathbf{c}_{\dagger} + \mathbf{v}_{d'}) d\mu(y) + \mathbf{h}_{\Pi_{\mathbb{S}}}(-\mathbf{v}_d - \mathbf{v}_{d'} - \lambda \mathbf{c}_{\dagger}) + \lambda p.$$

- If $\mathcal{F}_{\mathcal{T}}(\mathbb{S})$ is a **convex polytope**, there exists $\{\mathbf{c}_{k|d}\}_{k=1}^n$ such that

$$\Pi_{\mathbb{S}} = \left\{ \mathbf{x} \in \mathbb{R} \mid \mathbf{c}_{k|d}^{\top} \mathbf{x} \leq \mathbf{h}_{\Pi_{\mathbb{S}}}(\mathbf{c}_{k|d}) \forall k = 1, \dots, n \right\}.$$

In such cases, $U_d(p)$ has the following **“multiplier”** representation

$$U_d(p) = \inf_{\substack{\lambda \in \mathbb{R}, \\ \gamma \in \mathbb{R}_{\geq 0}^n}} \left\{ \int_{\mathcal{Y}} \mathbf{h}_{d(y)} \left((\kappa(y) - \lambda) \mathbf{c}_{\dagger} - \sum_{k=1}^n \gamma_{(k)} \mathbf{c}_{k|d} \right) d\mu(y) + \sum_{k=1}^n \gamma_{(k)} \mathbf{h}_{\Pi_{\mathbb{S}}}(\mathbf{c}_{k|d}) + \lambda p \right\}$$

→ In many cases, $n \ll |\mathcal{T}|$, leading to computational gains

A Simpler Approach?

A Simpler Approach?

For fixed p , **upper** bound is a *min* problem, **lower** bound is *max* problem

A Simpler Approach?

For fixed p , **upper** bound is a *min* problem, **lower** bound is *max* problem

→ adding constraints can only widen bounds, maintaining validity

A Simpler Approach?

For fixed p , **upper** bound is a *min* problem, **lower** bound is *max* problem

→ adding constraints can only widen bounds, maintaining validity

Imposing $v = 0$ gives upper bound $\inf_{\lambda \in \mathbb{R}} \left\{ \int_{\mathcal{Y}} h_{d(y)}((y - \lambda) c_{\dagger}) d\mu(y) + \lambda p \right\}$

A Simpler Approach?

For fixed p , **upper** bound is a *min* problem, **lower** bound is *max* problem

→ adding constraints can only widen bounds, maintaining validity

Imposing $\mathbf{v} = 0$ gives upper bound $\inf_{\lambda \in \mathbb{R}} \left\{ \int_{\mathcal{Y}} \mathbf{h}_{d(y)}((y - \lambda) \mathbf{c}_{\dagger}) d\mu(y) + \lambda p \right\}$

Support functions are **positive homogeneous**: $\mathbf{h}_C(\alpha \mathbf{c}) = \alpha \mathbf{h}_C(\mathbf{c}) \forall \alpha \geq 0$

A Simpler Approach?

For fixed p , **upper** bound is a *min* problem, **lower** bound is *max* problem

→ adding constraints can only widen bounds, maintaining validity

Imposing $\mathbf{v} = 0$ gives upper bound $\inf_{\lambda \in \mathbb{R}} \left\{ \int_{\mathcal{Y}} \mathbf{h}_{d(y)}((y - \lambda) \mathbf{c}_{\dagger}) d\mu(y) + \lambda p \right\}$

Support functions are **positive homogeneous**: $\mathbf{h}_C(\alpha \mathbf{c}) = \alpha \mathbf{h}_C(\mathbf{c}) \forall \alpha \geq 0$

→ Can write upper bound as

$$\inf_{\lambda \in \mathbb{R}} \left\{ \int_{-\infty}^{\lambda} (\lambda - y) \mathbf{h}_{d(y)}(-\mathbf{c}_{\dagger}) d\mu(y) + \int_{\lambda}^{\infty} (y - \lambda) \mathbf{h}_{d(y)}(\mathbf{c}_{\dagger}) d\mu(y) + \lambda p \right\}$$

A Simpler Approach?

For fixed p , **upper** bound is a *min* problem, **lower** bound is *max* problem

→ adding constraints can only widen bounds, maintaining validity

Imposing $\mathbf{v} = 0$ gives upper bound $\inf_{\lambda \in \mathbb{R}} \left\{ \int_{\mathcal{Y}} \mathbf{h}_{d(y)}((y - \lambda) \mathbf{c}_{\dagger}) d\mu(y) + \lambda p \right\}$

Support functions are **positive homogeneous**: $\mathbf{h}_C(\alpha \mathbf{c}) = \alpha \mathbf{h}_C(\mathbf{c}) \forall \alpha \geq 0$

→ Can write upper bound as

$$\begin{aligned} & \inf_{\lambda \in \mathbb{R}} \left\{ \int_{-\infty}^{\lambda} (\lambda - y) \mathbf{h}_{d(y)}(-\mathbf{c}_{\dagger}) d\mu(y) + \int_{\lambda}^{\infty} (y - \lambda) \mathbf{h}_{d(y)}(\mathbf{c}_{\dagger}) d\mu(y) + \lambda p \right\} \\ &= \inf_{\lambda \in \mathbb{R}} \left\{ \int_{-\infty}^{\lambda} y (-\mathbf{h}_{d(y)}(-\mathbf{c}_{\dagger})) d\mu(y) + \int_{\lambda}^{\infty} y \mathbf{h}_{d(y)}(\mathbf{c}_{\dagger}) d\mu(y) \right. \\ & \quad \left. \lambda \left(p - \left(\int_{-\infty}^{\lambda} -\mathbf{h}_{d(y)}(-\mathbf{c}_{\dagger}) d\mu(y) + \int_{\lambda}^{\infty} \mathbf{h}_{d(y)}(\mathbf{c}_{\dagger}) d\mu(y) \right) \right) \right\}. \end{aligned}$$

Simple Bounds: Solving for λ

Differentiating wrt λ and cancelling terms yields the following FOC

$$p = \int_{-\infty}^{\lambda} -h_{d(y)}(-c_{\dagger}) d\mu(y) + \int_{\lambda}^{\infty} h_{d(y)}(c_{\dagger}) d\mu(y) ,$$

Simple Bounds: Solving for λ

Differentiating wrt λ and cancelling terms yields the following FOC

$$p = \int_{-\infty}^{\lambda} -\mathbf{h}_{d(y)}(-\mathbf{c}_{\dagger}) \, d\mu(y) + \int_{\lambda}^{\infty} \mathbf{h}_{d(y)}(\mathbf{c}_{\dagger}) \, d\mu(y) \, ,$$

which implies

$$p - \int_{-\infty}^{\infty} -\mathbf{h}_{d(y)}(-\mathbf{c}_{\dagger}) \, d\mu(y) = \int_{\lambda}^{\infty} \mathbf{h}_{d(y)}(\mathbf{c}_{\dagger}) - (-\mathbf{h}_{d(y)}(-\mathbf{c}_{\dagger})) \, d\mu(y) \, ,$$

Simple Bounds: Solving for λ

Differentiating wrt λ and cancelling terms yields the following FOC

$$p = \int_{-\infty}^{\lambda} -h_{d(y)}(-c_{\dagger}) d\mu(y) + \int_{\lambda}^{\infty} h_{d(y)}(c_{\dagger}) d\mu(y) ,$$

which implies

$$p - \int_{-\infty}^{\infty} -h_{d(y)}(-c_{\dagger}) d\mu(y) = \int_{\lambda}^{\infty} h_{d(y)}(c_{\dagger}) - (-h_{d(y)}(-c_{\dagger})) d\mu(y) ,$$

→ Left side does not depend on λ , right side invertible as function of λ

Simple Bounds: Solving for λ

Differentiating wrt λ and cancelling terms yields the following FOC

$$p = \int_{-\infty}^{\lambda} -h_{d(y)}(-c_{\dagger}) d\mu(y) + \int_{\lambda}^{\infty} h_{d(y)}(c_{\dagger}) d\mu(y) ,$$

which implies

$$p - \int_{-\infty}^{\infty} -h_{d(y)}(-c_{\dagger}) d\mu(y) = \int_{\lambda}^{\infty} h_{d(y)}(c_{\dagger}) - (-h_{d(y)}(-c_{\dagger})) d\mu(y) ,$$

→ Left side does not depend on λ , right side invertible as function of λ

→ Apply inverse to both sides to get optimal λ as function of p

Simple Bounds: Solving for λ

Differentiating wrt λ and cancelling terms yields the following FOC

$$p = \int_{-\infty}^{\lambda} -h_{d(y)}(-c_{\dagger}) d\mu(y) + \int_{\lambda}^{\infty} h_{d(y)}(c_{\dagger}) d\mu(y) ,$$

which implies

$$p - \int_{-\infty}^{\infty} -h_{d(y)}(-c_{\dagger}) d\mu(y) = \int_{\lambda}^{\infty} h_{d(y)}(c_{\dagger}) - (-h_{d(y)}(-c_{\dagger})) d\mu(y) ,$$

- Left side does not depend on λ , right side invertible as function of λ
- Apply inverse to both sides to get optimal λ as function of p
- If Y is not continuous, can use suitable generalized inverse

“Generalized Trimming Bounds”: Setup

To formalize this “trimming” idea, need to introduce some new operators.

“Generalized Trimming Bounds”: Setup

To formalize this “trimming” idea, need to introduce some new operators.

For any non-negative Borel measurable function $f : \mathcal{Y} \rightarrow \mathbb{R}_{\geq 0}$, define

$$\mathfrak{F}_{\uparrow}^{-1}[f], \mathfrak{F}_{\downarrow}^{-1}[f] : \left[0, \int_{\mathcal{Y}} f(y) dy\right] \rightarrow \mathcal{Y}$$

“Generalized Trimming Bounds”: Setup

To formalize this “trimming” idea, need to introduce some new operators.

For any non-negative Borel measurable function $f : \mathcal{Y} \rightarrow \mathbb{R}_{\geq 0}$, define

$$\mathfrak{F}_{\uparrow}^{-1}[f], \mathfrak{F}_{\downarrow}^{-1}[f] : \left[0, \int_{\mathcal{Y}} f(y) dy\right] \rightarrow \mathcal{Y}$$

as follows, for every $0 \leq q \leq \int_{\mathcal{Y}} f(y) dy$,

$$\mathfrak{F}_{\uparrow}^{-1}[f](q) := \inf \left\{ x \in \mathcal{Y} \mid q \leq \int_{-\infty}^x f(y) dy \right\},$$

“Generalized Trimming Bounds”: Setup

To formalize this “trimming” idea, need to introduce some new operators.

For any **non-negative Borel measurable function** $f : \mathcal{Y} \rightarrow \mathbb{R}_{\geq 0}$, define

$$\mathfrak{F}_{\uparrow}^{-1}[f], \mathfrak{F}_{\downarrow}^{-1}[f] : \left[0, \int_{\mathcal{Y}} f(y) dy\right] \rightarrow \mathcal{Y}$$

as follows, for every $0 \leq q \leq \int_{\mathcal{Y}} f(y) dy$,

$$\begin{aligned}\mathfrak{F}_{\uparrow}^{-1}[f](q) &:= \inf \left\{ x \in \mathcal{Y} \mid q \leq \int_{-\infty}^x f(y) dy \right\}, \\ \mathfrak{F}_{\downarrow}^{-1}[f](q) &:= \sup \left\{ x \in \mathcal{Y} \mid q \leq \int_x^{\infty} f(y) dy \right\}.\end{aligned}$$

→ **If f is a density**, $\mathfrak{F}_{\uparrow}^{-1}[f]$ is **(generalized) quantile of CDF** associated with f , and $\mathfrak{F}_{\downarrow}^{-1}[f](q) = \mathfrak{F}_{\uparrow}^{-1}[f](1 - q)$

“Generalized Trimming Bounds”: Setup

To formalize this “trimming” idea, need to introduce some new operators.

For any **non-negative Borel measurable function** $f : \mathcal{Y} \rightarrow \mathbb{R}_{\geq 0}$, define

$$\mathfrak{F}_{\uparrow}^{-1}[f], \mathfrak{F}_{\downarrow}^{-1}[f] : \left[0, \int_{\mathcal{Y}} f(y) dy\right] \rightarrow \mathcal{Y}$$

as follows, for every $0 \leq q \leq \int_{\mathcal{Y}} f(y) dy$,

$$\begin{aligned}\mathfrak{F}_{\uparrow}^{-1}[f](q) &:= \inf \left\{ x \in \mathcal{Y} \mid q \leq \int_{-\infty}^x f(y) dy \right\}, \\ \mathfrak{F}_{\downarrow}^{-1}[f](q) &:= \sup \left\{ x \in \mathcal{Y} \mid q \leq \int_x^{\infty} f(y) dy \right\}.\end{aligned}$$

→ **If f is a density**, $\mathfrak{F}_{\uparrow}^{-1}[f]$ is **(generalized) quantile of CDF** associated with f , and $\mathfrak{F}_{\downarrow}^{-1}[f](q) = \mathfrak{F}_{\uparrow}^{-1}[f](1 - q)$

→ More generally, $\mathfrak{F}_{\downarrow}^{-1}[f](q) = \mathfrak{F}_{\uparrow}^{-1}[f](\int_{\mathcal{Y}} f(y) dy - q)$

Theorem 5 (Generalized Trimming Bounds)

Suppose $\mathbb{S}$, d , and $\mathcal{T}_\dagger$ are as in Theorem 3.

Define $\bar{f}_{\dagger|d}(y) := \mathbf{h}_{d(y)}(\mathbf{c}_\dagger)$ and $\underline{f}_{\dagger|d}(y) := -\mathbf{h}_{d(y)}(-\mathbf{c}_\dagger)$,

Theorem 5 (Generalized Trimming Bounds)

Suppose $\mathbb{S}$, d , and $\mathcal{T}_{\dagger}$ are as in Theorem 3.

Define $\bar{f}_{\dagger|d}(y) := \mathbf{h}_{d(y)}(\mathbf{c}_{\dagger})$ and $\underline{f}_{\dagger|d}(y) := -\mathbf{h}_{d(y)}(-\mathbf{c}_{\dagger})$, and

let $\bar{p}_{\dagger|d} := \int_{\mathcal{Y}} \bar{f}_{\dagger|d}(y) d\mu(y)$ and $\underline{p}_{\dagger|d} := \int_{\mathcal{Y}} \underline{f}_{\dagger|d}(y) d\mu(y)$.

Theorem 5 (Generalized Trimming Bounds)

Suppose $\mathbb{S}$, d , and $\mathcal{T}_\dagger$ are as in Theorem 3.

Define $\bar{f}_{\dagger|d}(y) := \mathbf{h}_{d(y)}(\mathbf{c}_\dagger)$ and $\underline{f}_{\dagger|d}(y) := -\mathbf{h}_{d(y)}(-\mathbf{c}_\dagger)$, and let $\bar{p}_{\dagger|d} := \int_{\mathcal{Y}} \bar{f}_{\dagger|d}(y) d\mu(y)$ and $\underline{p}_{\dagger|d} := \int_{\mathcal{Y}} \underline{f}_{\dagger|d}(y) d\mu(y)$.

Then, for any $d \in \mathcal{D}$,

$$\frac{\tilde{L}_d(p_L^*)}{p_L^*} \leq \mathbb{E} [\kappa(Y_d) \mid \mathbf{T} \in \mathcal{T}_\dagger] \leq \frac{\tilde{U}_d(p_U^*)}{p_U^*},$$

Theorem 5 (Generalized Trimming Bounds)

Suppose $\mathbb{S}$, d , and $\mathcal{T}_\dagger$ are as in Theorem 3.

Define $\bar{f}_{\dagger|d}(y) := \mathbf{h}_{d(y)}(\mathbf{c}_\dagger)$ and $\underline{f}_{\dagger|d}(y) := -\mathbf{h}_{d(y)}(-\mathbf{c}_\dagger)$, and let $\bar{p}_{\dagger|d} := \int_{\mathcal{Y}} \bar{f}_{\dagger|d}(y) d\mu(y)$ and $\underline{p}_{\dagger|d} := \int_{\mathcal{Y}} \underline{f}_{\dagger|d}(y) d\mu(y)$.

Then, for any $d \in \mathcal{D}$,

$$\frac{\tilde{L}_d(p_L^*)}{p_L^*} \leq \mathbb{E}[\kappa(Y_d) \mid \mathbf{T} \in \mathcal{T}_\dagger] \leq \frac{\tilde{U}_d(p_U^*)}{p_U^*},$$

where

$$\begin{aligned} \tilde{L}_d(p) &:= \int_{\mathcal{Y}} y \underline{f}_{\dagger}(y) d\mu(y) + \int_0^{p - \underline{p}_{\dagger|d}} \mathfrak{F}_\dagger^{-1}[\bar{f}_\dagger - \underline{f}_\dagger](q) d\nu(q), \\ \tilde{U}_d(p) &:= \int_{\mathcal{Y}} y \bar{f}_{\dagger}(y) d\mu(y) + \int_0^{p - \bar{p}_{\dagger|d}} \mathfrak{F}_\dagger^{-1}[\bar{f}_\dagger - \underline{f}_\dagger](q) d\nu(q), \end{aligned}$$

and p_L^* and p_U^* are as follows

Theorem 5 contd.

$$p_L^* := \begin{cases} \bar{p}_\dagger & \text{if } p_L > \bar{p}_\dagger \\ \underline{p}_\dagger & \text{if } p_L < \underline{p}_\dagger \\ p_L & \text{otherwise} \end{cases}, \text{ with } p_L := \inf \left\{ p \in [\underline{p}_{\dagger|d}, \bar{p}_{\dagger|d}] \mid \mathfrak{F}_\dagger^{-1} [\bar{f}_\dagger - \underline{f}_\dagger] (p - \underline{p}_{\dagger|d}) \geq \frac{L(p)}{p} \right\},$$

$$p_U^* := \begin{cases} \bar{p}_\dagger & \text{if } p_U > \bar{p}_\dagger \\ \underline{p}_\dagger & \text{if } p_U < \underline{p}_\dagger \\ p_U & \text{otherwise} \end{cases}, \text{ with } p_U := \inf \left\{ p \in [\underline{p}_{\dagger|d}, \bar{p}_{\dagger|d}] \mid \mathfrak{F}_\dagger^{-1} [\bar{f}_\dagger - \underline{f}_\dagger] (p - \underline{p}_{\dagger|d}) \leq \frac{U(p)}{p} \right\}.$$

Theorem 5 contd.

$$p_L^* := \begin{cases} \bar{p}_\dagger & \text{if } p_L > \bar{p}_\dagger \\ \underline{p}_\dagger & \text{if } p_L < \underline{p}_\dagger \\ p_L & \text{otherwise} \end{cases}, \text{ with } p_L := \inf \left\{ p \in [\underline{p}_{\dagger|d}, \bar{p}_{\dagger|d}] \mid \mathfrak{F}_\dagger^{-1} [\bar{f}_\dagger - \underline{f}_\dagger] (p - \underline{p}_{\dagger|d}) \geq \frac{L(p)}{p} \right\},$$

$$p_U^* := \begin{cases} \bar{p}_\dagger & \text{if } p_U > \bar{p}_\dagger \\ \underline{p}_\dagger & \text{if } p_U < \underline{p}_\dagger \\ p_U & \text{otherwise} \end{cases}, \text{ with } p_U := \inf \left\{ p \in [\underline{p}_{\dagger|d}, \bar{p}_{\dagger|d}] \mid \mathfrak{F}_\dagger^{-1} [\bar{f}_\dagger - \underline{f}_\dagger] (p - \underline{p}_{\dagger|d}) \leq \frac{U(p)}{p} \right\}.$$

In cases where $\underline{f}_\dagger(y) = 0$ for all $y \in \mathcal{Y}$, the bounds simplify to

$$\begin{aligned} \frac{1}{\underline{p}_\dagger} \int_0^{\underline{p}_\dagger} \mathfrak{F}_\dagger^{-1} [\bar{f}_\dagger] (q) \, d\nu(q) \\ \leq \mathbb{E} [Y_d \mid \mathbf{T} \in \mathcal{T}_\dagger] \\ \leq \frac{1}{\underline{p}_\dagger} \int_0^{\underline{p}_\dagger} \mathfrak{F}_\dagger^{-1} [\bar{f}_\dagger] (q) \, d\nu(q) \end{aligned}$$

Comments

- Bounds of Horowitz and Manski (1995), Lee (2009), Kitagawa (2021), and Marx (2024) can be obtained as **special cases of Theorem 5**
- Sharp in many cases, but not in general
 - Sharp if $\begin{bmatrix} \mathbf{R}_d \\ \mathbf{c}_\dagger^\top \end{bmatrix}$ *with duplicate and zero columns removed* is tall
 - In particular, sharp under the **Unordered Monotonicity** condition of Heckman and Pinto (2018)
 - Sharp for Examples 1 & 2, not sharp for Example 3

Remaining Content of Paper

1. **Identification:** Remaining identification results, including under some assumptions on outcomes
2. **Analytic bounds:** Automated derivation of analytic expressions for “point-wise” bounds
3. **Inference:** Support function based test statistic for inclusion tests
4. **Computation:** Primal-dual method for computing support function
 - feasible for large problems, exact solution in finitely many steps
 - can also provide fast approximations with deterministic error bounds

Conclusion

Summary

Framework for partial identification in IV treatment selection models

- multivalued treatments, discrete instruments
- sharp under random assignment of instrument
- tractable, not “black box”: analytic bounds
- sharp specification tests

Current and Future Work

- Formalizing inference and estimation
- Simulations and empirical applications
- Developing a Stata/R/Python package
- Additional guidance on when bounds are likely to be informative

Thank you!

Basic Example ($Z \in \{0, 1\}$, $D \in \{0, 1\}$, no selection restrictions)

For $d = 1$, $(\star)$ implies

$$\begin{bmatrix} 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \end{bmatrix} \boldsymbol{\xi}_1(y) = \mathbf{p}\mathbf{f}_1(y)$$

Basic Example ($Z \in \{0, 1\}$, $D \in \{0, 1\}$, no selection restrictions)

For $d = 1$, $(\star)$ implies

$$\begin{aligned} & \begin{bmatrix} 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \end{bmatrix} \boldsymbol{\xi}_1(y) = \mathbf{pf}_1(y) \\ \Rightarrow & \begin{bmatrix} 1 \\ 1 \end{bmatrix} P(\mathbf{T} = AT) f_{Y_1|\mathbf{T}=AT}(y) \leq \begin{bmatrix} P(D = 1|Z = 0) f_{Y|D=1,Z=0}(y) \\ P(D = 1|Z = 1) f_{Y|D=1,Z=1}(y) \end{bmatrix} \end{aligned}$$

Basic Example ($Z \in \{0, 1\}$, $D \in \{0, 1\}$, no selection restrictions)

For $d = 1$, $(\star)$ implies

$$\begin{bmatrix} 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \end{bmatrix} \boldsymbol{\xi}_1(y) = \mathbf{p}\mathbf{f}_1(y)$$
$$\Rightarrow \begin{bmatrix} 1 \\ 1 \end{bmatrix} P(\mathbf{T} = AT) f_{Y_1|\mathbf{T}=AT}(y) \leq \begin{bmatrix} P(D = 1|Z = 0) f_{Y|D=1,Z=0}(y) \\ P(D = 1|Z = 1) f_{Y|D=1,Z=1}(y) \end{bmatrix}$$

The support function for the 1-dimensional polytope (i.e. interval) defined by the non-negative solutions to the above system is simply the map

$$c \mapsto \begin{cases} c \left(\min_{z \in \{0,1\}} \left\{ P(D = 1|Z = z) f_{Y|D=1,Z=z}(y) \right\} \right) & \text{if } c \geq 0 \\ 0 & \text{if } c < 0 \end{cases}$$

Basic Example ($Z \in \{0, 1\}$, $D \in \{0, 1\}$, no selection restrictions)

$\Pi_{\mathcal{T}|1}$ consists of all $\pi \geq 0$ such that

$$\pi_{(AT)} \leq \int_{\mathcal{Y}} \min_{z \in \{0,1\}} \left\{ P(D=1|Z=z) f_{Y|D=1,Z=z}(y) \right\} d\mu(y) ,$$

$$\pi_{(AT)} + \pi_{(D)} = P(D=1|Z=0) = \int_{\mathcal{Y}} P(D=1|Z=0) f_{Y|D=1,Z=0}(y) d\mu(y) ,$$

$$\pi_{(AT)} + \pi_{(C)} = P(D=1|Z=1) = \int_{\mathcal{Y}} P(D=1|Z=1) f_{Y|D=1,Z=1}(y) d\mu(y) .$$

Similarly, $\Pi_{\mathcal{T}|0}$ consists of all $\pi \geq 0$ such that

$$\pi_{(NT)} \leq \int_{\mathcal{Y}} \min_{z \in \{0,1\}} \left\{ P(D=0|Z=z) f_{Y|D=0,Z=z}(y) \right\} d\mu(y) ,$$

$$\pi_{(C)} + \pi_{(NT)} = P(D=0|Z=0) ,$$

$$\pi_{(D)} + \pi_{(NT)} = P(D=0|Z=1) .$$

$\Pi_{\mathcal{T}}$ consists of all $\pi \geq 0$ that satisfy the 6 conditions above.

MTO Example: Sharp Bounds on $\pi_{[(\cdot, M)]}$

For the MTO example considered earlier, the following bounds are sharp

$$\max \left\{ \int_{\mathcal{Y}} \max \left\{ 0, \mathbf{p}\mathbf{f}_{M|Z=(1,1)}(y) - \mathbf{p}\mathbf{f}_{M|Z=(0,0)}(y) \right\} d\mu(y), \right. \\ \left. \mathbf{p}_H[(0,1)] - \mathbf{p}_H[(1,1)] \int_{\mathcal{Y}} \max \left\{ 0, \mathbf{p}\mathbf{f}_{L|Z=(0,0)}(y) - \mathbf{p}\mathbf{f}_{L|Z=(1,1)}(y) \right\} d\mu(y) \right\}$$

$$\leq \pi_{(HHM)} + \pi_{(HLM)} + \pi_{(LLM)} \leq$$

$$\min \left\{ \mathbf{p}_M[(1,1)] - \mathbf{p}_M[(0,1)], \right. \\ \left. \mathbf{p}_M[(1,1)] - \mathbf{p}_M[(0,0)] + \int_{\mathcal{Y}} \min \left\{ \mathbf{p}\mathbf{f}_{L|Z=(1,1)}(y), \mathbf{p}\mathbf{f}_{L|Z=(0,1)}(y) - \mathbf{p}\mathbf{f}_{L|Z=(0,0)}(y) \right\} d\mu(y) \right\}$$

Works Cited i

- Angrist, J. D. and G. W. Imbens (1995, June). Two-Stage Least Squares Estimation of Average Causal Effects in Models with Variable Treatment Intensity. *Journal of the American Statistical Association* 90(430), 431–442.
- Artstein, Z. and R. A. Vitale (1975, October). A Strong Law of Large Numbers for Random Compact Sets. *The Annals of Probability* 3(5), 879–882.
- Aumann, R. J. (1965, August). Integrals of set-valued functions. *Journal of Mathematical Analysis and Applications* 12(1), 1–12.
- Balke, A. and J. Pearl (1997). Bounds on Treatment Effects From Studies With Imperfect Compliance. *Journal of the American Statistical Association* 92(439), 1171–1176.

Works Cited ii

- Beresteanu, A., I. Molchanov, and F. Molinari (2012, January). Partial identification using random set theory. *Journal of Econometrics* 166(1), 17–32.
- Billera, L. J. and B. Sturmfels (1992). Fiber Polytopes. *Annals of Mathematics* 135(3), 527–549.
- Fang, Z., A. Santos, A. M. Shaikh, and A. Torgovitsky (2023). Inference for Large-Scale Linear Systems With Known Coefficients. *Econometrica* 91(1), 299–327.
- Fang, Z. and J. Seo (2021). A Projection Framework for Testing Shape Restrictions That Form Convex Cones. *Econometrica* 89(5), 2439–2458.
- Heckman, J. J. and R. Pinto (2018). Unordered Monotonicity. *Econometrica* 86(1), 1–35.

Works Cited iii

Heckman, J. J. and E. Vytlacil (2001). Policy-Relevant Treatment Effects. *American Economic Review* 91(2), 107–111.

Heckman, J. J. and E. J. Vytlacil (1999, April). Local instrumental variables and latent variable models for identifying and bounding treatment effects. *Proceedings of the National Academy of Sciences of the United States of America* 96(8), 4730–4734.

Heckman, J. J. and E. J. Vytlacil (2007a, January). Chapter 70 Econometric Evaluation of Social Programs, Part I: Causal Models, Structural Models and Econometric Policy Evaluation. In J. J. Heckman and E. E. Leamer (Eds.), *Handbook of Econometrics*, Volume 6, pp. 4779–4874. Elsevier.

Works Cited iv

- Heckman, J. J. and E. J. Vytlačil (2007b, January). Chapter 71
Econometric Evaluation of Social Programs, Part II: Using the Marginal
Treatment Effect to Organize Alternative Econometric Estimators to
Evaluate Social Programs, and to Forecast their Effects in New
Environments. In J. J. Heckman and E. E. Leamer (Eds.), *Handbook of
Econometrics*, Volume 6, pp. 4875–5143. Elsevier.
- Horowitz, J. L. and C. F. Manski (1995). Identification and Robustness
with Contaminated and Corrupted Data. *Econometrica* 63(2), 281–302.
- Kamat, V., S. Norris, and M. Pecenco (2023, July). Identification in
Multiple Treatment Models under Discrete Variation.
<https://arxiv.org/abs/2307.06174v1>.
- Kédagni, D. and I. Mourifié (2020, September). Generalized instrumental
inequalities: Testing the instrumental variable independence
assumption. *Biometrika* 107(3), 661–675.

Works Cited v

- Kirkebøen, L. J., E. Leuven, and M. Mogstad (2016, August). Field of Study, Earnings, and Self-Selection*. *The Quarterly Journal of Economics* 131(3), 1057–1111.
- Kitagawa, T. (2015). A Test for Instrument Validity. *Econometrica* 83(5), 2043–2063.
- Kitagawa, T. (2021, December). The identification region of the potential outcome distributions under instrument independence. *Journal of Econometrics* 225(2), 231–253.
- Kroft, K., I. Mourifié, and A. Vayalinkal (2024). Lee Bounds with Multilayered Sample Selection. *Unpublished Manuscript, University of Toronto*.
- Lee, D. S. (2009, July). Training, Wages, and Sample Selection: Estimating Sharp Bounds on Treatment Effects. *The Review of Economic Studies* 76(3), 1071–1102.

Works Cited vi

- Lee, S. and B. Salanié (2018). Identifying Effects of Multivalued Treatments. *Econometrica* 86(6), 1939–1963.
- Manski, C. F. (1989). Anatomy of the Selection Problem. *The Journal of Human Resources* 24(3), 343–360.
- Manski, C. F. (1990). Nonparametric Bounds on Treatment Effects. *The American Economic Review* 80(2), 319–323.
- Marx, P. (2024, January). Sharp bounds in the latent index selection model. *Journal of Econometrics* 238(2), 105561.
- Mogstad, M., A. Santos, and A. Torgovitsky (2018). Using Instrumental Variables for Inference About Policy Relevant Treatment Parameters. *Econometrica* 86(5), 1589–1619.
- Mountjoy, J. (2022, August). Community Colleges and Upward Mobility. *American Economic Review* 112(8), 2580–2630.

Works Cited vii

- Mourifié, I. and Y. Wan (2017, May). Testing Local Average Treatment Effect Assumptions. *The Review of Economics and Statistics* 99(2), 305–313.
- Noack, C. (2021, June). Sensitivity of LATE Estimates to Violations of the Monotonicity Assumption. <https://arxiv.org/abs/2106.06421v1>.
- Sun, Z. (2023, October). Instrument Validity for Heterogeneous Causal Effects. <http://arxiv.org/abs/2009.01995>.
- Torgovitsky, A. (2019). Partial identification by extending subdistributions. *Quantitative Economics* 10(1), 105–144.