

PHYS 502: Mathematical Physics II

Winter 2015, Homework #5

(Due March 11, 2015)

1. By first considering the behavior of the fundamental solution in the vicinity of $\mathbf{x} = \mathbf{x}'$, show that the Green's function $G(\mathbf{x}, \mathbf{x}')$ for the three-dimensional Helmholtz equation

$$\nabla^2 u + k^2 u = 0,$$

with the boundary condition that $u(\mathbf{x})e^{-i\omega t}$ represents outgoing waves at infinity, is

$$G(\mathbf{x}, \mathbf{x}') = -\frac{e^{ikr}}{4\pi r},$$

where $r = |\mathbf{x} - \mathbf{x}'|$.

2. In using the method of images to find the Dirichlet Green's function for Poisson's equation inside a sphere of radius a , it can be shown that the image of a point \mathbf{x} within the sphere is $\mathbf{x}_1 = \alpha\mathbf{x}$, with strength β , so that the Green's function is

$$G(\mathbf{x}, \mathbf{x}') = -\frac{1}{4\pi|\mathbf{x}' - \mathbf{x}|} + \frac{\beta}{4\pi|\mathbf{x}' - \mathbf{x}_1|}.$$

(a) The boundary condition on G is that $G(\mathbf{x}, \mathbf{x}') = 0$ if \mathbf{x}' lies on the surface of the sphere. By applying this condition to the two points where the diameter through \mathbf{x} intersects the surface of the sphere, show that $\beta = a/r$ and $\alpha = \beta^2$, where $r = |\mathbf{x}|$.

(b) Hence derive an expression for the solution $u(r, \theta, \phi)$ to Laplace's equation $\nabla^2 u = 0$ inside the sphere, subject to the boundary condition $u(a, \theta, \phi) = f(\theta, \phi)$.

(c) Compare this form of the solution with the series solution obtained by separation of variables within the sphere $r < a$.

3. As discussed in class, the Green's function solution to the inhomogeneous wave equation

$$\nabla^2 \phi - \frac{1}{c^2} \frac{\partial^2 \phi}{\partial t^2} = f(\mathbf{x}, t)$$

is the "retarded potential" solution

$$\phi(\mathbf{x}, t) = -\frac{c}{4\pi} \int d^3\mathbf{x}' dt' f(\mathbf{x}', t') \frac{\delta[|\mathbf{x} - \mathbf{x}'| - c(t - t')]}{|\mathbf{x} - \mathbf{x}'|},$$

where $\frac{1}{c}|\mathbf{x} - \mathbf{x}'|$ is the light travel time from the source point \mathbf{x}' to the field point \mathbf{x} .

Now suppose that the field is due to a moving point source of unit magnitude, so

$$f(\mathbf{x}', t') = \delta[\mathbf{x}' - \xi(t')],$$

that is, the trajectory of the source is given by $\mathbf{x}' = \xi(t')$. Use the above equation to find a closed-form expression (no integrals!) for the potential $\phi(\mathbf{x}, t)$.