

Index

A

Accuracy in estimation, 6
Addition rule, 125–27
 general addition rule, 127
Agresti–Coull method, 190
AIDS, 428–29
Alternative hypothesis, 154
Analysis of covariance (ANCOVA), 623–27
Analysis of deviance, 575
Analysis of variance, 464–66
 ANOVA table, 467
 assumptions, 474–75
 confidence interval for difference
 between means, planned,
 475–77, 489
 critical value for F , 472
 error mean square, 471, 485
 example of a linear model, 612–14
 fixed vs random effects, 481–82
 F -ratio, 471
 group mean square, 466, 467,
 470, 485
 how it works, 466
 hypotheses, 465–66
 linear regression, 560
 mean squares, 470–71
 multiple testing, 474–81
 nonparametric alternative,
 Kruskal–Wallis test, 475, 488
 planned comparisons, 476–77, 489
 R^2 , fraction of variation explained,
 473
 random effects, 481–82
 repeatability, 485–86
 robustness, 474
 sum of squares, partitioning,
 468–69
 test, 369
 transformations, 475
 Tukey–Kramer test, 477–78,
 479–81, 489–90
 two groups, 474
 unplanned comparisons, 477–78
 variance components, random
 effects, 484–85
“And” versus “or,” in probability, 130

Angular transformation, 382
ANOVA. *See* Analysis of variance
Antibodies, red-winged blackbird,
330–31
Arcsine transformation, 382–83, 567
Aspirin
 treatment of cancer, 235–36
 willow bark, 24
Astronaut height, 281–82

B

Babies
 birth day of the week, 204
 birth weight, normal distribution,
 274–75, 283–84
 growth, breast-fed vs formula,
 16, 202
 resemblance to mothers and
 fathers, 345–46
Bar graph, 31–32
 grouped bar graph, 41
Bats, vampire, 250–51
Bayes’ theorem, 138–40
Bell-shaped curve
 normal distribution, 274–76
Bias, 5–6
 reduce, in experimental design,
 428–31
 volunteer, 10–11
Binomial coefficient, 181
Binomial distribution, 180–83
 derivation, 191–92
 formula, 180–81
 normal approximation, 289–92
Binomial test, 185–88, 368
 normal approximation, 289–92
Biodiversity
 alien trout species, 342–43
 biomass in marine reserves,
 375–76
 fish species diversity in desert
 pools, 566
 plant species diversity and pond
 productivity, 570–71
 plants, South American, 47
 sexual conflict and insect
 diversification, 385–86

species-area relationship, 564
stability of plant biomass
 production, 557, 611
zooplankton diversity with fish, 612
Birds
 blue tit, inheritance of cap color,
 565
 Galápagos boobies being bad, 511
 Galápagos finch beaks, 14
 great tit, malaria, 40–42
 juncos, tail feather white and
 latitude, 563
 loggerhead shrike skewers lizards,
 335–36
 red-winged blackbird, testosterone
 and immunocompetence,
 330–31
 trematode parasite or egrets,
 herons, 244
Births
 baby weights, normal distribution,
 274–75, 283–84
 days of the week, 204
Bisphenol-A, effects on rat sperm
 production, 439–40
Bivariate normal distribution, 517–18
Blinding, in experimental design,
432–33
Blocking, in experimental design,
437–40
Blood
 hemoglobin concentration at high
 altitude, 44–46
 probability of O– and O+ blood
 types, 125–26
 smoking and high blood pressure,
 129–30
Body temperature, 310–14
 normal distribution, 275
Bonferroni correction, 460
Boobies, being bad in the Galápagos,
511
Bootstrap method, 647–54
 assumptions and limitations, 654
 confidence interval, 652
 multiple groups, 652–54
 standard error, 650–51

- Box plot, 76–77
 compared to cumulative frequency distribution, 82
- Brain
 and body size in mammals, 508
 Brodman's area 44, 648–49
 speech center in chimpanzees, 648–49
 undergraduate brain sizes, normal distribution, 275
- Breast cancer cells and genetics of tumors, 694–95
- Brown recluse spider, diet of dead crickets, 289–90
- C**
- Cancer
 aspirin treatment, 235–36
 breast cancer cells and tumor subtypes, 694–95
 lung, and smoking, 16–17
- Cannibalism, during mating in crickets, 389–90, 653
- Case-control study, 241–42
- Categorical variables, 11–12
- Cats, falling in New York, 3–4
- Causation doesn't lead to correlation, 24–26
- Censored data
 in survival analysis, 690–92
- Central limit theorem, 286–88
- Chance, equal, in random sample, 7
- χ^2 contingency test, 243–50
 assumptions, 249
 calculate expected frequencies, 249
 correction for continuity, 249–50
- χ^2 distribution, 207–9
 critical value, 210–12
 degrees of freedom, 209
- χ^2 goodness-of-fit test, 204–14
 assumptions, 212–13
 binomial test, 213–14
 P-value, calculating, 209–10
 with two categories, 213
- Chimpanzee
 brain speech center, 648–49
 humans are the third, 665
- Circadian clock, light behind knees, 464–65, 613
- Clinical trial, HIV transmission, 428–29
- Coefficient of variation, 70–71
 formula, 84
- Completely randomized design, 431
- Computer-intensive methods, 643–54
 bootstrap method, 647–54
 permutation test, 395–400
 simulation, 644–47
- Conditional probability, 135–38
- Confidence interval, 105–6
 Agresti–Coull method, 190
 correlation coefficient, 509–13
 difference between means, paired data, 333
 difference between means, two samples, 347
 difference between means, Welch's, 341–42
 difference between medians, 652–54
 linear regression slope, 553
 mean, 105–6, 307–9
 median, 652
 overlap, interpreting, 347–48
 planned comparisons, 475–77
 proportion, 189–91, 283, 289
 relation to hypothesis testing, 168–69
 standard deviation, 315–16
 survival analysis, 689, 692
 2SE rule of thumb, 106
 variance, 315–16
 Wald method, 191
- Confounding variables, 15, 16, 25
- Contingency analysis, 234
- χ^2 contingency test, 243–50
- contingency table, 40–41, 236, 239, 247
 Fisher's exact test, 250–52, 254
 mosaic plot, 41–42, 234–35
 odds ratio, 234, 237–43
 relative risk, 234–37, 241–43, 253
 two categorical variables, 234
- 2 × 2 tables, 234–37
- Contingency table, 40–41, 236, 239, 247
 observed and expected frequencies, 247
- Controls in experimental design, 429, 430–31
- Correlation, 508
 assumptions, 517–18
 attenuation, 523–24
 confidence interval, 513–14
 correlation coefficient (linear), 509–13, 518–19
 depends on data range, 519
 difference from regression, 507–8
 Fisher's *z*-transformation, 513–14, 526
 intraclass (repeatability), 485
 is not causation, 24–26
 nonparametric alternative, Spearman's rank correlation, 519–23
 standard error, 512–13
 transformations, 518
 t-test of zero correlation, 526
- Covariance, 510, 549
 formula, 525
- Cricket
 cannibalism during mating, 389–90, 653
 as diet of brown recluse spider, 289–90
- Critical value, definition, 210–12
- D**
- Data
 categorical, 32–35
 censored, 690–92
 displays of, 27–51
 dredging, 459–60
 incomplete, 690–92
 numerical, 35–39
 real world, 10–11
 spreadsheets, 50–51
 tables, 48–50
 time-to-event, 688
 types of, 11–13
- de Fermat, Pierre, 201
- Degrees of freedom, 209
 χ^2 distribution, 209
 F-distribution, 349
 t-distribution, 305–6
- Dependent events, 133–35
- Deserts
 fish species diversity in pools, 564–67
 locust swarms, 30
- Design of experiments, 427–48
 balanced sample sizes, 436
 blinding, 432–33
 blocking, 437–40
 clinical trials, 428–29
 completely randomized design, 431
 controls, 300–1, 430

- experimental studies, 25
 experimental unit, 429–30, 433, 434
 extreme treatments, 439–40
 factorial design, 441–43
 interspersed of treatments, 432, 435, 437
 observational studies, 25–26
 paired design, 437
 paired vs two independent samples, 328–29
 placebo, 300–301, 428–29, 431
 power and sample size
 (see Sample size planning)
 precision and sample size
 (see Sample size planning)
 randomization, 8–9, 431–32
 randomized block design, 438–39
 replication, 434–36
 sample size planning (see Sample size planning)
 sampling error, reduce effects of, 433–40
 which test to use?, 368–69
 Design of observational studies, 442–44
 adjust for known confounding variables, 443–44
 matching, 443–44
 sample size planning, 444–46
 Difference between means
 paired data, 330–31, 333
 two-sample data, 352
 Difference between medians, 652–54
 Disease
 AIDS, 428–29
 Ebola virus outbreak, 689–90
 Distribution. *See also* Normal distribution
 cumulative frequency, 80–82
 formula for, 276–78
 frequency, 13–15
 location and spread, 65, 77–78
 mode, 37
 normal, 14–15
 probability, 13–15, 119–21
 sampling distribution, 98–103
 Dose–response curve, 572–76
 LD50, 575
 slope and intercept, 574
 Down syndrome, 139–40
 Driving accidents, as effect of
 Toxoplasma, 241–42
- E**
 Ear length and age, 556
 Ebola outbreak, 689–90
 Ecstasy (MDMA), mistakes, 505
 Education spending, 31
 Efron, Bradley, 648
 Elephants
 density in area, 673
 tusk length, 117
 Error bars, 107–9
 Error mean square (ANOVA), 466, 467, 470–71
 Errors, Type I and Type II, 162–63
 Estimate, 1–2
 Estimation, 1–2, 97–118
 bias, 5–6
 precision, 5–6, 98
 Events, 119–20
 mutually exclusive, 122–23, 125–27
 Experimental artifacts, 25
 Experimental design. *See* Design of experiments
 Experimental mistakes, 505–506
 Experimental study, 15–16
 bias, reducing, 430–33
 clinical trials, 428–30
 data loss, planning for, 448
 factors, more than one, 440–42
 vs observational study, 442–44
 sample size, 444–48
 sampling error, reducing, 433–40
 Experimental unit, 429–30
 Explanatory variable, 13
 Extinction, marine invertebrates, in time, 217
- F**
 Factorial (!), 181
 Factorial design, 441–42, 449
 analyze, 618–23
 False discovery rate, 461
F-distribution, 467, 471–72, 488
 critical values, 472
 degrees of freedom, 472
 variance ratio, 471–72
 Fish
 cod collapse, 9
 diversity in desert pools, 565
 effects of alien brook trout on salmon, 342–43
 guppy attractiveness, 43
 guppy, cold tolerance, 572
 sockeye salmon body mass, 38–39
 stickleback fish and *Ectodysplasin* gene, 77–78
 trematode parasite of killifish, 244
 zooplankton diversity with fish, 616
 Fisher, Ronald, 25, 201, 250
 Fisher's exact test, 250–52
 formula, 254
 Fisher's *z*-transformation, 513–14, 526
 Fixed effects vs random effects, 481–82
 Flu, Spanish, 286–87
 Fly
 bristle number, normal distribution, 275–76
 long-proboscid fly pollination, 567
 stalk-eyed, staring matches, 307–8
 Fossil record, marine invertebrate mass extinction, 217
 Frequency
 cumulative frequency distribution, 80–82
 frequency distribution, 14, 32, 37
 relative frequency, 32
 Frequency table, 32–33, 35–36
 contingency table, 40–41
 mean and standard variation, measuring, 71–72
 observed and expected frequencies, 205, 247
 Frogs
 deformities (image), 427
 handedness in European toad, 154–55
 pesticide effects, 441–42
F-test
 analysis of variance, 482
 comparing two variances, 348–49
 formula, 353
 Fungi, mycorrhizal carbon transfer, 478–79
- G**
 Galápagos
 beak of the finch, 14
 boobies being bad, 511
 Galton, Francis, 201, 562
 General linear models. *See* Linear models

Genetics

- baby resemblance to mothers and fathers, 345–46
 - DNA fingerprinting, elephant dung, 673
 - Down syndrome, 139–40
 - gene content of X chromosome, 213
 - genetic testing, 139
 - handedness in flowers of mud plantain, 163–64, 182
 - Human Genome Project, 98
 - inbreeding, 49–50
 - inbreeding in wolves, 514–15
 - inheritance of cap color in blue tit, 565
 - inheritance of male guppy attractiveness, 43
 - inheritance of pod color in Mendel's peas, 130–31
 - length of human genes, 98–99
 - of schizoaffective disorder, chromosome 1, 666
 - spermatogenesis in mouse, 185–86
 - stickleback fish and *Ectodysplasin* gene, 77–78
 - tumor subtypes in breast cancer cells, 694–95
 - X chromosome and spermatogenesis in mouse, 185–86
- Geometric mean, 383–84
- Goodness-of-fit test
- assumptions, 212–13
 - χ^2 goodness-of-fit test, 204–14, 368, 369
 - Poisson distribution, 215–16
 - proportional model, 204–12
 - Shapiro–Wilk test, 377
- Gosset, William, 304
- Government spending, 31
- Graphs. *See also* Spreadsheets; Tables
- bar graph, 31–35
 - box plot, 76–77
 - chartjunk, 29, 30
 - for the color blind, 32
 - common mistakes, 28–29
 - cumulative frequency distribution, 80–81
 - effective graphs, 29–32
 - error bars, 107–9
 - funnel plot, 518
 - grouped bar graph, 41

- histogram, 35–39, 45–46
 - interaction plot, 441–42, 619, 625
 - line graph, 46–47
 - map, 47–48
 - mosaic plot, 41–42
 - multiple histograms, 44–46
 - pie chart, 35
 - residual plot, 564–65
 - scatter plot, 43
 - smoothing, 564, 571–72
 - strip chart, 32, 44–45
 - violin plot, 45
 - wedge, 27
 - zero baseline, 32
- Group mean square (ANOVA), 466, 467, 470, 485
- Guinness Brewing Company, 304
- Guppy
- cold tolerance, 572
 - male ornamentation and attractiveness, 43

H

- Handedness
- European toads, 154–55
 - mud plantain flowers, 163–64
- Haphazard choice vs random, 644–45
- Hazard rate, 695–99
- formula, 701, 702
- Hazard ratio, 694–97
- Hazard ratio calculation, 697–99
- Height of astronauts, 281–82
- Hemoglobin concentration at high altitude, 44–46
- Herbivory effect on intertidal algae, 619–20
- Histogram, 35–39
- bimodal, 37
 - compared to cumulative frequency distribution, 82
 - frequency distributions in, 38–39
 - mode, 37
 - multiple histograms, 44–46
 - shape, 37
- HIV transmission, 428–29
- Human Genome Project, 98
- Huxley, Thomas Henry, 665
- Hypergeometric distribution, 674
- Hypothesis testing, 2, 151–78
- alternative hypothesis, 154, 205–6
 - best practices, 161
 - confidence interval use, 161, 163, 168–69

- critical value, 210–12
- four steps, 155
- interpret nonsignificant result, 165–66
- null distribution, 157–58
- null hypothesis, 153, 205–6
- one-sided vs two-sided tests, 166–68
- power, 162–63
- P*-value, 158–60
- reject or do not reject, 154, 163–64
- significance level (α), 160–61
- state the hypotheses, 155
- statistical significance, 160–61
- statistical vs scientific hypotheses, 152–54
- test statistic, 155, 156
- Type I and Type II errors, 162–63

I

- Inbreeding, 49–50, 514–15
- Independence, 7
- independent events, 128–31
 - independent variables, 234
 - pseudoreplication, 117–18
 - using correct sampling units, 342–45
- Indian rope trick, 520
- Influenza, Spanish, 286–87
- Insects
- butterfly and rainfall, data dredging, 459–61
 - cabbage white butterfly, wasp parasite, 667–68
 - cockroach neurons, firing rate, 565
 - dead crickets eaten by spiders, 289–90
 - fly bristle number, normal distribution, 275–76
 - in tree holes, 438–39
 - jewel wasp sex ratio, 133–35
 - locust serotonin, 30, 107
 - long-proboscid fly pollinating iris, 567
 - sage cricket, cannibalism during mating, 389–90, 653
 - sexual conflict and speciation, 385–86
 - stalk-eyed fly, staring battles, 307–8
 - walking-stick, repeatability of femur measurements, 482–83

- wasp, parasite of butterfly eggs, 667–68
- Interaction between variables, 441
estimating and testing (*see* Linear models)
visualizing, 442, 612, 617, 621
- Interaction plot, 441–42, 619, 621
- Interquartile range, 75–76
compared with standard deviation, 80
effect of measurement scale, 85
- Intervals. *See also* Confidence interval
Sturges's rule of thumb, 39
- Intraclass correlation (repeatability), 485
- Invertebrates
insects (*see* Insects)
intertidal herbivory, 619–20
marine invertebrate extinctions in time, 217
population density and body mass in streams, 519
spiders (*see* Spiders)
trematode parasite of snails, 244
zooplankton diversity and fish, 616
- J**
- Juncos, tail feather whiteness vs latitude, 563
- K**
- Kaplan–Meier method, 691–95
assumptions, 693–94
hazard ratio, 695–96
hazard ratio calculation, 697–99
logrank test, 698–99
tumor subtypes in breast cancer cells, 694–95
- Knees who say night, 464–65, 613
- Kruskal–Wallis test, 369, 475, 487
- L**
- Lansoprazole treatment for ulcers, 300
- Levene's test
compare variances, 348–50
formula, 353–54
- Life expectancy and TVs per person, 24–25
- Likelihood, 664
bias, 676
binomial model, 668
definition, 664
estimate population size using mark-recapture, 673–76
hypergeometric model, 674
likelihood-based confidence interval, 671–72
log-likelihood, 669–70
log-likelihood curve, 671–72, 675
log-likelihood ratio test, 676–78
maximum likelihood estimate, 675–676
maximum likelihood estimation, 667–72
uses in biology, 665–67
- Line graph, 46–47
- Linear models
analysis of variance, 621
analyze covariates, 623–27
analyze experiment with blocking, 615
analyze factorial design, 618–23
analyze randomized block design, 615–18
assumptions, 627–28
categorical explanatory variables, 612–14
definition, 610
fit null model vs full model, 611
fixed vs random effects, 623
F-test, 613, 621
interaction term, 618–24, 625–27
linear regression, 610–13
multiple regression, 615
numeric explanatory variables, 612
residual plot, 628
- Linear regression. *See* Regression, linear
- Lions, black on nose, 546–47
- Lizard, horned, skewered by shrikes, 335–36
- Log transformation, 379–82, 566–67
- Logistic regression, 572–76
analysis of deviance, 573
*LD*50, 575
log-odds, 574
slope and intercept, 574
- Logit function, 574
- Log-likelihood ratio test, 676–78
- Log-odds, in logistic regression, 574
- Log-odds ratio. *See* Odds ratio
- Logrank test, 698–99
formula, 702
- M**
- Magic Indian rope trick, 520
- Malaria, great tit, 40–42
- Mammals
black noses, 546–47
cats falling in New York, 3–4
chimpanzee brain speech center, 648–49
elephant density, 673
fur seals stretch and shrink, 571–72
great apes and human ancestry, 665
inbreeding in wolves, 514–15
mole rats, castes, 624–25
mouse spermatogenesis genes, 185–86
rat sperm production and Bisphenol-A, 439–40
tigers, deaths from, 33–35
tusk length of elephants, 117
vampire bats attack cows in estrus, 250–51
- Mann–Whitney *U*-test, 369, 388–93
assumptions, 393–94
normal approximation, 393
power, 393
tied ranks, 392–93
- Map, 47–48
- Marine reserves, 375–76, 381
- Mass extinction, marine invertebrate fossil record, 217
- Maximum likelihood. *See* Likelihood
- Median survival time, 693
- Mean, 66–68
analysis of variance, 464–66
calculate from frequency table, 71–72
confidence interval, 307–9
effect of measurement scale, 72–73, 85
formula, 84
geometric mean, 383–84
mean vs median, 78–80
normal sampling distribution, 283–85
paired difference, 332, 334–35
population mean, 97
sample mean, 67–68, 83
sampling distribution, 283–85
standard error, 103–4, 284
t-test, one-sample, 310–14
t-test, paired, 333–34
t-test, two-sample, 338–40

- Mean squares
 in ANOVA, 460–71
 in linear regression, 553, 558, 559–60
- Measles outbreaks, 47–48
- Measurement error
 effect on correlation, 523–24
 effect on linear regression, 568
 estimate repeatability, 485–86
- Median, 73–75
 confidence interval, 652
 effect of measurement scale, 85
 formula, 84
 mean vs median, 78–80
 standard error, 651
- Memory and rope trick, 520
- Mendel, Gregor, 130–31
- Meta-analysis, 598–600
 confidence interval of effect size, 599
 define the question, 598
 degrees of freedom, 600
 effect size, 599
 estimates of sizes of errors, 600
 funnel plot, 518
 make your work accessible to meta-analysis, 600
 publication bias, 540–43
- Mistakes
 common graphical mistakes, 29–30
 experimental, 505–506
 fallacy of indirect comparison, 345–47
 regression fallacy, 562
 statistical, 505–506
 use the correct sampling units, 342–45
- Mode, 37
- Model, 610
- Model, probability. *See* Probability model
- Models, linear. *See* Linear models
- Mole rat castes, 624–65
- Mosaic plot, 41–42, 234–37, 242–45
- Mount Everest climbers and variables, 15–16
- Mouse
 genome, 185–86
 spermatogenesis genes on X chromosome, 185–86
- Multiple explanatory variables, analyzing, 610
- Multiple regression, 615
- Multiple testing
 Bonferroni correction, 460
 data dredging, 459–61
 false discovery rate, 461
 inflation of Type I error rate, 459–60
 Tukey–Kramer test, 477–81, 489–90
- Multiplication rule, 128–31
 general multiplication rule, 137
- Mutually exclusive events, 122–23, 125–27
- N**
- Newt, red-spotted, 442
- Nightingale, Florence, 27
- Nixon, Richard, 505
- Nominal variables, 11–12
- Nonlinear regression, 569–72
 asymptotic curve, 570–71
 formula-free curve fitting, 571
 Michaelis–Menten curve, 569–70
 parabola, 570
 quadratic curve, 570–71
 smoothing, 571–72
- Nonparametric methods, 384
 assumptions, 393–94
 bootstrap, 647–54
 Kruskal–Wallis test, 475, 488
 Mann–Whitney *U*-test, 388–93
 permutation test, 395–99
 power, 394–95
 sign test, 385–88
 Spearman's rank correlation, 519–23
 Type I and Type II error rates, 394–95
 Wilcoxon signed-rank test, 388
- Normal distribution, 14–15, 273–92
 approximation to binomial distribution, 289–92
 bell-shaped curve, 274–76
 bivariate normal distribution, 517–18
 critical values, 289
 formula, 276–78
 normal quantile plot, 373–75
 Shapiro–Wilk test, 377
 standard normal (*Z*) distribution, 278–83
- Null distribution, 156–58
- Null hypothesis, 153
- Null model, 611–14
- Numerical variables, 12–13
 and categorical variables, 43–45
 graphs for data, 39
- O**
- Observational study, 15–16, 25–26, 442–44
- Odds, 237–39
- Odds ratio, 238–43
 confidence interval, 240–41
 formula, 240
 vs relative risk, 241–43
 standard error, 240–41
- One-sample *t*-test, 310–14, 369
 assumptions, 314
 critical values, 312
 effect of larger sample, 313–14
 nonparametric alternative, sign test, 385–88
- One-sided vs two-sided tests, 166–68
- Ordinal variables, 12
- Outliers
 causes, 38
 data entry errors, 51
 definition, 38
 indicate non-normal distribution, 373, 386–87
 indicate use of nonparametric method, 372, 378, 383
 influence on linear regression, 563–64
 sensitivity of mean and median, 79–80
 violate assumption of bivariate normality, 518
 violate assumption of normality, 372, 378
- P**
- Paired measurements, 330–31
- Paired *t*-test, 333–34, 369
 assumptions, 334–35
 nonparametric alternative, sign test, 385–88
 nonparametric alternative, Wilcoxon signed-rank test, 388
- Parameters, 2, 99
- Pascal, Blaise, 201
- Pearson, Karl, 201, 508
- Percentiles, 80–81
 cumulative frequency distribution, 81–82

- Permutation test, 395–99
assumptions, 399
steps, 397
- Pesticides, effects on frogs, 441–42
- Phylogenetic non-independence, 640–41
- Phylogenetically independent contrasts, 641
- Phylogeny
human ancestry, 665–66
lilies, 639–40
- Phytoplankton growth and iron, 569
- Pie chart, 35
- Placebo effect, 300–301
- Planned comparisons, 475–77, 489
- Plants
biodiversity and stability, 557, 611
biodiversity, South American, 47
carbon transfer via fungi, 478–79
Harvard Forest, 8
inheritance of pea pod color, 130–31
intertidal algae and herbivory, 619–20
iris, pollination by long-proboscid flies, 567
lilies, habitat and flower type, 639–42
maize growth, 28–29
mud plantain, handedness, 163–64, 182
phytoplankton growth and iron, 569
species diversity and pond productivity, 570–71
tree-hole insects, 438–39
- Poisson distribution, 215–21
formula, 216
successes clumped or dispersed, 215–16
variance-to-mean ratio, 220–21
- Poisson, Siméon Denis, 216
- Polio, 151–52
vaccine, 10
- Pooled sample variance, 337–39
- Population, 4–5, 97
- Population size estimation using likelihood, 673–74
- Power, 162–63, 446–47
parametric vs nonparametric methods, 394–95
sample size planning (*see* Sample size planning)
- Precision, 5–6, 98
sample size planning (*see* Sample size planning)
- Probability, 119–21
addition rule, 125–27
“and” versus “or,” 130
Bayes’ theorem, 138–40
conditional probability, 135–38
dependent events, 133–35
events, 119–20
general addition rule, 127
general multiplication rule, 137
multiplication rule, 128–31
mutually exclusive events, 122–23, 125–27
random trial, 120
trees, 131–33, 134, 135
Venn diagram, 121–22
- Probability distribution, 122–25
binomial, 180–85, 214, 668
bivariate normal, 517–18
 χ^2 , 207–9
continuous, 123
discrete, 121
 F , 472
hypergeometric, 674
normal, 274–78
Poisson, 215–21
- Probability model, 203
binomial distribution, 180–85
Poisson distribution, 215–21
proportional model, 204–5
random in space or time, 215–21
- Proportion, 65, 82–83
binomial test, 185–88
confidence interval, 190–91
formula, 84
maximum likelihood estimate, 675
odds, 241–43
sampling distribution, 183–85
standard error, 184–85, 188–89
- Proportional hazards, 700
- Proportional model, 204–5
- Pseudoreplication, 117–18
using correct sampling units, 118, 342–45
- Publication bias, 540–44
- P -value, 158–60
best practices, 161
- Q**
Quadratic regression, 570
Quantiles, 80–81
cumulative frequency distribution, 81–82
interquartile range, 75–76
median, 73–75
normal quantile plot, 373–75
quartiles, 75–76
Quartiles, 75–76
- R**
 R^2
analysis of variance, 473
linear regression, 559–60
- Random
in space or time, 215–21
number generator, 8
numbers, inability of people to generate, 645
sample, 7–9
trial, in probability, 120
- Random-effects ANOVA, 482–86
- Randomization, in experimental design, 431–32
- Randomization test. *See* Permutation test
- Randomized block design, 438–40
analyze, 615–17
- Ranks, methods based on. *See* Nonparametric methods
- Rats
mole rats, castes, 624–65
sperm production and Bisphenol-A, 439–40
- Regression, 545–46
application to two study designs, 546
difference from correlation, 546, 557
fallacy, 562
regression toward the mean, 561–62
- Regression, linear, 546–53
ANOVA approach, 560
assumptions, 562–65
calculate slope and intercept, 549
confidence bands, 554–56
confidence interval, slope, 553
confidence in predictions, 554–56
detect nonlinearity, 564
detect non-normality, 564–65
detect unequal variances, 564–65

- Regression, linear (*continued*)
 effect of measurement error, 568
 formula for the line, 548
 least squares method, 548
 as a linear model, 610–11
 outliers, 563–64
 population regression, 551
 predicted values, 551–52
 prediction intervals, 555
 R^2 , 560
 regression toward the mean, 560–62
 residual plot, 564–65, 566–67
 residuals, 552–53
 standard error, slope, 553
 transformations, 566–67
 transform exponential function, 566
 transform power function, 566
 t -test of regression slope, 557–59
- Regression, logistic, 572–76
 analysis of deviance, 575
 LD_{50} , 575
 slope and intercept, 574
- Regression, multiple. *See* Linear models
- Regression, nonlinear, 569–72
 asymptotic curve, 569–71
 detecting, 564–65
 formula-free curve fitting, 571–72
 Michaelis–Menten curve, 569–70
 parabola, 570
 quadratic curve, 570–71
 relationships, 569–72
 smoothing, 571–72
- Relative risk, 234–37, 241–43, 253
 confidence interval, 253
 vs odds ratio, 241–43
 standard error, 253
- Repeatability, 485–86
 assumptions, 486
- Replication, 434–36
- Reptiles
 horned lizards skewered by shrikes, 335–36
 paradise tree snake, gliding, 66–67
- Residual plot, 564–65, 566–67
- Residuals, 565–68
- Response variable, 13
- Robust statistical procedures, 377
- Rope trick and memory of, 520
- Rounding, 70
- S**
- Salk, Jonas, 151
- Salmon
 bimodal body mass distribution in sockeye, 38–39
 chinook, effects of alien brook trout, 342–43
- Sample, 4–7
 of convenience, 9–10
 from a population, 97
 random, 7–9
 volunteer, 10–11
- Sample size planning
 binomial test, 451
 confidence interval of a log-odds ratio, 449–50
 confidence interval of a mean, 450
 confidence interval of a proportion, 449
 confidence interval of difference between two means, 450–51
 confidence interval of difference between two proportions, 450
 one-sample or paired t -test, 451–52
 plan for data loss, 448
 plan for power, 446–47, 451–52
 plan for precision, 444–46
 two-sample t -test, 452
 2×2 contingency test, 451
- Sampling, 3–11
 bias, 5–6
 sampling error, 5–6
 sampling unit, 4–5
 without replacement, 138
 with replacement, 138, 650
- Sampling distribution, 99–100
 bootstrap sampling distribution, 647–52
 of means, 283–85
 under the null hypothesis, 156
 proportion, 183–85
 sample mean, 67–68, 101–2
- Sampling error, 5–6
 reduce effects of, in experiment, 429, 433–40
- Scatter plot, 43
 smoothing, 563–64, 571–72
- Schizoaffective disorder, genetics, 666
- Seals that stretch and shrink, 571–72
- Serotonin, effect on locust behavior change, 30, 107
- Sex ratio
 jewel wasps, 133–34
 turtles, green sea, 189
 in two-child families, 131–33
- Sexual conflict and insect diversification, 385–86
- Shapiro–Wilk test, 369, 377
- Sign test, 369, 385–88
 power, 394–95
- Simulation
 hypothesis testing, 644–47
 null distribution, 645–47
- Skew, 38
- Smoking, 16–17
 and high blood pressure, 129–30
 and lung cancer, 16–17
- Snail, trematode parasite, 244
- Snakes
 paradise tree snake gliding, 66–67
- Spearman's rank correlation, 519–23
 assumptions, 523
 procedure for large n , 522–23
 test of no correlation, 520–22
- Species as data points, 639–42
- Sperm
 effect of Bisphenol-A, 439–40
 spermatogenesis genes in mouse, 185–86
- Spiders
 brown recluse, diet of dead crickets, 289–90
 distribution of, after Mount St. Helens eruption, 215
 pedipalp amputation and running speed, 74
- Spreadsheets, 50–51
- Square-root transformation, 379, 383, 567
- Stalk-eyed fly, 307–8
- Standard deviation, 68–70
 calculate from frequency table, 71–72
 compared with interquartile range, 80
 confidence interval, 315–16

- effect of measurement scale,
72–73, 85
formula, 84
Standard error, 103
bootstrap, 647–54
correlation coefficient, 509–13
difference between means, paired
data, 332, 351
difference between medians, 652–54
difference between two means,
two-sample data, 339, 351
linear regression slope, 553
mean, 103–4, 284
median, 651
proportion, 184–85, 188–89
Standard normal (*Z*) distribution. *See*
Normal distribution
Statistical hypothesis, 2
Statistical significance
vs biological importance, 177–78
significance level α , 160–61
Statistics, defined, 1–2
Stickleback fish and *Ectodysplasin*
gene, 77–78
Strip chart, 32, 39, 44–45
Student's *t* distribution. *See*
t-distribution
Study
case-control study, 241–43
experimental vs observational,
15–16, 442–44
Sturges's rule of thumb, 39
Sum of squares, 69
in ANOVA, 468–69
formula, 84
in linear regression, 553, 560, 578
Survival analysis
assumptions, 700
defined, 688
hazard ratio, 694
Kaplan–Meier curves, 688–94
logrank test, 698–99
logrank test formula, 702
proportional hazards, 700
survival curves, 688–94
- T**
Tables
contingency, 40
display, 48–50
effective, 48–51
frequency, 32–34, 35–36
standard normal (*Z*), 279
t-distribution, 304–7
critical values, 306
degrees of freedom, 305–6
Temperature, human body, 310–11
Testosterone and immunocompetence
in blackbirds, 330–31
Tigers, deaths from, 33–35
Time
random, in probability, 215–21
survival analysis, 687–700
time-to-event data, 688
Tit, blue, inheritance of cap color,
565
Tit, great, malaria, 40–42
Titanic, survival of men and women,
234, 235
Toads, handedness, 154–55
Toxoplasma, and driving accidents,
241–42
Transformations, 379–84
angular, 382
arcsine, 382–83, 567
confidence intervals, 383
correlation analysis, 518
log, 379–82, 566–67
square-root, 379, 383, 566–67
transform exponential function, 566
transform power function, 566
Tree, probability, 131–33, 134, 135
Trees and carbon transfer via fungi,
478–79
tree-hole insects, 438–39
Trematode, bird parasite, 244
Trout, effects on salmon, 342–43
t-test
correlation coefficient, 509–13,
518–19
mean, one-sample (*see*
One-sample *t*-test)
mean, paired-sample (*see* Paired
t-test)
means, two-sample (*see*
Two-sample *t*-test)
regression slope, 557–59
Tusk length of elephants, 117
Tukey–Kramer test, 477–81
assumptions, 481
Tumor subtypes in breast cancer cells,
694–95
Turtles, green sea, sex of, 189
Two-factor ANOVA
fixed effects, 623
random effects, 623
Two-sample *F*-test
comparing two variances, 348–49
Two-sample *t*-test, 338–40, 369
assumptions, 340–41
fallacy of indirect comparison,
345–47
formula, 353
nonparametric alternative,
Mann–Whitney *U*-test, 388–93
unequal standard deviations, 378
violations of assumptions,
377–78
Welch's *t*-test, 344–45
2SE rule of thumb, 106
2 × 2 contingency test, 451
2 × 2 tables, 234–37
Two-way ANOVA
fixed effects, 615
random effects, 623
Type I and Type II errors, 162–63
- U**
Ulcers, lansoprazole treatment, 300
Unplanned comparisons, 476,
477–81
U-test. *See* Mann–Whitney *U*-test
- V**
Vaccine
measles outbreaks, 46
polio, 10, 151–52
refuted link to autism, 46
Vampire bats, 250–51
Variables, 11–13
categorical, 11–12
confounding, 15, 16–17, 25, 443–44
explanatory, 13
interaction between, 441–42
multiple explanatory variables,
610
nominal, 11–12
numerical, 12–13
ordinal, 12
types of, 11–13
Variance, 68–70
calculate from frequency table,
71–72
components, random-effects
ANOVA, 484–85
confidence interval, 315–16
effect of measurement scale,
72–73, 85
formula, 84

Variance (*continued*)

- F*-test to compare two variances, 348–49, 353
- Levene's test of equal variances, 349, 353
- pooled sample, 337–38
- Variance components, random-effects ANOVA, 484–86
- Venn diagram, 121–22
- Violence and ice cream, 24–25
- Violin chart, 39, 44–45

W

- Wald method, 191
- Walking-stick, repeatability of femur measurements, 482–83
- Weighted average, 337

- Welch's confidence interval for difference between means, 344–45
 - formula, 352
- Welch's *t*-test, 341–42, 369
 - formula, 353
- Which test to use?, 368–69
- Wilcoxon rank-sum test. *See* Mann–Whitney *U*-test
- Wilcoxon signed-rank test, 388
- Willow bark and pain relief, 24
- Wolves, inbreeding, 514–15
- Wood wide web, 478

X

- χ^2 . *See* section “C”
- X chromosome

- gene content, 213
- spermatogenesis genes in mouse, 185–86

Y

- Yates correction for continuity, 249–50

Z

- Z distribution. *See* Normal distribution: standard normal
- Zika virus infection effects on fetuses, 36–37
- Zooplankton, diversity with fish, 616