UC San Diego

Trilinear estimates for quadratic dispersive equations


Vitor Borges
includes work with Ioan Bejenaru and Tiklung Chan
University of California San Diego



UC Berkeley – HADES
October 27, 2026

Introduction

What is dispersion?



Different frequencies move at different speeds, and the wave disperses!

Quadratic semilinear equations

Schematically, a quadratic and dispersive equation has the form

$$(i\partial_t + L(D))u = N(u),$$
$$u(0) = u_0$$

Examples: Schrödinger (Δ), Klein–Gordon ($\langle D \rangle$), wave ($|D|$).

Physics: optics, quantum mechanics, gravitation, water waves…

Motivating question

For which initial data do we get global-in-time solutions? How do these solutions behave as $t \to \infty$?

Quadratic semilinear equations

Schematically, a quadratic and dispersive equation has the form

$$(i\partial_t + L(D))u = N(u),$$
$$u(0) = u_0$$

Examples: Schrödinger (Δ), Klein–Gordon ($\langle D \rangle$), wave ($|D|$).

Physics: optics, quantum mechanics, gravitation, water waves…

Motivating question

For which initial data do we get global-in-time solutions? How do these solutions behave as $t \to \infty$?

Quadratic semilinear equations

Schematically, a quadratic and dispersive equation has the form

$$(i\partial_t + L(D))u = N(u),$$
$$u(0) = u_0$$

Examples: Schrödinger (Δ), Klein–Gordon ($\langle D \rangle$), wave ($|D|$).

Physics: optics, quantum mechanics, gravitation, water waves…

Motivating question

For which initial data do we get global-in-time solutions? How do these solutions behave as $t \to \infty$?

Quadratic semilinear equations

Schematically, a quadratic and dispersive equation has the form

$$(i\partial_t + L(D))u = N(u),$$
$$u(0) = u_0$$

Examples: Schrödinger (Δ), Klein–Gordon ($\langle D\rangle$), wave ($|D|$).

Physics: optics, quantum mechanics, gravitation, water waves…

Motivating question

Under what conditions do small initial data u_0 give rise to global-in-time solutions that asymptotically approach free waves?

$$\|u(t) - e^{itL(D)}u_*\|_{H^s_x} \longrightarrow 0 \quad \text{as} \quad t \to \infty.$$

where $u_{\text{free}}(t) = e^{itL(D)}u_*$ solves $(i\partial_t + L(D))u_{\text{free}} = 0$ with $u_{\text{free}}(0) = u_*$.

Why should one expect scattering?

$$(i\partial_t - \Delta)u = \underbrace{N(u)}_{\text{nonlinear forcing}}$$

Small accumulated forcing $\longrightarrow$ small correction to free evolution.

Linear dispersion

$$(i\partial_t - \Delta)u_{\text{free}} = 0$$

$$\|u_{\text{free}}(t)\|_{L^\infty_x} \lesssim (1+t)^{-n/2}.$$

Decay weakens quadratic forcing.

Possible ODE buildup

$$\partial_t v = v^2$$

$$v(t) = \frac{c}{T-t}.$$

Blows up as $t \longrightarrow T^-$.

> **Can dispersion win?** For small data, hope the late-time forcing becomes negligible and the solution approaches a free wave.

Why should one expect scattering?

$$(i\partial_t - \Delta)u = \underbrace{N(u)}_{\text{nonlinear forcing}}$$

Small accumulated forcing $\longrightarrow$ small correction to free evolution.

Linear dispersion

$$(i\partial_t - \Delta)u_{\text{free}} = 0$$

$$\|u_{\text{free}}(t)\|_{L^\infty_x} \lesssim (1+t)^{-n/2}.$$

Decay weakens quadratic forcing.

Possible ODE buildup

$$\partial_t v = v^2$$

$$v(t) = \frac{c}{T-t}.$$

Blows up as $t \longrightarrow T^-$.

> **Can dispersion win?** For small data, hope the late-time forcing becomes negligible and the solution approaches a free wave.

Why should one expect scattering?

$$(i\partial_t - \Delta)u = \underbrace{N(u)}_{\text{nonlinear forcing}}$$

Small accumulated forcing $\longrightarrow$ small correction to free evolution.

Linear dispersion

$$(i\partial_t - \Delta)u_{\text{free}} = 0$$

$$\|u_{\text{free}}(t)\|_{L_x^\infty} \lesssim (1+t)^{-n/2}.$$

Decay weakens quadratic forcing.

Possible ODE buildup

$$\partial_t v = v^2$$

$$v(t) = \frac{c}{T-t}.$$

Blows up as $t \longrightarrow T^-$.

Can dispersion win? For small data, hope the late-time forcing becomes negligible and the solution approaches a free wave.

Why should one expect scattering?

$$(i\partial_t - \Delta)u = \underbrace{N(u)}_{\text{nonlinear forcing}}$$

Small accumulated forcing $\longrightarrow$ small correction to free evolution.

Linear dispersion

$$(i\partial_t - \Delta)u_{\text{free}} = 0$$

$$\|u_{\text{free}}(t)\|_{L^\infty_x} \lesssim (1+t)^{-n/2}.$$

Decay weakens quadratic forcing.

Possible ODE buildup

$$\partial_t v = v^2$$

$$v(t) = \frac{c}{T-t}.$$

Blows up as $t \longrightarrow T^-$.

Can dispersion win? For small data, hope the late-time forcing becomes negligible and the solution approaches a free wave.

Literature

Schrödinger equations

Colliander–Keel–Staffilani–Takaoka–Tao (2002)
Germain–Masmoudi–Shatah (2009)
Bejenaru–Ionescu–Kenig–Tataru (2011)
Herr–Tataru–Tzvetkov (2011)
Ifrim–Tataru (2015)
Ifrim–Tataru (2023)
Ifrim–Tataru (2024)

Klein–Gordon equations

Shatah (1985)
Bejenaru–Herr (2017)
Chae–Oh (2017)
Ionescu–Pausader (2019)
Bejenaru–Borges (2026)
Borges–Chan (2026)

Wave equations and GR

Christodoulou–Klainerman (1993)
Tao (2001)
Tataru (2001)
Rodnianski–Tao (2004)
Lindblad–Rodnianski (2010)
Krieger–Sterbenz–Tataru (2015)
Krieger–Tataru (2017)
Ionescu–Pausader (2022)
Klainerman–Szeftel (2023)

Water waves

Wu (2011)
Germain–Masmoudi–Shatah (2012)
Alazard–Delort (2015)
Ionescu–Pusateri (2015)
Ifrim–Tataru (2016)
Deng–Ionescu–Pausader–Pusateri (2017)
Ionescu–Pusateri (2018)

Main result

Theorem (Bejenaru–B.–Chan)

The Schrödinger equation

$$(i\partial_t - \Delta)u = c_1 u^2 + c_2 \bar{u}^2, \qquad c_1, c_2 \in \mathbb{C}, \tag{2NLS}$$

with small initial data $u_0 \in \dot{H}^{-\frac{1}{2}}(\mathbb{R}^3)$ in the scaling-critical space is globally well-posed, scatters to free solutions, and has Lipschitz dependence on its initial data. The solution is also unique in the resolution space $\dot{X}^{-\frac{1}{2}} \subset C_t \dot{H}_x^{-\frac{1}{2}}$.

Roadmap & recap: multilinear estimates

> **Goal:** Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

Roadmap & recap: multilinear estimates

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

Roadmap & recap: multilinear estimates

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

Roadmap & recap: multilinear estimates

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

(iii) Harmonic analysis: geometry of interacting waves

Roadmap & recap: multilinear estimates

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

(iii) Harmonic analysis: geometry of interacting waves

(iv) From harmonic analysis to PDEs

Roadmap & recap: multilinear estimates

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

(iii) Harmonic analysis: geometry of interacting waves

(iv) From harmonic analysis to PDEs

(v) Everything coming together

Contraction argument

$$(i\partial_t + L(D))u = F \qquad (F = N(u))$$

Goal: Find a complete space $X \subset C_t H^s_x$, and a forcing space Y such that

- If $\|u(0)\|_{H^s_x}$ is small, then $e^{itL(D)}u_0 \in X$
- Linear estimate: $\|u\|_X \lesssim \|u_0\|_{H^s_x} + \|F\|_Y$
- Nonlinear estimate: $\|N(u) - N(v)\|_Y \lesssim (\|u\|_X + \|v\|_X)^{p-1} \|u - v\|_X$

Contraction argument

$$(i\partial_t + L(D))u = F \qquad (F = N(u))$$

Goal: Find a complete space $X \subset C_t H^s_x$, and a forcing space Y such that

- If $\|u(0)\|_{H^s_x}$ is small, then $e^{itL(D)}u_0 \in X$
- Linear estimate: $\|u\|_X \lesssim \|u_0\|_{H^s_x} + \|F\|_Y$
- Nonlinear estimate: $\|N(u) - N(v)\|_Y \lesssim (\|u\|_X + \|v\|_X)^{p-1} \|u - v\|_X$

Contraction argument

$$(i\partial_t + L(D))u = F \qquad (F = N(u))$$

Goal: Find a complete space $X \subset C_t H^s_x$, and a forcing space Y such that

- If $\|u(0)\|_{H^s_x}$ is small, then $e^{itL(D)}u_0 \in X$
- Linear estimate: $\|u\|_X \lesssim \|u_0\|_{H^s_x} + \|F\|_Y$
- Nonlinear estimate: $\|N(u) - N(v)\|_Y \lesssim (\|u\|_X + \|v\|_X)^{p-1} \|u - v\|_X$

Contraction argument

$$(i\partial_t + L(D))u = F \qquad (F = N(u))$$

Goal: Find a complete space $X \subset C_t H^s_x$, and a forcing space Y such that

- If $\|u(0)\|_{H^s_x}$ is small, then $e^{itL(D)}u_0 \in X$
- Linear estimate: $\|u\|_X \lesssim \|u_0\|_{H^s_x} + \|F\|_Y$
- Nonlinear estimate: $\|N(u) - N(v)\|_Y \lesssim (\|u\|_X + \|v\|_X)^{p-1} \|u - v\|_X$

Contraction argument

$$(i\partial_t + L(D))u = F \qquad (F = N(u))$$

Goal: Find a complete space $X \subset C_t H^s_x$, and a forcing space Y such that

- If $\|u(0)\|_{H^s_x}$ is small, then $e^{itL(D)}u_0 \in X$
- Linear estimate: $\|u\|_X \lesssim \|u_0\|_{H^s_x} + \|F\|_Y$
- Nonlinear estimate: $\|N(u) - N(v)\|_Y \lesssim (\|u\|_X + \|v\|_X)^{p-1} \|u - v\|_X$

Contraction argument

$$(i\partial_t + L(D))u = F \qquad (F = N(u))$$

Goal: Find a complete space $X \subset C_t H^s_x$, and a forcing space Y such that

- If $\|u(0)\|_{H^s_x}$ is small, then $e^{itL(D)}u_0 \in X$
- Linear estimate: $\|u\|_X \lesssim \|u_0\|_{H^s_x} + \|F\|_Y$
- Nonlinear estimate: $\|N(u) - N(v)\|_Y \lesssim (\|u\|_X + \|v\|_X)^{p-1} \|u - v\|_X$

For small initial data, it is enough to establish

$$\|u\|_X \lesssim \|u_0\|_{H^s_x} + \|N(u)\|_Y, \quad \text{and} \quad \|N(u)\|_Y \lesssim \|u\|_X^p \tag{1}$$

Contraction argument

$$(i\partial_t + L(D))u = F \qquad (F = N(u))$$

Goal: Find a complete space $X \subset C_t H^s_x$, and a forcing space Y such that


- If $\|u(0)\|_{H^s_x}$ is small, then $e^{itL(D)}u_0 \in X$
- Linear estimate: $\|u\|_X \lesssim \|u_0\|_{H^s_x} + \|F\|_Y$
- Nonlinear estimate: $\|N(u) - N(v)\|_Y \lesssim \left(\|u\|_X + \|v\|_X\right)^{p-1} \|u - v\|_X$


For small initial data, it is enough to establish

$$\|u\|_X \lesssim \|u_0\|_{H^s_x} + \|N(u)\|_Y, \quad \text{and} \quad \|N(u)\|_Y \lesssim \|u\|_X^p \tag{1}$$

In fact, if X is a *reasonable* space, then (1) establishes both global well-posedness and scattering.


Higher dimensions are easier






More dimensions = more dispersion

Higher dimensions are easier




More dimensions = more dispersion

Higher dimensions are easier



More dimensions = more dispersion

Higher dimensions are easier



More dimensions = more dispersion

Linear Schrödinger: Strichartz estimates

Free evolution: $\|e^{-it\Delta}u_0\|_{L_t^q L_x^r} \lesssim \|u_0\|_{L_x^2}, \qquad \frac{2}{q}+\frac{n}{r}=\frac{n}{2}, \quad 2 \le q, r \le \infty$

With forcing: if $(i\partial_t - \Delta)u = F$, then

$$\|u\|_{L_{t,x}^p} \lesssim \|u_0\|_{L_x^2} + \|F\|_{L_{t,x}^{p'}}, \qquad p = \frac{2(n+2)}{n}.$$

Linear Schrödinger: Strichartz estimates

Free evolution: $\|e^{-it\Delta}u_0\|_{L_t^q L_x^r} \lesssim \|u_0\|_{L_x^2}, \qquad \frac{2}{q}+\frac{n}{r}=\frac{n}{2}, \quad 2 \le q, r \le \infty$

With forcing: if $(i\partial_t - \Delta)u = F$, then

$$\|u\|_{L_{t,x}^p} \lesssim \|u_0\|_{L_x^2} + \|F\|_{L_{t,x}^{p'}}, \qquad p = \frac{2(n+2)}{n}.$$

Toy model: quadratic Schrödinger in dimension 4

$$(i\partial_t + \Delta)u = u^2, \quad u : \mathbb{R}_t \times \mathbb{R}^n_x \to \mathbb{C}.$$

Goal: Construct spaces X and Y such that $\|u^2\|_Y \lesssim \|u\|_X^2$.
By duality, it is enough to establish

$$|\langle u^2, w\rangle| \lesssim \|u\|_X^2 \|w\|_{Y^*}, \qquad w \in Y^*.$$

Strichartz estimates: $\|u\|_{L^p_{t,x}} \lesssim \|u_0\|_{L^2_x} + \|u^2\|_{L^{p'}_{t,x}}$, $p = \frac{2(2+n)}{n}$

$n = 4$	$p = 3$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$p = \frac{10}{3}$	$L^{10/3} \cdot L^{10/3} \cdot L^{10/3} \not\subset L^1_{t,x}$	✗
$n = 2$	$p = 4$	$L^4 \cdot L^4 \cdot L^4 \not\subset L^1_{t,x}$	✗

> **Takeaway:** Linear information is not sufficient to establish a contraction for $n \leq 3$.

Toy model: quadratic Schrödinger in dimension 4

$$(i\partial_t + \Delta)u = u^2, \quad u : \mathbb{R}_t \times \mathbb{R}^n_x \to \mathbb{C}.$$

Goal: Construct spaces X and Y such that $\|u^2\|_Y \lesssim \|u\|_X^2$.

By duality, it is enough to establish

$$|\langle u^2, w\rangle| \lesssim \|u\|_X^2 \|w\|_{Y^*}, \qquad w \in Y^*.$$

Strichartz estimates: $\|u\|_{L^p_{t,x}} \lesssim \|u_0\|_{L^2_x} + \|u^2\|_{L^{p'}_{t,x}}$, $p = \frac{2(2+n)}{n}$

$n = 4$	$p = 3$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$p = \frac{10}{3}$	$L^{10/3} \cdot L^{10/3} \cdot L^{10/3} \not\subset L^1_{t,x}$	✗
$n = 2$	$p = 4$	$L^4 \cdot L^4 \cdot L^4 \not\subset L^1_{t,x}$	✗

> **Takeaway:** Linear information is not sufficient to establish a contraction for $n \leq 3$.

Toy model: quadratic Schrödinger in dimension 4

$$(i\partial_t + \Delta)u = u^2, \quad u : \mathbb{R}_t \times \mathbb{R}^n_x \to \mathbb{C}.$$

Goal: Construct spaces X and Y such that $\|u^2\|_Y \lesssim \|u\|_X^2$.
By duality, it is enough to establish

$$|\langle u^2, w\rangle| \lesssim \|u\|_X^2 \|w\|_{Y^*}, \qquad w \in Y^*.$$

Strichartz estimates: $\|u\|_{L^p_{t,x}} \lesssim \|u_0\|_{L^2_x} + \|u^2\|_{L^{p'}_{t,x}}$, $p = \frac{2(2+n)}{n}$

$n = 4$	$p = 3$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$p = \frac{10}{3}$	$L^{10/3} \cdot L^{10/3} \cdot L^{10/3} \not\subset L^1_{t,x}$	✗
$n = 2$	$p = 4$	$L^4 \cdot L^4 \cdot L^4 \not\subset L^1_{t,x}$	✗

> **Takeaway:** Linear information is not sufficient to establish a contraction for $n \leq 3$.

Toy model: quadratic Schrödinger in dimension 4

$$(i\partial_t + \Delta)u = u^2, \quad u : \mathbb{R}_t \times \mathbb{R}^n_x \to \mathbb{C}.$$

Goal: Construct spaces X and Y such that $\|u^2\|_Y \lesssim \|u\|_X^2$.
By duality, it is enough to establish

$$|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad w \in Y^*.$$

Strichartz estimates: $\|u\|_{L^p_{t,x}} \lesssim \|u_0\|_{L^2_x} + \|u^2\|_{L^{p'}_{t,x}}$, $p = \frac{2(2+n)}{n}$

$n = 4$	$p = 3$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$p = \frac{10}{3}$	$L^{10/3} \cdot L^{10/3} \cdot L^{10/3} \not\subset L^1_{t,x}$	✗
$n = 2$	$p = 4$	$L^4 \cdot L^4 \cdot L^4 \not\subset L^1_{t,x}$	✗

> **Takeaway:** Linear information is not sufficient to establish a contraction for $n \leq 3$.

Toy model: quadratic Schrödinger in dimension 4

$$(i\partial_t + \Delta)u = u^2, \quad u : \mathbb{R}_t \times \mathbb{R}^n_x \to \mathbb{C}.$$

Goal: Construct spaces X and Y such that $\|u^2\|_Y \lesssim \|u\|_X^2$.
By duality, it is enough to establish

$$|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad w \in Y^*.$$

Strichartz estimates: $\|u\|_{L^p_{t,x}} \lesssim \|u_0\|_{L^2_x} + \|u^2\|_{L^{p'}_{t,x}}$, $p = \frac{2(2+n)}{n}$

$n = 4$	$p = 3$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$p = \frac{10}{3}$	$L^{10/3} \cdot L^{10/3} \cdot L^{10/3} \not\subset L^1_{t,x}$	✗
$n = 2$	$p = 4$	$L^4 \cdot L^4 \cdot L^4 \not\subset L^1_{t,x}$	✗

> **Takeaway:** Linear information is not sufficient to establish a contraction for $n \leq 3$.

Toy model: quadratic Schrödinger in dimension 4

$$(i\partial_t + \Delta)u = u^2, \quad u : \mathbb{R}_t \times \mathbb{R}^n_x \to \mathbb{C}.$$

Goal: Construct spaces X and Y such that $\|u^2\|_Y \lesssim \|u\|_X^2$.
By duality, it is enough to establish

$$|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad w \in Y^*.$$

Strichartz estimates: $\|u\|_{L^p_{t,x}} \lesssim \|u_0\|_{L^2_x} + \sup |\langle u^2, w\rangle|$ with $\|w\|_{(L^{p'}_{t,x})^*} = 1$, $p = \frac{2(2+n)}{n}$

$n = 4$	$p = 3$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$p = \frac{10}{3}$	$L^{10/3} \cdot L^{10/3} \cdot L^{10/3} \not\subset L^1_{t,x}$	✗
$n = 2$	$p = 4$	$L^4 \cdot L^4 \cdot L^4 \not\subset L^1_{t,x}$	✗

> **Takeaway:** Linear information is not sufficient to establish a contraction for $n \leq 3$.

Toy model: quadratic Schrödinger in dimension 4

$$(i\partial_t + \Delta)u = u^2, \quad u : \mathbb{R}_t \times \mathbb{R}^n_x \to \mathbb{C}.$$

Goal: Construct spaces X and Y such that $\|u^2\|_Y \lesssim \|u\|_X^2$.
By duality, it is enough to establish

$$|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad w \in Y^*.$$

Strichartz estimates: $\|u\|_{L^p_{t,x}} \lesssim \|u_0\|_{L^2_x} + \sup |\langle u^2, w\rangle|$ with $\|w\|_{L^p_{t,x}} = 1$, $p = \frac{2(2+n)}{n}$

$n = 4$	$p = 3$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$p = \frac{10}{3}$	$L^{10/3} \cdot L^{10/3} \cdot L^{10/3} \not\subset L^1_{t,x}$	✗
$n = 2$	$p = 4$	$L^4 \cdot L^4 \cdot L^4 \not\subset L^1_{t,x}$	✗

> **Takeaway:** Linear information is not sufficient to establish a contraction for $n \leq 3$.

Toy model: quadratic Schrödinger in dimension 4

$$(i\partial_t + \Delta)u = u^2, \quad u : \mathbb{R}_t \times \mathbb{R}^n_x \to \mathbb{C}.$$

Goal: Construct spaces X and Y such that $\|u^2\|_Y \lesssim \|u\|_X^2$.
By duality, it is enough to establish

$$\Big| \int_{\mathbb{R}^{1+n}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad w \in Y^*.$$

Strichartz estimates: $\|u\|_{L^p_{t,x}} \lesssim \|u_0\|_{L^2_x} + \sup |\langle u^2, w\rangle|$ with $\|w\|_{L^p_{t,x}} = 1$, $p = \frac{2(2+n)}{n}$

$n = 4$	$p = 3$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$p = \frac{10}{3}$	$L^{10/3} \cdot L^{10/3} \cdot L^{10/3} \not\subset L^1_{t,x}$	✗
$n = 2$	$p = 4$	$L^4 \cdot L^4 \cdot L^4 \not\subset L^1_{t,x}$	✗

> **Takeaway:** Linear information is not sufficient to establish a contraction for $n \leq 3$.

Toy model: quadratic Schrödinger in dimension 4

$$(i\partial_t + \Delta)u = u^2, \quad u : \mathbb{R}_t \times \mathbb{R}^n_x \to \mathbb{C}.$$

Goal: Construct spaces X and Y such that $\|u^2\|_Y \lesssim \|u\|_X^2$.
By duality, it is enough to establish

$$\Big| \int_{\mathbb{R}^{1+n}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_{L^p_{t,x}} \|v\|_{L^p_{t,x}} \|w\|_{L^p_{t,x}}$$

Strichartz estimates: $\|u\|_{L^p_{t,x}} \lesssim \|u_0\|_{L^2_x} + \sup |\langle u^2, w\rangle|$ with $\|w\|_{L^p_{t,x}} = 1$, $p = \frac{2(2+n)}{n}$

$n = 4$	$p = 3$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$p = \frac{10}{3}$	$L^{10/3} \cdot L^{10/3} \cdot L^{10/3} \not\subset L^1_{t,x}$	✗
$n = 2$	$p = 4$	$L^4 \cdot L^4 \cdot L^4 \not\subset L^1_{t,x}$	✗

> **Takeaway:** Linear information is not sufficient to establish a contraction for $n \leq 3$.

Toy model: quadratic Schrödinger in dimension 4

$$(i\partial_t + \Delta)u = u^2, \quad u : \mathbb{R}_t \times \mathbb{R}^n_x \to \mathbb{C}.$$

Goal: Construct spaces X and Y such that $\|u^2\|_Y \lesssim \|u\|_X^2$.
By duality, it is enough to establish

$$\Big| \int_{\mathbb{R}^{1+n}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_{L^p_{t,x}} \|v\|_{L^p_{t,x}} \|w\|_{L^p_{t,x}}$$

Strichartz estimates: $\|u\|_{L^p_{t,x}} \lesssim \|u_0\|_{L^2_x} + \sup |\langle u^2, w\rangle|$ with $\|w\|_{L^p_{t,x}} = 1$, $p = \frac{2(2+n)}{n}$

$n = 4$	$p = 3$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$p = \frac{10}{3}$	$L^{10/3} \cdot L^{10/3} \cdot L^{10/3} \not\subset L^1_{t,x}$	✗
$n = 2$	$p = 4$	$L^4 \cdot L^4 \cdot L^4 \not\subset L^1_{t,x}$	✗

> **Takeaway:** Linear information is not sufficient to establish a contraction for $n \leq 3$.

Toy model: quadratic Schrödinger in dimension 4

$$(i\partial_t + \Delta)u = u^2, \quad u : \mathbb{R}_t \times \mathbb{R}^n_x \to \mathbb{C}.$$

Goal: Construct spaces X and Y such that $\|u^2\|_Y \lesssim \|u\|_X^2$.
By duality, it is enough to establish

$$\Big| \int_{\mathbb{R}^{1+n}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_{L^p_{t,x}} \|v\|_{L^p_{t,x}} \|w\|_{L^p_{t,x}}$$

Strichartz estimates: $\|u\|_{L^p_{t,x}} \lesssim \|u_0\|_{L^2_x} + \sup |\langle u^2, w\rangle|$ with $\|w\|_{L^p_{t,x}} = 1$, $p = \frac{2(2+n)}{n}$

$n = 4$	$p = 3$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$p = \frac{10}{3}$	$L^{10/3} \cdot L^{10/3} \cdot L^{10/3} \not\subset L^1_{t,x}$	✗
$n = 2$	$p = 4$	$L^4 \cdot L^4 \cdot L^4 \not\subset L^1_{t,x}$	✗

Takeaway: Linear information is not sufficient to establish a contraction for $n \leq 3$.

Toy model: quadratic Schrödinger in dimension 4

$$(i\partial_t + \Delta)u = u^2, \quad u : \mathbb{R}_t \times \mathbb{R}^n_x \to \mathbb{C}.$$

Goal: Construct spaces X and Y such that $\|u^2\|_Y \lesssim \|u\|_X^2$.
By duality, it is enough to establish

$$\Big| \int_{\mathbb{R}^{1+n}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_{L^p_{t,x}} \|v\|_{L^p_{t,x}} \|w\|_{L^p_{t,x}}$$

Strichartz estimates: $\|u\|_{L^p_{t,x}} \lesssim \|u_0\|_{L^2_x} + \sup |\langle u^2, w\rangle|$ with $\|w\|_{L^p_{t,x}} = 1$, $p = \frac{2(2+n)}{n}$

$n = 4$	$p = 3$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$p = \frac{10}{3}$	$L^{10/3} \cdot L^{10/3} \cdot L^{10/3} \not\subset L^1_{t,x}$	✗
$n = 2$	$p = 4$	$L^4 \cdot L^4 \cdot L^4 \not\subset L^1_{t,x}$	✗

> **Takeaway:** Linear information is not sufficient to establish a contraction for $n \leq 3$.

Toy model: quadratic Schrödinger in dimension 4

$$(i\partial_t + \Delta)u = u^2, \quad u : \mathbb{R}_t \times \mathbb{R}^n_x \to \mathbb{C}.$$

Goal: Construct spaces X and Y such that $\|u^2\|_Y \lesssim \|u\|_X^2$.
By duality, it is enough to establish

$$\Big| \int_{\mathbb{R}^{1+n}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_{L^p_{t,x}} \|v\|_{L^p_{t,x}} \|w\|_{L^p_{t,x}}$$

Strichartz estimates: $\|u\|_{L^p_{t,x}} \lesssim \|u_0\|_{L^2_x} + \sup |\langle u^2, w\rangle|$ with $\|w\|_{L^p_{t,x}} = 1$, $p = \frac{2(2+n)}{n}$

$n = 4$	$p = 3$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$p = \frac{10}{3}$	$L^{10/3} \cdot L^{10/3} \cdot L^{10/3} \not\subset L^1_{t,x}$	✗
$n = 2$	$p = 4$	$L^4 \cdot L^4 \cdot L^4 \not\subset L^1_{t,x}$	✗

Takeaway: Linear information is not sufficient to establish a contraction for $n \leq 3$.

First nonlinear iteration

> **Guiding principle:** nonlinear interactions carry more information than linear theory suggests

Consider the equation

$$(i\partial_t - \Delta)u = vw$$

where $v = e^{-it\Delta}g$, $w = e^{-it\Delta}h$ are free solutions localized at frequency 1.

In (spacetime) Fourier variables

$$\widehat{u}(\tau, \xi) = \frac{\widehat{(vw)}}{\tau - |\xi|^2}$$

This is well-defined when $\widehat{v} * \widehat{w}$ is supported *away* from the surface $\tau = |\xi|^2$. Such interactions are called *nonresonant*.

First nonlinear iteration

> **Guiding principle:** nonlinear interactions carry more information than linear theory suggests

Consider the equation

$$(i\partial_t - \Delta)u = vw$$

where $v = e^{-it\Delta}g$, $w = e^{-it\Delta}h$ are free solutions localized at frequency 1.

In (spacetime) Fourier variables

$$\widehat{u}(\tau, \xi) = \frac{\widehat{(vw)}}{\tau - |\xi|^2}$$

This is well-defined when $\widehat{v} * \widehat{w}$ is supported *away* from the surface $\tau = |\xi|^2$. Such interactions are called *nonresonant*.

First nonlinear iteration

> **Guiding principle:** nonlinear interactions carry more information than linear theory suggests

Consider the equation

$$u = \frac{vw}{i\partial_t - \Delta}$$

where $v = e^{-it\Delta}g$, $w = e^{-it\Delta}h$ are free solutions localized at frequency 1.

In (spacetime) Fourier variables

$$\widehat{u}(\tau,\xi) = \frac{\widehat{(vw)}}{\tau - |\xi|^2}$$

This is well-defined when $\widehat{v} * \widehat{w}$ is supported *away* from the surface $\tau = |\xi|^2$. Such interactions are called *nonresonant*.

First nonlinear iteration

> **Guiding principle:** nonlinear interactions carry more information than linear theory suggests

Consider the equation

$$u = \frac{vw}{i\partial_t - \Delta}$$

where $v = e^{-it\Delta}g$, $w = e^{-it\Delta}h$ are free solutions localized at frequency 1.

In (spacetime) Fourier variables

$$\hat{u}(\tau, \xi) = \frac{\widehat{(vw)}}{\tau - |\xi|^2}$$

This is well-defined when $\hat{v} * \hat{w}$ is supported *away* from the surface $\tau = |\xi|^2$. Such interactions are called *nonresonant*.

First nonlinear iteration

> **Guiding principle:** nonlinear interactions carry more information than linear theory suggests

Consider the equation

$$u = \frac{vw}{i\partial_t - \Delta}$$

where $v = e^{-it\Delta}g$, $w = e^{-it\Delta}h$ are free solutions localized at frequency 1.

In (spacetime) Fourier variables

$$\widehat{u}(\tau, \xi) = \frac{\widehat{v} * \widehat{w}}{\tau - |\xi|^2}$$

This is well-defined when $\widehat{v} * \widehat{w}$ is supported *away* from the surface $\tau = |\xi|^2$. Such interactions are called *nonresonant*.

First nonlinear iteration

> **Guiding principle:** nonlinear interactions carry more information than linear theory suggests

Consider the equation

$$u = \frac{vw}{i\partial_t - \Delta}$$

where $v = e^{-it\Delta}g$, $w = e^{-it\Delta}h$ are free solutions localized at frequency 1.

In (spacetime) Fourier variables

$$\widehat{u}(\tau, \xi) = \frac{\widehat{v} * \widehat{w}}{\tau - |\xi|^2}$$

This is well-defined when $\widehat{v} * \widehat{w}$ is supported *away* from the surface $\tau = |\xi|^2$. Such interactions are called *nonresonant*.

First nonlinear iteration

> **Guiding principle:** nonlinear interactions carry more information than linear theory suggests

Consider the equation

$$u = \frac{vw}{i\partial_t - \Delta}$$

where $v = e^{-it\Delta}g$, $w = e^{-it\Delta}h$ are free solutions localized at frequency 1.

In (spacetime) Fourier variables

$$\widehat{u}(\tau, \xi) = \frac{\widehat{v} * \widehat{w}}{\tau - |\xi|^2}$$

This is well-defined when $\widehat{v} * \widehat{w}$ is supported *away* from the surface $\tau = |\xi|^2$. Such interactions are called *nonresonant*.

Bilinear $L^2_{t,x}$

Let $Q_j u$ denote the localization of u to the region $|\tau - \xi^2| \leq 2^j$, $j \in \mathbb{Z}$.


By Plancharel,

$$\|Q_j u\|_{L^2_{t,x}} = \|Q_j (i\partial_t - \Delta)^{-1}(vw)\|_{L^2_{t,x}} = 2^{-j}\|vw\|_{L^2_{t,x}}.$$

Compared to the pure Strichartz approach, now we have access to $L^2_{t,x}$:

$$\left| \int_{\mathbb{R}^{1+3}} Q_j u v \bar{w} \, dx \, dt \right| \lesssim \|Q_j u\|_{L^2_{t,x}} \|vw\|_{L^2_{t,x}}$$

$n = 4$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$L^2 \cdot L^{10/3} \cdot L^{10/3} \subset L^{10/11}_{t,x}$	✓
$n = 2$	$L^2 \cdot L^4 \cdot L^4 \subset L^1_{t,x}$	✓



Takeaway: Nonlinear information supplies a more complete picture

Bilinear $L^2_{t,x}$

Let $Q_j u$ denote the localization of u to the region $|\tau - \xi^2| \leq 2^j$, $j \in \mathbb{Z}$.
By Plancharel,

$$\|Q_j u\|_{L^2_{t,x}} = \|Q_j (i\partial_t - \Delta)^{-1}(vw)\|_{L^2_{t,x}} = 2^{-j}\|vw\|_{L^2_{t,x}}.$$


Compared to the pure Strichartz approach, now we have access to $L^2_{t,x}$:

$$\left| \int_{\mathbb{R}^{1+3}} Q_j u v \bar{w} \, dx \, dt \right| \lesssim \|Q_j u\|_{L^2_{t,x}} \|vw\|_{L^2_{t,x}}$$

$n = 4$ $\quad L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$ ✓

$n = 3$ $\quad L^2 \cdot L^{10/3} \cdot L^{10/3} \subset L^{10/11}_{t,x}$ ✓

$n = 2$ $\quad L^2 \cdot L^4 \cdot L^4 \subset L^1_{t,x}$ ✓



Takeaway: Nonlinear information supplies a more complete picture

Bilinear $L^2_{t,x}$

Let $Q_j u$ denote the localization of u to the region $|\tau - \xi^2| \leq 2^j$, $j \in \mathbb{Z}$.
By Plancharel,

$$\|Q_j u\|_{L^2_{t,x}} = \|Q_j (i\partial_t - \Delta)^{-1}(vw)\|_{L^2_{t,x}} = 2^{-j}\|vw\|_{L^2_{t,x}}.$$


Compared to the pure Strichartz approach, now we have access to $L^2_{t,x}$:

$$\left| \int_{\mathbb{R}^{1+3}} Q_j u v \bar{w}\, dx\, dt \right| \lesssim \|Q_j u\|_{L^2_{t,x}} \|vw\|_{L^2_{t,x}}$$

$n = 4$ $\quad L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$ ✓

$n = 3$ $\quad L^2 \cdot L^{10/3} \cdot L^{10/3} \subset L^{10/11}_{t,x}$ ✓

$n = 2$ $\quad L^2 \cdot L^4 \cdot L^4 \subset L^1_{t,x}$ ✓



Takeaway: Nonlinear information supplies a more complete picture

Bilinear $L^2_{t,x}$

Let $Q_j u$ denote the localization of u to the region $|\tau - \xi^2| \leq 2^j$, $j \in \mathbb{Z}$.
By Plancharel,

$$\|Q_j u\|_{L^2_{t,x}} = \|Q_j (i\partial_t - \Delta)^{-1}(vw)\|_{L^2_{t,x}} = 2^{-j}\|vw\|_{L^2_{t,x}}.$$


Compared to the pure Strichartz approach, now we have access to $L^2_{t,x}$:

$$\left| \int_{\mathbb{R}^{1+3}} Q_j u v \bar{w}\, dx\, dt \right| \lesssim \|Q_j u\|_{L^2_{t,x}} \|vw\|_{L^2_{t,x}}$$

$n = 4$ $\quad L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$ ✓

$n = 3$ $\quad L^2 \cdot L^{10/3} \cdot L^{10/3} \subset L^{10/11}_{t,x}$ ✓

$n = 2$ $\quad L^2 \cdot L^4 \cdot L^4 \subset L^1_{t,x}$ ✓



Takeaway: Nonlinear information supplies a more complete picture

Bilinear $L^2_{t,x}$

Let $Q_j u$ denote the localization of u to the region $|\tau - \xi^2| \leq 2^j$, $j \in \mathbb{Z}$.
By Plancharel,

$$\|Q_j u\|_{L^2_{t,x}} = \|Q_j (i\partial_t - \Delta)^{-1}(vw)\|_{L^2_{t,x}} = 2^{-j}\|vw\|_{L^2_{t,x}}.$$


Compared to the pure Strichartz approach, now we have access to $L^2_{t,x}$:

$$\left| \int_{\mathbb{R}^{1+3}} Q_j u v \bar{w} \, dx \, dt \right| \lesssim \|Q_j u\|_{L^2_{t,x}} \|vw\|_{L^2_{t,x}}$$

$n = 4$ $\quad L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$ ✓

$n = 3$ $\quad L^2 \cdot L^{10/3} \cdot L^{10/3} \subset L^{10/11}_{t,x}$ ✓

$n = 2$ $\quad L^2 \cdot L^4 \cdot L^4 \subset L^1_{t,x}$ ✓



Takeaway: Nonlinear information supplies a more complete picture

Bilinear $L^2_{t,x}$


Let $Q_j u$ denote the localization of u to the region $|\tau - \xi^2| \leq 2^j$, $j \in \mathbb{Z}$.
By Plancharel,

$$\|Q_j u\|_{L^2_{t,x}} = \|Q_j (i\partial_t - \Delta)^{-1}(vw)\|_{L^2_{t,x}} = 2^{-j}\|vw\|_{L^2_{t,x}}.$$

Compared to the pure Strichartz approach, now we have access to $L^2_{t,x}$:

$$\left| \int_{\mathbb{R}^{1+3}} Q_j u v \bar{w}\, dx\, dt \right| \lesssim \|Q_j u\|_{L^2_{t,x}} \|vw\|_{L^2_{t,x}}$$

$n = 4$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$L^2 \cdot L^{10/3} \cdot L^{10/3} \subset L^{10/11}_{t,x}$	✓
$n = 2$	$L^2 \cdot L^4 \cdot L^4 \subset L^1_{t,x}$	✓



Takeaway: Nonlinear information supplies a more complete picture

Bilinear $L^2_{t,x}$


Let $Q_j u$ denote the localization of u to the region $|\tau - \xi^2| \leq 2^j$, $j \in \mathbb{Z}$.
By Plancharel,

$$\|Q_j u\|_{L^2_{t,x}} = \|Q_j (i\partial_t - \Delta)^{-1}(vw)\|_{L^2_{t,x}} = 2^{-j}\|vw\|_{L^2_{t,x}}.$$

Compared to the pure Strichartz approach, now we have access to $L^2_{t,x}$:

$$\left| \int_{\mathbb{R}^{1+3}} Q_j u v \bar{w}\, dx\, dt \right| \lesssim \|Q_j u\|_{L^2_{t,x}} \|vw\|_{L^2_{t,x}}$$

$n = 4$	$L^3 \cdot L^3 \cdot L^3 \subset L^1_{t,x}$	✓
$n = 3$	$L^2 \cdot L^{10/3} \cdot L^{10/3} \subset L^{10/11}_{t,x}$	✓
$n = 2$	$L^2 \cdot L^4 \cdot L^4 \subset L^1_{t,x}$	✓



Takeaway: Nonlinear information supplies a more complete picture

Nonresonant Dirac–Klein–Gordon in 2D

Theorem (Bejenaru–B. ’26)

Let $M, m > 0$ with $2M > m > 0$. For small initial data

$$(\psi_0, \phi_0, \phi_1) \in H_x^{\frac{1}{2}+\epsilon}(\mathbb{R}^2) \times H_x^{1+\epsilon}(\mathbb{R}^2) \times H_x^{\epsilon}(\mathbb{R}^2),$$

the *nonresonant* system

$$(-i\gamma^\mu \partial_\mu + M)\psi = \phi\psi$$
$$(\Box + m^2)\phi = \langle \psi, \gamma^0 \psi \rangle$$

is globally well-posed and scatters.

Recap (trilinear formulation and resonance decomposition)

Our goal is to find spaces X and Y such that $|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad w \in Y^*.$

- In high dimensions $n \geq 4$, Strichartz estimate on $L^3_{t,x}$ suffices.
- In low dimensions, we decompose u into $u = u_{\text{nonr}} + u_{\text{res}}$ and study
$$\langle uv, w\rangle = \langle u_{\text{res}} v, w\rangle + \langle u_{\text{nonr}} v, w\rangle$$
separately.
- Since u_{nonr} is not a free solution, we have access to extra integrability.

Since u_{res} is close to a free solution, we first consider the case where u, v, w are free waves.

Recap (trilinear formulation and resonance decomposition)

Our goal is to find spaces X and Y such that $|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad w \in Y^*.$

- In high dimensions $n \geq 4$, Strichartz estimate on $L^3_{t,x}$ suffices.
- In low dimensions, we decompose u into $u = u_{\text{nonr}} + u_{\text{res}}$ and study
$$\langle uv, w\rangle = \langle u_{\text{res}} v, w\rangle + \langle u_{\text{nonr}} v, w\rangle$$
separately.
- Since u_{nonr} is not a free solution, we have access to extra integrability.

Since u_{res} is close to a free solution, we first consider the case where u, v, w are free waves.

Recap (trilinear formulation and resonance decomposition)

Our goal is to find spaces X and Y such that $|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad w \in Y^*.$


- In high dimensions $n \geq 4$, Strichartz estimate on $L^3_{t,x}$ suffices.
- In low dimensions, we decompose u into $u = u_{\text{nonr}} + u_{\text{res}}$ and study
$$\langle uv, w\rangle = \langle u_{\text{res}} v, w\rangle + \langle u_{\text{nonr}} v, w\rangle$$
separately.
- Since u_{nonr} is not a free solution, we have access to extra integrability.

Since u_{res} is close to a free solution, we first consider the case where u, v, w are free waves.

Recap (trilinear formulation and resonance decomposition)

Our goal is to find spaces X and Y such that $|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad w \in Y^*.$

- In high dimensions $n \geq 4$, Strichartz estimate on $L^3_{t,x}$ suffices.
- In low dimensions, we decompose u into $u = u_{\text{nonr}} + u_{\text{res}}$ and study
$$\langle uv, w\rangle = \langle u_{\text{res}} v, w\rangle + \langle u_{\text{nonr}} v, w\rangle$$
separately.
- Since u_{nonr} is not a free solution, we have access to extra integrability.




Since u_{res} is close to a free solution, we first consider the case where u, v, w are free waves.

Recap (trilinear formulation and resonance decomposition)

Our goal is to find spaces X and Y such that $|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad w \in Y^*.$

- In high dimensions $n \geq 4$, Strichartz estimate on $L^3_{t,x}$ suffices.
- In low dimensions, we decompose u into $u = u_{\text{nonr}} + u_{\text{res}}$ and study
$$\langle uv, w\rangle = \langle u_{\text{res}} v, w\rangle + \langle u_{\text{nonr}} v, w\rangle$$
separately.
- Since u_{nonr} is not a free solution, we have access to extra integrability.




Since u_{res} is close to a free solution, we first consider the case where u, v, w are free waves.

Recap (trilinear formulation and resonance decomposition)

Our goal is to find spaces X and Y such that $|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad w \in Y^*.$


- In high dimensions $n \geq 4$, Strichartz estimate on $L^3_{t,x}$ suffices.
- In low dimensions, we decompose u into $u = u_{\text{nonr}} + u_{\text{res}}$ and study
$$\langle uv, w\rangle = \langle u_{\text{res}} v, w\rangle + \langle u_{\text{nonr}} v, w\rangle$$
separately.
- Since u_{nonr} is not a free solution, we have access to extra integrability.




Since u_{res} is close to a free solution, we first consider the case where u, v, w are free waves.

Wave packets

Let $u = e^{-it\Delta} f$ be a free wave. In Fourier space, $\widehat{u}$ is supported on the paraboloid $\tau = |\xi|^2$:







u is approximately constant on tubes of dimensions $(\delta^{-1})^3 \times \delta^{-2}$ oriented along N.

Wave packets

Let $u = e^{-it\Delta} f$ be a free wave. In Fourier space, $\widehat{u}$ is supported on the paraboloid $\tau = |\xi|^2$:





u is approximately constant on tubes of dimensions $(\delta^{-1})^3 \times \delta^{-2}$ oriented along N.

Two-wave interactions

Question: In 3D, Hölder + Strichartz give

$$\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^{5/3}_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}.$$

Can we do better?

If N_1 and N_2 are *transversal*, $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, $\frac{3}{2} < p < \frac{5}{3}$

Two-wave interactions

Question: In 3D, Hölder + Strichartz give

$$\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^{5/3}_{t,x}} \lesssim \|e^{-it\Delta} f\|_{L^{10/3}_{t,x}} \|e^{-it\Delta} g\|_{L^{10/3}_{t,x}}.$$

Can we do better?

If N_1 and N_2 are *transversal*, $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, $\frac{3}{2} < p < \frac{5}{3}$

Two-wave interactions

Question: In 3D, Hölder + Strichartz give

$$\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^{5/3}_{t,x}} \lesssim \|e^{-it\Delta} f\|_{L^{10/3}_{t,x}} \|e^{-it\Delta} g\|_{L^{10/3}_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}.$$

Can we do better?

If N_1 and N_2 are *transversal*, $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}, \quad \frac{3}{2} < p < \frac{5}{3}$

Two-wave interactions

Question: In 3D, Hölder + Strichartz give

$$\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^{5/3}_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}.$$

Can we do better?


If N_1 and N_2 are *transversal*, $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, $\frac{3}{2} < p < \frac{5}{3}$


Two-wave interactions

Question: In 3D, Hölder + Strichartz give

$$\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^{5/3}_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}.$$

Can we do better?






If N_1 and N_2 are *transversal*, $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, $\frac{3}{2} < p < \frac{5}{3}$


Two-wave interactions

Question: In 3D, Hölder + Strichartz give

$$\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^{5/3}_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}.$$

Can we do better?







If N_1 and N_2 are *transversal*, $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, $\frac{3}{2} < p < \frac{5}{3}$

Why is transversality important?

Heuristic: “transversal waves interact in smaller regions and for less time”







If the normals satisfy $|N_1 \wedge N_2| \geq \nu > 0$, then

$$\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}} \lesssim \nu^{3-\frac{5}{p}} \|f\|_{L^2_x} \|g\|_{L^2_x}, \qquad \frac{3}{2} < p \leq 2.$$

Why is transversality important?

Heuristic: “transversal waves interact in smaller regions and for less time”





If the normals satisfy $|N_1 \wedge N_2| \geq \nu > 0$, then

$$\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}} \lesssim \nu^{3-\frac{5}{p}} \|f\|_{L^2_x} \|g\|_{L^2_x}, \qquad \frac{3}{2} < p \leq 2.$$

Bilinear estimates and the remaining resonances

For three free Schrödinger waves in dimension 3, the direct Hölder route is

$$\left|\int uv\bar{w}\ dx\,dt\right| \leq \|uv\|_{L^q_{t,x}} \|w\|_{L^{10/3}_{t,x}}, \qquad \frac{1}{q}+\frac{3}{10}=1.$$

We need a bilinear estimate with $q=\frac{10}{7}$, however bilinear restriction requires $q>\frac{3}{2}>\frac{10}{7}$

> **Takeaway:** the bilinear restriction range is not enough to control resonant quadratic interactions!

Bilinear suffices in the radial class: In that case, the correct Strichartz space is $L^{14/5}_{t,x}$, and so

$$\left|\int uv\bar{w}\ dx\,dt\right| \leq \|uv\|_{L^q_{t,x}} \|w\|_{L^{14/5}_{t,x}},$$

and the dual space $q=\frac{14}{9}>\frac{3}{2}$ belongs to the bilinear restriction range!

Bilinear estimates and the remaining resonances

For three free Schrödinger waves in dimension 3, the direct Hölder route is

$$\left|\int uv\bar{w}\ dx\,dt\right| \leq \|uv\|_{L^q_{t,x}} \|w\|_{L^{10/3}_{t,x}}, \qquad \frac{1}{q}+\frac{3}{10}=1.$$

We need a bilinear estimate with $q=\frac{10}{7}$, however bilinear restriction requires $q>\frac{3}{2}>\frac{10}{7}$

> **Takeaway:** the bilinear restriction range is not enough to control resonant quadratic interactions!

Bilinear suffices in the radial class: In that case, the correct Strichartz space is $L^{14/5}_{t,x}$, and so

$$\left|\int uv\bar{w}\ dx\,dt\right| \leq \|uv\|_{L^q_{t,x}} \|w\|_{L^{14/5}_{t,x}},$$

and the dual space $q=\frac{14}{9}>\frac{3}{2}$ belongs to the bilinear restriction range!

Bilinear estimates and the remaining resonances

For three free Schrödinger waves in dimension 3, the direct Hölder route is

$$\left| \int uv\bar{w} \, dx \, dt \right| \leq \|uv\|_{L^q_{t,x}} \|w\|_{L^{10/3}_{t,x}}, \qquad \frac{1}{q} + \frac{3}{10} = 1.$$

We need a bilinear estimate with $q = \frac{10}{7}$, however bilinear restriction requires $q > \frac{3}{2} > \frac{10}{7}$

> **Takeaway:** the bilinear restriction range is not enough to control resonant quadratic interactions!

Bilinear suffices in the radial class: In that case, the correct Strichartz space is $L^{14/5}_{t,x}$, and so

$$\left| \int uv\bar{w} \, dx \, dt \right| \leq \|uv\|_{L^q_{t,x}} \|w\|_{L^{14/5}_{t,x}},$$

and the dual space $q = \frac{14}{9} > \frac{3}{2}$ belongs to the bilinear restriction range!

Bilinear estimates and the remaining resonances

For three free Schrödinger waves in dimension 3, the direct Hölder route is

$$\left| \int uv\bar{w} \, dx \, dt \right| \leq \|uv\|_{L^q_{t,x}} \|w\|_{L^{10/3}_{t,x}}, \qquad \frac{1}{q} + \frac{3}{10} = 1.$$

We need a bilinear estimate with $q = \frac{10}{7}$, however bilinear restriction requires $q > \frac{3}{2} > \frac{10}{7}$

> **Takeaway:** the bilinear restriction range is not enough to control resonant quadratic interactions!

Bilinear suffices in the radial class: In that case, the correct Strichartz space is $L^{14/5}_{t,x}$, and so

$$\left| \int uv\bar{w} \, dx \, dt \right| \leq \|uv\|_{L^q_{t,x}} \|w\|_{L^{14/5}_{t,x}},$$

and the dual space $q = \frac{14}{9} > \frac{3}{2}$ belongs to the bilinear restriction range!

The 3D Klein–Gordon–Schrödinger system

Theorem (B.–Chan, ‘26)


$$(i\partial_t + \Delta)u = un, \qquad (\Box + 1)n = \pm|u|^2.$$

For any $0 < \epsilon \ll 1$, sufficiently small **radial** data


$$(u_0, n_0, n_1) \in L^2_x(\mathbb{R}^3) \times H^{-\frac{1}{2}+\epsilon}_x(\mathbb{R}^3) \times H^{-\frac{3}{2}+\epsilon}_x(\mathbb{R}^3)$$

give a global solution which scatters to free solutions as $t \to \pm\infty$.

Schrödinger equation



Klein–Gordon equation



Color key

- Blue: nonresonant
- Yellow: bilinear restriction
- Gold: radial

Can we do better than bilinear estimates?

Hölder tells you how to estimate $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}}$ using only *linear* information.

Bilinear estimates use nonlinear information (transversality) to improve integrability $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}}$ with some $q < p$.

Question: Do *trilinear* interactions capture any extra information compared to bilinear?

Yes!

$$\Big\| \prod_{j=1}^{3} e^{-it\Delta} f_j \Big\|_{L^p_{t,x}} \lesssim \prod_{j=1}^{3} \|f_j\|_{L^2_x}, \quad p > \frac{14}{15}$$

provided that the normals N_1, N_2, N_3 are transversal $|N_1 \wedge N_2 \wedge N_3| \geq \nu > 0$.

Caveat: For general curves, transversality does not improve the range below $p = 1$.

Can we do better than bilinear estimates?

Hölder tells you how to estimate $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}}$ using only *linear* information.

Bilinear estimates use nonlinear information (transversality) to improve integrability $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}}$ with some $q < p$.

Question: Do *trilinear* interactions capture any extra information compared to bilinear?

Yes!

$$\Big\| \prod_{j=1}^{3} e^{-it\Delta} f_j \Big\|_{L^p_{t,x}} \lesssim \prod_{j=1}^{3} \|f_j\|_{L^2_x}, \quad p > \frac{14}{15}$$

provided that the normals N_1, N_2, N_3 are transversal $|N_1 \wedge N_2 \wedge N_3| \geq \nu > 0$.

Caveat: For general curves, transversality does not improve the range below $p = 1$.

Can we do better than bilinear estimates?

Hölder tells you how to estimate $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}}$ using only *linear* information.

Bilinear estimates use nonlinear information (transversality) to improve integrability $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}}$ with some $q < p$.

Question: Do *trilinear* interactions capture any extra information compared to bilinear?

Yes!

$$\Big\| \prod_{j=1}^{3} e^{-it\Delta} f_j \Big\|_{L^p_{t,x}} \lesssim \prod_{j=1}^{3} \|f_j\|_{L^2_x}, \quad p > \frac{14}{15}$$

provided that the normals N_1, N_2, N_3 are transversal $|N_1 \wedge N_2 \wedge N_3| \geq \nu > 0$.

Caveat: For general curves, transversality does not improve the range below $p = 1$.

Can we do better than bilinear estimates?

Hölder tells you how to estimate $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}}$ using only *linear* information.

Bilinear estimates use nonlinear information (transversality) to improve integrability $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}}$ with some $q < p$.

Question: Do *trilinear* interactions capture any extra information compared to bilinear?

Yes!

$$\Big\| \prod_{j=1}^{3} e^{-it\Delta} f_j \Big\|_{L^p_{t,x}} \lesssim \prod_{j=1}^{3} \|f_j\|_{L^2_x}, \quad p > \frac{14}{15}$$

provided that the normals N_1, N_2, N_3 are transversal $|N_1 \wedge N_2 \wedge N_3| \geq \nu > 0$.

Caveat: For general curves, transversality does not improve the range below $p = 1$.

Can we do better than bilinear estimates?

Hölder tells you how to estimate $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}}$ using only *linear* information.

Bilinear estimates use nonlinear information (transversality) to improve integrability $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}}$ with some $q < p$.

Question: Do *trilinear* interactions capture any extra information compared to bilinear?

Yes!

$$\Big\| \prod_{j=1}^{3} e^{-it\Delta} f_j \Big\|_{L^p_{t,x}} \lesssim \prod_{j=1}^{3} \|f_j\|_{L^2_x}, \quad p > \frac{14}{15}$$

provided that the normals N_1, N_2, N_3 are transversal $|N_1 \wedge N_2 \wedge N_3| \geq \nu > 0$.

Caveat: For general curves, transversality does not improve the range below $p = 1$.

Can we do better than bilinear estimates?

Hölder tells you how to estimate $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^p_{t,x}}$ using only *linear* information.

Bilinear estimates use nonlinear information (transversality) to improve integrability $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}}$ with some $q < p$.

Question: Do *trilinear* interactions capture any extra information compared to bilinear?

Yes!

$$\Big\| \prod_{j=1}^{3} e^{-it\Delta} f_j \Big\|_{L^p_{t,x}} \lesssim \prod_{j=1}^{3} \|f_j\|_{L^2_x}, \quad p > \frac{14}{15}$$

provided that the normals N_1, N_2, N_3 are transversal $|N_1 \wedge N_2 \wedge N_3| \geq \nu > 0$.

Caveat: For general curves, transversality does not improve the range below $p = 1$.

What does curvature buy?

Curves on $\mathbb{R}^{1+1}$



Transversality gives bilinear $L^2_{t,x}$.

Surfaces on $\mathbb{R}^{1+2}$

Transversality gives bilinear $L^2_{t,x}$.

Curvature along the intersection gives bilinear $L^{5/3+}_{t,x}$.

Hypersurfaces on $\mathbb{R}^{1+3}$

In $\mathbb{R}^3$: transversality gives trilinear $L^{1+}_{t,x}$.

In $\mathbb{R}^4$: extra curvature gives trilinear $L^{14/15+}_{t,x}$.


What does curvature buy?

Curves on $\mathbb{R}^{1+1}$



Transversality gives bilinear $L^2_{t,x}$.

Surfaces on $\mathbb{R}^{1+2}$



Transversality gives bilinear $L^2_{t,x}$.

Curvature along the intersection gives bilinear $L^{5/3+}_{t,x}$.

Hypersurfaces on $\mathbb{R}^{1+3}$

In $\mathbb{R}^3$: transversality gives trilinear $L^{1+}_{t,x}$.

In $\mathbb{R}^4$: extra curvature gives trilinear $L^{14/15+}_{t,x}$.


What does curvature buy?

Curves on $\mathbb{R}^{1+1}$



Transversality gives bilinear $L^2_{t,x}$.


Surfaces on $\mathbb{R}^{1+2}$



Transversality gives bilinear $L^2_{t,x}$.

Curvature along the intersection gives bilinear $L^{5/3+}_{t,x}$.

Hypersurfaces on $\mathbb{R}^{1+3}$



In $\mathbb{R}^3$: transversality gives trilinear $L^{1+}_{t,x}$.

In $\mathbb{R}^4$: extra curvature gives trilinear $L^{14/15+}_{t,x}$.

Roadmap: harmonic analysis

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

(iii) Harmonic analysis: geometry of interacting waves

(iv) From harmonic analysis to PDEs

(v) Everything coming together

Roadmap: harmonic analysis

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability

(iii) Harmonic analysis: geometry of interacting waves

(iv) From harmonic analysis to PDEs

(v) Everything coming together

Roadmap: harmonic analysis

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

(iv) From harmonic analysis to PDEs

(v) Everything coming together

Roadmap: harmonic analysis

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

- Linear wave packets

(iv) From harmonic analysis to PDEs

(v) Everything coming together

Roadmap: harmonic analysis

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

- Linear wave packets
- Bilinear range is very good in the radial case

(iv) From harmonic analysis to PDEs

(v) Everything coming together

Roadmap: harmonic analysis

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

- Linear wave packets
- Bilinear range is very good in the radial case
- Transversality improves integrability

(iv) From harmonic analysis to PDEs

(v) Everything coming together

Roadmap: harmonic analysis

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

- Linear wave packets
- Bilinear range is very good in the radial case
- Transversality improves integrability
- Trilinear transversality is better than bilinear transversality

(iv) From harmonic analysis to PDEs

(v) Everything coming together

Transference spaces

Recall that our original goal was to find spaces X and Y satisfying:

- Linear estimate: $\|u\|_X \lesssim \|u(0)\|_{L^2_x} + \|F\|_Y$,
- Multilinear estimate: $|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}$

We have discussed two types of multilinear estimates:

$$|\langle vw, u_{\text{nonr}}\rangle| \lesssim \|v\|_X \|w\|_X \|u_{\text{nonr}}\|_{L^2_{t,x}},$$

$$|\langle vw, u_{\text{res}}\rangle| \lesssim \|vw\|_{\text{Bil}} \|u_{\text{res}}\|_{\text{Bil}^*},$$

in the case where u_{res} is a free solution.

Question: How do we combine all this information into spaces X and Y?

Transference spaces

Recall that our original goal was to find spaces X and Y satisfying:

- Linear estimate: $\|u\|_X \lesssim \|u(0)\|_{L^2_x} + \|F\|_Y$,
- Multilinear estimate: $|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}$

We have discussed two types of multilinear estimates:

$$|\langle vw, u_{\text{nonr}}\rangle| \lesssim \|v\|_X \|w\|_X \|u_{\text{nonr}}\|_{L^2_{t,x}},$$

$$|\langle vw, u_{\text{res}}\rangle| \lesssim \|vw\|_{\text{Bil}} \|u_{\text{res}}\|_{\text{Bil}^*},$$

in the case where u_{res} is a free solution.

Question: How do we combine all this information into spaces X and Y?

Transference spaces

Recall that our original goal was to find spaces X and Y satisfying:

- Linear estimate: $\|u\|_X \lesssim \|u(0)\|_{L^2_x} + \|F\|_Y$,
- Multilinear estimate: $|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}$

We have discussed two types of multilinear estimates:

$$|\langle vw, u_{\text{nonr}}\rangle| \lesssim \|v\|_X \|w\|_X \|u_{\text{nonr}}\|_{L^2_{t,x}},$$

$$|\langle vw, u_{\text{res}}\rangle| \lesssim \|v\|_X \|w\|_X \|u_{\text{res}}\|_{\text{Bil}^*},$$

in the case where u_{res} is a free solution.

Question: How do we combine all this information into spaces X and Y?

Transference spaces

Recall that our original goal was to find spaces X and Y satisfying:

- Linear estimate: $\|u\|_X \lesssim \|u(0)\|_{L^2_x} + \|F\|_Y$,
- Multilinear estimate: $|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}$

We have discussed two types of multilinear estimates:

$$|\langle vw, u_{\text{nonr}}\rangle| \lesssim \|v\|_X \|w\|_X \|u_{\text{nonr}}\|_{L^2_{t,x}},$$

$$|\langle vw, u_{\text{res}}\rangle| \lesssim \|v\|_X \|w\|_X \|u_{\text{res}}\|_{\text{Bil}^*},$$

in the case where u_{res} is a free solution.

Question: How do we combine all this information into spaces X and Y?

Candidate 1: Strichartz spaces – revisited

Let

$$X = L^2_t L^6_x \cap L^\infty_t L^2_x, \quad Y = L^2_t L^{6/5}_x + L^1_t L^2_x.$$

Since $Y^* = X$, the trilinear estimate reduces to

$$\Big| \int_{\mathbb{R}^{1+3}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_X \|v\|_X \|w\|_X$$

Pros

- Easy to define
- Easy to establish duality

Cons

- Only considers linear information
- Not enough to close the contraction when $n \leq 3$
- It does not allow for (bilinear) $L^2_{t,x}$

Candidate 1: Strichartz spaces – revisited

Let

$$X = L_t^2 L_x^6 \cap L_t^\infty L_x^2, \quad Y = L_t^2 L_x^{6/5} + L_t^1 L_x^2.$$

Since $Y^* = X$, the trilinear estimate reduces to

$$\Big| \int_{\mathbb{R}^{1+3}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_X \|v\|_X \|w\|_X$$

Pros

- Easy to define
- Easy to establish duality

Cons

- Only considers linear information
- Not enough to close the contraction when $n \leq 3$
- It does not allow for (bilinear) $L_{t,x}^2$

Candidate 1: Strichartz spaces – revisited

Let

$$X = L_t^2 L_x^6 \cap L_t^\infty L_x^2, \quad Y = L_t^2 L_x^{6/5} + L_t^1 L_x^2.$$

Since $Y^* = X$, the trilinear estimate reduces to

$$\Big| \int_{\mathbb{R}^{1+3}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_X \|v\|_X \|w\|_X$$

Pros

- Easy to define
- Easy to establish duality

Cons

- Only considers linear information
- Not enough to close the contraction when $n \leq 3$
- It does not allow for (bilinear) $L_{t,x}^2$

Candidate 1: Strichartz spaces – revisited

Let

$$X = L_t^2 L_x^6 \cap L_t^\infty L_x^2, \quad Y = L_t^2 L_x^{6/5} + L_t^1 L_x^2.$$

Since $Y^* = X$, the trilinear estimate reduces to

$$\Big| \int_{\mathbb{R}^{1+3}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_X \|v\|_X \|w\|_X$$

Pros

- Easy to define
- Easy to establish duality

Cons

- Only considers linear information
- Not enough to close the contraction when $n \leq 3$
- It does not allow for (bilinear) $L_{t,x}^2$

Candidate 1: Strichartz spaces – revisited

Let

$$X = L_t^2 L_x^6 \cap L_t^\infty L_x^2, \quad Y = L_t^2 L_x^{6/5} + L_t^1 L_x^2.$$

Since $Y^* = X$, the trilinear estimate reduces to

$$\Big| \int_{\mathbb{R}^{1+3}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_X \|v\|_X \|w\|_X$$

Pros

- Easy to define
- Easy to establish duality

Cons

- Only considers linear information
- Not enough to close the contraction when $n \leq 3$
- It does not allow for (bilinear) $L_{t,x}^2$

Candidate 1: Strichartz spaces – revisited

Let

$$X = L_t^2 L_x^6 \cap L_t^\infty L_x^2, \quad Y = L_t^2 L_x^{6/5} + L_t^1 L_x^2.$$

Since $Y^* = X$, the trilinear estimate reduces to

$$\Big| \int_{\mathbb{R}^{1+3}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_X \|v\|_X \|w\|_X$$

Pros

- Easy to define
- Easy to establish duality

Cons

- Only considers linear information
- Not enough to close the contraction when $n \leq 3$
- It does not allow for (bilinear) $L_{t,x}^2$

Candidate 1: Strichartz spaces – revisited

Let

$$X = L^2_t L^6_x \cap L^\infty_t L^2_x, \quad Y = L^2_t L^{6/5}_x + L^1_t L^2_x.$$

Since $Y^* = X$, the trilinear estimate reduces to

$$\Big| \int_{\mathbb{R}^{1+3}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_X \|v\|_X \|w\|_X$$

Pros

- Easy to define
- Easy to establish duality

Cons

- Only considers linear information
- Not enough to close the contraction when $n \leq 3$
- It does not allow for (bilinear) $L^2_{t,x}$

Candidate 1: Strichartz spaces – revisited

Let

$$X = L_t^2 L_x^6 \cap L_t^\infty L_x^2, \quad Y = L_t^2 L_x^{6/5} + L_t^1 L_x^2.$$

Since $Y^* = X$, the trilinear estimate reduces to

$$\Big| \int_{\mathbb{R}^{1+3}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_X \|v\|_X \|w\|_X$$

Pros

- Easy to define
- Easy to establish duality

Cons

- Only considers linear information
- Not enough to close the contraction when $n \leq 3$
- It does not allow for (bilinear) $L_{t,x}^2$

Candidate 1: Strichartz spaces – revisited

Let

$$X = L_t^2 L_x^6 \cap L_t^\infty L_x^2, \quad Y = L_t^2 L_x^{6/5} + L_t^1 L_x^2.$$

Since $Y^* = X$, the trilinear estimate reduces to

$$\Big| \int_{\mathbb{R}^{1+3}} uv\bar{w}\, dx\, dt \Big| \lesssim \|u\|_X \|v\|_X \|w\|_X$$

Pros

- Easy to define
- Easy to establish duality

Cons

- Only considers linear information
- Not enough to close the contraction when $n \leq 3$
- It does not allow for (bilinear) $L_{t,x}^2$

Candidate 2: The atomic space U^a_Δ

$$e^{-it\Delta} f = \mathbb{1}_{[0,\infty)}(t) e^{-it\Delta} f$$




Candidate 2: The atomic space U^a_Δ

$$e^{-it\Delta} f = \mathbb{1}_{[0,\infty)}(t) e^{-it\Delta} f$$




Candidate 2: The atomic space U^a_Δ

$$e^{-it\Delta} f = \mathbb{1}_{[0,\infty)}(t) e^{-it\Delta} f$$




Candidate 2: The atomic space U^a_Δ

$$e^{-it\Delta} f = \mathbb{1}_{[0,\infty)}(t) e^{-it\Delta} f$$




Candidate 2: The atomic space U^a_Δ

$$e^{-it\Delta} f = \mathbb{1}_{[0,\infty)}(t) e^{-it\Delta} f$$



Candidate 2: The atomic space U^a_Δ

$$e^{-it\Delta} f = \mathbb{1}_{[0,\infty)}(t) e^{-it\Delta} f$$



A U^a_Δ-*atom* has the form


$$u = \sum_{j=1}^N \mathbb{1}_{I_j}(t) e^{-it\Delta} f_j$$

with normalization

$$\sum_{j=1}^N \|f_j\|^a_{L^2_x} = 1.$$

Candidate 2: The atomic space U^a_Δ

$$e^{-it\Delta} f = \mathbb{1}_{[0,\infty)}(t) e^{-it\Delta} f$$



A U^a_Δ-*atom* has the form

$$u = \sum_{j=1}^N \mathbb{1}_{I_j}(t) e^{-it\Delta} f_j$$


with normalization

$$\sum_{j=1}^N \|f_j\|^a_{L^2_x} = 1.$$

$$U^a_\Delta = \left\{ u = \sum_{J=1}^\infty \lambda_J u_J : u_J \text{ is an atom, } (\lambda_J) \in \ell^1_J \right\}$$

Candidate 2: The atomic space U^2_Δ

$$e^{-it\Delta} f = \mathbb{1}_{[0,\infty)}(t) e^{-it\Delta} f$$



A U^2_Δ-*atom* has the form

$$u = \sum_{j=1}^N \mathbb{1}_{I_j}(t) e^{-it\Delta} f_j$$

with normalization

$$\sum_{j=1}^N \|f_j\|^2_{L^2_x} = 1.$$

$$U^2_\Delta = \left\{ u = \sum_{J=1}^\infty \lambda_J u_J : u_J \text{ is an atom, } (\lambda_J) \in \ell^1_J \right\}$$

Transference from free solutions

U^2_Δ-functions are "almost free solutions" and most estimates for free solutions also hold in this space:

- Linear: if $\|e^{-it\Delta} f\|_{L^p_t L^q_x} \lesssim \|f\|_{L^2_x}$ for all $f \in L^2_x$, then $\|u\|_{L^p_t L^q_x} \lesssim \|u\|_{U^2_\Delta}$ for all $u \in U^2_\Delta$.
- Bilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, then $\|uv\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta}$
- Trilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot e^{-it\Delta} h\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$, then $\|uvw\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$.

U^2_Δ is almost the perfect space... Except for duality.

$$|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad U^2_\Delta \subset V^2_\Delta$$

Estimate we need

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}$$

Estimate we can transfer

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}.$$

Transference from free solutions

U^2_Δ-functions are "almost free solutions" and most estimates for free solutions also hold in this space:

- Linear: if $\|e^{-it\Delta} f\|_{L^p_t L^q_x} \lesssim \|f\|_{L^2_x}$ for all $f \in L^2_x$, then $\|u\|_{L^p_t L^q_x} \lesssim \|u\|_{U^2_\Delta}$ for all $u \in U^2_\Delta$.
- Bilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, then $\|uv\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta}$
- Trilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot e^{-it\Delta} h\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$, then $\|uvw\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$.

U^2_Δ is almost the perfect space... Except for duality.

$$|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad U^2_\Delta \subset V^2_\Delta$$

Estimate we need

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}$$

Estimate we can transfer

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}.$$

Transference from free solutions

U^2_Δ-functions are "almost free solutions" and most estimates for free solutions also hold in this space:

- Linear: if $\|e^{-it\Delta} f\|_{L^p_t L^q_x} \lesssim \|f\|_{L^2_x}$ for all $f \in L^2_x$, then $\|u\|_{L^p_t L^q_x} \lesssim \|u\|_{U^2_\Delta}$ for all $u \in U^2_\Delta$.
- Bilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, then $\|uv\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta}$
- Trilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot e^{-it\Delta} h\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$, then $\|uvw\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$.

U^2_Δ is almost the perfect space... Except for duality.

$$|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad U^2_\Delta \subset V^2_\Delta$$

Estimate we need

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}$$

Estimate we can transfer

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}.$$

Transference from free solutions

U^2_Δ-functions are "almost free solutions" and most estimates for free solutions also hold in this space:

- Linear: if $\|e^{-it\Delta} f\|_{L^p_t L^q_x} \lesssim \|f\|_{L^2_x}$ for all $f \in L^2_x$, then $\|u\|_{L^p_t L^q_x} \lesssim \|u\|_{U^2_\Delta}$ for all $u \in U^2_\Delta$.
- Bilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, then $\|uv\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta}$
- Trilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot e^{-it\Delta} h\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$, then $\|uvw\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$.

U^2_Δ is almost the perfect space... Except for duality.

$$|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad U^2_\Delta \subset V^2_\Delta$$

Estimate we need

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}$$

Estimate we can transfer

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}.$$

Transference from free solutions

U^2_Δ-functions are "almost free solutions" and most estimates for free solutions also hold in this space:

- Linear: if $\|e^{-it\Delta} f\|_{L^p_t L^q_x} \lesssim \|f\|_{L^2_x}$ for all $f \in L^2_x$, then $\|u\|_{L^p_t L^q_x} \lesssim \|u\|_{U^2_\Delta}$ for all $u \in U^2_\Delta$.
- Bilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, then $\|uv\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta}$
- Trilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot e^{-it\Delta} h\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$, then $\|uvw\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$.

U^2_Δ is almost the perfect space... Except for duality.

$$|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad U^2_\Delta \subset V^2_\Delta$$

Estimate we need

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}$$

Estimate we can transfer

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}.$$

Transference from free solutions

U^2_Δ-functions are "almost free solutions" and most estimates for free solutions also hold in this space:

- Linear: if $\|e^{-it\Delta} f\|_{L^p_t L^q_x} \lesssim \|f\|_{L^2_x}$ for all $f \in L^2_x$, then $\|u\|_{L^p_t L^q_x} \lesssim \|u\|_{U^2_\Delta}$ for all $u \in U^2_\Delta$.
- Bilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, then $\|uv\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta}$
- Trilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot e^{-it\Delta} h\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$, then $\|uvw\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$.

U^2_Δ is almost the perfect space... Except for duality.

$$|\langle uv, w\rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad U^2_\Delta \subset V^2_\Delta$$

Estimate we need

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}$$

Estimate we can transfer

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}.$$

Transference from free solutions

U^2_Δ-functions are "almost free solutions" and most estimates for free solutions also hold in this space:

- Linear: if $\|e^{-it\Delta} f\|_{L^p_t L^q_x} \lesssim \|f\|_{L^2_x}$ for all $f \in L^2_x$, then $\|u\|_{L^p_t L^q_x} \lesssim \|u\|_{U^2_\Delta}$ for all $u \in U^2_\Delta$.
- Bilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, then $\|uv\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta}$
- Trilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot e^{-it\Delta} h\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$, then $\|uvw\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$.

U^2_Δ is almost the perfect space... Except for duality.

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}, \qquad U^2_\Delta \subset V^2_\Delta$$

Estimate we need

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}$$

Estimate we can transfer

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}.$$

Transference from free solutions

U^2_Δ-functions are "almost free solutions" and most estimates for free solutions also hold in this space:

- Linear: if $\|e^{-it\Delta} f\|_{L^p_t L^q_x} \lesssim \|f\|_{L^2_x}$ for all $f \in L^2_x$, then $\|u\|_{L^p_t L^q_x} \lesssim \|u\|_{U^2_\Delta}$ for all $u \in U^2_\Delta$.
- Bilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, then $\|uv\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta}$
- Trilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot e^{-it\Delta} h\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$, then $\|uvw\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$.

U^2_Δ is almost the perfect space... Except for duality.

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{Y^*}, \qquad U^2_\Delta \subset V^2_\Delta$$

Estimate we need

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}$$

Estimate we can transfer

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}.$$

Transference from free solutions

U^2_Δ-functions are "almost free solutions" and most estimates for free solutions also hold in this space:

- Linear: if $\|e^{-it\Delta} f\|_{L^p_t L^q_x} \lesssim \|f\|_{L^2_x}$ for all $f \in L^2_x$, then $\|u\|_{L^p_t L^q_x} \lesssim \|u\|_{U^2_\Delta}$ for all $u \in U^2_\Delta$.
- Bilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, then $\|uv\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta}$
- Trilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot e^{-it\Delta} h\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$, then $\|uvw\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$.

U^2_Δ is almost the perfect space... Except for duality.

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}, \qquad U^2_\Delta \subset V^2_\Delta$$

Estimate we need

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}$$

Estimate we can transfer

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}.$$

Transference from free solutions

U^2_Δ-functions are "almost free solutions" and most estimates for free solutions also hold in this space:

- Linear: if $\|e^{-it\Delta} f\|_{L^p_t L^q_x} \lesssim \|f\|_{L^2_x}$ for all $f \in L^2_x$, then $\|u\|_{L^p_t L^q_x} \lesssim \|u\|_{U^2_\Delta}$ for all $u \in U^2_\Delta$.
- Bilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, then $\|uv\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta}$
- Trilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot e^{-it\Delta} h\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$, then $\|uvw\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$.

U^2_Δ is almost the perfect space... Except for duality.

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}, \qquad U^2_\Delta \subset V^2_\Delta$$

Estimate we need

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}$$

Estimate we can transfer

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}.$$

Transference from free solutions

U^2_Δ-functions are "almost free solutions" and most estimates for free solutions also hold in this space:

- Linear: if $\|e^{-it\Delta} f\|_{L^p_t L^q_x} \lesssim \|f\|_{L^2_x}$ for all $f \in L^2_x$, then $\|u\|_{L^p_t L^q_x} \lesssim \|u\|_{U^2_\Delta}$ for all $u \in U^2_\Delta$.
- Bilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x}$, then $\|uv\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta}$
- Trilinear: if $\|e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot e^{-it\Delta} h\|_{L^q_{t,x}} \lesssim \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$, then $\|uvw\|_{L^q_{t,x}} \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$.

U^2_Δ is almost the perfect space... Except for duality.

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}, \qquad U^2_\Delta \subset V^2_\Delta$$

Estimate we need

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{V^2_\Delta}$$

Estimate we can transfer

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}.$$

Candidate 3: the 2-variation space V_Δ^2

$$V_\Delta^2 = \left\{ e^{-it\Delta} v(t) : \sup_{\{t_j\} \text{ partitions}} \sum_j \|v(t_{j+1}) - v(t_j)\|_{L_x^2}^2 < \infty \right\}$$

When it comes to transference, V_Δ^2 is *almost* as good as U_Δ^2:

- $U_\Delta^2 \subset V_\Delta^2 \subset U_\Delta^a, \quad a > 2.$
- if $\|e^{-it\Delta} f\|_{L_t^p L_x^q} \lesssim \|f\|_{L_x^2}$, then $\|v\|_{L_t^p L_x^q} \lesssim \|v\|_{V_\Delta^2}$, provided $p > 2$.

Recovering these almost-endpoint losses often make V_Δ^2 hard to use efficiently.

Variation-space duality

But while harder to prove, V_Δ^2 duality is perfect for PDEs: if $X = V_\Delta^2$, then

$$|\langle uv, w \rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}$$

So **if** we can show V_Δ^2 transference $\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$, we solve (2NLS)!

Candidate 3: the 2-variation space V_Δ^2

$$V_\Delta^2 = \left\{ e^{-it\Delta} v(t) : \sup_{\{t_j\} \text{ partitions}} \sum_j \|v(t_{j+1}) - v(t_j)\|_{L_x^2}^2 < \infty \right\}$$

When it comes to transference, V_Δ^2 is *almost* as good as U_Δ^2:

- $U_\Delta^2 \subset V_\Delta^2 \subset U_\Delta^a, \quad a > 2.$
- if $\|e^{-it\Delta} f\|_{L_t^p L_x^q} \lesssim \|f\|_{L_x^2}$, then $\|v\|_{L_t^p L_x^q} \lesssim \|v\|_{V_\Delta^2}$, provided $p > 2$.

Recovering these almost-endpoint losses often make V_Δ^2 hard to use efficiently.

Variation-space duality

But while harder to prove, V_Δ^2 duality is perfect for PDEs: if $X = V_\Delta^2$, then

$$|\langle uv, w \rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}$$

So **if** we can show V_Δ^2 transference $\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$, we solve (2NLS)!

Candidate 3: the 2-variation space V_Δ^2

$$V_\Delta^2 = \left\{ e^{-it\Delta} v(t) : \sup_{\{t_j\} \text{ partitions}} \sum_j \|v(t_{j+1}) - v(t_j)\|_{L_x^2}^2 < \infty \right\}$$

When it comes to transference, V_Δ^2 is *almost* as good as U_Δ^2:

- $U_\Delta^2 \subset V_\Delta^2 \subset U_\Delta^a, \quad a > 2.$
- if $\|e^{-it\Delta} f\|_{L_t^p L_x^q} \lesssim \|f\|_{L_x^2}$, then $\|v\|_{L_t^p L_x^q} \lesssim \|v\|_{V_\Delta^2}$, provided $p > 2$.

Recovering these almost-endpoint losses often make V_Δ^2 hard to use efficiently.

Variation-space duality

But while harder to prove, V_Δ^2 duality is perfect for PDEs: if $X = V_\Delta^2$, then

$$|\langle uv, w \rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}$$

So **if** we can show V_Δ^2 transference $\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$, we solve (2NLS)!

Candidate 3: the 2-variation space V_Δ^2

$$V_\Delta^2 = \left\{ e^{-it\Delta} v(t) : \sup_{\{t_j\} \text{ partitions}} \sum_j \|v(t_{j+1}) - v(t_j)\|_{L_x^2}^2 < \infty \right\}$$

When it comes to transference, V_Δ^2 is *almost* as good as U_Δ^2:

- $U_\Delta^2 \subset V_\Delta^2 \subset U_\Delta^a, \quad a > 2.$
- if $\|e^{-it\Delta} f\|_{L_t^p L_x^q} \lesssim \|f\|_{L_x^2}$, then $\|v\|_{L_t^p L_x^q} \lesssim \|v\|_{V_\Delta^2}$, provided $p > 2$.

Recovering these almost-endpoint losses often make V_Δ^2 hard to use efficiently.

Variation-space duality

But while harder to prove, V_Δ^2 duality is perfect for PDEs: if $X = V_\Delta^2$, then

$$|\langle uv, w \rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}$$

So **if** we can show V_Δ^2 transference $\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$, we solve (2NLS)!

Candidate 3: the 2-variation space V_Δ^2

$$V_\Delta^2 = \left\{ e^{-it\Delta} v(t) : \sup_{\{t_j\} \text{ partitions}} \sum_j \|v(t_{j+1}) - v(t_j)\|_{L_x^2}^2 < \infty \right\}$$

When it comes to transference, V_Δ^2 is *almost* as good as U_Δ^2:

- $U_\Delta^2 \subset V_\Delta^2 \subset U_\Delta^a, \quad a > 2.$
- if $\|e^{-it\Delta} f\|_{L_t^p L_x^q} \lesssim \|f\|_{L_x^2}$, then $\|v\|_{L_t^p L_x^q} \lesssim \|v\|_{V_\Delta^2}$, provided $p > 2$.

Recovering these almost-endpoint losses often make V_Δ^2 hard to use efficiently.

Variation-space duality

But while harder to prove, V_Δ^2 duality is perfect for PDEs: if $X = V_\Delta^2$, then

$$|\langle uv, w \rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}$$

So **if** we can show V_Δ^2 transference $\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$, we solve (2NLS)!

Candidate 3: the 2-variation space V_Δ^2

$$V_\Delta^2 = \left\{ e^{-it\Delta} v(t) : \sup_{\{t_j\} \text{ partitions}} \sum_j \|v(t_{j+1}) - v(t_j)\|_{L_x^2}^2 < \infty \right\}$$

When it comes to transference, V_Δ^2 is *almost* as good as U_Δ^2:

- $U_\Delta^2 \subset V_\Delta^2 \subset U_\Delta^a, \quad a > 2.$
- if $\|e^{-it\Delta} f\|_{L_t^p L_x^q} \lesssim \|f\|_{L_x^2}$, then $\|v\|_{L_t^p L_x^q} \lesssim \|v\|_{V_\Delta^2}$, provided $p > 2$.

Recovering these almost-endpoint losses often make V_Δ^2 hard to use efficiently.

Variation-space duality

But while harder to prove, V_Δ^2 duality is perfect for PDEs: if $X = V_\Delta^2$, then

$$|\langle uv, w \rangle| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}$$

So **if** we can show V_Δ^2 transference $\left| \int u v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$, we solve (2NLS)!

Candidate 3: the 2-variation space V_Δ^2

$$V_\Delta^2 = \left\{ e^{-it\Delta} v(t) : \sup_{\{t_j\} \text{ partitions}} \sum_j \|v(t_{j+1}) - v(t_j)\|_{L_x^2}^2 < \infty \right\}$$

When it comes to transference, V_Δ^2 is *almost* as good as U_Δ^2:

- $U_\Delta^2 \subset V_\Delta^2 \subset U_\Delta^a, \quad a > 2.$
- if $\|e^{-it\Delta} f\|_{L_t^p L_x^q} \lesssim \|f\|_{L_x^2}$, then $\|v\|_{L_t^p L_x^q} \lesssim \|v\|_{V_\Delta^2}$, provided $p > 2$.

Recovering these almost-endpoint losses often make V_Δ^2 hard to use efficiently.

Variation-space duality

But while harder to prove, V_Δ^2 duality is perfect for PDEs: if $X = V_\Delta^2$, then

$$\left| \int u v \bar{w} \, dx \, dt \right| \lesssim \|u\|_X \|v\|_X \|w\|_{Y^*}$$

So **if** we can show V_Δ^2 transference $\left| \int u v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$, we solve (2NLS)!

Candidate 3: the 2-variation space V_Δ^2

$$V_\Delta^2 = \left\{ e^{-it\Delta} v(t) : \sup_{\{t_j\} \text{ partitions}} \sum_j \|v(t_{j+1}) - v(t_j)\|_{L_x^2}^2 < \infty \right\}$$

When it comes to transference, V_Δ^2 is *almost* as good as U_Δ^2:

- $U_\Delta^2 \subset V_\Delta^2 \subset U_\Delta^a, \quad a > 2.$
- if $\|e^{-it\Delta} f\|_{L_t^p L_x^q} \lesssim \|f\|_{L_x^2}$, then $\|v\|_{L_t^p L_x^q} \lesssim \|v\|_{V_\Delta^2}$, provided $p > 2$.

Recovering these almost-endpoint losses often make V_Δ^2 hard to use efficiently.

Variation-space duality

But while harder to prove, V_Δ^2 duality is perfect for PDEs: if $X = V_\Delta^2$, then

$$\left| \int u v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{U_\Delta^2}$$

So **if** we can show V_Δ^2 transference $\left| \int u v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$, we solve (2NLS)!

Candidate 3: the 2-variation space V^2_Δ

$$V^2_\Delta = \left\{ e^{-it\Delta}v(t) : \sup_{\{t_j\} \text{ partitions}} \sum_j \|v(t_{j+1}) - v(t_j)\|^2_{L^2_x} < \infty \right\}$$

When it comes to transference, V^2_Δ is *almost* as good as U^2_Δ:

- $U^2_\Delta \subset V^2_\Delta \subset U^a_\Delta, \quad a > 2.$
- if $\|e^{-it\Delta} f\|_{L^p_t L^q_x} \lesssim \|f\|_{L^2_x}$, then $\|v\|_{L^p_t L^q_x} \lesssim \|v\|_{V^2_\Delta}$, provided $p > 2$.

Recovering these almost-endpoint losses often make V^2_Δ hard to use efficiently.

Variation-space duality

But while harder to prove, V^2_Δ duality is perfect for PDEs: if $X = V^2_\Delta$, then

$$\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{U^2_\Delta}$$

So **if** we can show V^2_Δ transference $\left| \int uv\bar{w}\, dx\, dt \right| \lesssim \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$, we solve (2NLS)!

Recap: transference

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

- Linear wave packets
- Bilinear range is very good in the radial case
- Transversality improves integrability
- Trilinear transversality is better than bilinear transversality

(iv) From harmonic analysis to PDEs

(v) The fun part: everything coming together

Recap: transference

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

- Linear wave packets
- Bilinear range is very good in the radial case
- Transversality improves integrability
- Trilinear transversality is better than bilinear transversality

(iv) From harmonic analysis to PDEs

- Strichartz spaces are good for linear transference

(v) The fun part: everything coming together

Recap: transference

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

- Linear wave packets
- Bilinear range is very good in the radial case
- Transversality improves integrability
- Trilinear transversality is better than bilinear transversality

(iv) From harmonic analysis to PDEs

- Strichartz spaces are good for linear transference
- U^2_Δ is great for multilinear transference

(v) The fun part: everything coming together

Recap: transference

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

- Linear wave packets
- Bilinear range is very good in the radial case
- Transversality improves integrability
- Trilinear transversality is better than bilinear transversality

(iv) From harmonic analysis to PDEs

- Strichartz spaces are good for linear transference
- U^2_Δ is great for multilinear transference
- U^2_Δ has bad duality

(v) The fun part: everything coming together

Recap: transference

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

- Linear wave packets
- Bilinear range is very good in the radial case
- Transversality improves integrability
- Trilinear transversality is better than bilinear transversality

(iv) From harmonic analysis to PDEs

- Strichartz spaces are good for linear transference
- U^2_Δ is great for multilinear transference
- U^2_Δ has bad duality
- V^2_Δ has good duality, but transference is harder

(v) The fun part: everything coming together

Main result: revisited

Theorem (Bejenaru–B.–Chan)

The Schrödinger equation

$$(i\partial_t - \Delta)u = c_1 u^2 + c_2 \bar{u}^2, \qquad c_1, c_2 \in \mathbb{C}, \tag{2NLS}$$

with small initial data $u_0 \in \dot{H}^{-\frac{1}{2}}(\mathbb{R}^3)$ in the scaling-critical space is globally well-posed, scatters to free solutions, and has Lipschitz dependence on its initial data. The solution is also unique in the resolution space $\dot{X}^{-\frac{1}{2}} \subset C_t \dot{H}_x^{-\frac{1}{2}}$.

In order to prove this, we need to:

- Classify the resonances
- Check whether resonant interactions are transversal
- Split $u = u_{\text{res}} + u_{\text{nonr}}$ and establish

$$\left| \int_{\mathbb{R}^{1+3}} u_{\text{nonr}} v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}, \qquad \left| \int_{\mathbb{R}^{1+3}} u_{\text{res}} v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$$

Main result: revisited

Theorem (Bejenaru–B.–Chan)

The Schrödinger equation

$$(i\partial_t - \Delta)u = c_1 u^2 + c_2 \bar{u}^2, \qquad c_1, c_2 \in \mathbb{C}, \tag{2NLS}$$

with small initial data $u_0 \in \dot{H}^{-\frac{1}{2}}(\mathbb{R}^3)$ in the scaling-critical space is globally well-posed, scatters to free solutions, and has Lipschitz dependence on its initial data. The solution is also unique in the resolution space $\dot{X}^{-\frac{1}{2}} \subset C_t \dot{H}_x^{-\frac{1}{2}}$.

In order to prove this, we need to:

- Classify the resonances
- Check whether resonant interactions are transversal
- Split $u = u_{\text{res}} + u_{\text{nonr}}$ and establish

$$\left| \int_{\mathbb{R}^{1+3}} u_{\text{nonr}} v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}, \qquad \left| \int_{\mathbb{R}^{1+3}} u_{\text{res}} v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$$

Main result: revisited

Theorem (Bejenaru–B.–Chan)

The Schrödinger equation

$$(i\partial_t - \Delta)u = c_1 u^2 + c_2 \bar{u}^2, \qquad c_1, c_2 \in \mathbb{C}, \tag{2NLS}$$

with small initial data $u_0 \in \dot{H}^{-\frac{1}{2}}(\mathbb{R}^3)$ in the scaling-critical space is globally well-posed, scatters to free solutions, and has Lipschitz dependence on its initial data. The solution is also unique in the resolution space $\dot{X}^{-\frac{1}{2}} \subset C_t \dot{H}_x^{-\frac{1}{2}}$.

In order to prove this, we need to:

- Classify the resonances
- Check whether resonant interactions are transversal
- Split $u = u_{\text{res}} + u_{\text{nonr}}$ and establish

$$\left| \int_{\mathbb{R}^{1+3}} u_{\text{nonr}} v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}, \qquad \left| \int_{\mathbb{R}^{1+3}} u_{\text{res}} v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$$

Main result: revisited

Theorem (Bejenaru–B.–Chan)

The Schrödinger equation

$$(i\partial_t - \Delta)u = c_1 u^2 + c_2 \bar{u}^2, \qquad c_1, c_2 \in \mathbb{C}, \tag{2NLS}$$

with small initial data $u_0 \in \dot{H}^{-\frac{1}{2}}(\mathbb{R}^3)$ in the scaling-critical space is globally well-posed, scatters to free solutions, and has Lipschitz dependence on its initial data. The solution is also unique in the resolution space $\dot{X}^{-\frac{1}{2}} \subset C_t \dot{H}_x^{-\frac{1}{2}}$.

In order to prove this, we need to:

- Classify the resonances
- Check whether resonant interactions are transversal
- Split $u = u_{\text{res}} + u_{\text{nonr}}$ and establish

$$\left| \int_{\mathbb{R}^{1+3}} u_{\text{nonr}} v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}, \qquad \left| \int_{\mathbb{R}^{1+3}} u_{\text{res}} v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$$

Main result: revisited

Theorem (Bejenaru–B.–Chan)

The Schrödinger equation

$$(i\partial_t - \Delta)u = c_1 u^2 + c_2 \bar{u}^2, \qquad c_1, c_2 \in \mathbb{C}, \tag{2NLS}$$

with small initial data $u_0 \in \dot{H}^{-\frac{1}{2}}(\mathbb{R}^3)$ in the scaling-critical space is globally well-posed, scatters to free solutions, and has Lipschitz dependence on its initial data. The solution is also unique in the resolution space $\dot{X}^{-\frac{1}{2}} \subset C_t \dot{H}_x^{-\frac{1}{2}}$.

In order to prove this, we need to:

- Classify the resonances
- Check whether resonant interactions are transversal
- Split $u = u_{\text{res}} + u_{\text{nonr}}$ and establish

$$\left| \int_{\mathbb{R}^{1+3}} u_{\text{nonr}} v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}, \qquad \left| \int_{\mathbb{R}^{1+3}} u_{\text{res}} v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$$

Main result: revisited

Theorem (Bejenaru–B.–Chan)

The Schrödinger equation

$$(i\partial_t - \Delta)u = c_1 u^2 + c_2 \bar{u}^2, \qquad c_1, c_2 \in \mathbb{C}, \tag{2NLS}$$

with small initial data $u_0 \in \dot{H}^{-\frac{1}{2}}(\mathbb{R}^3)$ in the scaling-critical space is globally well-posed, scatters to free solutions, and has Lipschitz dependence on its initial data. The solution is also unique in the resolution space $\dot{X}^{-\frac{1}{2}} \subset C_t \dot{H}_x^{-\frac{1}{2}}$.

In order to prove this, we need to:

- Classify the resonances
- Check whether resonant interactions are transversal
- Split $u = u_{\text{res}} + u_{\text{nonr}}$ and establish

$$\left| \int_{\mathbb{R}^{1+3}} u_{\text{nonr}} v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}, \qquad \left| \int_{\mathbb{R}^{1+3}} u_{\text{res}} v \bar{w} \, dx \, dt \right| \lesssim \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$$

Resonances for $(i\partial_t - \Delta)u = u^2$

$$\int_{\mathbb{R}^{1+3}} uv\bar{w}\,dx\,dt = \left(\hat{u} * \hat{v} * \hat{\bar{w}}\right)(0) = \int \hat{u}(\xi,\tau)\hat{v}(\eta,\tilde{\tau})\overline{\hat{w}}(\xi+\eta,\tau+\tilde{\tau})\,d\xi\,d\eta\,d\tau\,d\tilde{\tau}$$

Resonant interactions occur exactly when u, v, and w are free waves, which are (Fourier) supported on the paraboloid $\tau = \xi^2$. So resonances occur when $\xi^2 + \eta^2 = (\xi+\eta)^2$, that is, when $\xi \cdot \eta = 0$.

To check transversality, we compute $|N_\xi \wedge N_\eta \wedge N_{\xi+\eta}| \approx \min\{|\xi|, |\eta|\}$, which degenerates at low frequencies!

Resonances for $(i\partial_t - \Delta)u = u^2$

$$\int_{\mathbb{R}^{1+3}} uv\bar{w}\,dx\,dt = \left(\hat{u} * \hat{v} * \hat{\bar{w}}\right)(0) = \int \hat{u}(\xi,\tau)\hat{v}(\eta,\tilde{\tau})\overline{\hat{w}}(\xi+\eta,\tau+\tilde{\tau})\,d\xi\,d\eta\,d\tau\,d\tilde{\tau}$$

Resonant interactions occur exactly when u, v, and w are free waves, which are (Fourier) supported on the paraboloid $\tau = \xi^2$. So resonances occur when $\xi^2 + \eta^2 = (\xi+\eta)^2$, that is, when $\xi \cdot \eta = 0$.

To check transversality, we compute $|N_\xi \wedge N_\eta \wedge N_{\xi+\eta}| \approx \min\{|\xi|, |\eta|\}$, which degenerates at low frequencies!


Resonances for $(i\partial_t - \Delta)u = u^2$


$$\int_{\mathbb{R}^{1+3}} uv\bar{w}\,dx\,dt = \left(\hat{u} * \hat{v} * \hat{\bar{w}}\right)(0) = \int \hat{u}(\xi,\tau)\hat{v}(\eta,\tilde{\tau})\overline{\hat{w}}(\xi+\eta,\tau+\tilde{\tau})\,d\xi\,d\eta\,d\tau\,d\tilde{\tau}$$

Resonant interactions occur exactly when u, v, and w are free waves, which are (Fourier) supported on the paraboloid $\tau = \xi^2$. So resonances occur when $\xi^2 + \eta^2 = (\xi+\eta)^2$, that is, when $\xi \cdot \eta = 0$.

To check transversality, we compute $|N_\xi \wedge N_\eta \wedge N_{\xi+\eta}| \approx \min\{|\xi|, |\eta|\}$, which degenerates at low frequencies!

Three-wave interactions





Nonresonant estimates

We can assume that v, w have frequency 1 and u has frequency 2^k, with $k \leq 0$.
Let $Q_j u$ localize u to the region $|\tau - \xi^2| \approx 2^j, j \in \mathbb{Z}$.

High modulation on u

High modulation on v


Schematic frequency–modulation slices; hatched regions have $|\tau - |\xi|^2| \gtrsim 2^{2k}$.

Claim: $\left|\int_{\mathbb{R}^{1+3}} (Q_{\geq 2k}u) v \bar{w}\, dx\, dt\right| + \left|\int_{\mathbb{R}^{1+3}} u (Q_{\geq 2k}v) \bar{w}\, dx\, dt\right| \lesssim \langle k \rangle^3 \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$


Nonresonant estimates

We can assume that v, w have frequency 1 and u has frequency 2^k, with $k \leq 0$.
Let $Q_j u$ localize u to the region $|\tau - \xi^2| \approx 2^j$, $j \in \mathbb{Z}$.

High modulation on u



High modulation on v




Schematic frequency–modulation slices; hatched regions have $|\tau - |\xi|^2| \gtrsim 2^{2k}$.

Claim: $\left|\int_{\mathbb{R}^{1+3}} (Q_{\geq 2k}u) v \bar{w}\, dx\, dt\right| + \left|\int_{\mathbb{R}^{1+3}} u (Q_{\geq 2k}v) \bar{w}\, dx\, dt\right| \lesssim \langle k \rangle^3 \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$


Nonresonant estimates

We can assume that v, w have frequency 1 and u has frequency 2^k, with $k \leq 0$.
Let $Q_j u$ localize u to the region $|\tau - \xi^2| \approx 2^j$, $j \in \mathbb{Z}$.

High modulation on u



High modulation on v



Schematic frequency–modulation slices; hatched regions have $|\tau - |\xi|^2| \gtrsim 2^{2k}$.

Claim: $\left|\int_{\mathbb{R}^{1+3}} (Q_{\geq 2k}u) v \bar{w}\, dx\, dt\right| + \left|\int_{\mathbb{R}^{1+3}} u (Q_{\geq 2k}v) \bar{w}\, dx\, dt\right| \lesssim \langle k \rangle^3 \|u\|_{V_\Delta^2} \|v\|_{V_\Delta^2} \|w\|_{V_\Delta^2}$

Techniques: bilinear $L^2_{t,x}$, $\dot{X}^{0,1/2}$, Fourier cube decomposition

Resonant estimates

Goal: $\left| \int_{\mathbb{R}^{1+3}} Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\, dx\, dt \right| \leq C_\epsilon 2^{\epsilon k} \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$

Step 1: Establish the free wave estimate

$$\left| \int_{\mathbb{R}^{1+3}} e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot \overline{e^{-it\Delta} h}\, dx\, dt \right| \leq C_\epsilon 2^{-k\epsilon} \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$$

- Bejenaru '22: Free wave estimate for functions at frequency 1.
- Bejenaru–Chan '26 (companion paper): Polynomial partitioning, induction on scales, ϵ-removal.

Step 2: Transfer the estimate from free waves to V^2_Δ functions

$$\left| \int_{\mathbb{R}^{1+3}} Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\, dx\, dt \right| \leq C_\epsilon 2^{\epsilon k} \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$$

Resonant estimates

Goal: $\left| \int_{\mathbb{R}^{1+3}} Q_{<2k}u \cdot Q_{<2k}v \cdot \overline{Q_{<2k}w}\, dx\, dt \right| \leq C_\epsilon 2^{\epsilon k} \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$

Step 1: Establish the free wave estimate

$$\left| \int_{\mathbb{R}^{1+3}} e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot \overline{e^{-it\Delta} h}\, dx\, dt \right| \leq C_\epsilon 2^{-k\epsilon} \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$$

- Bejenaru '22: Free wave estimate for functions at frequency 1.
- Bejenaru–Chan '26 (companion paper): Polynomial partitioning, induction on scales, ϵ-removal.

Step 2: Transfer the estimate from free waves to V^2_Δ functions

$$\left| \int_{\mathbb{R}^{1+3}} Q_{<2k}u \cdot Q_{<2k}v \cdot \overline{Q_{<2k}w}\, dx\, dt \right| \leq C_\epsilon 2^{\epsilon k} \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$$

Resonant estimates

Goal: $\left| \int_{\mathbb{R}^{1+3}} Q_{<2k}u \cdot Q_{<2k}v \cdot \overline{Q_{<2k}w}\, dx\, dt \right| \leq C_\epsilon 2^{\epsilon k} \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$

Step 1: Establish the free wave estimate

$$\left| \int_{\mathbb{R}^{1+3}} e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot \overline{e^{-it\Delta} h}\, dx\, dt \right| \leq C_\epsilon 2^{-k\epsilon} \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$$

- Bejenaru '22: Free wave estimate for functions at frequency 1.
- Bejenaru–Chan '26 (companion paper): Polynomial partitioning, induction on scales, ϵ-removal.

Step 2: Transfer the estimate from free waves to V^2_Δ functions

$$\left| \int_{\mathbb{R}^{1+3}} Q_{<2k}u \cdot Q_{<2k}v \cdot \overline{Q_{<2k}w}\, dx\, dt \right| \leq C_\epsilon 2^{\epsilon k} \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$$

Resonant estimates

Goal: $\left| \int_{\mathbb{R}^{1+3}} Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\, dx\, dt \right| \leq C_\epsilon 2^{\epsilon k} \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$

Step 1: Establish the free wave estimate

$$\left| \int_{\mathbb{R}^{1+3}} e^{-it\Delta} f \cdot e^{-it\Delta} g \cdot \overline{e^{-it\Delta} h}\, dx\, dt \right| \leq C_\epsilon 2^{-k\epsilon} \|f\|_{L^2_x} \|g\|_{L^2_x} \|h\|_{L^2_x}$$

- Bejenaru '22: Free wave estimate for functions at frequency 1.
- Bejenaru–Chan '26 (companion paper): Polynomial partitioning, induction on scales, ϵ-removal.

Step 2: Transfer the estimate from free waves to V^2_Δ functions

$$\left| \int_{\mathbb{R}^{1+3}} Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\, dx\, dt \right| \leq C_\epsilon 2^{\epsilon k} \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$$

Trilinear transference to V^2_Δ

Goal:

$$\Big| \int_{\mathbb{R}^{1+3}} Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\, dx\, dt \Big| \leq C_\epsilon 2^{\epsilon k} \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$$

V^2_Δ does not have an atomic structure, and it is hard to work with directly. So instead we show that

$$\|Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\|_{\ell^p L^1(\mathbb{R}^{1+3})} \leq C_\epsilon 2^{\epsilon k} \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$$
$$\|Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\|_{\ell^\infty L^1(\mathbb{R}^{1+3})} \leq C_\epsilon 2^{\epsilon k} \|u\|_{U^a_\Delta} \|v\|_{U^a_\Delta} \|w\|_{U^a_\Delta},$$

for some $\frac{14}{15} < p < 1$ and $a > 2$. Recall that $U^2_\Delta \subset V^2_\Delta \subset U^a_\Delta$, and so V^2_Δ-transference follows from interpolation.

The fact that we can prove an estimate for $p < 1$ is crucial for the interpolation, which means that the curvature of the paraboloid plays a key role in this result.

Trilinear transference to V^2_Δ

Goal:

$$\Big| \int_{\mathbb{R}^{1+3}} Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\, dx\, dt \Big| \leq C_\epsilon 2^{\epsilon k} \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$$

V^2_Δ does not have an atomic structure, and it is hard to work with directly. So instead we show that

$$\|Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\|_{\ell^p L^1(\mathbb{R}^{1+3})} \leq C_\epsilon 2^{\epsilon k} \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$$
$$\|Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\|_{\ell^\infty L^1(\mathbb{R}^{1+3})} \leq C_\epsilon 2^{\epsilon k} \|u\|_{U^a_\Delta} \|v\|_{U^a_\Delta} \|w\|_{U^a_\Delta},$$

for some $\frac{14}{15} < p < 1$ and $a > 2$. Recall that $U^2_\Delta \subset V^2_\Delta \subset U^a_\Delta$, and so V^2_Δ-transference follows from interpolation.

The fact that we can prove an estimate for $p < 1$ is crucial for the interpolation, which means that the curvature of the paraboloid plays a key role in this result.

Trilinear transference to V^2_Δ

Goal:

$$\Big| \int_{\mathbb{R}^{1+3}} Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\, dx\, dt \Big| \leq C_\epsilon 2^{\epsilon k} \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$$

V^2_Δ does not have an atomic structure, and it is hard to work with directly. So instead we show that

$$\|Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\|_{\ell^p L^1(\mathbb{R}^{1+3})} \leq C_\epsilon 2^{\epsilon k} \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$$
$$\|Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\|_{\ell^\infty L^1(\mathbb{R}^{1+3})} \leq C_\epsilon 2^{\epsilon k} \|u\|_{U^a_\Delta} \|v\|_{U^a_\Delta} \|w\|_{U^a_\Delta},$$

for some $\frac{14}{15} < p < 1$ and $a > 2$. Recall that $U^2_\Delta \subset V^2_\Delta \subset U^a_\Delta$, and so V^2_Δ-transference follows from interpolation.

The fact that we can prove an estimate for $p < 1$ is crucial for the interpolation, which means that the curvature of the paraboloid plays a key role in this result.

Trilinear transference to V^2_Δ

Goal:

$$\Big| \int_{\mathbb{R}^{1+3}} Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\, dx\, dt \Big| \leq C_\epsilon 2^{\epsilon k} \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$$

V^2_Δ does not have an atomic structure, and it is hard to work with directly. So instead we show that

$$\|Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\|_{\ell^p L^1(\mathbb{R}^{1+3})} \leq C_\epsilon 2^{\epsilon k} \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$$
$$\|Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\|_{\ell^\infty L^1(\mathbb{R}^{1+3})} \leq C_\epsilon 2^{\epsilon k} \|u\|_{U^a_\Delta} \|v\|_{U^a_\Delta} \|w\|_{U^a_\Delta},$$

for some $\frac{14}{15} < p < 1$ and $a > 2$. Recall that $U^2_\Delta \subset V^2_\Delta \subset U^a_\Delta$, and so V^2_Δ-transference follows from interpolation.

The fact that we can prove an estimate for $p < 1$ is crucial for the interpolation, which means that the curvature of the paraboloid plays a key role in this result.

Trilinear transference to V^2_Δ

Goal:

$$\Big| \int_{\mathbb{R}^{1+3}} Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\, dx\, dt \Big| \leq C_\epsilon 2^{\epsilon k} \|u\|_{V^2_\Delta} \|v\|_{V^2_\Delta} \|w\|_{V^2_\Delta}$$

V^2_Δ does not have an atomic structure, and it is hard to work with directly. So instead we show that

$$\|Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\|_{\ell^p L^1(\mathbb{R}^{1+3})} \leq C_\epsilon 2^{\epsilon k} \|u\|_{U^2_\Delta} \|v\|_{U^2_\Delta} \|w\|_{U^2_\Delta}$$
$$\|Q_{<2^k} u \cdot Q_{<2^k} v \cdot \overline{Q_{<2^k} w}\|_{\ell^\infty L^1(\mathbb{R}^{1+3})} \leq C_\epsilon 2^{\epsilon k} \|u\|_{U^a_\Delta} \|v\|_{U^a_\Delta} \|w\|_{U^a_\Delta},$$

for some $\frac{14}{15} < p < 1$ and $a > 2$. Recall that $U^2_\Delta \subset V^2_\Delta \subset U^a_\Delta$, and so V^2_Δ-transference follows from interpolation.

The fact that we can prove an estimate for $p < 1$ is crucial for the interpolation, which means that the curvature of the paraboloid plays a key role in this result.

Full recap

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

- Linear wave packets
- Bilinear range is very good in the radial case
- Transversality improves integrability
- Trilinear transversality is better than bilinear transversality

(iv) From harmonic analysis to PDEs

- Strichartz spaces are good for linear transference
- U^2_Δ is great for multilinear transference
- U^2_Δ has bad duality
- V^2_Δ has good duality, but transference is harder

(v) Everything coming together

Full recap

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

- Linear wave packets
- Bilinear range is very good in the radial case
- Transversality improves integrability
- Trilinear transversality is better than bilinear transversality

(iv) From harmonic analysis to PDEs

- Strichartz spaces are good for linear transference
- U^2_Δ is great for multilinear transference
- U^2_Δ has bad duality
- V^2_Δ has good duality, but transference is harder

(v) Everything coming together

- Resonance analysis

Full recap

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

- Linear wave packets
- Bilinear range is very good in the radial case
- Transversality improves integrability
- Trilinear transversality is better than bilinear transversality

(iv) From harmonic analysis to PDEs

- Strichartz spaces are good for linear transference
- U^2_Δ is great for multilinear transference
- U^2_Δ has bad duality
- V^2_Δ has good duality, but transference is harder

(v) Everything coming together

- Resonance analysis
- Check that trilinear interactions are transversal

Full recap

Goal: Explore the proof of scattering for $(i\partial_t - \Delta)u = u^2$

(i) The building blocks: contraction theorems and high-dimensional waves

(ii) Why do simpler approaches fail when studying (2NLS)?

- Slow dispersion in 3d, Insufficient time integrability
- Bilinear range is insufficient in general

(iii) Harmonic analysis: geometry of interacting waves

- Linear wave packets
- Bilinear range is very good in the radial case
- Transversality improves integrability
- Trilinear transversality is better than bilinear transversality

(iv) From harmonic analysis to PDEs

- Strichartz spaces are good for linear transference
- U^2_Δ is great for multilinear transference
- U^2_Δ has bad duality
- V^2_Δ has good duality, but transference is harder

(v) Everything coming together

- Resonance analysis
- Check that trilinear interactions are transversal
- Use interpolation to transfer free estimates to V^2_Δ

TL;DR

- Multilinear wave interactions give contain extra information beyond the linear theory
- This information comes from transversality and curvature properties of the characteristic surface
- Using multilinear estimates beyond free solutions requires an transferring them to larger function spaces
- V^2_Δ is a very suitable transference space in a lot of cases

TL;DR

- Multilinear wave interactions give contain extra information beyond the linear theory
- This information comes from transversality and curvature properties of the characteristic surface
- Using multilinear estimates beyond free solutions requires an transferring them to larger function spaces
- V^2_Δ is a very suitable transference space in a lot of cases

TL;DR

- Multilinear wave interactions give contain extra information beyond the linear theory
- This information comes from transversality and curvature properties of the characteristic surface
- Using multilinear estimates beyond free solutions requires an transferring them to larger function spaces
- V^2_Δ is a very suitable transference space in a lot of cases

TL;DR

- Multilinear wave interactions give contain extra information beyond the linear theory
- This information comes from transversality and curvature properties of the characteristic surface
- Using multilinear estimates beyond free solutions requires an transferring them to larger function spaces
- V^2_Δ is a very suitable transference space in a lot of cases

TL;DR

- Multilinear wave interactions give contain extra information beyond the linear theory
- This information comes from transversality and curvature properties of the characteristic surface
- Using multilinear estimates beyond free solutions requires an transferring them to larger function spaces
- V^2_Δ is a very suitable transference space in a lot of cases

Thank you!

Bibliography (1/3)

[GMS09] Pierre Germain, Nader Masmoudi, and Jalal Shatah.
Global Solutions for 3D Quadratic Schrödinger Equations.
International Mathematics Research Notices 2009(3), 414–432 (2009).
doi:10.1093/imrn/rnn135.

[CKSTT02] James Colliander, Markus Keel, Gigliola Staffilani, Hideo Takaoka, and Terence Tao.
A Refined Global Well-Posedness Result for Schrödinger Equations with Derivative.
SIAM Journal on Mathematical Analysis 34(1), 64–86 (2002).
doi:10.1137/S0036141001394541; arXiv:math/0110026.

[BIKT11] Ioan Bejenaru, Alexandru D. Ionescu, Carlos E. Kenig, and Daniel Tataru.
Global Schrödinger Maps in Dimensions $d \geq 2$: Small Data in the Critical Sobolev Spaces.
Annals of Mathematics 173(3), 1443–1506 (2011).
doi:10.4007/annals.2011.173.3.5; arXiv:0807.0265.

[Sha85] Jalal Shatah.
Normal Forms and Quadratic Nonlinear Klein–Gordon Equations.
Communications on Pure and Applied Mathematics 38(5), 685–696 (1985).
doi:10.1002/cpa.3160380516.

[CO17] Myeongju Chae and Sung-Jin Oh.
Small Data Global Existence and Decay for Relativistic Chern–Simons Equations.
Annales Henri Poincaré 18, 2123–2198 (2017).
doi:10.1007/s00023-016-0547-9; arXiv:1512.03039.

[BB26] Ioan Bejenaru and Vitor Borges.
Global Well-Posedness and Scattering for the Massive Dirac–Klein–Gordon System in Two Dimensions.
Communications in Mathematical Physics 407, Article 7 (2026).
doi:10.1007/s00220-025-05499-9; arXiv:2501.03963.
Published online December 6, 2025.

[BC26] Vitor Borges and Tiklung Chan.
Scattering for the Klein–Gordon–Schrödinger System in Three Dimensions with Radial Data.
Preprint (2026).
arXiv:2604.10366.

[Tat01] Daniel Tataru.
On Global Existence and Scattering for the Wave Maps Equation.
American Journal of Mathematics 123(1), 37–77 (2001).
doi:10.1353/ajm.2001.0005.

[Tao01] Terence Tao.
Global Regularity of Wave Maps II. Small Energy in Two Dimensions.
Communications in Mathematical Physics 224(2), 443–544 (2001).
doi:10.1007/PL00005588; arXiv:math/0011173.

[KT17] Joachim Krieger and Daniel Tataru.
Global Well-Posedness for the Yang–Mills Equation in $4+1$ Dimensions. Small Energy.
Annals of Mathematics 185(3), 831–893 (2017).
doi:10.4007/annals.2017.185.3.3; arXiv:1509.00751.

Bibliography (2/3)

[CK93] Demetrios Christodoulou and Sergiu Klainerman.
The Global Nonlinear Stability of the Minkowski Space.
Princeton Mathematical Series, vol. 41. Princeton University Press, 1993.
doi:10.1515/9781400863174.
DOI for later electronic edition.

[LR10] Hans Lindblad and Igor Rodnianski.
The Global Stability of Minkowski Space-Time in Harmonic Gauge.
Annals of Mathematics 171(3), 1401–1477 (2010).
doi:10.4007/annals.2010.171.1401; arXiv:math/0411109.

[GMS12] Pierre Germain, Nader Masmoudi, and Jalal Shatah.
Global Solutions for the Gravity Water Waves Equation in Dimension 3.
Annals of Mathematics 175(2), 691–754 (2012).
doi:10.4007/annals.2012.175.2.6; arXiv:0906.5343.

[IT16] Mihaela Ifrim and Daniel Tataru.
Two Dimensional Water Waves in Holomorphic Coordinates II: Global Solutions.
Bulletin de la Société Mathématique de France 144(2), 369–394 (2016).
doi:10.24033/bsmf.2717; arXiv:1404.7583.

[FX18] Chenjie Fan and Weijun Xu.
Global Well-Posedness for the Mass-Critical Stochastic Nonlinear Schrödinger Equation on $\mathbb{R}$: Small Initial Data.
Preprint (2018).
arXiv:1803.03257.

[HTT11] Sebastian Herr, Daniel Tataru, and Nikolay Tzvetkov.
Global Well-Posedness of the Energy-Critical Nonlinear Schrödinger Equation with Small Initial Data in $H^1(\mathbb{T}^3)$.
Duke Mathematical Journal 159(2), 329–349 (2011).
doi:10.1215/00127094-1415889; arXiv:1005.2832.

[IT15] Mihaela Ifrim and Daniel Tataru.
Global Bounds for the Cubic Nonlinear Schrödinger Equation (NLS) in One Space Dimension.
Nonlinearity 28(8), 2661–2675 (2015).
doi:10.1088/0951-7715/28/8/2661; arXiv:1404.7581.

[IT23] Mihaela Ifrim and Daniel Tataru.
Global Solutions for 1D Cubic Defocusing Dispersive Equations: Part I.
Forum of Mathematics, Pi 11, e31 (2023).
doi:10.1017/fmp.2023.30; arXiv:2205.12212.

[IT24] Mihaela Ifrim and Daniel Tataru.
Global Solutions for Cubic Quasilinear Schrödinger Flows in Two and Higher Dimensions.
Preprint (2024), subsequently revised.
arXiv:2404.09970.

[BH17] Ioan Bejenaru and Sebastian Herr.
On Global Well-Posedness and Scattering for the Massive Dirac–Klein–Gordon System.
Journal of the European Mathematical Society 19(8), 2445–2467 (2017).
doi:10.4171/JEMS/721; arXiv:1409.1778.

Bibliography (3/3)

[KST15] Joachim Krieger, Jacob Sterbenz, and Daniel Tataru.
Global Well-Posedness for the Maxwell–Klein–Gordon Equation in $4+1$ Dimensions. Small Energy.
Duke Mathematical Journal 164(6), 973–1040 (2015).
doi:10.1215/00127094-2885982; arXiv:1211.3527.

[RT04] Igor Rodnianski and Terence Tao.
Global Regularity for the Maxwell–Klein–Gordon Equation with Small Critical Sobolev Norm in High Dimensions.
Communications in Mathematical Physics 251(2), 377–426 (2004).
doi:10.1007/s00220-004-1152-1; arXiv:math/0309353.

[IP19] Alexandru D. Ionescu and Benoît Pausader.
On the Global Regularity for a Wave–Klein–Gordon Coupled System.
Acta Mathematica Sinica, English Series 35(6), 933–986 (2019).
doi:10.1007/s10114-019-8413-6; arXiv:1703.02846.

[IP22] Alexandru D. Ionescu and Benoît Pausader.
The Einstein–Klein–Gordon Coupled System: Global Stability of the Minkowski Solution.
Annals of Mathematics Studies, vol. 213. Princeton University Press, 2022.
doi:10.1515/9780691233031; arXiv:1911.10652.

[KS23] Sergiu Klainerman and Jérémie Szeftel.
Kerr Stability for Small Angular Momentum.
Pure and Applied Mathematics Quarterly 19(3), 791–1678 (2023).
doi:10.4310/PAMQ.2023.v19.n3.a1; arXiv:2104.11857.

[Wu11] Sijue Wu.
Global Wellposedness of the 3-D Full Water Wave Problem.
Inventiones Mathematicae 184(1), 125–220 (2011).
doi:10.1007/s00222-010-0288-1; arXiv:0910.2473.

[AD15] Thomas Alazard and Jean-Marc Delort.
Global Solutions and Asymptotic Behavior for Two Dimensional Gravity Water Waves.
Annales scientifiques de l'École normale supérieure 48(5), 1149–1238 (2015).
doi:10.24033/asens.2268.

[IP15] Alexandru D. Ionescu and Fabio Pusateri.
Global Solutions for the Gravity Water Waves System in 2D.
Inventiones Mathematicae 199(3), 653–804 (2015).
doi:10.1007/s00222-014-0521-4; arXiv:1303.5357.

[IP18] Alexandru D. Ionescu and Fabio Pusateri.
Global Regularity for 2D Water Waves with Surface Tension.
Memoirs of the American Mathematical Society 256(1227) (2018).
doi:10.1090/memo/1227; arXiv:1408.4428.

[DIPP17] Yu Deng, Alexandru D. Ionescu, Benoît Pausader, and Fabio Pusateri.
Global Solutions of the Gravity-Capillary Water-Wave System in Three Dimensions.
Acta Mathematica 219(2), 213–402 (2017).
doi:10.4310/ACTA.2017.v219.n2.a1; arXiv:1601.05685.