



Forward and inverse modelling of eco-evolutionary dynamics



in biological and economic systems

Victor Boussange

Forward and inverse modelling of eco-evolutionary dynamics in biological and economic systems

September 10, 2022

Cover picture: Top: forest in Sorapiss, Dolomites, Italy. Bottom: New York City, USA. @ Luca
Bravo / PhotoSpirit

The document format is based on the *Clean Thesis* style developed by Ricardo Langner.

Contents

| | | |
|----------|---------------------|-----------|
| 1 | Introduction | 1 |
| 2 | CV | 15 |

Introduction

” Concepts without precepts are empty, and precepts without concepts are blind.

— Immanuel Kant
Critique of Pure Reason

Biological and economic systems as complex adaptive systems

What are the similarities between biological and economic systems? Both are complex adaptive systems (CAS) [1]: they are composed of heterogeneous entities structured at different levels of organizations, that interact in nonlinear ways and experience evolutionary processes. Interaction and evolutionary processes take many different forms and operate at different organizational level [Levin1998] (see Fig. 1.1). Surprisingly, the variety of processes involved and their couplings do not necessarily lead to unpredictable, chaotic, or erratic structures and dynamics [Olff2009]. Instead, emergent structural properties and behavior often arise [mitchell2009complexity]. In biological systems, those include patterns of species richness, where for instance montane regions are often associated with a disproportionately high number of species [2]. In economic systems, those include the distribution of international income, where some countries have systematically developed much more rapidly than others [acemoglu2001colonial]. A common direction on the research agenda in Biology and Economics is to comprehend the set of interaction and evolutionary processes that determine these emergent properties [3], and how do they do so. In biological systems, the nature of the processes of interaction and evolution is identified, and the current challenge is to comprehend the mechanisms resulting from their couplings. In economic systems, we still do not exactly understand the nature of those processes, and how are they involved.

Ecological and evolutionary processes drive the dynamics of biological systems

In biological systems, interaction processes are generally designated as ecological processes, and involve fluxes of energy and matter across space and time, encompassing the processes of interaction between organisms (biotic interactions) and between




Fig. 1.1: Schematic diagram of proposed organisational levels in biological and economic systems. A is inspired from [Hendry+2016]

organisms and their environment (abiotic interactions), and dispersal processes (movement of individual across space) [4]. Evolutionary processes designate those processes responsible for the change of heritable characteristics (DNA, genes, phenotypes) over successive generations [Hall2013]. The coupling between ecological and evolutionary processes is acknowledged since the very birth of the theory of Evolution, when Darwin realised a link between the different ecological opportunities across the Galápagos islands and the different beak shapes in the finches he found in each island [darwin2004origin], during his voyage on the Beagle. He reasoned that the variations in ecological opportunities lead to a differential in survival for certain phenotypes, which over time resulted in the evolution of different beak shapes. Since then, we know that ecological processes directly affect evolutionary response [Ezard2009]. In the recent years, the idea that not only ecological processes can affect evolutionary response, but also that evolutionary processes could affect ecological processes, has developed [XXX]. Empirical studies have demonstrated that evolution can be rapid and occur on similar time scales as ecology [Hairston2005, 5] and have quantifiable effect on ecological dynamics [Ezard2009], leading to feedbacks between ecological and evolutionary processes, so-called eco-evolutionary feedbacks [Schoener2011, 5]. Eco-evolutionary feedbacks involve situations where an ecological property influences evolutionary change, which then feeds back to an ecological property, or vice versa [Govaert2019a]. Examples are feedbacks

between population density (ecological property) and trait evolution (evolutionary change), which can lead to evolutionary branching through the effect of competition [Doebeli1999]. Eco-evolutionary feedbacks are also involved in adaptation mechanisms [Doebeli1999], where species disperse and phenotypic variations allow to adapt to local environments [XXX]. Those feedbacks may greatly influence the mechanisms driving the dynamics of ecosystems [6], but our quantitative understanding of their modality is limited [Lion2022]. In particular, eco-evolutionary feedbacks are expected to play a critical role in the evolution of the biosphere in the coming decades [7], as ecosystems are being rapidly affected by anthropogenic pressure and with climate change [8, 9]. In order to mitigate the consequences of human development, it is of utmost urgency to better understand eco-evolutionary feedbacks [7], and develop mechanistic models embedding this knowledge [6]. This will in turn provide more reliable forecasts of ecosystem states [Clark2001], to help designing adequate management of ecosystem services [6].

Drivers of economic change

The processes that determine economic change is controversial in economics [Nelson2014]. To explain economic development, mainstream economic theory assumes that economic systems are in equilibrium, in the sense that the demand and supply of goods and services are balanced on all relevant markets. Firms are rational in maximizing profits by adapting to demand and supply, and the observed economic change is driven by exogenous forces, such as technological change [Romer1986]. Evolutionary economics, promoted by the seminal work of [Nelson2014], criticizes this view and seeks to explain economic change by focusing on endogenous forces. Evolutionary economics suggests that interactions between firms and economic activities, and evolutionary processes acting upon them, are major processes contributing to economic change [Hodgson2019]. For instance, firms or economic activities may interact positively or negatively [Wernerfelt1989, Pistorius2007Ozman2009, Saavedra2009a, Cohendet2018, Menon2015], spread across space [RogersEverettM2003DoI, Zahra2000], and adapt [Cordes2006] or transform into new economic institutions [Freeman2002, Hodgson2004, Aldrich2008], affecting economic development at the regional and national scale. Because these processes are analogous to eco-evolutionary processes driving the dynamics of biological systems, a number of modelling approaches have recently borrowed concepts and methods from biology, aiming at better understanding the processes underlying emergent properties in economic systems [Tacchella2018, Saavedra2009a, Scholl2020, Zhang2018, Modis1997, Saavedra2014, Farmer1999, Michalakelis2011, Marasco2016, Gatabazi2019, Cauwels56, Applegate2021, Suweis2015]. For instance, [Saavedra2009a] has

successfully used a model of mutualistic interaction to explain structural patterns in industrial cooperation. Also, [Scholl2020] uses the concepts of foodwebs and density dependence to explain market malfunctions and excess volatility in financial markets. However, those studies did not seek to understand how these processes may affect economic development, and whether eco-evolutionary processes could explain differences in economic development across countries remains to be tested.

Forward modelling of eco-evolutionary processes

The complex interplay between ecological and evolutionary processes, acting at different scales of time and space and organization, can hardly be studied with experimental approaches [Hagen2022, 10]. As such, a deductive approach, relying on forward modelling, has traditionally been put forward to advance our understanding of the mechanisms underlying [Brummitt2020]. Along this approach, hypotheses about causal processes are embedded in a model, which forward integration generates emergent properties. Such emergent behavior may be seen as predictions from the consideration of the causal processes [May2004]. The role of the modeller is to point at the mechanisms by which the properties emerge, disentangling the underlying interplay between the processes. In the early 1930s to 1940s, by formulating tractable mathematical models implementing the processes of reproduction, dispersal and mutations, the work of Fisher, Wright and Haldane has greatly contributed to the modern synthesis of evolutionary biology [huxley1942evolution], generally accepted as the basis of our current understanding of evolutionary dynamics. The mechanistic models commonly take the form of differential equations (DE), and express how the processes under investigation affect the rate of change of the population characteristics, such as the proportion of a given allele. However, the requirement of tractable mathematical models (DEs that yield analytical solutions) has involved strong assumptions on the processes investigated, that are poorly representative of the complexity of eco-evolutionary feedbacks in nature [Govaert2019a]. In particular, ecological scenarios have been strongly simplified, and did not take into account how evolution could affect population dynamics [Lion2022].

With the increase in computational capacity, novel modelling approaches relying on individual based models (IBMs) have appeared [deangelis2005individual]. IBMs allow to capture processes acting at the individual level, requiring less simplifying assumptions than traditional mathematical models [deangelis2005individual]. Capturing more realistic scenarios by allowing the forward integration of complex hypothesis, the lack of analytical tractability of IBMs may nonetheless occult the mechanisms underlying emergent properties [May2004, 11]. The recent development of mathematical techniques, such moment closure approximations

[law1999moment, Gandhi2000, 12, 11], adaptive dynamics theory [Metz1995], and probability theory [13], are generating novel pathways by filling the gap between IBMs and mathematical models. Analogous to renormalisation group analysis developed in quantum and statistical physics [Sayama], they form a toolbox to rigorously derive how emergent properties are influenced by processes operating at different organizational levels. As such, they allow an analytical underpinning to IBM simulations, and can generate a general understanding of the key mechanisms at stake [11].

The combination of numerical simulations and, e.g., adaptive dynamics theory, has successfully shed new lights on the emergence of evolutionary branching under frequency-dependent selection [14, 15]. An other example is the work of [Debarre2013a, 16, 17], that has provided new insights on the effect of habitat heterogeneity on population dynamics. However, our current understanding of eco-evolutionary feedbacks omits potentially significant factors, such as the structuration of populations over complex spatial structures [XX] and highly dimensional phenotypic space [XXX].

The consideration of such details is important to advance our understanding, but raises challenging methodological issues. In particular, complex models may hinder the fundamental mechanisms underlying the emergence of a pattern. Also, the consideration of multiple traits leads to an increase in the dimensionality of the associated DE problem, which in turn leads to an exponential increase in computational cost [XXX]. In order to better understand eco-evolutionary feedbacks, we need to investigate more realistic scenarios, which will, in turn, require the development of novel numerical methods that can cope with the extra computational cost.

Inverse modelling

Diametrically opposed to forward modelling, inverse modelling consists in using observation data to infer causal processes [18]. Inverse modelling has recently seen an increased attention, thanks to the increased computational power and availability of datasets [Csillary2010]. It involves the use of inference methods to estimate model parameters, such as bayesian or maximum likelihood inference methods [19]. Those methods proceed by defining a distance between the model simulation and the observation data, which relates to the probability of the parameters given the model and the data [19]. The most likely parameters are associated with the minimum distance, obtained using ad-hoc algorithms. Provided that they are inferred together with uncertainties, parameters can be interpreted to better understand the strengths and effects of the embedded processes [10]. For instance, [20, 21] infers

and analyse the parameters of population dynamic models to understand processes involved in ecosystem functions. Because the parameter estimation problem is equivalent to finding the maximum probability of the model given the observation data, inference methods can also be used to discriminate between candidate mechanisms embedded in alternative models [22, 23]. For instance, [Skeels2022] shows that temperature-dependent evolutionary speed is the most likely mechanism to explain variations in biodiversity patterns, using inference methods to discriminate between alternative dynamic models embedding different hypotheses. Nonetheless, the use of inverse modelling in evolutionary biology has been limited because of a number of issues, some of which specifically relating to eco-evolutionary models. Central to these issues are the high computational cost of the forward integration of eco-evolutionary models [24], the large number of parameters involved [25], and their strong nonlinearities[26, 27, 28]. Advances in the field of artificial intelligence could circumvent these issues.

Machine learning to leverage forward and inverse modelling

In the recent years, the field of artificial intelligence (AI) has made enormous progresses in computer vision [XXX] and natural language processing [XXX]. At the backbone of this success are key computational techniques that are increasingly applied to leverage research in other scientific fields [Rolnick2023, 29, 30, 31], providing unique opportunities to investigate causal processes in CAS. In particular, computer vision and natural language processing rely on deep learning methods, that allow neural networks to learn abstract representation of mechanisms from large datasets [32]. Neural network abstractions are hardly interpretable by humans [XXX], and because their prediction ability is limited by the information contained in the training datasets, they are ill-suited for extrapolating beyond observed trends [33, 6]. While neural networks cannot be used *per se* to gain scientific insights and extrapolate, their traditional applications and associated methods have been flourishingly imported in various research fields for this purpose.

For example, neural networks have been used to reduce the cost of the forward integration of climate models, by learning more efficient representations of physical mechanisms [XXX]. They have also been used to approximate the solution of partial differential equations (PDEs) [34], with the major advantage of approximating high dimensional problems at a lower computational cost than traditional methods. Notoriously, [Arnulf] has developed methods consisting in training neural networks through a correspondence between PDEs and stochastic processes. [Arnulf] have mathematically proved and numerically showed that their methods can efficiently approximate very high dimensional PDEs (10 000 dimensions), which could allow

to include more realism in eco-evolutionary models. Nonetheless, PDE models for CAS critically involve non-local terms, which capture non-local interactions between microscopic agents. The extension of such techniques to non-local PDEs could leverage the forward modelling of biological and economic systems.

Also, neural networks have been employed to leverage inverse modelling techniques. For instance, [35, 31] have developed innovative parameter estimation techniques, where neural networks are trained to approximate DEs which parameters are free. Extra constraints, provided by incorporating observation data, allow to recover the parameter values during the training. Underlying this approach is the technique of automatic differentiation (AD), which allows to generate the gradient of the loss function, used in the training process, with respect to any input parameters [XXX]. This technique involves several variants [XXX] which can be generalised to differentiate any numerical function, allowing to train not only neural networks but also any scientific model against data [36]. With the ability to handle computational models characterised by a very large number of parameters, AD, together with the development of other training techniques [XXX], have the potential to leverage inverse modelling [36]. Overall, the use of AI techniques offers unique opportunities for the investigation of causal processes in CAS [Frank2022], but requires further developments for adapting to the specificities of CAS models.

Programming languages

Combining ML techniques with scientific models requires computational environments that allow to easily develop scientific models, ensure simulation performance, and provide composability between ML and other scientific libraries [36]. However, performance and composability are features that are poorly represented in mainstream programming languages used by the scientific community, such as Python, Matlab or R. Those languages are naturally attractive because they are dynamically typed [XX], allowing convenient development iterations. Nonetheless, prototypes written in Python, Matlab or R need to be rewritten in low level, compiled languages such as C, C++ or Fortran for speed and predictable mapping to hardware [Perkel2019, Bezanson2017]. This conversion requires significant involvement, leading to a problem commonly designated as the "two language problem" [Bezanson2017]. In order to circumvent issues of performance, most libraries in Python, Matlab or R rely on bindings with low level languages. For instance, the most used deep learning libraries in Python, TensorFlow and PyTorch, are internally written in C. However, bindings with low level languages come with major negative externalities, such as restricting the understandability of their internals

to computer scientists – prohibiting potential development contributions from the scientific community –, and preventing the composability of libraries [XXX].

Julia is a programming language that was launched in 2012 to address the issue of the two-language problem [Bezanson2017, Bezanson2018]. Julia was built over a type-specializing, just-in-time compiler, which makes it easy to generate highly performant programs, while preserving the essential features of Python, Matlab or R, such as dynamic typing and automatic memory management. Importantly, it relies on multiple dispatch, which allows to generate highly generic code with good performance. This permits to write libraries in pure Julia, guaranteeing productivity and composability. As such, the internals of any Julia library can be understood by non computer-scientists, who can further use his expertise to participate to its development. As a matter of fact, many Julia libraries benefit from a high number of contributions of independent users (see, e.g., github.com/DifferentialEquations.jl). Multiple dispatch also allows to automatically generate the gradient of any Julia program without any modification [ForwardDiff.jl, Zygote.jl]. This means that any scientific library in Julia, such as differential equation solvers, can be combined with deep learning tools, with unique opportunities for forward and inverse modelling problems [Frank2022]. As such, Julia allows to prototype a program which is readily generic and can directly be shared to the research community. Further, by granting the composability of libraries, it allows to blend ML techniques with scientific models. This makes Julia a promising computational environment to accelerate research in CAS.

Thesis outline

In summary, while it is increasingly acknowledged that feedbacks between ecological and evolutionary processes play an important role in the dynamics of biological systems [5, 6], our understanding of the mechanisms in which they are involved has been limited to simplified scenarios. Further, while analogous processes have been suggested to influence the dynamics of economic systems [Hodgson2019], a quantification of their effect is missing. Under increasing anthropogenic pressure, these research directions become essential [6], but raise challenging methodological issues. Here, I present novel forward and inverse modelling approaches to advance our understanding of eco-evolutionary dynamics in biological and economic systems, and utilise them to shed light on the underlying processes and resulting mechanisms.

In ??, I investigate how eco-evolutionary processes, in combination with complex habitat spatial structures, influence the trait distribution of biological populations. I proceed using a forward modelling approach, building a stochastic eco-evolutionary IBM where individuals are structured over a spatial graph, and experience the

fundamental processes of reproduction, competition, mutation and migration. I seek to understand how those microscopic forces result in trait differentiation at the population level. I derive DE approximations of the IBM that, together with extensive numerical simulations, provide analytical insights into how the graph properties affect the population size and trait differentiation. In particular, I show that three main graph properties relating to landscape connectivity and heterogeneity, as well as to habitat spatial auto-correlation, shape the trait differentiation of the biological population. These results establish mechanistic links between landscape features and the eco-evolutionary dynamics of biological populations.

In ??, I develop an inverse modelling method to estimate the parameters of highly nonlinear population dynamic models. The method is based on a machine learning framework and involves AI techniques together with a novel learning strategy. The learning strategy consists in training the model against mini-batches of data with short time horizon, which I analytically show to regularize the learning problem. I implement the ML framework in the Julia library **MiniBatchInference.jl**, and demonstrate through numerical experiments that it can efficiently estimate model parameters and provide statistical evidences for causal processes from noisy, incomplete and independent time series. Altogether, the proposed ML framework is a workhorse for inverse modelling and can elucidate mechanistic pathways in biological and economic systems.

In ??, I quantify the effect of eco-evolutionary processes on the dynamics of economic systems. I employ the ML framework developed in ?? to investigate how alternative eco-evolutionary population dynamic models can explain the dynamics of economic activities in the richest 100 countries, relying on 59 years of economic data. The models investigated embed the processes of ecological interactions between economic activities, spatial transfers, and economic activity transformations, which support is compared to a simple logistic growth model, taken as a null model. I find strong statistical evidence for positive interactions between national economic activities, and spatial transfers across countries. To my knowledge, this is the first study that provides quantitative evidences that similar processes may influence the dynamics of biological and economic systems.

In ??, I extend two recent methods to solve high dimensional PDEs, in order to handle non-local nonlinear terms. The first method relies on Picard iterations, while the second is based on machine learning and involves neural networks to approximate PDE solutions. The numerical difficulties arising due to the non-local term are avoided by using a plain vanilla Monte Carlo integration. I implement the methods in the Julia library **HighDimPDE.jl**, and evaluate their performance on high dimensional PDE models arising in physics and biology, including population dynamic eco-evolutionary models. For all models, the methods yield good results

with short run times, offering the possibility to include more realism in future eco-evolutionary models.

References

- [1] S. A. Levin. “Complex adaptive systems: Exploring the known, the unknown and the unknowable”. In: *Bulletin of the American Mathematical Society* 40.01 (2002), pp. 3–20. DOI: 10.1090/S0273-0979-02-00965-5.
- [2] C. Rahbek et al. “Humboldt’s enigma: What causes global patterns of mountain biodiversity?” In: *Science* 365.6458 (2019), pp. 1108–1113. DOI: 10.1126/science.aax0149.
- [3] J. M. Nordbotten et al. “Ecological and evolutionary dynamics of interconnectedness and modularity”. In: *Proc. Natl. Acad. Sci. USA* 115.4 (2018), pp. 750–755. DOI: 10.1073/pnas.1716078115.
- [4] M. Vellend. “Conceptual Synthesis in Community Ecology”. In: *The Quarterly Review of Biology* 85.2 (2010), pp. 183–206. DOI: 10.1086/652373.
- [5] F. Pelletier, D. Garant, and A. Hendry. “Eco-evolutionary dynamics”. eng. In: *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 364.1523 (2009), pp. 1483–1489. DOI: 10.1098/rstb.2009.0027.
- [6] M. C. Urban et al. “Improving the forecast for biodiversity under climate change”. In: *Science* 353.6304 (2016). DOI: 10.1126/science.aad8466.
- [7] J. Norberg et al. “Eco-evolutionary responses of biodiversity to climate change”. In: *Nature Climate Change* 2.10 (2012), pp. 747–751. DOI: 10.1038/nclimate1588.
- [8] E. C. Ellis. “Anthropogenic transformation of the terrestrial biosphere”. In: *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 369.1938 (2011), pp. 1010–1035. DOI: 10.1098/rsta.2010.0331.
- [9] G. Midgley and L. Hannah. “Extinction Risk from Climate Change”. In: *Biodiversity and Climate Change*. Yale University Press, 2019, pp. 294–296. DOI: 10.2307/j.ctv8jnzw1.37.
- [10] M. Pontarp, Å. Bränström, and O. L. Petchey. “Inferring community assembly processes from macroscopic patterns using dynamic ecoevolutionary models and Approximate Bayesian Computation (ABC)”. In: *Methods in Ecology and Evolution* 10.4 (2019). Ed. by T. Poisot, pp. 450–460. DOI: 10.1111/2041-210X.13129.
- [11] S. Lion. “Moment equations in spatial evolutionary ecology”. In: *Journal of Theoretical Biology* 405 (2016), pp. 46–57. DOI: 10.1016/j.jtbi.2015.10.014.

- [12] J. M. Nordbotten et al. “The dynamics of trait variance in multi-species communities”. In: *R. Soc. Open Sci.* 7.8 (2020), Article No. 200321, 20 pp. DOI: 10.1098/rsos.200321.
- [13] N. Champagnat, R. Ferrière, and S. Méléard. “Unifying evolutionary dynamics: From individual stochastic processes to macroscopic models”. In: *Theoretical Population Biology* 69.3 (2006), pp. 297–321. DOI: 10.1016/j.tpb.2005.10.004.
- [14] U. Dieckmann and M. Doebeli. “On the origin of species by sympatric speciation”. In: *Nature* 400.6742 (1999), pp. 354–357. DOI: 10.1038/22521.
- [15] M. Doebeli and U. Dieckmann. “Speciation along environmental gradients”. In: *Nature* 421.6920 (2003), pp. 259–264. DOI: 10.1038/nature01274.
- [16] G. Meszéna, I. Czibula, and S. Geritz. “Adaptive Dynamics in a 2-Patch Environment: A Toy Model for Allopatric and Parapatric Speciation”. In: *Journal of Biological Systems* 05.02 (1997), pp. 265–284. DOI: 10.1142/S0218339097000175.
- [17] S. Mirrahimi and S. Gandon. “Evolution of specialization in heterogeneous environments: equilibrium between selection, mutation and migration”. In: *Genetics* 214.2 (2020), pp. 479–491. DOI: 10.1534/genetics.119.302868.
- [18] C. Peng et al. “Integrating models with data in ecology and palaeoecology: advances towards a model-data fusion approach”. In: *Ecology Letters* 14.5 (2011), pp. 522–536. DOI: 10.1111/j.1461-0248.2011.01603.x.
- [19] M. Schartau et al. “Reviews and syntheses: parameter identification in marine planktonic ecosystem modelling”. In: *Biogeosciences* 14.6 (2017), pp. 1647–1701. DOI: 10.5194/bg-14-1647-2017.
- [20] S. I. Higgins, S. Scheiter, and M. Sankaran. “The stability of African savannas: insights from the indirect estimation of the parameters of a dynamic model”. In: *Ecology* 91.6 (2010), pp. 1682–1692. DOI: 10.1890/08-1368.1.
- [21] A. Curtsdotter et al. “Ecosystem function in predator-prey food webs—confronting dynamic models with empirical data”. In: *Journal of Animal Ecology* 88.2 (2019). Ed. by D. Stouffer, pp. 196–210. DOI: 10.1111/1365-2656.12892.
- [22] D. R. A. Kenneth P. Burnham and Model. *Model Selection and Multimodel Inference*. Ed. by K. P. Burnham and D. R. Anderson. New York, NY: Springer New York, 2002. DOI: 10.1007/b97636.

- [23] J. B. Johnson and K. S. Omland. “Model selection in ecology and evolution”. In: *Trends in Ecology & Evolution* 19.2 (2004), pp. 101–108. DOI: 10.1016/j.tree.2003.10.013.
- [24] R. A. Fisher et al. “Vegetation demographics in Earth System Models: A review of progress and priorities”. In: *Global Change Biology* 24.1 (2018), pp. 35–54. DOI: 10.1111/gcb.13910.
- [25] I. L. Boyd. “The Art of Ecological Modeling”. In: *Science* 337.6092 (2012), pp. 306–307. DOI: 10.1126/science.1225049.
- [26] A. Hastings et al. “Chaos in Ecology: Is Mother Nature a Strange Attractor?” In: *Annual Review of Ecology and Systematics* 24.1 (1993), pp. 1–33. DOI: 10.1146/annurev.es.24.110193.000245.
- [27] J. Huisman and F. J. Weissing. “Biodiversity of plankton by species oscillations and chaos”. In: *Nature* 402.6760 (1999), pp. 407–410. DOI: 10.1038/46540.
- [28] E. Benincà et al. “Chaos in a long-term experiment with a plankton community”. In: *Nature* 451.7180 (2008), pp. 822–825. DOI: 10.1038/nature06512.
- [29] K. Kashinath et al. “Physics-informed machine learning: Case studies for weather and climate modelling”. In: *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 379.2194 (2021). DOI: 10.1098/rsta.2020.0093.
- [30] T. Schneider et al. “Earth System Modeling 2.0: A Blueprint for Models That Learn From Observations and Targeted HighResolution Simulations”. In: *Geophysical Research Letters* 44.24 (2017), pp. 12,396–12,417. DOI: 10.1002/2017GL076101. arXiv: 1709.00037.
- [31] A. Yazdani et al. “Systems biology informed deep learning for inferring parameters and hidden dynamics”. In: *PLOS Computational Biology* 16.11 (2020). Ed. by V. Hatzimanikatis, e1007575. DOI: 10.1371/journal.pcbi.1007575.
- [32] Y. LeCun, Y. Bengio, and G. Hinton. “Deep learning”. In: *Nature* 521.7553 (2015), pp. 436–444. DOI: 10.1038/nature14539.
- [33] A. D. Barnosky et al. “Approaching a state shift in Earth’s biosphere”. In: *Nature* 486.7401 (2012), pp. 52–58. DOI: 10.1038/nature11018. arXiv: 9605103 [cs].
- [34] J. Sirignano and K. Spiliopoulos. “DGM: A deep learning algorithm for solving partial differential equations”. In: *J. Comput. Phys.* 375 (2018), pp. 1339–1364.

- [35] M. Raissi, P. Perdikaris, and G. Karniadakis. “Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations”. In: *Journal of Computational Physics* 378 (2019), pp. 686–707. doi: 10.1016/j.jcp.2018.10.045.
- [36] C. Rackauckas et al. “Universal Differential Equations for Scientific Machine Learning”. In: *arXiv:2001.04385v3* (2020), 18 pages.

CV

2

Personal Information

| | |
|-------------|-----------------------|
| Residence | Zürich, Switzerland |
| E-mail | bvictor@ethz.ch |
| Website | vbuossange.github.io |
| Github | github.com/vbuossange |
| Age | Born 1995 (age 27) |
| Citizenship | France citizen |

Personal skills

| | |
|-----------------------|---|
| Languages | English (fluent) French (native) Spanish (B2) German (B1) |
| Programming languages | Julia Python C++ Java Matlab R Bash VBA |
| Sports | Ski mountaineering Alpinism Rock climbing Enduro mountainbiking Surfing |
| Alpine CV | [vbuossange.github.io/pages/alpine_cv/] |

Education

- 10.2022 **Ph.D in Environmental Sciences**, Swiss Federal Institute for Forest, Snow and Landscape (WSL | Swiss Federal Institute of Technology Zurich, ETH), Switzerland
Forward and inverse modelling of eco-evolutionary processes. Under the guidance of Prof. Dr. Loïc Pellissier.
- 06.2017 **Full year academic exchange**, University of New South Wales (UNSW Sydney), Australia
- 06.2017 **Master thesis in theoretical geomechanics**, UNSW Sydney | CSIRO, Australia
Numerical continuation and bifurcation analysis for unconventional geomechanics. Under the guidance of Dr. Thomas Poulet.
- 08.2018 **M.S. in Energy and Environmental Engineering**, Institut National Des Sciences Appliquées de Lyon (INSA Lyon), France
Three-year undergraduate engineering course in Energy and Environmental Systems, focused on Advanced Energy Systems and Efficiency.
- 08.2018 **B.S. in Mathematics and Physics**, Institut National Des Sciences Appliquées de Lyon (INSA Lyon), France
Ranking : 21/650 students.

Professional appointments

- 08.2018 **R&D intern**, Compagnie National du Rhône (CNR), France
03.2018 Development of an Energy Management System based on various optimisation techniques for optimal production of renewable resources. Applications to EU sponsored projects: **Jupiter1000** (power-to-gas), **Move in pure** (vehicle-to-grid), **Marie-Galante island** (micro-grid)

Publications

Peer-reviewed

1. **Boussange, V.** & Pellissier, L., *Eco-evolutionary model on spatial graphs reveals how habitat structure affects phenotypic differentiation*. *Commun Biol* 5, 668 (2022). [[bioRxiv](#)]

Preprints

1. **Boussange, V.**, Vilimelis-Aceituno, P., Pellissier, L., *Mini-batching ecological data to improve ecosystem models with machine learning* [[bioRxiv](#)] (2022), 46 pages. In review.
2. **Boussange, V.**, Becker, S., Jentzen, A., Kuckuck, B., Pellissier, L., *Deep learning approximations for non-local nonlinear PDEs with Neumann boundary conditions*. [[arXiv](#)] (2022), 59 pages. Revision requested from Partial Differential Equations and Applications.

Proceedings

1. Poulet, T., Alevizos, S., Veveakis, M., **Boussange, V.**, Regenauer-Lieb, K., *Episodic mineralising fluid injection through chemical shear zones*, ASEG Extended Abstracts (2018), 5 pages.

In preparation

1. **Boussange, V.**, Sornette, D., Lischke, H., Pellissier, L., *Analogous forces to ecological interactions, dispersal and mutations shape the dynamics of economic activities*.

Talks

- 07.2022 **Speaker**, HIGHDIMPDE.JL: A Julia package for solving high-dimensional PDEs, JuliaCon2022, online. [youtube.com/watch?v=4sXqGhhknT4](https://www.youtube.com/watch?v=4sXqGhhknT4)
- 06.2022 **Speaker**, Interpretable machine learning for forecasting dynamical processes in ecosystems, World Biodiversity Forum, Davos, Switzerland.
- 06.2022 **Invited speaker**, Investigating empirical patterns of biodiversity with mechanistic eco-evolutionary models, Seminar at the Theoretical Ecology and Evolution group, Universität Bern.
- 11.2021 **Invited speaker**, Numerical approximations of solutions of highly dimensional, non-local nonlinear PDEs, StAMBio seminar, St Andrews, UK.
- 10.2021 **Speaker**, Graph topology and habitat assortativity drive phenotypic differentiation in an eco-evolutionary model, Conference on Complex Systems, Lyon, France.
- 10.2021 **Speaker**, Using graph-based metrics to assess the effect of landscape topography on diversification, ECBC, Amsterdam, Netherlands.
- 09.2021 **Speaker**, Solving non-local nonlinear Partial Differential Equations in high dimensions with HighDimPDE.jl, International Conference on Computational Methods in Systems Biology, Bordeaux, France.
- 04.2021 **Speaker**, Responses of neutral and adaptive diversity to complex geographic population structure, Mathematical Population Dynamics, Ecology and Evolution, CIRM Marseille, France.

Softwares

- 2022 **MiniBatchInference.jl** Julia
github.com/vboussange/MiniBatchInference.jl
A Julia package for maximum likelihood estimation and model selection of strongly nonlinear dynamical models.
- 2021 **HighDimPDE.jl** Julia
github.com/vboussange/HighDimPDE.jl
A Julia package that breaks down the curse of dimensionality in solving non local, non linear PDEs.
- 2021 **EvoId.jl** Julia
2019 github.com/vboussange/EvoId.jl

Evolutionary individual based modelling, mathematically grounded.

| | | |
|------|---------------------------------------|--|
| 2018 | OptiVPP <i>confidential</i> | Python, GAMS |
| | | Energy Management System for Virtual Power Plants. |

Open source software contributions

SciML
DiffEqFlux.jl
CUDA.jl
Flux.jl
LightGraphs.jl

Teaching and supervision

12.2020 **701-3001-00L Environmental Systems Data Science**, ETH Zürich, D-USYS,
09.2020 Switzerland

06.2020 **262-0100-00L Lab rotation**, ETH Zürich, D-BSSE, Switzerland
04.2020

12.2020 **Taste of research internship**, Polytech Nice-Sophia, France
09.2020

Reviews

2022 **Journal of Open Source Software**
2019 **Journal of Theoretical Biology**

