


# Forward and inverse modelling of eco-evolutionary dynamics

in biological and economic systems

Victor BOUSSANGE



**Victor Boussange**

*Forward and inverse modelling of eco-evolutionary dynamics in biological and economic systems*

September 17, 2022

Cover picture: Top: forest in Sorapiss, Dolomites, Italy. Bottom: New York City, USA. @ Luca  
Bravo / PhotoSpirit

The document format is based on the *Clean Thesis* style developed by Ricardo Langner.

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Context . . . . .	1
1.1.1	Biological and economic systems as complex adaptive systems	1
1.1.2	Ecological and evolutionary processes drive the dynamics of biological systems . . . . .	2
1.1.3	Drivers of economic change . . . . .	3
1.2	Modeling eco-evolutionary dynamics . . . . .	4
1.2.1	Forward modelling of eco-evolutionary processes . . . . .	4
1.2.2	Inverse modelling . . . . .	6
1.2.3	Machine learning to leverage forward and inverse modelling .	7
1.2.4	Programming languages . . . . .	8
1.3	Thesis outline . . . . .	9
<b>2</b>	<b>Discussion</b>	<b>13</b>
2.1	Contributions . . . . .	13
2.1.1	Advancing our fundamental understanding of eco-evolutionary processes in biological and economic systems . . . . .	13
2.1.2	Leveraging forward and inverse modelling with ML . . . . .	17
2.2	Limitations . . . . .	20
2.2.1	Forward modelling . . . . .	20
2.2.2	Inverse modelling . . . . .	21
2.3	Perspectives . . . . .	22
2.4	Concluding remarks . . . . .	25
<b>3</b>	<b>CV</b>	<b>27</b>
<b>List of Figures</b>		<b>41</b>



# Introduction

„ Nature loves to hide.

— Heraclitus (c.6th-5th century BCE)

## 1.1 Context

### 1.1.1 Biological and economic systems as complex adaptive systems

What are the similarities between the dynamics of biological and economic systems? Think of a biological system as community of interacting biological organisms ([chapin2002principles](#)), and think of an economic system as a community of interacting economic agents (Dopfer and Potts, 2007). *A priori*, the processes underlying the dynamics of an economic system strongly differ from that of biological systems, because the behavior of economic agents is motivated by rationality ([XXX](#)), where economic agents maximize utility. Nonetheless, economic agents are faced with uncertainty ([Foster2012](#)) and their rationality is bounded ([HerbertSimon](#)). As a result, economic agents adopt a variety of behavioral rules (e.g. technological, organizational, institutional, [Foster2012](#)) through trial-and-errors, which are subject to natural selection through competition processes ([Schumpeter](#)). In this perspective, both biological and economic systems are complex adaptive systems (CAS) (Levin, 2002), composed of heterogeneous entities that interact in nonlinear ways and experience evolutionary processes. Interestingly, the stochasticity of the processes involved, and their couplings, do not necessarily lead to unpredictable structures and dynamics ([Olff2009](#)), but rather induce organised structural properties and invariant patterns ([Olff2009](#); [mitchell2009complexity](#)). In biological systems, invariant patterns include patterns of species richness, where for instance montane regions are often associated with a disproportionately high number of species (Rahbek et al., 2019b). In economic systems, invariant patterns include the bimodal shape of the distribution of international income, where some countries have systematically developed much more rapidly than others ([ace moglu2001colonial](#)). The processes of interaction and evolution involved take many different forms and operate at different organizational level ([Levin1998](#)), from genes to ecosystems, and from



**Fig. 1.1:** Schematic diagram of proposed organisational levels in biological and economic systems. A is inspired from (Hendry+2016)

organizational routines to economies (see Fig. 1.1). A common direction on the research agenda to understand the dynamics of biological and economic systems is to identify the fundamental dynamical processes involved, and the associated mechanisms (Fig. 1.1,Nordbotten et al., 2018). In biological systems, the fundamental processes resulting in patterns of species richness are identified (Hagen; Rahbek et al., 2019a; Rangel et al., 2018), and the current challenge is to underpin the mechanisms resulting from their couplings (Fig. 1.2). In economic systems, we still do not exactly understand the fundamental processes at stake.

### 1.1.2 Ecological and evolutionary processes drive the dynamics of biological systems

In biological systems, interaction processes are more commonly designated as ecological processes, and encompass the processes of interaction between organisms (biotic interactions) and between organisms and their environment (abiotic interactions), as well as dispersal processes (movement of individual across space) (Vellend, 2010). Evolutionary processes designate those processes responsible for the change of heritable characteristics (DNA, genes, phenotypes) over successive generations (Hall2013). The coupling between ecological and evolutionary processes is acknowl-

edged since the very birth of the theory of evolution. During his voyage on the Beagle, Darwin documented a link between the different ecological opportunities across the Galápagos Islands and the different beak shapes in the finches he found on each island ([darwin2004origin](#)). He reasoned that the variations in ecological opportunities lead to a differential in survival for certain phenotypes, which over time resulted in the evolution of different beak shapes. Since then, we know that ecological processes interact with evolutionary processes, and together shape the long term dynamics of biological systems ([Hagen](#); Rahbek et al., 2019a; Rangel et al., 2018). Empirical studies have now demonstrated that evolution can be rapid and occur on similar time scales as ecology ([Hairston2005](#); Pelletier et al., 2009) and have quantifiable effects on ecological dynamics ([Ezard2009](#)), leading to feedbacks between ecological and evolutionary processes, so-called eco-evolutionary feedbacks ([Schoener2011](#); [Govaert2019](#); Pelletier et al., 2009). Eco-evolutionary feedbacks involve situations where an ecological process (e.g., replication, competition, dispersal) influences an evolutionary process (e.g. phenotypic change), which then feeds back to an ecological process, or vice versa ([Govaert2019a](#)). Examples are feedbacks between population dynamics (replication and competition) and trait evolution (phenotypic change), which can lead to evolutionary branching through the effect of competition ([Doebeli1999](#)). In spatially structured populations, another classical example of eco-evolutionary feedbacks is the mechanism of local adaptation (Savolainen et al., 2007), where feedbacks between population dynamics, dispersal and trait evolution can facilitate or prevent populations to adapt to local environmental conditions (Meszéna et al., 1997; Doebeli and Dieckmann, 2003). Importantly, the eco-evolutionary feedbacks involved in adaptation mechanisms are expected to affect the dynamics of the biosphere in the coming decades (Norberg et al., 2012; Urban et al., 2016), because of rapid changes in local environmental conditions experienced in ecosystems, through the effects of anthropogenic pressure and climate change (Ellis, 2011; Midgley and Hannah, 2019). Nevertheless, our understanding of their effect and modality under realistic biological scenarios is limited ([Lion2022](#)). Eco-evolutionary feedbacks may greatly influence the dynamics of ecosystems (Norberg et al., 2012; Urban et al., 2016), calling for a better understanding of their effect in realistic biological scenarios ([Lion2022](#)).

### 1.1.3 Drivers of economic change

In economic systems, the fundamental processes determining economic change are controversial ([Dopfer](#); [Nelson2014](#); Hodgson, 2019). To explain economic development, mainstream economic theory (Lawson, 2013) assumes that economic systems are in equilibrium, in the sense that the demand and supply of goods and

services are balanced on all relevant markets. Firms are rational in maximizing profits by adapting to demand and supply, and the observed economic change is driven by exogenous forces, such as technological change (**Romer1986**). Evolutionary economics, promoted by the seminal work of**Nelson2014**, criticizes this view and seeks to explain economic change by focusing on endogenous forces. Evolutionary economics suggests that interactions between economic agents, firms and economic activities, and evolutionary processes acting upon them, are major processes contributing to economic change (Hodgson, 2019). For instance, firms or economic activities may interact positively through supply chains Ozman, 2009; Saavedra et al., 2009; Van Der Panne, 2004, compete in markets (Wernerfelt, 1989), spread across space through business expansions (Rogers, 2003; Zahra et al., 2000), and adapt (Cordes, 2006) or transform into new economic institutions (Freeman, 2002; Hodgson and Knudsen, 2004; Aldrich et al., 2008), affecting economic development at the local, regional, national, and international scale. Because these processes are analogous to eco-evolutionary processes driving the dynamics of biological systems, a number of modelling approaches have borrowed concepts and methods from biology, aiming at underpinning the processes underlying invariant patterns in economic systems (Tacchella et al., 2018; Saavedra et al., 2009; Scholl et al., 2021; Zhang et al., 2018; Modis, 1997; Saavedra et al., 2014; Farmer and Lo, 1999; Michalakelis et al., 2011; Marasco et al., 2016; Gatabazi et al., 2019; Cauwels and Sornette, 2012; Applegate and Lampert, 2021; Suweis et al., 2015). For instance, (Saavedra et al., 2009) has successfully used a model of mutualistic interaction to explain structural patterns in industrial cooperation. Also, (Scholl et al., 2021) uses the concepts of food webs and density dependence to explain market malfunctions and excess volatility in financial markets. However, those studies did not seek to understand how eco-evolutionary processes may affect economic development at the national scale. Recent modelling approaches developed in evolutionary biology may help to disentangle whether eco-evolutionary processes could explain differences in economic development across countries.

## 1.2 Modeling eco-evolutionary dynamics


### 1.2.1 Forward modelling of eco-evolutionary processes

The complex interplay between ecological and evolutionary processes can hardly be studied with experimental approaches (**Hagen2022**; Pontarp et al., 2019). As such, a deductive approach, relying on forward modelling, has traditionally been put forward to underpin the mechanisms underlying invariant patterns in biological systems (Brummitt et al., 2020). Along this approach, hypotheses about causal processes

are embedded in a model, whose forward integration generates emergent (non-anticipated) properties. Emergent properties may be seen as predictions from the consideration of the causal processes (**May2004**), and the role of the modeler is to underpin the underlying mechanisms, i.e. to disentangle how the interplay between the processes generate the observed behavior. In the early 1930s to 1940s, by formulating tractable mathematical models implementing the processes of reproduction, dispersal and mutations, the work of Fisher, Wright and Haldane has greatly contributed to the modern synthesis of evolutionary biology (**huxley1942evolution**), generally accepted as the basis of our current understanding of evolutionary dynamics. In order to obtain tractable mathematical model, Fisher, Wright and Haldane have neglected eco-evolutionary feedbacks (**Govaert2019a**). In particular, ecological scenarios have been strongly simplified, and did not take into account how evolutionary processes could affect population dynamics (**Lion2022**).

With the increase in computational capacity, novel modelling approaches relying on individual based models (IBMs) have appeared (**deangelis2005individual**). These models require less simplifying assumptions than traditional mathematical models (**deangelis2005individual**), and can underpin more realistic mechanisms by allowing to capture processes acting at the individual level. However, the lack of analytical tractability of IBMs is a shortcoming, because it challenges the ability of the modeler to underpin with generality the mechanisms resulting from the underlying hypotheses (**May2004**; Lion, 2016). The recent development of mathematical techniques, such moment closure approximations (**law1999moment**; **Gandhi2000**; Nordbotten et al., 2020; Lion, 2016), adaptive dynamics theory (**Metz1995**), and probability theory (Champagnat et al., 2006), are generating novel pathways by filling the gap between IBMs and mathematical models. Analogous to renormalization group analysis developed in quantum and statistical physics (**Sayama**), they form a toolbox to rigorously derive how emergent properties are influenced by processes operating at different organizational levels. As such, these mathematical techniques allow an analytical underpinning to IBM simulations, and can generate a general understanding of the key mechanisms at stake (Lion, 2016).

The combination of numerical simulations and, e.g., adaptive dynamics theory, has successfully shed new lights on the emergence of evolutionary branching under frequency-dependent selection (Dieckmann and Doebeli, 1999; Doebeli and Dieckmann, 2003). An other example is the work of **Debarre2013a**; Meszéna et al., 1997; Mirrahimi and Gandon, 2020, that has provided new insights on the effect of habitat heterogeneity on population dynamics. However, our current understanding of eco-evolutionary feedbacks neglects specificities of real biological populations that may significantly alter the resulting mechanisms, such as the structuration



**Fig. 1.2: Forward and inverse modelling approaches for the understanding of CAS.** A forward modelling approach starts with assuming a set of processes, that are embedded in a model, and seeks to understand how their interplay transforms in mechanisms that are associated with a macroscopic pattern (macroscopic invariance). An inverse modelling approach integrates empirical observation to constrain the model output, generating knowledge by inferring the probability and the strength of the processes investigated, given the empirical observations. Similarly to forward modelling, mechanisms may be deduced from the processes inferred, leading to the undersanding of invariances (pattern).

of populations over complex spatial structures (Nowak and Komarova, 2001) and highly dimensional phenotypic space (Doebeli and Ispolatov, 2010).

The consideration of such factors is important to advance our understanding, but raises challenging methodological issues. In particular, adding complexity in eco-evolutionary models may hinder the fundament mechanisms underlying the emergence of a pattern. Also, the consideration of multiple traits leads to an increase in the dimensionality of the model, which in turn leads to an exponential increase in computational cost (Bellman, 2010). In order to better understand eco-evolutionary feedbacks, we need to investigate more realistic scenarios, which will, in turn, require developments in order to cope with the extra complexity and computational cost.

## 1.2.2 Inverse modelling

Another approach to underpin processes and mechanisms in biological systems consists in inverse modelling, where processes and mechanisms are inferred from empirical data through an induction process (Fig. 1.2). Inverse modelling can take

the form of parameter estimation (Schartau et al., 2017) or model selection (Johnson and Omland, 2004), both involving the use of inference methods to estimate, respectively, the most probable model parameter value, or the most probable model among candidates, given empirical data. In parameter estimation, provided that they are inferred together with uncertainties, parameters can be interpreted to better understand the strengths and effects of the associated processes (Pontarp et al., 2019). For instance, Higgins et al., 2010; Curtsdotter et al., 2019 infer the parameters of population dynamic models to understand the processes involved in ecosystem functions. In model selection, candidate models embedding competing hypotheses about processes and mechanisms are derived, and the relative support of each model given the data is computed to discriminate between the hypothesis (Johnson and Omland, 2004). For instance, using inverse modelling and alternative eco-evolutionary models, (Skeels2022) shows that temperature-dependent evolutionary speed is the most likely hypothesis explaining variations in biodiversity patterns, among alternative hypotheses embedded in the models.

The computation of the most probable model parameter values, or the computation of the different model supports, critically involves inference methods. Inference methods commonly demand many forward integration of the model, resulting in a computational cost that can be prohibitively expensive (Schneider et al., 2017). The number of forward integration required may dramatically increase with the number of model parameters (Csillary2010), and the number of model parameters, together with the model nonlinearities, can eventually lead to false estimates of the most probable model parameter values (XXX). Consequently, inverse modelling methods have mostly been used with simple evolutionary models (Csillary2010). Eco-evolutionary models are dependent on numerous parameters (Boyd, 2012), are strongly nonlinear (Hastings et al., 1993; Huisman and Weissing, 1999; Benincà et al., 2008), and their integration is computationally expensive (Fisher et al., 2018), challenging the use of inverse modelling to underpin eco-evolutionary processes. Advances in the field of artificial intelligence could circumvent these issues, allowing to advance our knowledge of eco-evolutionary feedbacks in empirical systems (XXX).

### 1.2.3 Machine learning to leverage forward and inverse modelling

In the recent years, the field of artificial intelligence (AI) has made enormous progresses in computer vision (XXX) and natural language processing (XXX). At the backbone of this success are key computational techniques that could leverage the forward and inverse modelling of CAS. Computer vision and natural language processing rely on deep learning methods, that allow neural networks to learn abstract representation of mechanisms from large datasets (LeCun et al., 2015).

These abstractions are hardly interpretable by humans (XXX), and their prediction ability is limited by the information contained in the training datasets. As such, neural networks cannot be used *per se* to gain scientific insights and extrapolate beyond observed trends (Barnosky et al., 2012; Urban et al., 2016). Nevertheless, their traditional applications and associated methods have been successfully derived in other scientific fields for this purpose (Rolnick2023; Kashinath et al., 2021; Schneider et al., 2017; Yazdani et al., 2020).

Neural networks have been used in forward modelling, to reduce the cost of the forward integration of climate models, by learning more efficient representations of physical mechanisms (XXX). They have also been used to approximate the solution of partial differential equations (PDEs) (Sirignano and Spiliopoulos, 2018), with the major advantage of approximating high dimensional problems at a lower computational cost than traditional methods. Underlying the training of neural network is the technique of backpropagation (XXX). This technique can be generalised to any scientific model against data (Rackauckas et al., 2020a), with the potential to leverage inverse modelling techniques. As such, the derivation of AI techniques to investigate causal processes in CAS offers unique opportunities (Frank2022).

## 1.2.4 Programming languages

Combining ML techniques with scientific models requires computational environments that allow to easily develop scientific models, ensure simulation performance, and provide composability between ML and other scientific libraries (Rackauckas et al., 2020a). Unfortunately, performance and composability are features that are poorly represented in mainstream programming languages used by the scientific community, such as Python, Matlab or R. Those languages are naturally attractive because they are dynamically typed (XX), allowing convenient development iterations. Nonetheless, prototypes written in Python, Matlab or R need to be rewritten in low level, compiled languages such as C, C++ or Fortran for speed and predictable mapping to hardware (Perkel2019; Bezanson2017). This conversion requires significant involvement, leading to a problem commonly designated as the "two language problem" (Bezanson2017). In order to circumvent issues of performance, most libraries in Python, Matlab or R rely on bindings with low level languages. For instance, the most used deep learning libraries in Python, TensorFlow and PyTorch, are internally written in C. However, bindings with low level languages come with major negative externalities, such as restricting the understandability of their internals to computer scientists – prohibiting potential development contributions from the scientific community –, and preventing the composability of libraries (XXX).

Julia is a programming language that was launched in 2012 to address the issue of the two-language problem (Bezanson2017; Bezanson2018). Julia was built over a type-specializing, just-in-time compiler, which makes it easy to generate highly performant programs, while preserving the essential features of Python, Matlab or R, such as dynamic typing and automatic memory management. Importantly, it relies on multiple dispatch, which allows to generate highly generic code with good performance. This permits to write libraries in pure Julia, guaranteeing productivity and composability. As such, the internals of any Julia library can be understood by non computer-scientists, who can further use his expertise to participate to its development. Many Julia libraries benefit from a high number of contributions of independent users (see, e.g., [github.com/DifferentialEquations.jl](https://github.com/DifferentialEquations.jl)). Multiple dispatch also allows to automatically generate the gradient of any Julia program without any modification (**ForwardDiff.jl**; **Zygote.jl**). This means that any scientific library in Julia, such as differential equation solvers, can be combined with deep learning tools, with unique opportunities for forward and inverse modelling problems (Frank2022). As such, Julia allows to prototype a program which is readily generic and can directly be shared to the research community. Further, by granting the composability of libraries, it allows to blend ML techniques with scientific models. This makes Julia a promising computational environment to accelerate research in CAS.

## 1.3 Thesis outline

In summary, while it is increasingly acknowledged that feedbacks between ecological and evolutionary processes play an important role in the dynamics of biological systems (Pelletier et al., 2009; Urban et al., 2016), our understanding of the mechanisms in which they are involved has been limited to simplified scenarios. Further, while analogous processes have been suggested to influence the dynamics of economic systems (Hodgson, 2019), a quantification of their effect is missing. Under increasing anthropogenic pressure, these research directions become essential (Urban et al., 2016), but raise challenging methodological issues. Here, I present novel forward and inverse modelling approaches to advance our understanding of eco-evolutionary dynamics in biological and economic systems, and utilise them to shed light on the underlying processes and resulting mechanisms.

In ??, I investigate how eco-evolutionary processes, in combination with complex habitat spatial structures, influence the trait distribution of biological populations. I proceed using a forward modelling approach, building a stochastic eco-evolutionary IBM where individuals are structured over a spatial graph, and experience the fundamental processes of reproduction, competition, mutation and migration. I

seek to understand how those microscopic forces result in trait differentiation at the population level. I derive DE approximations of the IBM that, together with extensive numerical simulations, provide analytical insights into how the graph properties affect the population size and trait differentiation. In particular, I show that three main graph properties, measuring landscape connectivity, heterogeneity in connectivity, and habitat spatial auto-correlation, shape the trait differentiation of the biological population. These results establish mechanistic links between landscape features and the eco-evolutionary dynamics of biological populations.

In ??, I develop an inverse modelling method to estimate the parameters of highly nonlinear population dynamic models. The method is based on a machine learning framework and involves AI techniques together with a novel learning strategy. This learning strategy consists in training the model against mini-batches of data with short time horizon, which I analytically show to regularize the learning problem. I implement the ML framework in the Julia library **MiniBatchInference.jl**, and demonstrate through numerical experiments that it can efficiently estimate model parameters and provide statistical evidences for causal processes from noisy, incomplete and independent time series. Altogether, the proposed ML framework is a workhorse for inverse modelling and can elucidate mechanistic pathways in biological and economic systems.

In ??, I quantify the effect of eco-evolutionary processes on the dynamics of economic systems. I employ the ML framework developed in ?? to investigate how alternative eco-evolutionary population dynamic models can explain the dynamics of economic activities in the richest 100 countries, relying on 59 year of economic data. The models embed the processes of ecological interactions between economic activities, spatial transfers, and economic activity transformations, which support is compared to a simple logistic growth model, taken as a null model. I find strong statistical evidence for positive interactions between national economic activities, and spatial transfers across countries. To my knowledge, this is the first study that provides quantitative evidences that similar processes may influence the dynamics of biological and economic systems.

In ??, I extend two recent methods to solve high dimensional PDEs, in order to handle non-local nonlinear terms. The first method relies on Picard iterations, while the second is based on machine learning and involves neural networks to approximate PDE solutions. The numerical difficulties arising due to the non-local term are avoided by using a plain vanilla Monte Carlo integration. I implement the methods in the Julia library **HighDimPDE.jl**, and evaluate their performance on high dimensional PDE models arising in physics and biology, including population dynamic eco-evolutionary models. For all models, the methods yield good results

with short run times, offering the possibility to include more realism in future eco-evolutionary models.



# Discussion

”

*To be but one with all living things, to return, by  
a radiant self-forgetfulness, to the All of Nature.*

— Friedrich Hölderlin (1770-1843)

## 2.1 Contributions

Understanding biological and economic systems involves the underpinning of the elemental processes underlying the systems' macroscopic features, and establishing the associated mechanisms (Levin, 2002, Fig. 1.2). Following this approach, my work aimed at advancing our understanding of how ecological and evolutionary processes, and their interplay, shape the dynamics of biological and economic systems. In particular, this thesis contributed to

- (i) a fundamental understanding of the role of eco-evolutionary processes in shaping the dynamics of biological populations structured in complex landscapes ??,
- (ii) the quantification of the effect of mechanisms associated with eco-evolutionary processes in economic systems ??,
- (iii) methodological advances in the forward and inverse modelling of eco-evolutionary dynamics ??????.

In the following, I discuss the chapters of my thesis collectively within the broader context of our current understanding of the dynamics of biological and economic systems, and the accompanying modelling paradigm.


### 2.1.1 Advancing our fundamental understanding of eco-evolutionary processes in biological and economic systems

#### Linking eco-evolutionary processes to patterns of differentiation

Spatial patterns of biodiversity result from the processes of selection, mutation and dispersal, that act upon individual organisms ([hamilton2021population](#)). In finite size populations, mutations result in "drift" ([Slatkin1987a](#)), causing stochastic

variations in the allelic proportions and phenotypes of biological populations. In geographically structured population, drift results in patterns of neutral differentiation (**Slatkin1987a**), where isolated populations are characterised by differentiated allelic proportions and phenotypes. Dispersal tends to reduce neutral differentiation (**Slatkin1987a**), and this effect is modulated by landscape connectivity (Wright, 1943; McRae, 2006; McRae and Beier, 2007) through the mechanism of "isolation by limited dispersal" (Orsini et al., 2013). By increasing the dispersal ability of organisms, landscape connectivity decreases neutral differentiation (Lande, 1991). When landscapes present heterogeneous habitats, natural selection can supplement the effect of genetic drift and increase the sole effect of stochasticity on differentiation (**fisher1958genetical**). Under this scenario, local environment conditions select individuals with traits that provide them higher fitness (**Gaither2018**). At the population level, this results in populations adapting to their local environment, a mechanism coined "local adaptation" (Kawecki and Ebert, 2004) and resulting in patterns of "adaptive differentiation". Adaptive differentiation is hindered by dispersal, which prevents local adaptation by bringing maladapted individuals, that destabilise the evolution of traits towards the optimal (Meszéna et al., 1997; Débarre et al., 2013; Mirrahimi and Gandon, 2020). While adaptive differentiation concerns traits under selection, it indirectly affects the differentiation of neutral traits, that are co-evolving with traits under selection through linkages (Billiard et al., 2015; Lepers et al., 2021). This results in turn to the mechanism of "isolation by adaptation", where habitat heterogeneity, rather than landscape connectivity, increases neutral differentiation (**nosil2008**). Simple mechanisms resulting in neutral and adaptive differentiation are identified, but how they are modulated by eco-evolutionary feedbacks and landscape complexity is unclear.

In ??, I demonstrate a novel mechanism, involving the process of intra-specific competition, that considerably affects neutral differentiation. Through the creation of unbalanced migration fluxes which increases the intensity of competition in highly connected populations, heterogeneity in connectivity reduces gene flow and reinforces neutral differentiation. I also investigate how the mechanism of local adaptation in complex landscapes, where habitats are arranged in a realistic fashion. I show that the complexity of habitat spatial distribution can be reduced to a measure of habitat spatial auto-correlation, coined the "habitat assortativity". Landscapes characterised by a high habitat assortativity systematically support populations that are locally better adapted than in landscape with low assortativity, resulting in higher adaptive differentiation. Specifically, I provide an analytical condition for local adaptation that sheds light on how it relates to dispersal intensity, selection strength, habitat heterogeneity, and habitat assortativity.



**Fig. 2.1:** Summary of the causal pathways involved in neutral and adaptive differentiation, disentangled in ?? . Ecological processes are displayed in green boxes, evolutionary processes are displayed in red boxes.

Because habitat assortativity affects local adaptation, it must also affect neutral differentiation through the mechanism of isolation by adaptation. Closing the loop, I demonstrate that habitat assortativity affects population differentiation through two antagonistic effects. By favoring local adaptation, it promotes isolation by adaptation, therefore increasing neutral differentiation. In parallel, it also favors gene flow within clusters of similar environmental conditions, decreasing isolation by limited dispersal. This results in habitat assortativity decreasing neutral differentiation for low dispersal intensity, and increasing neutral differentiation for high dispersal intensity. This complex feedback is essential to understand population differentiation in complex landscapes.

I provide a summary of the processes and resulting mechanisms shaping neutral and adaptive differentiation in Fig. 2.1. Overall, ?? links processes to patterns, establishing an extensive map of causal pathways involved in the phenotypic differentiation of populations.

### Linking economic patterns to eco-evolutionary processes

?? provides a novel understanding of the role of eco-evolutionary processes in economic systems. Neoclassical economics and evolutionary economics seek to explain

economic change with formal modelling, focusing on the relationships between economic variables such as output, employment and productivity (Boschma and Frenken, 2005). In particular, evolutionary economics is concerned with explaining economic change by endogenous forces, such as interactions between firms and economic activities, and evolutionary processes acting upon them (Metcalfe2006; Hodgson, 2019). In opposition to this approach, complexity economics (Hidalgo, 2021) seeks to predict variations in national income by using fine-grained data of economic activity outputs and dimensionality reduction techniques (Mitchell, n.d.), without making assumptions on the underlying processes (Hidalgo). While a current concern in evolutionary economics is to obtain an agreement between the mechanisms proposed and empirical observations (XXX), complexity economics seeks to understand the causal processes underlying the success of the dimensionality reduction technique (XXX).

By being agnostic to economic variables yet providing a causal link with the most probable processes underlying economic development, ?? bridges evolutionary economics and economic complexity. Our approach relies on the simple observation that the dynamics of economic activities contains signatures from the many complex processes that underpin them. These signatures consist in peculiar temporal variations and couplings that can be conveniently modelled by biologically inspired population dynamic models, which use are additionally justified by deep analogies between economic activities and biological populations. Analogously to biological populations that are characterised by genes, economic activities are characterised by organizational routines (NelsonWinter), which experience evolutionary processes and define how they engage in ecological processes (NelsonWinter). As a result, and similarly to the size of biological populations, the capital growth of economic activites is determined by the ensemble of organizational routines that characterise them (Boschma and Frenken, 2005). Similarly to approaches used in ecology and evolution (Skeels), inverse modelling approaches can then be used to disentangle, from their signatures (Skeels) left on the dynamics of economic activities, the role of the eco-evolutionary processes acting upon them.

Specifically, ?? explores the effect of eco-evolutionary processes, proposed in the evolutionary economics and geography economics literature, on the dynamics of economic activities at the national scale. In particular, ?? seeks to test whether the dynamics of economic activities can be explained by different types of interdependencies, including positive (XXX) and negative interactions (XXX), spatial transfers (XXX), and economic transformations (XXX). Using population dynamic models capturing the different interdependencies, ?? provides empirical evidence that economic activities engage in positive interactions, and benefit from spatial transfers of knowledge and routines. Positive interactions may arise from a variety of processes

proposed in the evolutionary economic literature, such as supply chains (Ozman, 2009; Saavedra et al., 2009) and knowledge spillovers (Menon, 2015). Its support implies that diversity promotes economic development (**Hidalgo2018**), as economic activities promote each other. The support for spatial transfers implies that, besides international patent laws (**XXX**), transfers of knowledge and organisational routines have a considerable effect on economic activities. Nevertheless, discrepancies in the strength-of-evidence obtained for spatial transfers across countries highlight that some countries are overall more akin to spatial transfers than others, which may be explained by differences in cognitive, organizational, social, institutional or geographic proximities across countries (**XXX**). I provide a summary of the mechanisms evidenced in economic systems in ???. Overall, ?? bridges approaches in economics and biology to understand the mechanisms shaping the endogenous dynamics of economic systems.

### 2.1.2 Leveraging forward and inverse modelling with ML

#### Advances in the modelling of realistic spatial and phenotypic structures

???? provide new tools to better understand mechanisms resulting from ecological and evolutionary processes, and their interplay, in the context of realistic population structures and phenotypic distributions. Evolutionary dynamics have been traditionally studied in the context of regular population structures (Lieberman et al., 2005). For instance, Slatkin, 1973; Slatkin, 1978; Kirkpatrick and Barton, 1997; Polechová and Barton, 2015; Polechová, 2018; AndradeRestrepo et al., 2019; Doebeli and Dieckmann, 2003; Meszéna et al., 1997; Yeaman and Otto, 2011; Débarre et al., 2013; Mirrahimi and Gandon, 2020 consider regular spatial structures to investigate differentiation in biological populations, missing the effect of spatial complexity on the underlying mechanisms. Biological habitats differ in their connectivity (**XXX**), and economic entities are structured through complex networks (**XXX**). Lieberman et al., 2005 and subsequent studies of "evolutionary dynamics on graphs" (**XXX**) show that this complexity affects the interplay between selection and drift. However, evolutionary dynamics on graphs does not consider eco-evolutionary feedbacks (**Govaert2019a**). Thus far, models that include frequency dependence together with realistic, complex population structures were missing.

While a vast majority of the work on eco-evolutionary feedbacks has focused on the evolution of scalar phenotypes (**Doebeli2011**), in most organisms, many phenotypic properties combine in complicated ways to determine ecological processes (**Doebeli2014**). For instance, **Doebeli2011** shows that the consideration of multiple traits is likely to generate more diversity than expected with one dimensional models. Trade-offs in traits is also an essential feature shaping the evolutionary dynamics

of biological populations, with consequences on the dynamics of e.g. cancer cell evolution (XXX) and plankton dynamics (XXX). While there is overall a genuine need to better understand evolutionary dynamics in high dimensional spaces, the simulation of high dimensional models is tremendously difficult, since the numerical cost of traditional methods grows exponentially in the number of dimensions of the phenotypic space.

?? develops a generic modelling framework to capture the effect of eco-evolutionary processes on biological populations structured in complex landscapes, and ?? provides tools to efficiently simulate the resulting high-dimensional models. The IBM presented in ?? involves the combination of graphs and highly dimensional phenotypic spaces, together with eco-evolutionary feedbacks, to model population structures. The modelling framework presented can readily be generalised, and the code associated to the numerical experiments in ?? include a Julia library, **Evoid.jl**, that implements a more general version of the model. As such, the modelling framework presented in ?? may be used to investigate other questions involving complex population structures and the co-evolution of characterisits. Reproducing the discrete and stochastic nature of ecological and evolutionary processes (Champagnat et al., 2006), simulations of the IBM may not provide a general understanding of system investigated (XXX), and cannot be scaled to simulate large systems involving millions of individuals (XXX). Nevertheless, the IBM proposed in ?? is mathematically tractable under simplifying assumptions, and can be efficiently simulated with a deterministic PDE approximation. Tractability allows to obtain analytical insights on how structural properties affect macroscopic population under simplifying assumptions (??). The PDE approximation, combined with the numerical methods presented in ??, further allow efficient simulations. I have implemented the numerical methods for simulating high dimensional models in the Julia library **HighDimPDE.jl** (**HighDimPDE**), a registered Julia package belonging to the SciML organisation (XXX). The user interface respects standards from the SciML organisation, meaning that Julia users can easily adopt it. The package aims at hosting many more solver algorithms that break down the curse of dimensionality, and is currently receiving contributions from developers to implement the DeepBSDE scheme (Han et al., 2018).

The combination of analytical insights and numerical simulations can help to elucidate the links between microscopic processes and macroscopic properties (Levin). ???? provide novel analytical and numerical tools to better understand how complex spatio-evolutionary structures affect the interplay between eco-evolutionary processes and generate macroscopic patterns.

## **Advances in inference methods for the investigation of eco-evolutionary processes**

???? develop and test a novel inverse modelling method that allows to infer highly nonlinear dynamical processes from observation data, opening up new venues to advance our understanding and prediction of the dynamics of biological and economic systems. The most developed inference methods in current use for inverse modelling are Bayesian inference methods with Markov Chain Monte Carlo (XXX) and variational methods (XXX). Bayesian inference methods require a large number of forward model integrations (Schneider et al., 2017), and are highly affected by the number of model parameters (Csillary2010). Variational methods require the model sensitivity to its parameters (XXX) and are prone to converge to local minima, especially with complex models (XXX). Those central issues likely explain the very poor number of studies that have used inverse modelling to further our knowledge on eco-evolutionary dynamics (XXX).

?? presents a novel inference framework that can efficiently recover the most probable parameter values of eco-evolutionary models, given temporal data. The framework is based on a variational method, but resolves its main shortcomings by including key ingredients in the recipe, including automatic differentiation (Rackauckas et al., 2020b), state-of-the-art optimizers (Kingma and Ba, 2014), and a learning strategy based on a mini-batch method. The use of automatic differentiation simply eliminates the effort required to obtain the model sensitivity to its parameters, and the state-of-the-art optimizers, together with the mini-batch method, ensure the efficiency and reliability of the method in handling highly nonlinear models. ?? is part of an ongoing effort to blend ML and traditional models to gain scientific understanding and extrapolability (XXX). In physical systems such as ocean and atmospheric systems (XXX), invariant laws are known, and ML is mostly used to improve model forecast skill. In contrast, models of biological and economic systems are yet to be formulated, methods such as the the ML framework presented in ?? can greatly contribute to gain scientific knowledge. By contrasting competing hypotheses embedded in alternative models, ???? provide concrete examples, both with synthetic and empirical data, that the ML framework can successfully elucidate eco-evolutionary mechanistic pathways. The proposed method is also relevant for improving the forecast of predictive eco-evolutionary model Urban et al., 2016, and integrates the practical constraints of datasets including short time series with partial, noisy, shallow and independent observations (Dornelas et al., 2018).

The ML framework is implemented in the multi-purpose Julia package **MiniBatchInference.jl** (**MiniBatchInference**), readily available to the scientific community. **MiniBatchInference.jl** is built around the celebrated differential equation solver

`DifferentialEquations.jl` and the deep learning library `Flux.jl`. As such, the use of `MiniBatchInference.jl` requires very limited efforts to any user familiar with those libraries.

Overall, the method proposed in ?? successfully blends ML methods with mechanistic ecosystem models to improve our gain scientific knowledge from observation data. Concrete case examples in ???? show that it enables the testing of eco-evolutionary theories against data, advancing our understanding and the improvement of current mechanistic models, with the potential to provide better forecasts of ecosystems states (Urban et al., 2016).

## 2.2 Limitations

### 2.2.1 Forward modelling

Alternatives to the methods presented in ???? may be more appropriate for the forward modelling of eco-evolutionary dynamics. While IBMs are interesting tools to investigate stochastic drift in finite size populations, the Gillespie algorithm (Gillespie, 1976) used to simulate the IBM in ?? is computationally intensive, and requires to compute the fitness of all individuals at each time step, which depends on the characteristics of all the other individuals. The resulting computational complexity grows polynomially with the number of individuals ( $\mathcal{O}(N^2)$ ). While it is an interesting tool to investigate stochastic drift in finite size populations, it cannot be used to model large populations. On the other hand, PDEs can provide deterministic approximations of IBMs (Champagnat et al., 2006), while demanding less computational power. The methods presented in ?? can approximate PDEs in high dimensions (up to 10 traits, which seems more than enough in practice (XXX)), but still suffer from a number of issues that may prevent their practical use. In particular, the MLP method can only provide the population density for one single trait value, and as such, cannot characterise the total population density with a reasonable computational complexity. On the other hand, the ML based approximation method can provide the full population density, but involves the training of many neural networks, one at each time step. This is worrying, since the training of a neural network is numerically costly, and that long simulation times may be required by practitioners. An other problem with the numerical methods proposed in ?? is that they involve the tuning of meta parameters, including the choice of a kernel for the integration of the nonlocal term. This choice is critical for the success of the approximation, but how to determine it is not clear. Together, the methods proposed in ?? may require further development to be used in practice by practitioners. Yet irrespective of the numerical method used, solving PDEs inevitably

requires a considerable computational effort, because PDEs track the evolution of the full phenotypic density of populations. Nevertheless, only the first three moments of the population density are usually of interest to investigate eco-evolutionary dynamics (population size, trait mean and trait variance, see **XXX**). Instead of seeking to numerically approximate the full phenotypic density, moment closure approximation methods (**Wickman2021; Lion2022**; Nordbotten et al., 2020) may be, so far, more appropriate tools. Those approaches consist in approximating the population density with a gaussian distribution. This, in turn, allows to transform the PDE problem into a system of coupled differential equations involving the time evolution of the population size (1 variable for a single species population), the mean trait value in each dimension of the trait space ( $d$  variables), and the variance-covariance matrix of the trait density ( $d^2$  variables). As such, the computational cost of this method scales only polynomially with the number of dimension ( $\mathcal{O}(d^2)$ ), while providing the exact information required to investigate eco-evolutionary dynamics in high dimensional spaces. It is worth noting that instead of neural networks, gaussian functions could easily be used with the ML-based approximation method to simulate PDEs. Equivalent to the assumption taken in the moment closure methods presented in **Wickman2021; Lion2022**; Nordbotten et al., 2020, we expect that this approach would greatly improve the computational efficiency of the ML-based approximation method in simulating eco-evolutionary models, while solving the problem of the choice of a kernel for the integration of the nonlocal term. Using Gaussian functions may considerably lower the number of iterations required in the training process, while reducing the computational cost, as they involve less parameters ( $d(d + 1) + 1$ ) than neural networks (*xx* in ??).

### 2.2.2 Inverse modelling

The inverse modelling framework proposed in ?? and used in ?? also present shortcomings, which may favor the use of other methods to infer eco-evolutionary processes from data. First, the mini-batching learning strategy requires the choice of a minibatch size to ensure the convergence to the maximum likelihood estimate. This choice is arbitrary, but may affect the model selection process. Reducing the batch size implies that the model is fitted on the short term dynamics of the data, but because the model is likely to only characterise some of the features of the data, the resulting support could differ, were the model rather fitted to long term dynamics. Theoretical developments will be required to better understand the assumptions behind the choice of the mini-batch size, which may provide guidance to the choice of this meta parameter. Second, the proposed method may fail to find the maximum likelihood estimate of complex models, since the associated likelihood landscape is

harder to navigate than simpler ones. This may lead to a bias in the model selection process towards simpler models. Third, the information criterion-based model selection procedure used in ???? is uniquely based on a trade-off between goodness-of-fit and number of parameters of the model, which may not be satisfactory to characterise the complexity of dynamical models. For instance, XXX shows that the logistic map, which consists of only two parameters, can be fitted to any pattern. Other criterion, involving the complexity of the dynamical behavior of the model (such as, e.g., its Lyapunov exponent), could as such be developed. Fourth, the ML framework developed in ?? requires a differentiable model, a strong prerequisite that may not be met by stochastic models (XXX). Fifth, the ML framework provides a single point estimate of the posterior distribution, which is subsequently used for model selection. Yet the models' posterior distributions may be multimodal, where the alternative modes carry valuable information to consider in the model selection process (**Daniels2015**). In this case, fully Bayesian method may be required to characterise the full posterior distribution. Alternatively, **Skeels2022** employs a more flexible approach for model selection, which does not require differentiability. The method consists in aggregating model simulation outputs into summary statistics, which are used to train classifier algorithms in recognizing the generating model. The classifier algorithms, such as random forests and neural networks, are further used on summary statistics obtained from the empirical data, recovering the most likely models and associated hypotheses. This approach is similar to Approximation Bayesian Computation methods **Csillary2010**, and requires summary statistics that can correctly discriminate between models. This is a strong requirement to grant the success of the model selection procedure, but it has the major advantage of clarifying why the model is more likely.

Overall, the methods presented in ???? solve major problems arising in the forward and inverse modelling of eco-evolutionary dynamics, but suffer from shortcomings. These limitations may direct practitioners to alternative methods, but also invite to further developments.

## 2.3 Perspectives

### Development opportunities in inverse modelling

The mini-batch method developed in ?? and the ML-based approximation method developed in ?? offer unique development opportunities to leverage inverse modelling. The mini-batch method is relevant beyond the ML framework presented in ??, and could be used within a fully Bayesian framework (XXX), where the full posterior distribution of the model is estimated. While Bayesian inference with

MCMC chains methods may not be appropriate for eco-evolutionary models (see ??), automatic differentiation variational inference (ADVI,(XXX)) offers an appealing alternative. With ADVI, the posterior distribution is approximated by a gaussian distribution (XXX), significantly reducing the number of model integration (Gosh et al., 2021). Improving the ML framework presented in ??, ADVI could capture multimodality in the model posterior distribution (by approximating the multimodal distribution with a gaussian distribution with large variance). This, in turn, could improve the model selection procedure (??), and provide uncertainties measures to the value of the parameters inferred. Providing uncertainty estimates while ensuring computational efficiency, Bayesian Learning via Stochastic Gradient Langevin Dynamics (**Welling2011BayesianIV**) could also be implemented in the ML framework. This method builds upon recent advances in Bayesian Deep Learning (**Wilson2020**) and interprets the iterative gradient-based optimization procedure as a Markov chain with an equilibrium distribution over the posterior distribution of the model parameters. It therefore comes with the scalability of variational methods and the interpretability of Bayesian methods, and can provide good estimates of uncertainty errors for complex models.

The ML-based approximation method for high dimensional PDEs, presented in ??, could be used for inverse modelling. In ??, the parameters of the PDE model are assumed fixed, but could be set as free parameters, analogously to the parameters of the neural networks used for approximating the solution. The loss function in ?? would then take the PDE model parameters as additional arguments, and include an additional term, involving the distance between the PDE model solution and the data. This term, analogous to ?? in ??, would constrain the PDE parameters, similarly to the training of physics informed neural networks (Raissi et al., 2019; Yazdani et al., 2020). In contrast to Raissi et al., 2019; Yazdani et al., 2020, a major advantage of this approach it to efficiently perform inverse modelling with high-dimensional dynamical models. Because Julia is a programming language with pervasive AD, this development would require little effort with the Julia library **HighDimPDE.jl**. Together, the ML methods developed in ???? offer unique development opportunities to bring more robustness and efficiency to inverse modelling methods, providing uncertainty estimation and the possibility to handle high dimensional models.

### Confronting eco-evolutionary model on spatial graphs and empirical data

The confrontation of the predictions from ?? with empirical data, and the use of inference methods with the proposed eco-evolutionary model on spatial graphs, could advance our understanding of eco-evolutionary dynamics in empirical systems. ?? proposed topology metrics that should correlate with standard population

differentiation metrics ( $Q_{ST}$  metrics). Because real landscapes can be projected on spatial graphs ((XXX)and ??), the topology metrics, together with empirical data on population differentiation (e.g., Fluerin), could be used to verify our predictions. Discrepancies may indicate that other important processes may be involved in empirical patterns. On the other hand, a validation of our predictions could help to predict population differentiation at a global scale. These predictions could, in turn, be linked to patterns of species richness, in order to underpin how population genetics may lead to speciation over time (XXX). In the same direction, the use of the eco-evolutionary model on spatial graphs, together with paleo-climatic data (HagenXXX) and inference methods, could help addressing fundamental questions on the processes involved in current biodiversity patterns. ?? succinctly tests whether our predictions hold for a more general setting involving trait-based competition. Trait-based competition may be ubiquitous in biological systems (XXX), and similarly to the process of environmental filtering (XXX), can lead to diversification. An important question on the research agenda is to underpin how competition may mediate environmental filtering and promote and hamper diversification over time. This fundamental question could be addressed by embedding the competing hypotheses in alternative models, which support could be tested against data. Along shorter time scales, the eco-evolutionary model on spatial graph could be calibrated on empirical data of species distribution (XXX)with the inference method proposed in ??, and combined with climate scenarios (XXX)to better predict how biological populations will adapt to climate change (Norberg et al., 2012; Urban et al., 2016). Together, the model developed in ?? and the resulting predictions, together with the ML framework presented in ??, could bring insights on the actual mechanisms involved in empirical systems, and help to predict their responses to climate change.

### **Econobiology, a new venue to understand economic systems, and design more appropriate governance**

The current understanding of key processes and patterns in biological systems may provide insights into organization principles in economic systems, and provide guidance for economic policies. The biologically inspired eco-evolutionary model presented in ?? was successful in characterizing the dynamics of economic activities. ?? therefore evidences deep connections between processes acting upon economic and biological systems. On top of encouraging the characterization of these processes at a finer level, this calls for investigating parallels in the dynamics and organizational principles between biological and economic systems. There may be connections between nutrient cycles in ecosystems (Veldhuis2018) and capital cycles in economic systems (King1993), where ecological compartments and financial intermediaries

develop to improve the efficiency of nutrient flows and capital allocations (Odum, 1969). **Veldhuis2018; King1993** show that this efficiency sustains productivity and growth, but the developmental process may fail, where ecosystems stay in a state of lower maturity (**Margalef1963**) or technological lock-in (**Hidalgo2009**). What are the links between the evolution of nutrient and capital cycles, and the fundamental processes acting upon lower organization levels? Future directions should seek to understand at a finer level the processes evidenced in ??, and understand how they may result in organizational principles at the economy level. **Veldhuis2018** provides a synthesis of our understanding of how ecosystem organization emerges through self-reinforcing mechanisms, promoted by ecological and evolutionary processes acting upon producers, consumers, and decomposers. This synthesis provides an interesting roadmap, and analogous self-reinforcing mechanisms could play important roles in determining fluxes of capital in economic systems. The quest for finding common organizational principles may help us distinguishing important differences between economic systems. Since biological systems have undergone evolution for a much longer time than economic systems, they may be more effective and resilient than economic systems. These difference may inspire economic policies. Altogether, biology may eventually help us understanding organizational principles in economic systems, and provide guidance for improving economic regulations.

## 2.4 Concluding remarks

It is, more than ever before, urgent to better understand and forecast the dynamics of life on Earth. While we know that the global temperature on Earth will rise by 2 to 4 degrees in 2100, predictions on ecosystem states are much more uncertain. Yet obtaining more realistic ecosystem forecasts is critical to help designing adequate management of ecosystem services. Blending scientific knowledge, formalised in mathematical model, with machine learning, provides unique opportunities to quickly improve our understanding of ecosystem dynamics and provide forecasts. Machine learning is part of the astonishing technological progress experienced by humankind in the last centuries. This progress has led us to think that we are rational beings, that have crafted singular socio-economic to create wealth. Yet it may well be that socio-economic systems are analogous to, but less efficient and resilient than, biological systems. Leaving aside our navel gazing, we could surely get insights from organizational principles in ecosystems, that have survived major environmental crisis for more than 3.5 billion years. To conclude, let us cite Hölderlin: "*To be but one with all living things, to return, by a radiant self-forgetfulness, to the All of Nature*".



# CV

## Personal Information

Residence	Zürich, Switzerland
E-mail	bvictor@ethz.ch
Website	vboussange.github.io
Github	github.com/vboussange
Age	Born 1995 (age 27)
Citizenship	France citizen

## Personal skills

Languages	English (fluent) French (native) Spanish (B2) German (B1)
Programming languages	Julia Python C++ Java Matlab R Bash VBA
Sports	Ski mountaineering Alpinism Rock climbing Enduro mountainbiking Surfing
Alpine CV	[vboussange.github.io/pages/alpine_cv/]

## Education

- 10.2022 **Ph.D in Environmental Sciences**, Swiss Federal Institute for Forest, Snow and Landscape (WSL | Swiss Federal Institute of Technology Zurich, ETH), Switzerland  
*Forward and inverse modelling of eco-evolutionary processes in biological and economic systems.* Under the guidance of Prof. Dr. Loïc Pellissier.
- 06.2017 **Full year academic exchange**, University of New South Wales (UNSW Sydney), Australia
- 06.2017 **Master thesis in theoretical geomechanics**, UNSW Sydney | CSIRO, Australia  
*Numerical continuation and bifurcation analysis for unconventional geomechanics.* Under the guidance of Dr. Thomas Poulet.
- 08.2018 **M.S. in Energy and Environmental Engineering**, Institut National Des Sciences Appliquées de Lyon (INSA Lyon), France  
Three-year undergraduate engineering course in Energy and Environmental Systems, focused on Advanced Energy Systems and Efficiency.
- 08.2018 **B.S. in Mathematics and Physics**, Institut National Des Sciences Appliquées de Lyon (INSA Lyon), France  
Ranking : 21/650 students.

## Professional appointments

- 08.2018 **R&D intern**, Compagnie National du Rhône (CNR), France  
03.2018 Development of an Energy Management System based on various optimisation techniques for optimal production of renewable resources. Applications to EU sponsored projects: **Jupiter1000** (power-to-gas), **Move in pure** (vehicle-to-grid), **Marie-Galante island** (micro-grid)

## Publications

### Peer-reviewed

1. **Boussange, V.** & Pellissier, L., *Eco-evolutionary model on spatial graphs reveals how habitat structure affects phenotypic differentiation*. *Commun Biol* 5, 668 (2022). [[bioRxiv](#)]

### Preprints

1. **Boussange, V.**, Vilimelis-Aceituno, P., Pellissier, L., *Mini-batching ecological data to improve ecosystem models with machine learning* [[bioRxiv](#)] (2022), 46 pages. In review.
2. **Boussange, V.**, Becker, S., Jentzen, A., Kuckuck, B., Pellissier, L., *Deep learning approximations for non-local nonlinear PDEs with Neumann boundary conditions*. [[arXiv](#)] (2022), 59 pages. Revision requested from Partial Differential Equations and Applications.

### Proceedings

1. Poulet, T., Alevizos, S., Veveakis, M., **Boussange, V.**, Regenauer-Lieb, K., *Episodic mineralising fluid injection through chemical shear zones*, ASEG Extended Abstracts (2018), 5 pages.

### In preparation

1. **Boussange, V.**, Sornette, D., Lischke, H., Pellissier, L., *Analogous forces to ecological interactions, dispersal and mutations shape the dynamics of economic activities*.

## Talks

- 07.2022 **Speaker**, HIGHDIMPDE.JL: A Julia package for solving high-dimensional PDEs, JuliaCon2022, online. [youtube.com/watch?v=4sXqGhhknT4](https://www.youtube.com/watch?v=4sXqGhhknT4)
- 06.2022 **Speaker**, Interpretable machine learning for forecasting dynamical processes in ecosystems, World Biodiversity Forum, Davos, Switzerland.
- 06.2022 **Invited speaker**, Investigating empirical patterns of biodiversity with mechanistic eco-evolutionary models, Seminar at the Theoretical Ecology and Evolution group, Universität Bern.
- 11.2021 **Invited speaker**, Numerical approximations of solutions of highly dimensional, non-local nonlinear PDEs, StAMBio seminar, St Andrews, UK.
- 10.2021 **Speaker**, Graph topology and habitat assortativity drive phenotypic differentiation in an eco-evolutionary model, Conference on Complex Systems, Lyon, France.
- 10.2021 **Speaker**, Using graph-based metrics to assess the effect of landscape topography on diversification, ECBC, Amsterdam, Netherlands.
- 09.2021 **Speaker**, Solving non-local nonlinear Partial Differential Equations in high dimensions with HighDimPDE.jl, International Conference on Computational Methods in Systems Biology, Bordeaux, France.
- 04.2021 **Speaker**, Responses of neutral and adaptive diversity to complex geographic population structure, Mathematical Population Dynamics, Ecology and Evolution, CIRM Marseille, France.

## Softwares

- 2022 **MiniBatchInference.jl** Julia  
[github.com/vboussange/MiniBatchInference.jl](https://github.com/vboussange/MiniBatchInference.jl)  
A Julia package for maximum likelihood estimation and model selection of strongly nonlinear dynamical models.
- 2021 **HighDimPDE.jl** Julia  
[github.com/vboussange/HighDimPDE.jl](https://github.com/vboussange/HighDimPDE.jl)  
A Julia package that breaks down the curse of dimensionality in solving non local, non linear PDEs.
- 2021 **EvoId.jl** Julia  
2019 [github.com/vboussange/EvoId.jl](https://github.com/vboussange/EvoId.jl)

Evolutionary individual based modelling, mathematically grounded.

2018	<b>OptiVPP</b> <i>confidential</i>	Python, GAMS
		Energy Management System for Virtual Power Plants.

## Open source software contributions

SciML  
DiffEqFlux.jl  
CUDA.jl  
Flux.jl  
LightGraphs.jl

## Teaching and supervision

12.2020 **701-3001-00L Environmental Systems Data Science**, ETH Zürich, D-USYS,  
09.2020 Switzerland

06.2020 **262-0100-00L Lab rotation**, ETH Zürich, D-BSSE, Switzerland  
04.2020

12.2020 **Taste of research internship**, Polytech Nice-Sophia, France  
09.2020

## Reviews

2022 **Journal of Open Source Software**  
2019 **Journal of Theoretical Biology**



# Bibliography

- Aldrich, Howard E., Geoffrey M. Hodgson, David L. Hull, et al. (2008). “In defence of generalized Darwinism”. In: *Journal of Evolutionary Economics* 18.5, pp. 577–596 (cit. on p. 4).
- AndradeRestrepo, Martín, Nicolas Champagnat, and Régis Ferrière (2019). “Local adaptation, dispersal evolution, and the spatial ecoevolutionary dynamics of invasion”. In: *Ecology Letters* 22.5. Ed. by Vincent Calcagno, pp. 767–777 (cit. on p. 17).
- Applegate, J. M. and Adam Lampert (2021). “Firm size populations modeled through competition-colonization dynamics”. In: *Journal of Evolutionary Economics* 31.1, pp. 91–116 (cit. on p. 4).
- Barnosky, Anthony D., Elizabeth A. Hadly, Jordi Bascompte, et al. (2012). “Approaching a state shift in Earth’s biosphere”. In: *Nature* 486.7401, pp. 52–58. arXiv: 9605103 [cs] (cit. on p. 8).
- Bellman, Richard (2010). *Dynamic Programming*. Princeton Landmarks in Mathematics. Reprint of the 1957 edition. Princeton University Press, Princeton, NJ, pp. xxx+340 (cit. on p. 6).
- Benincà, Elisa, Jef Huisman, Reinhard Heerkloss, et al. (2008). “Chaos in a long-term experiment with a plankton community”. In: *Nature* 451.7180, pp. 822–825 (cit. on p. 7).
- Billiard, Sylvain, Régis Ferrière, Sylvie Méléard, and Viet Chi Tran (2015). “Stochastic dynamics of adaptive trait and neutral marker driven by eco-evolutionary feedbacks”. In: *Journal of Mathematical Biology* 71.5, pp. 1211–1242. arXiv: 1310.6274 (cit. on p. 14).
- Boschma, Ron A and Koen Frenken (2005). “Why is economic geography not an evolutionary science? Towards an evolutionary economic geography”. In: (cit. on p. 16).
- Boyd, Ian L. (2012). “The Art of Ecological Modeling”. In: *Science* 337.6092, pp. 306–307 (cit. on p. 7).
- Brummitt, Charles D., Andrés Gómez-Liévano, Ricardo Hausmann, and Matthew H. Bonds (2020). “Machine-learned patterns suggest that diversification drives economic development”. In: *Journal of The Royal Society Interface* 17.162, p. 20190283. arXiv: 1812.03534 (cit. on p. 4).
- Cauwels, Peter and Didier Sornette (2012). “Quis Predit Ipsa Pretia: Facebook Valuation and Diagnostic of a Bubble Based on Nonlinear Demographic Dynamics”. In: *The Journal of Portfolio Management* 38.2, pp. 56–66 (cit. on p. 4).
- Champagnat, Nicolas, Régis Ferrière, and Sylvie Méléard (2006). “Unifying evolutionary dynamics: From individual stochastic processes to macroscopic models”. In: *Theoretical Population Biology* 69.3, pp. 297–321 (cit. on pp. 5, 18, 20).

- Cordes, Christian (2006). "Darwinism in economics: from analogy to continuity". In: *Journal of Evolutionary Economics* 16.5, pp. 529–541 (cit. on p. 4).
- Curtsdotter, Alva, H. Thomas Banks, John E. Banks, et al. (2019). "Ecosystem function in predator-prey food webs: confronting dynamic models with empirical data". In: *Journal of Animal Ecology* 88.2. Ed. by Daniel Stouffer, pp. 196–210 (cit. on p. 7).
- Débarre, F., O. Ronce, and S. Gandon (2013). "Quantifying the effects of migration and mutation on adaptation and demography in spatially heterogeneous environments". In: *Journal of Evolutionary Biology* 26.6, pp. 1185–1202 (cit. on pp. 14, 17).
- Dieckmann, Ulf and Michael Doebeli (1999). "On the origin of species by sympatric speciation". In: *Nature* 400.6742, pp. 354–357 (cit. on p. 5).
- Doebeli, Michael and Ulf Dieckmann (2003). "Speciation along environmental gradients". In: *Nature* 421.6920, pp. 259–264 (cit. on pp. 3, 5, 17).
- Doebeli, Michael and Iaroslav Ispolatov (2010). "Complexity and diversity". In: *Science* 328.5977, pp. 494–497 (cit. on p. 6).
- Dopfer, Kurt and Jason Potts (2007). *The General Theory of Economic Evolution*. Vol. 4. Routledge, pp. 2–4 (cit. on p. 1).
- Dornelas, Maria, Laura H. Antão, Faye Moyes, et al. (2018). "BioTIME: A database of biodiversity time series for the Anthropocene". In: *Global Ecology and Biogeography* 27.7, pp. 760–786 (cit. on p. 19).
- Ellis, Erle C. (2011). "Anthropogenic transformation of the terrestrial biosphere". In: *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 369.1938, pp. 1010–1035 (cit. on p. 3).
- Farmer, J. D. and A. W. Lo (1999). "Frontiers of finance: Evolution and efficient markets". In: *Proceedings of the National Academy of Sciences* 96.18, pp. 9991–9992 (cit. on p. 4).
- Fisher, Rosie A., Charles D. Koven, William R.L. Anderegg, et al. (2018). "Vegetation demographics in Earth System Models: A review of progress and priorities". In: *Global Change Biology* 24.1, pp. 35–54 (cit. on p. 7).
- Freeman, Christopher (2002). *As time goes by : from the industrial revolutions to the information revolution*. Oxford: Oxford University Press (cit. on p. 4).
- Gatabazi, P., J.C. Mba, E. Pindza, and C. Labuschagne (2019). "Grey LotkaVolterra models with application to cryptocurrencies adoption". In: *Chaos, Solitons & Fractals* 122, pp. 47–57 (cit. on p. 4).
- Gillespie, Daniel T. (1976). "A general method for numerically simulating the stochastic time evolution of coupled chemical reactions". In: *Journal of Computational Physics* 22.4, pp. 403–434 (cit. on p. 20).
- Gosh, Sanmitra, Paul Birrell, and Daniela De Angelis (2021). "Variational inference for non-linear ordinary differential equations". In: *Proceedings of The 24th International Conference on Artificial Intelligence and Statistics* 130.29, pp. 2719–2727 (cit. on p. 23).

- Han, Jiequn, Arnulf Jentzen, and Weinan E (2018). “Solving high-dimensional partial differential equations using deep learning”. In: *Proc. Natl. Acad. Sci. USA* 115.34, pp. 8505–8510 (cit. on p. 18).
- Hastings, Alan, Carole L. Hom, Stephen Ellner, Peter Turchin, and H. Charles J. Godfray (1993). “Chaos in Ecology: Is Mother Nature a Strange Attractor?” In: *Annual Review of Ecology and Systematics* 24.1, pp. 1–33 (cit. on p. 7).
- Hidalgo, César A. (2021). “Economic complexity theory and applications”. In: *Nature Reviews Physics* 3.2, pp. 92–113 (cit. on p. 16).
- Higgins, Steven I., Simon Scheiter, and Mahesh Sankaran (2010). “The stability of African savannas: insights from the indirect estimation of the parameters of a dynamic model”. In: *Ecology* 91.6, pp. 1682–1692 (cit. on p. 7).
- Hodgson, Geoffrey M. (2019). *Evolutionary Economics*. Vol. 66. Cambridge University Press, pp. 37–39 (cit. on pp. 3, 4, 9, 16).
- Hodgson, Geoffrey M. and Thorbjorn Knudsen (2004). “The firm as an interactor: firms as vehicles for habits and routines”. In: *Journal of Evolutionary Economics* 14.3, pp. 281–307 (cit. on p. 4).
- Huisman, Jef and Franz J. Weissing (1999). “Biodiversity of plankton by species oscillations and chaos”. In: *Nature* 402.6760, pp. 407–410 (cit. on p. 7).
- Johnson, Jerald B. and Kristian S. Omland (2004). “Model selection in ecology and evolution”. In: *Trends in Ecology & Evolution* 19.2, pp. 101–108 (cit. on p. 7).
- Kashinath, K., M. Mustafa, A. Albert, et al. (2021). “Physics-informed machine learning: Case studies for weather and climate modelling”. In: *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 379.2194 (cit. on p. 8).
- Kawecki, Tadeusz J. and Dieter Ebert (2004). “Conceptual issues in local adaptation”. In: *Ecology Letters* 7.12, pp. 1225–1241 (cit. on p. 14).
- Kingma, Diederik P. and Jimmy Ba (2014). “Adam: A Method for Stochastic Optimization”. In: *arXiv:1412.6980*, 15 pp. arXiv: 1412.6980 (cit. on p. 19).
- Kirkpatrick, Mark and N. H. Barton (1997). “Evolution of a Species’ Range”. In: *The American Naturalist* 150.1, pp. 1–23 (cit. on p. 17).
- Lande, R. (1991). “Isolation by distance in a quantitative trait”. In: *Genetics* 128.2, pp. 443–452 (cit. on p. 14).
- Lawson, Tony (2013). “What is this school’ called neoclassical economics?” In: *Cambridge Journal of Economics* 37.5, pp. 947–983 (cit. on p. 3).
- LeCun, Yann, Yoshua Bengio, and Geoffrey Hinton (2015). “Deep learning”. In: *Nature* 521.7553, pp. 436–444 (cit. on p. 7).
- Lepers, Clotilde, Sylvain Billiard, Matthieu Porte, Sylvie Méléard, and Viet Chi Tran (2021). “Inference with selection, varying population size, and evolving population structure: application of ABC to a forwardbackward coalescent process with interactions”. In: *Heredity* 126.2, pp. 335–350. arXiv: 1910.10201 (cit. on p. 14).

- Levin, Simon A. (2002). "Complex adaptive systems: Exploring the known, the unknown and the unknowable". In: *Bulletin of the American Mathematical Society* 40.01, pp. 3–20 (cit. on pp. 1, 13).
- Lieberman, Erez, Christoph Hauert, and Martin A. Nowak (2005). "Evolutionary dynamics on graphs". In: *Nature* 433.7023, pp. 312–316 (cit. on p. 17).
- Lion, Sébastien (2016). "Moment equations in spatial evolutionary ecology". In: *Journal of Theoretical Biology* 405, pp. 46–57 (cit. on p. 5).
- Marasco, A., A. Picucci, and A. Romano (2016). "Market share dynamics using LotkaVolterra models". In: *Technological Forecasting and Social Change* 105, pp. 49–62 (cit. on p. 4).
- McRae, Brad H. (2006). "ISOLATION BY RESISTANCE". In: *Evolution* 60.8, p. 1551 (cit. on p. 14).
- McRae, Brad H. and Paul Beier (2007). "Circuit theory predicts gene flow in plant and animal populations". In: *Proceedings of the National Academy of Sciences* 104.50, pp. 19885–19890 (cit. on p. 14).
- Menon, Carlo (2015). "Spreading big ideas? The effect of top inventing companies on local inventors \*". In: *Journal of Economic Geography* 15.4, pp. 743–768 (cit. on p. 17).
- Meszéna, Géza, István Czibula, and Stefan Geritz (1997). "Adaptive Dynamics in a 2-Patch Environment: A Toy Model for Allopatric and Parapatric Speciation". In: *Journal of Biological Systems* 05.02, pp. 265–284 (cit. on pp. 3, 5, 14, 17).
- Michalakelis, Christos, Thomas Sphicopoulos, and Dimitris Varoutas (2011). "Modeling competition in the telecommunications market based on concepts of population biology". In: *IEEE Transactions on Systems, Man and Cybernetics Part C: Applications and Reviews* 41.2, pp. 200–210 (cit. on p. 4).
- Midgley, Guy and Lee Hannah (2019). "Extinction Risk from Climate Change". In: *Biodiversity and Climate Change*. Yale University Press, pp. 294–296 (cit. on p. 3).
- Mirrahimi, Sepideh and Sylvain Gandon (2020). "Evolution of specialization in heterogeneous environments: equilibrium between selection, mutation and migration". In: *Genetics* 214.2, pp. 479–491 (cit. on pp. 5, 14, 17).
- Mitchell, Melanie (n.d.). "Analogy-Making as a Complex Adaptive System". In: () (cit. on p. 16).
- Modis, Theodore (1997). "Genetic re-engineering of corporations". In: *Technological Forecasting and Social Change* 56.2, pp. 107–118 (cit. on p. 4).
- Norberg, Jon, Mark C. Urban, Mark Vellend, Christopher A. Klausmeier, and Nicolas Loeuille (2012). "Eco-evolutionary responses of biodiversity to climate change". In: *Nature Climate Change* 2.10, pp. 747–751 (cit. on pp. 3, 24).
- Nordbotten, Jan M., Simon A. Levin, Eörs Szathmáry, and Nils C. Stenseth (2018). "Ecological and evolutionary dynamics of interconnectedness and modularity". In: *Proc. Natl. Acad. Sci. USA* 115.4, pp. 750–755 (cit. on p. 2).

- Nordbotten, Jan Martin, Folmer Bokma, Jo Skeie Hermansen, and Nils Chr Stenseth (2020). “The dynamics of trait variance in multi-species communities”. In: *R. Soc. Open Sci.* 7.8, Article No. 200321, 20 pp. (Cit. on pp. 5, 21).
- Nowak, Martin A and Natalia L Komarova (2001). “Towards an evolutionary theory of language”. In: *Trends in Cognitive Sciences* 5.7, pp. 288–295 (cit. on p. 6).
- Odum, Eugene P. (1969). “The strategy of ecosystem development”. In: *Science* 164.3877, pp. 262–270 (cit. on p. 25).
- Orsini, Luisa, Joost Vanoverbeke, Ine Swillen, Joachim Mergeay, and Luc De Meester (2013). “Drivers of population genetic differentiation in the wild: Isolation by dispersal limitation, isolation by adaptation and isolation by colonization”. In: *Molecular Ecology* 22.24, pp. 5983–5999 (cit. on p. 14).
- Ozman, M. (2009). “Inter-firm networks and innovation: a survey of literature”. In: *Economics of Innovation and New Technology* 18.1, pp. 39–67 (cit. on pp. 4, 17).
- Pelletier, F., D. Garant, and A.P. Hendry (2009). “Eco-evolutionary dynamics”. eng. In: *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 364.1523, pp. 1483–1489 (cit. on pp. 3, 9).
- Polechová, Jitka (2018). “Is the sky the limit? On the expansion threshold of a species’ range”. In: *PLoS Biology* 16.6, pp. 1–18 (cit. on p. 17).
- Polechová, Jitka and Nicholas H. Barton (2015). “Limits to adaptation along environmental gradients”. In: *Proceedings of the National Academy of Sciences of the United States of America* 112.20, pp. 6401–6406 (cit. on p. 17).
- Pontarp, Mikael, Åke Bränström, and Owen L. Petchey (2019). “Inferring community assembly processes from macroscopic patterns using dynamic ecoevolutionary models and Approximate Bayesian Computation (ABC)”. In: *Methods in Ecology and Evolution* 10.4. Ed. by Timothée Poisot, pp. 450–460 (cit. on pp. 4, 7).
- Rackauckas, Christopher, Yingbo Ma, Julius Martensen, et al. (2020a). “Universal Differential Equations for Scientific Machine Learning”. In: *arXiv:2001.04385v3*, 18 pages (cit. on p. 8).
- Rackauckas, Christopher, Yingbo Ma, Julius Martensen, et al. (2020b). “Universal Differential Equations for Scientific Machine Learning”. In: *arXiv: 2001.04385* (cit. on p. 19).
- Rahbek, Carsten, Michael K. Borregaard, Alexandre Antonelli, et al. (2019a). “Building mountain biodiversity: Geological and evolutionary processes”. In: *Science* 365.6458, pp. 1114–1119 (cit. on pp. 2, 3).
- Rahbek, Carsten, Michael K. Borregaard, Robert K. Colwell, et al. (2019b). “Humboldt’s enigma: What causes global patterns of mountain biodiversity?” In: *Science* 365.6458, pp. 1108–1113 (cit. on p. 1).
- Raiissi, M., P. Perdikaris, and G.E. Karniadakis (2019). “Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations”. In: *Journal of Computational Physics* 378, pp. 686–707 (cit. on p. 23).

- Rangel, Thiago F., Neil R. Edwards, Philip B. Holden, et al. (2018). "Modeling the ecology and evolution of biodiversity: Biogeographical cradles, museums, and graves". In: *Science* 361.6399 (cit. on pp. 2, 3).
- Rogers, Everett M (2003). *Diffusion of Innovations*. 5th ed. Riverside: Free Press (cit. on p. 4).
- Saavedra, S., R. P. Rohr, L. J. Gilarranz, and J. Bascompte (2014). "How structurally stable are global socioeconomic systems?" In: *Journal of The Royal Society Interface* 11.100, pp. 20140693–20140693 (cit. on p. 4).
- Saavedra, Serguei, Felix Reed-tsochas, and Brian Uzzi (2009). "A simple model of bipartite cooperation for ecological and organizational networks". In: *Nature* 457.7228, pp. 436–466 (cit. on pp. 4, 17).
- Savolainen, Outi, Tanja Pyhäjärvi, and Timo Knürr (2007). "Gene Flow and Local Adaptation in Trees". In: *Annual Review of Ecology, Evolution, and Systematics* 38.1, pp. 595–619 (cit. on p. 3).
- Schartau, Markus, Philip Wallhead, John Hemmings, et al. (2017). "Reviews and syntheses: parameter identification in marine planktonic ecosystem modelling". In: *Biogeosciences* 14.6, pp. 1647–1701 (cit. on p. 7).
- Schneider, Tapio, Shiwei Lan, Andrew Stuart, and João Teixeira (2017). "Earth System Modeling 2.0: A Blueprint for Models That Learn From Observations and Targeted HighResolution Simulations". In: *Geophysical Research Letters* 44.24, pp. 12,396–12,417. arXiv: 1709.00037 (cit. on pp. 7, 8, 19).
- Scholl, Maarten Peter, Anisoara Calinescu, and J. Doyne Farmer (2021). "How market ecology explains market malfunction". In: *Proceedings of the National Academy of Sciences* 118.26, e2015574118. arXiv: 2009.09454 (cit. on p. 4).
- Sirignano, Justin and Konstantinos Spiliopoulos (2018). "DGM: A deep learning algorithm for solving partial differential equations". In: *J. Comput. Phys.* 375, pp. 1339–1364 (cit. on p. 8).
- Slatkin, Montgomery (1973). "GENE FLOW AND SELECTION IN A CLINE". In: *Genetics* 75.4, pp. 733–756 (cit. on p. 17).
- (1978). "Spatial patterns in the distributions of polygenic characters". In: *Journal of Theoretical Biology* 70.2, pp. 213–228 (cit. on p. 17).
- Suweis, Samir, Joel A Carr, Amos Maritan, Andrea Rinaldo, and Paolo D'Odorico (2015). "Resilience and reactivity of global food security". In: *Proceedings of the National Academy of Sciences* 112.22, pp. 6902–6907 (cit. on p. 4).
- Tacchella, A., D. Mazzilli, and L. Pietronero (2018). "A dynamical systems approach to gross domestic product forecasting". In: *Nature Physics* 14.8, pp. 861–865 (cit. on p. 4).
- Urban, M. C., G. Bocedi, A. P. Hendry, et al. (2016). "Improving the forecast for biodiversity under climate change". In: *Science* 353.6304 (cit. on pp. 3, 8, 9, 19, 20, 24).
- Van Der Panne, Gerben (2004). "Agglomeration externalities: Marshall versus Jacobs". In: *Journal of Evolutionary Economics* 14.5, pp. 593–604 (cit. on p. 4).

- Vellend, Mark (2010). "Conceptual Synthesis in Community Ecology". In: *The Quarterly Review of Biology* 85.2, pp. 183–206 (cit. on p. 2).
- Wernerfelt, Birger (1989). "From Critical Resources to Corporate Strategy". In: *Journal of General Management* 14.3, pp. 4–12 (cit. on p. 4).
- Wright, Sewal (1943). "Isolation By Distance". In: *Genetics* (cit. on p. 14).
- Yazdani, Alireza, Lu Lu, Maziar Raissi, and George Em Karniadakis (2020). "Systems biology informed deep learning for inferring parameters and hidden dynamics". In: *PLOS Computational Biology* 16.11. Ed. by Vassily Hatzimanikatis, e1007575 (cit. on pp. 8, 23).
- Yeaman, Sam and Sarah P. Otto (2011). "Establishment and maintenance of adaptive genetic divergence under migration, selection, and drift". In: *Evolution* 65.7, pp. 2123–2129 (cit. on p. 17).
- Zahra, Shaker A., R. Duane Ireland, and Michael A. Hitt (2000). "International Expansion by New Venture Firms: International Diversity, Mode of Market Entry, Technological Learning, and Performance". In: *Academy of Management Journal* 43.5, pp. 925–950 (cit. on p. 4).
- Zhang, Guanglu, Daniel A. McAdams, Venkatesh Shankar, and Milad Mohammadi Darani (2018). "Technology Evolution Prediction Using LotkaVolterra Equations". In: *Journal of Mechanical Design* 140.6, pp. 1–9 (cit. on p. 4).



# List of Figures

1.1 Schematic diagram of proposed organisational levels in biological and economic systems. A is inspired from (Hendry+2016) . . . . .	2
1.2 <b>Forward and inverse modelling approaches for the understanding of CAS.</b> A forward modelling approach starts with assuming a set of processes, that are embedded in a model, and seeks to understand how their interplay transforms in mechanisms that are associated with a macroscopic pattern (macroscopic invariance). An inverse modelling approach integrates empirical observation to constrain the model output, generating knowledge by inferring the probability and the strength of the processes investigated, given the empirical observations. Similarly to forward modelling, mechanisms may be deduced from the processes inferred, leading to the undersanding of invariances (pattern). . . . .	6
2.1 <b>Summary of the causal pathways involved in neutral and adaptive differentiation, disentangled in ??.</b> Ecological processes are displayed in green boxes, evolutionary processes are displayed in red boxes. . . . .	15

