


In-house liver imaging dataset




Demographic Summary


Patient count: 3,277




1. Gender



2. Age Groups



3. Ethnicity (Merged Categories)



Ethnicity merged from detailed source categories.

In-house liver imaging dataset

Patients: 85* Unique accession numbers: 281* 423 unique StudyInstanceUID CT series: 6399
imaging / scan_dir + imaging_metadata.csv accession numbers mapping: 268/281 found, 80 unique personid
imaging / scan_dir + imaging_transfer.csv person_id mapping: 86 patients, 1 lack date information.

Current processed data structure

```
PERSON_ID_1/  
├── YYYYMMDD__AccessionNumber_1/  
│   ├── CT_S005__Arterial__000030_000013/  
│   │   ├── 000001.dcm  
│   │   ├── 000002.dcm  
│   │   └── ...  
│   ├── CT_S006__Venous__000030_000014/  
│   │   └── ...  
│   ├── CT_S200__Locator__000030_000015/  
│   │   └── ...  
│   ├── REG_S003__NA__000030_000016/  
│   │   └── ...  
│   └── SR_S001__Dose_Report__000030_000017/  
│       └── ...  
├── YYYYMMDD__AccessionNumber_2/  
│   └── ...  
└── YYYYMMDD__AccessionNumber_3/  
    └── ...  
PERSON_ID_2/  
└── ...
```

Current statistics

Phase / category	Series	Unique patients	Unique scan dates	Unique accessions
Unknown	2,504	84	273	279
Arterial	895	81	258	265
Delayed	565	77	232	237
Portal venous	342	60	118	122
Localizer	208	58	138	140
Other venous	178	40	55	55
Dose report	45	25	42	42
Non-contrast	28	10	14	14


*Lower bounds from the supplied summary; overall deduplicated totals are not provided. Counts overlap across categories.

Source: current processed folder structure and CT phase summary supplied by the team.

PLC-CECT

361 patients. CT imaging with four acquisition phases.

Patients by diagnosis



HCC: hepatocellular carcinoma ICC: intrahepatic cholangiocarcinoma
cHCC-CCA: combined hepatocellular-cholangiocarcinoma

MODALITY

CT: 361 patients
NC / arterial / venous / delayed

CONTROL GROUP

83 people with no disease
of the liver or biliary tract

BIOMARKERS

Blood / molecular markers:
not reported in release


ACCESS

Public Science Data Bank
Download not verified here

MCT-LTDiag

517 patients. Every patient has four-phase CT.

Patients by diagnosis



CRLM: colorectal liver metastases. BCLM: breast cancer liver metastases.
Diagnosis follows pathological or clinical assessment.

Source: Wu et al., Scientific Data. HCC 103 + ICC 100 + CRLM 103 + BCLM 115 + hemangioma 96 = 517.

MODALITY

CT: 517 patients, 2,068 volumes
NC / arterial / portal / delayed

DIAGNOSIS LABELS

HCC, ICC, CRLM and BCLM
Hemangioma

BIOMARKERS

Blood measures not reported
Public lab table unconfirmed


ACCESS

Public release on Dataverse
Download permission unverified

WAW-TACE

233 patients. All patients have HCC (100%).

Patients by AFP level (ng/mL)



AFP = alpha-fetoprotein. AFP ≥400 ng/mL: 55/233 patients (23.6%).
The 400 ng/mL threshold describes the distribution within this HCC cohort.

MODALITY

Multiphase CT: 233 patients
NC / arterial / portal / delayed

COHORT / PURPOSE

All 233 patients have HCC
Segmentation and TACE research

BIOMARKERS

AFP, albumin, bilirubin,
creatinine, INR, ALT: 233 each

ACCESS

Public Zenodo release
CT, masks and clinical data

Source: WAW-TACE clinical table. Baseline AFP is available for all 233 patients before first TACE.

AMPLIFAI

584 unique patients. 590 CT studies in the training and validation release.

LI-RADS labels and study counts

Label	Meaning	Studies
LR-1	Definitely benign	3
LR-2	Probably benign	3
LR-3	Intermediate probability of HCC	16
LR-4	Probably HCC	47
LR-5	Definitely HCC by imaging	258
LR-TIV	Definite tumor in a vein	64
LR-M	Malignant, not specific for HCC	129
Unlisted*	Category not tabulated in the paper	70

*The seven LI-RADS counts total 520 studies. The remaining 70 are unlisted in that table. 590 studies represent 584 unique patients.

DATASET PURPOSE

Predict LI-RADS categories from multiphase liver CT

WHAT DOES LR MEAN?

LI-RADS = Liver Imaging Reporting and Data System

CT PHASES (STUDIES)

Non-contrast: 424 / 590

Arterial: 590 / 590

Portal venous: 586 / 590

Delayed: 451 / 590


Counts overlap across phases.
Access requires registration.

Source: AMPLIFAI, Tables 2–3. Phase counts refer to 590 CT studies from 584 patients; four parent datasets overlap.

TCGA-LIHC

97 imaging patients across the TCGA-LIHC collection. CT, MRI and PET are included.

Patients by collection diagnosis



MODALITIES

CT, MRI and PET
97 patients across all modalities

DIAGNOSIS LABELS

Hepatocellular carcinoma
Linked pathology and grading

BIOMARKERS

Clinical and genomic data
Match availability by patient ID

ACCESS

Public TCIA imaging
Genomic access varies by file

97 is the total imaging cohort, not the CT-only count.


Patient counts by modality and CT-biomarker intersections are not verified.

Source: TCIA TCGA-LIHC collection, 97 subjects. Modalities may overlap within patients.

AMOS-MM + LEAVS

Unique patient count not reported. The dataset contains 2,000+ CT scans.

Liver findings in the 200-scan test subset



Diffuse findings can include cirrhosis. Focal findings can include masses or cysts.
These labels describe the pattern of abnormality, not a specific cancer diagnosis.

DATASET PURPOSE

Generate reports from CT
Detect abnormal abdominal organs

LABEL MEANING

Normal: no reported abnormality
Diffuse: widespread liver changes

LABEL MEANING

Focal: localized lesion(s)
Both: diffuse and focal findings

CT ACQUISITION PHASE

Not specified in the cited sources
No phase-specific counts reported

LiTS

201 patients. One contrast-enhanced CT volume per patient.

Liver and tumor segmentation benchmark

INPUT IMAGES

201 CT volumes; one portal venous phase per case.

SEGMENTATION LABELS

131 training cases: liver and tumor masks.

70 test cases: reference masks are hidden.

The tumor masks are available. A uniform patient-level HCC / ICC / metastasis classification is not provided, so diagnostic subtype counts cannot be reported.

Sources: LiTS benchmark (cohort / masks); MCT-LTDiag, Table 1 (portal venous phase).
The original LiTS benchmark does not provide a per-case acquisition-timing inventory.

MODALITY / CT PHASE

Contrast-enhanced 3D CT
Portal venous phase (single phase)

TUMOR INFORMATION

Includes primary and
metastatic liver tumors

BIOMARKERS

Blood / molecular markers:
not reported in release


ACCESS

Public CT dataset
131 public tumor-mask sets

LiMT (LiMT)

150 patients / cases. Arterial-phase contrast-enhanced CT.

Patients with each diagnosis label



A patient may have multiple lesion types. Category counts overlap and must not be added to estimate the number of patients.

Source: Liu et al., LiMT 2025. Normal 50, cyst 49, hemangioma 37, HCC 31, metastases 20.

MODALITY

CT: 150 arterial-phase volumes
About 35 s after injection

DIAGNOSIS LABELS

HCC, metastases, cysts
Hemangioma and normal cases

BIOMARKERS

Blood / molecular markers:
not reported in release

ACCESS

Public Google Drive release
Dataset used in our project

HCC-TACE-Seg

105 patients with HCC. CT phases: non-contrast, arterial and portal venous. Scans were acquired before and after TACE.

CT acquisition phases and timing

CT phase	Reported acquisition timing
Non-contrast	Before contrast injection
Arterial	17 s after peak aortic enhancement using bolus tracking
Portal venous	60 s after contrast injection

Additional data: segmentation masks, clinical variables and treatment-response / survival outcomes.

IMAGING MODALITY

CT only in this collection
DICOM SEG stores annotations

DATASET PURPOSE

Liver / tumor segmentation
Predict response to TACE

BIOMARKERS / DIAGNOSIS

AFP: 105/105, all HCC
Differentiation grade: 83/105

SCAN TIME POINTS

Before TACE: 1–12 weeks
After TACE: within 14 weeks