



Nutrient Removal Values - Phosphorus

	Projected phosphorus removal (as P ₂ O ₅), "wet" or "as is" basis		% Moisture, Dry matter				Number of samples in survey *
			Median*		Survey range*		
	lb P ₂ O ₅ per yield unit *	Survey range*	- moisture -	----- dry matter -----			
Grains, oilseeds, fiber, sugar crops							
Barley, grain	0.35 / bu (48 lb)	¹	0.32 - 0.40	11.1	88.9	87.2 - 90.0	197
Beans (dry)	0.62 / cwt (100 lb)	²	---	10.0	90.0	---	---
Beans (navy)	1.24 / cwt (100 lb)	³	---	10.0	90.0	---	---
Buckwheat, grain	0.38 / bu (52 lb)	²	---	12.0	88.0	---	---
Canola, grain	0.86 / bu (60 lb)	⁴	---	8.0	92.0	---	---
Corn, grain (dry shelled)	0.31 / bu (56 lb)	¹	0.28 - 0.33	14.2	85.9	84.9 - 86.7	4,032
	0.55 / cwt (100 lb)	¹	0.51 - 0.59	14.2	85.9	84.9 - 86.7	4,032
Corn, grain (high moisture)	0.26 / bu (56 lb)	²	0.24 --- 0.28	29.7	70.3	67.4 - 72.9	4,820
	0.46 / cwt (100 lb)	²	0.43 --- 0.50	29.7	70.3	67.4 - 72.9	4,820
Cotton (seed, lint, trash) (≈1300 lb seed cotton yields 480 lb lint)	11.2 / 1300 lb (bale)	³	---	---	---	---	---
Cotton seeds	1.33 / cwt (100 lb)	²	---	8.0	92.0	---	---
Field pea, seeds	0.56 / bu (60 lb)	²	---	11.5	88.5	---	---
Flax, grain	0.84 / bu (56 lb)	⁵	---	8.7	91.3	---	---
Millet grain	0.36 / bu (50 lb)	²	---	13.0	87.0	---	---
Oats, grain	0.23 / bu (32 lb)	¹	0.22 - 0.27	11.0	89.0	87.8 - 90.6	91
Peanuts, whole	0.55 / cwt	⁶	---	10.0	90.0	---	---
Popcorn	0.38 / bu (56 lb)	³	---	10.1	89.9	---	---
Potato tuber, fresh	0.13 / cwt (100 lb)	³	---	77.0	23.0	---	---
Rice, grain	0.29 / bu (45 lb)	²	---	11.0	89.0	---	---
Rye, grain	0.62 / bu (56 lb)	²	---	11.0	89.0	---	---
Safflower seeds	0.55 / bu (38 lb)	⁴	---	8.0	92.0	---	---
Sorghum, grain (milo)	0.32 / bu (56 lb)	¹	0.30 - 0.36	13.0	87.0	85.8 - 88.2	820
	0.58 / cwt (100 lb)	¹	0.54 - 0.63	13.0	87.0	85.8 - 88.2	820
Sorghum, grain (high moisture)	0.28 / bu (56 lb)	¹	0.24 - 0.29	30.5	69.5	65.2 - 74.5	97
	0.49 / cwt (100 lb)	¹	0.43 - 0.52	30.5	69.5	65.2 - 74.5	97
Soybeans, whole	0.78 / bu (60 lb)	⁴	---	12.0	88.0	---	---
Spelt, grain	0.32 / bu (40 lb)	²	---	12.0	88.0	---	---
Sugarbeet (root)	2.19 / ton	³	---	80.0	20.0	---	---
Sugarcane	1.20 / ton	⁷	---	74.0	26.0	---	---
Sunflower seeds	1.37 / cwt (100 lb)	⁴	---	9.8	90.3	---	---
Triticale, grain	0.47 / bu (60 lb)	²	---	11.0	89.0	---	---
Wheat, grain	0.46 / bu (60 lb)	¹	0.42 - 0.51	11.3	88.7	87.5 - 89.9	355
Hay, legumes							
Alfalfa hay	10.0 / ton	¹	8.4 - 11.1	12.7	87.3	85.0 - 89.1	8,360
Clover, hay alsike clover	9.7 / ton	²	---	12.6	87.4	---	---
Clover hay, ladino clover	13.1 / ton	²	---	10.0	90.0	---	---

"Phosphorus" is expressed in terms of fertilizer equivalent phosphate as " P_2O_5 ", where $P \times 2.28 = P_2O_5$.

To estimate the equivalent hay or silage yield based on grain yield or hay yield based on stocking rates as animal units, see Crop File 1.02.022, Yield Goal Equivalents: Silage, Hay, and Pasture

To estimate the equivalent hay or silage yield based on grain yield or hay yield based on stocking rates as animal units, see Crop File 1.02.022, Yield Goal Equivalents: Silage, Hay, and Pasture	Projected phosphorus removal (as P ₂ O ₅), "wet" or "as is" basis		% Moisture, Dry matter				Number of samples in survey *	
			Median*		Survey range*			
	lb P ₂ O ₅ per yield unit	*	Survey range*	- moisture -	----- dry matter -----			
Hay, legumes (continued)								
Clover hay, red clover	10.0	/ ton	2	---	12.0	88.0	---	---
Clover hay, sweetclover	10.4	/ ton	2	---	9.0	91.0	---	---
Cowpea hay	12.5	/ ton	2	---	9.6	90.4	---	---
Lespedeza hay	9.2	/ ton	6	---	8.0	92.0	---	---
Peanut hay	3.7	/ ton	1	3.3 - 4.9	10.8	89.2	86.4 - 90.8	56
Sanfoin hay	n/a			---	---	---	---	---
Soybean hay	6.7	/ ton	1	4.4 - 8.0	13.0	87.1	83.2 - 90.4	76
Trefoil hay, birdsfoot trefoil	9.7	/ ton	2	---	11.0	89.0	---	---
Vetch hay	13.8	/ ton	2	---	11.0	89.0	---	---
Hay, grasses								
Bahiagrass	8.2	/ ton	2	---	10.0	90.0	---	---
Bermudagrass hay	8.6	/ ton	1	7.0 - 10.3	10.0	90.0	87.4 - 91.8	350
Bluegrass hay	10.7	/ ton	7	---	11.0	89.1	---	---
Bluestem hay	5.1	/ ton	3	---	9.9	90.1	---	---
Bromegrass hay	7.2	/ ton	1	5.6 - 8.4	11.8	88.2	86.3 - 89.9	241
CRP hay	4.4	/ ton	1	-	11.4	88.6	85.7 - 90.6	---
Fescue hay	10.4	/ ton	2	---	12.2	87.8	---	---
Meadow hay	7.3	/ ton	1	6.1 - 8.6	10.6	89.4	87.0 - 90.9	215
Mixed grass hay	7.1	/ ton	1	4.9 - 8.6	10.7	89.3	86.5 - 91.8	2,951
Native grass hay	4.4	/ ton	1	3.2 - 6.4	12.9	87.1	67.5 - 90.5	227
Orchardgrass hay	12.0	/ ton	2	---	12.0	88.0	---	---
Prairie hay	4.8	/ ton	2	3.6 --- 6.1	11.4	88.6	86.6 - 90.2	295
Reed canarygrass hay	10.4	/ ton	2	---	9.0	91.0	---	---
Ryegrass hay	12.3	/ ton	2	---	8.5	91.5	---	---
Switchgrass hay	5.9	/ ton	3	---	4.6	95.4	---	---
Teff hay	n/a			---	---	---	---	---
Timothy hay	8.0	/ ton	2	---	12.0	88.0	---	---
Wheatgrass hay	8.4	/ ton	2	---	11.7	88.4	---	---
Hay, small grains								
Barley hay	9.5	/ ton	1	7.9 - 11.1	13.2	86.8	84.1 - 88.7	757
Millet hay	7.5	/ ton	1	5.9 - 9.5	13.1	86.9	83.3 - 89.4	399
Oat hay	9.6	/ ton	1	8.4 - 11.2	12.1	87.9	84.9 - 90.1	919
Rye hay	10.8	/ ton	1	8.2 - 12.8	12.4	87.6	84.6 - 90.2	71
Triticale hay	9.8	/ ton	1	8.2 - 11.8	10.4	89.6	86.0 - 92.0	204
Wheat hay	8.9	/ ton	1	6.9 - 10.9	11.5	88.5	84.9 - 90.7	684
Hay, annual sorghums								
Canex hay	n/a			---	---	---	---	---
Forage sorghum hay	6.1	/ ton	1	5.0 - 8.0	16.1	83.9	76.6 - 90.0	101
Haygrazer hay	6.4	/ ton	1	5.1 - 8.7	17.2	82.8	54.6 - 88.9	139
Milo (grain sorghum) hay	5.8	/ ton	1	4.0 - 7.2	21.0	79.0	61.3 - 85.9	128
Sorghum hay	6.4	/ ton	1	5.3 - 8.3	17.1	82.9	74.7 - 87.5	294
Sorghum-sudan hay	6.9	/ ton	1	5.7 - 8.8	16.3	83.7	74.7 - 88.0	156
Sudan hay	6.5	/ ton	1	5.4 - 8.8	16.2	83.8	77.3 - 87.8	370
Sumac hay	n/a			---	---	---	---	---

"Phosphorus" is expressed in terms of fertilizer equivalent phosphate as "P₂O₅", where P x 2.28 = P₂O₅.

To estimate the equivalent hay or silage yield based on grain yield or hay yield based on stocking rates as animal units, see Crop File 1.02.022, Yield Goal Equivalents: Silage, Hay, and Pasture

	Projected phosphorus removal (as P ₂ O ₅), "wet" or "as is" basis		% Moisture, Dry matter				Number of samples in survey *
			Median*		Survey range*		
	lb P ₂ O ₅ per yield unit	*	Survey range*	- moisture -	----- dry matter -----		
Hay, miscellaneous							
Cane hay	6.4 / ton	¹	4.9 - 7.9	17.0	83.0	77.4 - 86.8	610
Corn hay	6.6 / ton	¹	3.5 - 8.3	14.3	85.7	77.9 - 89.7	108
Kochia hay	8.2 / ton	²	---	12.6	87.4	---	---
Weed hay	8.6 / ton	¹	6.5 - 10.7	12.2	87.9	84.4 - 90.3	58
Silage, annual crops							
Barley silage	4.5 / ton	¹	3.8 - 5.1	66.3	33.7	30.8 - 38.3	387
Cane silage	2.3 / ton	¹	2.1 - 2.6	71.4	28.6	24.9 - 33.3	231
Corn silage	3.4 / ton	¹	2.9 - 3.8	66.3	33.7	30.2 - 37.8	9,910
Corn silage (sweet corn)	n/a		---	---	---	---	---
Forage sorghum silage	2.4 / ton	³	---	70.2	29.8	---	---
Milo silage (grain sorghum)	3.8 / ton	¹	3.1 - 4.2	64.9	35.1	28.9 - 41.4	103
Oat silage	4.2 / ton	¹	3.7 - 5.1	64.8	35.3	29.1 - 42.8	115
Oatlage	4.1 / ton	¹	3.5 - 4.9	66.3	33.7	28.9 - 43.0	145
Rye silage	4.5 / ton	¹	3.8 - 5.7	68.9	31.1	26.0 - 38.6	214
Small grain silage	4.3 / ton	¹	3.8 - 4.8	65.1	34.9	30.7 - 38.7	597
Sorghum silage	2.6 / ton	¹	2.2 - 3.1	70.5	29.5	26.2 - 33.1	1,750
Sorghum-sudan silage	2.2 / ton	¹	---	68.5	31.5	---	---
Sudangrass silage	3.2 / ton	³	2.3 - 3.8	71.0	29.1	24.5 - 36.3	71
Triticale silage	4.3 / ton	¹	3.7 - 5.1	67.3	32.7	28.1 - 38.8	346
Wheatlage	4.1 / ton	¹	3.6 - 4.8	63.9	36.1	31.5 - 41.7	881
Wheat silage	4.2 / ton	¹	3.6 - 4.9	65.5	34.5	30.3 - 39.9	410
Silage, perennial crops							
Alfalfa haylage	5.4 / ton	¹	4.7 - 6.2	59.1	40.9	34.1 - 48.1	752
Alfalfa silage	4.4 / ton	¹	3.9 - 5.3	64.0	36.0	31.3 - 43.2	168
Bermudagrass silage	2.4 / ton	²	---	74.0	26.0	---	---
Bromegrass silage	4.8 / ton	²	---	65.0	35.0	---	---
Mixed grass silage	3.3 / ton	²	---	70.0	30.0	---	---
Ryegrass silage	5.5 / ton	²	---	68.0	32.0	---	---
Timothy silage	4.2 / ton	²	---	66.0	34.0	---	---
Crop residues							
Barley straw	3.3 / ton	³	---	10.0	90.0	---	---
(72 lb straw/bu)	0.12 / bu harvested	³	---	---	---	---	---
Buckwheat straw	4.1 / ton	³	---	11.3	88.7	---	---
Canola straw (236 lb straw/cwt)	n/a		---	---	---	---	---
Corn cobs	1.6 / ton	²	---	6.4	93.6	---	---
Corn stalks	3.5 / ton	¹	2.7 - 4.7	15.0	85.0	79.9 - 88.1	132
Corn stover	4.9 / ton	¹	3.2 - 6.9	11.3	88.7	83.3 - 92.1	100
(70 lb stover/bu)	0.14 / bu harvested	³	---	---	---	---	---
Cotton stover							
(1300 lb seed cotton yields 480 lb lint)	6.7 / bale	²	---	7.7	92.3	---	---
Flax straw (80 lb straw/bu)	0.14 / bu harvested	³	---	7.7	93.0	---	---
Grain sorghum (milo) stalks	n/a		---	---	---	---	---
Millet straw	4.3 / ton	²	---	10.0	90.0	---	---
(84 lb straw/bu)	0.18 / bu harvested	³	---	---	---	---	---

"Phosphorus" is expressed in terms of fertilizer equivalent phosphate as "P₂O₅", where P x 2.28 = P₂O₅.

To estimate the equivalent hay or silage yield based on grain yield or hay yield based on stocking rates as animal units, see Crop File 1.02.022, Yield Goal Equivalents: Silage, Hay, and Pasture

To estimate the equivalent hay or silage yield based on grain yield or hay yield based on stocking rates as animal units, see Crop File 1.02.022, Yield Goal Equivalents: Silage, Hay, and Pasture

	Projected phosphorus removal (as P ₂ O ₅), "wet" or "as is" basis		% Moisture, Dry matter				Number of samples in survey *	
			Median*		Survey range*			
	lb P ₂ O ₅ per yield unit	*	Survey range*	- moisture -	----- dry matter -----			
Crop residues								
Oat straw (44 lb straw/bu)	2.9	/ ton	2	---	9.0	90.3	---	---
	0.06	/ bu harvested	3	---	---	---	---	---
Peanut stover	6.8	/ ton	2	---	---	---	---	---
Popcorn stover (56 lb stover/bu)	30.8	/ ton	3	---	12.7	87.3	---	---
	0.86	/ bu harvested	3	---	---	---	---	---
Potato vines	2.0	/ ton	2	---	85.0	15.0	---	---
Rice straw (100 lb straw/cwt)	3.3	/ ton	3	---	9.2	90.8	---	---
	0.17	/ cwt harvested	3	---	---	---	---	---
Rye straw (87 lb straw/bu)	3.0	/ ton	7	---	9.1	90.9	---	---
	0.17	/ bu harvested	3	---	---	---	---	---
Sorghum stover (56 lb stover/bu)	4.8	/ ton	2	---	8.1	91.9	---	---
	0.16	/ bu harvested	3	---	---	---	---	---
Soybean stover (96 lb stover/bu)	1.1	/ ton	2	---	13.9	86.2	83.1 88.2	69
	0.11	/ bu harvested	3	---	---	---	---	---
Sugarbeet top	1.9	/ ton	2	---	81.0	19.0	---	---
Sunflower stover (150 lb stover/cwt)	2.0	/ ton	7	---	13.3	86.7	---	---
	0.24	/ cwt harvested	3	---	---	---	---	---
Wheat straw (92 lb straw/bu)	2.9	/ ton	1	2.5 - 4.9	9.8	90.3	86.5 94.0	187
	0.13	/ bu harvested	3	---	---	---	---	---

Fresh forage, vegetative stage - annuals

Cane plants, fresh	2.3 / ton	3	---	73.7	26.4	---	---
Corn plants, fresh	2.6 / ton	1	2.3 - 3.2	74.0	26.1	21.3 - 33.8	121
Forage sorghum plants, fresh	2.3 / ton	1	---	71.5	28.5	---	---
Millet plants, fresh	3.4 / ton	3	2.6 - 4.1	70.0	30.1	22.2 - 40.2	56
Milo plants, fresh (grain sorghum)	2.5 / ton	3	---	66.6	33.4	---	---
Oat plants, fresh	4.0 / ton	3	---	70.4	29.6	---	---
Rye plants, fresh	3.5 / ton	3	---	68.9	31.2	---	---
Sorghum plants, fresh	2.6 / ton	1	1.9 - 3.6	71.9	28.1	23.3 - 34.1	112
Sorghum-sudan plants, fresh	2.3 / ton	3	---	72.4	27.6	---	---
Sudangrass plants, fresh	3.0 / ton	2	---	75.1	24.9	---	---
Triticale plants, fresh	4.4 / ton	3	---	68.0	32.0	---	---
Weeds, fresh	n/a		---	---	---	---	---
Wheat plants, fresh	4.4 / ton	1	3.5 - 5.4	63.2	36.9	27.5 - 47.3	92

Fresh forage, vegetative stage - perennial grasses

Bermudagrass, fresh	2.6 / ton	3	---	71.0	29.0	---	---
Bluestem, fresh	3.3 / ton	2	---	39.0	61.0	---	---
Bromegrass, fresh	4.7 / ton	2	---	70.0	30.0	---	---
Fescue, fresh	4.9 / ton	2	---	71.0	29.0	---	---
Mixed grass, fresh	3.9 / ton	1	1.9 - 5.1	30.5	69.6	46.2 - 86.9	2,870
Native grass, fresh	3.4 / ton	1	2.3 - 4.5	58.5	41.6	33.5 - 52.1	149
Orchardgrass, fresh	4.3 / ton	2	---	76.0	24.0	---	---
Timothy, fresh	3.3 / ton	2	---	74.0	26.0	---	---
Wheatgrass (crested), fresh	5.4 / ton	2	---	63.0	37.0	---	---

"Phosphorus" is expressed in terms of fertilizer equivalent phosphate as "P₂O₅", where $P \times 2.28 = P_2O_5$.

To estimate the equivalent hay or silage yield based on grain yield or hay yield based on stocking rates as animal units, see Crop File 1.02.022, Yield Goal Equivalents: Silage, Hay, and Pasture

Projected phosphorus removal (as P ₂ O ₅), "wet" or "as is" basis		% Moisture, Dry matter		Number of samples in survey *
		Median*	Survey range*	
lb P ₂ O ₅ per yield unit	*	Survey range*	- moisture - ----- dry matter -----	

Fresh forage, vegetative stage - legumes

Alfalfa, fresh	4.3	/ ton	¹	3.8 - 4.9	72.4	27.6	20.8 - 39.5	63
Clover (ladino), fresh	3.3	/ ton	²	---	81.0	19.0	---	---
Clover (red), fresh	3.3	/ ton	²	---	76.0	24.0	---	---
Lespedeza, fresh	2.7	/ ton	²	---	75.0	25.0	---	---
Trefoil (birdsfoot) fresh	2.5	/ ton	²	---	78.0	22.0	---	---

Removal by animal type^{8,9}

Beef cattle	0.0051 lb P ₂ O ₅ per pound of weight gain (based on average 20% protein in meat, bones, offal, etc.)
Poultry	0.0257 lb P ₂ O ₅ per 5 lb market-weight broiler
Dairy cows	0.00217 lb P ₂ O ₅ per cwt (100 lb) of milk

"Phosphorus" is expressed in terms of fertilizer equivalent phosphate as "P₂O₅", where $P \times 2.28 = P_2O_5$.

(Crop residue weights per unit of yield harvested were derived from NRCS PLANTS database.)

* References and Resources

- Servi-Tech Laboratories. 2014. In-house data survey of feed analysis results, 2006-2013. Results reported as "Median" or 50th percentile. "Typical range" includes the middle 50% of all samples, so 25% of samples are above range and 25% below. Statistical calculations based on listed number of samples of particular feedsuff that was included in data survey.
- BEEF. 2014 Feed Composition Table. BEEF magazine, Minneapolis MN(<http://beefmagazine.com/datasheet/2014-feed-composition-table>, 25 June 2014)
- USDA, NRCS. 2007. The PLANTS Database . National Plant Data Center, Baton Rouge, LA(<http://plants.usda.gov>, 25 June 2014)
- Walker, J. 2007. Oilseed Crops in Beef Cattle Rations. Extension Extra 2058. South Dakota State Univ. Coop. Ext. Serv., Brookings SD. (http://pubstorage.sdstate.edu/agbio_publications/articles/exex2058.pdf, 25 June 2014)
- Dahnke, W.C., L.J. Swenson, and A. Johnson. Crop Nutrient Removals vs. Fertilizer Additions in North Dakota. http://library.ndsu.edu/tools/dspace/load/?file=/repository/bitstream/handle/10365/8084/farm_44_01_03.pdf
- Osmond, D.L. and J. Kang. 2008. Nutrient Removal by Crops in North Carolina. Soil Facts #AG-439-16W. North Carolina State Univ., Coop. Ext. Serv., Raleigh NC.
- IPNI. IPNI Estimates of Nutrient Uptake and Removal. International Plant Food Institute, Peachtree Corners, GA(<http://www.ipni.net/article/IPNI-3296>, 25 June 2014)
- Smolen, M.D., P.L. Kenkel, and D.E. Storm. 1995. Nitrogen and phosphorus distribution in Oklahoma: A mass balance of animal feeds, animal manures, crops and fertilizer nutrients. Proceedings of "Innovations and New Horizons in Livestock and Poultry Manure Management", Austin, Texas. 6-7 Sept. 1995 pg. 39-50.
- Phosphorus nutrition and excretion by dairy animals. CIR849. Coop. Ext. Serv., Univ. of Florida. 2003.<http://edis.ifas.ufl.edu/ds165>

Albers, D.W., S. Hefner, and D. Klobe. 1993. Fertility Management of Cotton. Pub. G4256. Univ. of Missouri Extension, Columbia MO. (<http://extension.missouri.edu/p/G4256>, 25 June 2014)

Pettygrove, S. and I. Bay. 2009. Manure Technical Bulletin Series: Crop Nutrient Harvest Removal. Div. of Agric. and Nat.Res., Univ. of California, Davis CA (<http://manuremanagement.ucdavis.edu/files/134365.pdf>, 25 June 2014)

Rochester, J.J., G.A. Constable, D.M. Oosterhuis, and M. Errington. 2012. "Nutritional Requirements of Cotton During Flowering and Fruiting" in Flowering and Fruiting in Cotton. The Cotton Foundation, Cordova TN. pg. 41. (www.cotton.org/foundation/upload/F-F-Chapter-4.pdf, 25 June 2014)

Stichler, C. and M. McFarland. 2001. Crop Nutrient Needs in South and Southwest Texas. Pub. No. B-6053. Texas Agric. Ext. Serv., College Station TX.