


# Developing Low Temperature Contacts for Monolayer TMD Heterostructure Devices

---

Vanessa Kwong

Dr. Andrew Joe

University of California, Riverside, Department of Physics and Astronomy





# Background

Semiconductor transition metal dichalcogenide (TMD) devices bring efficiency to electronics.

- Silicon-based chip
- **Direct band gap** in the monolayer limit, by which a photon can be emitted without the aid of a phonon


**WSe<sub>2</sub>** and device quality through Photoluminescence (PL)




# Device Fabrication Method

## Mechanical Exfoliation

- Scotch tape cleaves bulk material at where vdW forces < chemical bonds
- An array of replicated, sheared bulk material pressed down onto substrate.
- **O<sub>2</sub> plasma** *organic materials*
- **Heating** of tapes *varied adhesion*





Graphite, h-BN (illustrated), and WSe<sub>2</sub> onto wafer chips




WSe<sub>2</sub> monolayers (Measurements scaled at 100x on 90nm SiO<sub>2</sub>)

## Dry Transfer

- Stacking of samples through picking up and dropping off with **PDMS/PC** stamp on microscope slide
- Heating / cooling of stamp for adhesiveness




# Patterning



1  $\mu\text{m}$

*Hall bar design on KLayout*


- Alignment marks and pre-patterned contacts
- Unique pattern superimposed on alignment marks with transferred on with identical coordinates





# Optical (PL) data for gate tunability




- Single-gated device with graphite contact to prepattern



- Au vs Pt

Applied voltage values → Wavelength and intensity of light emitted (PL)



- Electron doping





$X_0$  = neutral exciton emission

$X_T$  = charged exciton emission


$X_D$  = dark exciton, out-of-plane  
transition dipole moment




# Horizontal linecuts of PL plots



Exciton activity  
(broader)  
Wavelength ~750nm



Distinct peaks



Doped region



# Linecut fitted to Lorentzian function

**Line width**  
*Wavelengths  
in a spectral  
line; width at  
half max*

**Narrow**

identical exciton  
behavior (ideal)




$w = 8.4 \text{ meV}$

**Broad**

Various wavelengths  
emitted; Defects  
from cleanliness or  
temperature

$w = 2 - 4 \text{ meV}$   
at 4K  
(Zhou, et al.  
*Nat. Nanotech.*)



$w = 34.5 \text{ meV}$



# Conclusion


- With current data, we continue with the fabrication of a device with Hall bar patterning
- Current data determine quality of  $\text{WSe}_2$
- Guidance for PL

Our reference brings us closer to the ultimate goal of our research of measuring charged transport in developed contacts for  $\text{WSe}_2$



# Acknowledgements

*University of California, Riverside, Department of Physics and  
Astronomy*



Prof. Andrew Joe

Zhihan Wu    Hongyu Yao  
SM Umayer    Martin Ochoa  
Louis Lin

**Center for Nanoscale  
Science & Engineering**

Dong Yan  
Mohammad Alghamdi

**Jing Shi Research  
Group**

Jia-Mou Chen



*External*

**National Institute for  
Materials Science**

*2D materials - hBN*  
Takashi Taniguchi  
Kenji Watanabe

# References

- Barbone, Matteo, et al. "Charge-tunable biexciton complexes in monolayer WSe<sub>2</sub>." *Nature Communications*, **9**, 3721, Sept. 2018. DOI: <https://doi.org/10.1038/s41467-018-05632-4>
- Chen, Xiaotong, et al. "Excitonic Complexes in Two-Dimensional Transition Metal Dichalcogenides." *Nature Communications*, **14**, 8233, Dec. 2023. DOI: <https://doi.org/10.1038/s41467-023-44119-9>
- Fallahazad, Babak, et al. "Shubnikov-de Haas oscillations of high mobility holes in monolayer and bilayer WSe<sub>2</sub>: Landau level degeneracy, effective mass, and negative compressibility." *Physical Review Letters*, **116**, Feb. 2016. DOI: <https://doi.org/10.1103/PhysRevLett.116.086601>.
- Joe, Anderew Y., et al. "Transport Study of Charge Carrier Scattering in Monolayer WSe<sub>2</sub>." *Physical Review Letters*, **132**, Jan 2024. DOI: <https://doi.org/10.1103/PhysRevLett.132.056303>.
- Liu, Erfu, et al. "Valley-selective chiral phonon replicas of dark excitons and trions in monolayer WSe<sub>2</sub>." *Physical Review Research*, **1**, 3, Oct. 2019. DOI: <https://doi.org/10.1103/PhysRevResearch.1.032007>
- Tuan, Dinh Van, et al. "Six-Body and Eight-Body Exciton States in Monolayer WSe<sub>2</sub>." *Physical Review Letters*, **129**, Aug. 2022. DOI: <https://doi.org/10.1103/PhysRevLett.129.076801>
- Zhou, You, et al. "Probing dark excitons in atomically thin semiconductors via near-field to surface plasmon polaritons." *Nature Nanotechnology*, **106**, Jun. 2017. DOI: <https://doi.org/10.1038/nnano.2017.106>