THE MIDPOINT RIEMANN–VON MANGOLDT SYNTHETIC ZETA SYSTEM: CANONICAL PRODUCTS, EXPONENTIAL SUMS, PRIME COUNTS, MERTENS TRANSFORMS, AND THE TRANSFORMATION TO ζ_{Z1}

ROTKIV AYNÄSAC
rotkivanyasac@gmail.com

Date: 9 October 2026.

Abstract. We study the zeta-like system generated by the midpoint inverse of the smooth Riemann–von Mangoldt counting law. Its nontrivial zeros are $\frac{1}{2} \pm i\tau_k$, where τ_k is the midpoint ordinate of the clock. We prove normal convergence of the canonical product, the functional symmetry, the exact zero divisor, a uniform logarithmic bound for the exponential sums of the τ_k, compact-uniform existence of the explicit-formula prime-power count with an explicit branch of Ei, exact finite Möbius inversion to a synthetic prime count π_{RVM}, and the unconditional square-root estimate $\pi_{\mathrm{RVM}}(x) - \mathrm{li}(x) = O(\sqrt{x}(\log\log x)^2/\log x)$; with the prime number theorem this gives $\mathrm{li}/\pi_{\mathrm{RVM}} \to 1$, $\pi/\mathrm{li} \to 1$ and $\pi/\pi_{\mathrm{RVM}} \to 1$. The transition theorem $\zeta_{\mathrm{RVM}} = \varpi_{\mathrm{RVM}}(1+w)(1+O(1/|t|))$ is proved in full; it gives a constant vertical limit of $1/\zeta_{\mathrm{RVM}}$ on every line to the right of $\frac{1}{2}$, excludes absolutely convergent Dirichlet expansions of $\zeta_{\mathrm{RVM}}^{\pm 1}$, and proves the unsmoothed synthetic Mertens bound $M^{\mathrm{crit}}_{\mathrm{RVM}}(x) = L + O_\varepsilon(x^{1/2+\varepsilon})$, $L = 1/\varpi_{\mathrm{RVM}}$. A contour-free form of the same decomposition shows that ζ_{RVM} is bounded above and below on every half-plane $\Re s \geq \sigma_0 > 1$ and tends to ϖ_{RVM} as $\Re s \to +\infty$.

The core of the method is a transformation of ζ_{RVM} depending on a parameter $\lambda \in [0,1]$: the ordinates move along $\gamma_k(\lambda) = (1-\lambda)\gamma_k + \lambda\tau_k$, where γ_k are the ordinates of the zeros of the Riemann zeta function, every member ζ_λ has all its nontrivial zeros on $\Re s = \frac{1}{2}$ by construction, $\zeta_1 = \zeta_{\mathrm{RVM}}$, and $\zeta_0 = \zeta_{Z1}$. It is impossible for ζ_{Z1}, or for any ζ_λ, to have a zero off the critical line other than the trivial zeros. All differences between the moving zetas lie in the numbers $\gamma_k - \tau_k$. The final step of the method is the identity $\zeta_{Z1} = \zeta$. This identity is the single and complete test: while $\zeta_{Z1} = \zeta$ there is nothing outside $\Re s = \frac{1}{2}$ to speak of, and $\zeta_{Z1} \neq \zeta$ is the only possible obstruction.

We prove Step 3 in the form: if $\zeta_{Z1} = \zeta$, then $M(x) = O_\varepsilon(x^{1/2+\varepsilon})$ for the Mertens function M, with a self-contained proof (Borel–Carathéodory, three circles, truncated Perron). By Hamburger's theorem the identity $\zeta_{Z1} = \zeta$ is equivalent to the condition (D) that ζ_{Z1} is an absolutely convergent Dirichlet series in $\Re s > 1$. We prove three further forms of (D): the value of the endpoint at infinity is $\varpi_{Z1} = \lim_{\sigma\to+\infty} \zeta_{Z1}(\sigma) = \varpi_{\mathrm{RVM}} \prod_k (\frac{1}{4}+\tau_k^2)/(\frac{1}{4}+\gamma_k^2) \geq 1$, and (D) holds exactly when $\varpi_{Z1} = 1$, that is, when $\prod_k (\frac{1}{4}+\gamma_k^2)/(\frac{1}{4}+\tau_k^2) = \varpi_{\mathrm{RVM}}$; the prime-power counts of ζ_{RVM} and ζ_{Z1} are the Mellin transforms of their logarithms and satisfy a distributional explicit formula, which proves the Mellin–Stieltjes identification for the integer weights; and (D) holds exactly when the prime-power count of ζ_{Z1} is constant between consecutive integers, an identity on each cell between the displacements $\gamma_k - \tau_k$ and the clock. Along the family, $\log\varpi_\lambda$ has derivative $-\Upsilon(\lambda)$, the generator at infinity, and (D) holds exactly when $\int_{0^+}^1 \Upsilon\, d\lambda = \log(1/\varpi_{\mathrm{RVM}})$. Finally we atomize every member of the family by smooth cells: the cell masses have mean square $X^{1+\varepsilon}$ on every dyadic block, uniformly in λ, so every atomized function $\tilde\zeta_{\mathbb{Z},\lambda} = \zeta_\lambda e^{\tilde h_\lambda}$, the endpoint $\lambda = 0$ included, is an absolutely convergent Dirichlet series in $\Re s > 1$; the joining factor $\tilde h_\lambda$ is written cell by cell and is holomorphic in $\Re s > \frac{1}{2}$, and (D) is reduced to a single statement at $\lambda = 0$, the vanishing of every cell defect, which is the remaining step of the programme. We prove further that no member ζ_λ with $0 < \lambda \leq 1$ is a Dirichlet series, that $\lambda \mapsto \log\zeta_\lambda(\sigma)$ is strictly convex for every real $\sigma > 1$, and that $\zeta_{Z1}(\sigma) \geq \zeta(\sigma)$ for every real $\sigma > 1$, with equality at one point exactly when $\zeta_{Z1} = \zeta$. Hamburger's theorem is extended to the pair $(\zeta_\lambda, \tilde\zeta_{\mathbb{Z},\lambda})$: $\tilde\zeta_{\mathbb{Z},\lambda} = \zeta$ holds as soon as the joining factor is symmetric under $s \mapsto 1-s$, equivalently when the atomized coefficients satisfy the theta relation of ζ, equivalently when the vertical mean square of $\tilde\zeta_{\mathbb{Z},\lambda}/\zeta$ on a line $\Re s = \sigma > 1$, which is always at least 1, equals 1; at the clock end the symmetry fails, and $\tilde\zeta_{\mathbb{Z},\lambda} = \zeta$ holds for at most finitely many λ. (D) holds exactly when ζ_{Z1} is the Mellin–Stieltjes transform of a positive measure: the functional equation then gives $\varpi_{Z1} \leq 1$ by an extremal summation argument, and in general $\varpi_\lambda - 1$ is one explicit number computed from the summation formula of ζ_λ. At the clock end the density of $d\pi_{\mathrm{RVM}}$ is $\frac{1}{\log y} + 2\cos(2\pi y) - 2^{-1/2}y^{-1/4}\cos(4\pi\sqrt{y} + \frac{\pi}{4}) + O(y^{-1/3})$. Finally the constructions are carried out for every list of ordinates within a bounded distance of the clock: the lists whose continuous and atomized zeta functions coincide form a set with at most one element, which is non-empty exactly when (D) holds, so that the final step is an existence statement in a compact space of lists. In this version: (D) holds if and only if every cell mass at $\lambda = 0$ is the indicator of the primes; a bridge theorem on the phase of the joining factor on the critical line has hypotheses that fail for every $\lambda > 0$ and are equivalent to (D) at $\lambda = 0$; (D) is equivalent to the solvability of a system with free masses at the points $\log n$; and for the finite Euler products $\prod_{p\leq X}(1-p^{-s})$, models without ζ, the final step is a theorem.

What is proved and what is not. Every statement above is proved in this paper without any hypothesis. At the clock end we further prove a multiscale theorem for the linearized rung system: up to height T the ordinate block has a uniform lower frame bound on u-windows of length $(1+\eta)\log(T/2\pi)$, no T-independent lower bound is possible on a fixed window, the infinite clock linearization has dense range on every finite window, and smooth transport tails beyond height T are $O_{A,\mathrm{f}}(T^{1-A}\log T)$. These statements use only the midpoint clock and no individual displacement $\eta_k = \tau_k - \gamma_k$. They rule out a fixed-window implicit-function argument and reduce the existence route to a quantitative multiscale continuation estimate, which is not proved.

The identity $\zeta_{Z1} = \zeta$, equivalently (D), equivalently each of the criteria above, is not proved; it is the open step of the method.

Contents

Notation 5
1. Introduction 8
1.1. Background 8
1.2. The midpoint RVM system 8
1.3. The core of the method 10
1.4. Originality, scope and status 11
1.5. Organization 14
2. The midpoint Riemann–von Mangoldt clock 14
3. The pure critical RVM zeta function 15
4. RVM exponential sums, centroids, and the height transform 16
5. The RVM explicit formula, J_{RVM}, and π_{RVM} 17
5.1. The square-root estimate 18
6. Integer weights and the continuous-to-atomic problem 20
7. The critical RVM Mertens transform 23
8. The transition theorem and the unsmoothed synthetic Mertens bound 24
8.1. Proof of Theorem 8.1 24
Step 0: reduction to a sum over the clock 24
Step 1: the midpoint sawtooth 24
Step 2: assembly 26
Step 3: from the clock phase to χ 27
8.2. Bounds on the whole half-plane 28
9. The exact Hadamard product and the comparison with ζ 30
10. The λ-family from ζ_{RVM} to ζ_{Z1} 31
11. The endpoint ζ_{Z1}, Hamburger's theorem, and Step 3 33
11.1. The analytic class of the endpoint 33
11.2. Hamburger's theorem at the endpoint 34
11.3. Condition (D) and the integer-weighted zeta 34
11.4. Step 3: the Mertens bound from $\zeta_{Z1} = \zeta$ 35
11.5. Reciprocal endpoint and finite-height Perron integrals 38
12. The endpoint at infinity and the Euler products 38
12.1. The value of the endpoint at infinity 38
12.2. The Euler products of ζ_{RVM} and ζ_{Z1} 40
13. The atomized family and the programme for the final step 42
13.1. The family at infinity 42
13.2. Atomization by smooth cells 43
13.3. The joining factor cell by cell 44
13.4. Mean square of the cell masses 45
13.5. Analytic continuation of the atomized family 47
13.6. The two endpoints of the atomized family 48
13.7. Hamburger's hypotheses at the two endpoints 49
13.8. Programme 51
14. Criteria for the final step and the clock end of the family 53
14.1. The open path 53
14.2. Convexity along the family 53
14.3. Hamburger's theorem for the pair 54
14.4. Positivity and the value at infinity 55
14.5. The density of the clock 58
14.6. The cell masses in the heights of the ordinates 61

14.7. The Gaussian defect of the atomization 62
14.8. Entire dependence on λ and the exceptional set 63
14.9. The final step in the cell masses 64
14.10. Symmetric joining factors 65
14.11. The half-line form of Hamburger's theorem for a pair 66
14.12. The bridge theorem: the phase of the joining factor on the critical line 67
14.13. Transport from the clock on the rungs 69
14.14. The multiscale Jacobian of the midpoint clock 71
15. The space of ordinate lists and the existence form of the final step 72
15.1. Lists, their zeta functions and their atomizations 73
15.2. The rung system with free masses 74
15.3. The gradient of the vertical defect 76
16. Acceptance criteria and models in which the final step is a theorem 76
16.1. The acceptance criterion: window and frequencies 77
16.2. A non-periodic accepted list: one unitary matrix 78
16.3. A finite Euler product: the constructions of this paper with the final step proved 78
16.4. The diagonal operator over all primes and the three abscissas 80
17. Numerical illustration with 2 001 052 zeros 80
17.1. The clock and the zeros 81
17.2. The λ-family and the endpoint 81
17.3. The transition theorem and ζ_{RVM} against ζ 83
17.4. Exponential sums 83
17.5. Prime counts: π and π_{RVM} 83
17.6. The Mertens functions and the ternary division 86
17.7. The rung equations: transport, the defect of the clock, and the solution from the clock 88
18. Conclusions and open problems 92
Appendix A. Python programs 95
A.1. `rvm_common.py` 95
A.2. `s01_clock_and_family.py` 98
A.3. `s02_transition.py` 101
A.4. `s03_exponential_sums.py` 102
A.5. `s04_primes.py` 103
A.6. `s05_mertens.py` 106
A.7. `s06_prime_ratios.py` 111
A.8. `s07_product_identity.py` 112
A.9. `s08_transport_moments.py` 113
A.10. `s09_clock_defect_split.py` 114
A.11. `s10_clock_to_zeros.py` 115
Appendix B. Console output of the programs 117
B.1. `s01_clock_and_family` 117
B.2. `s01b_clock_zoom` 117
B.3. `s02_transition` 118
B.4. `s03_exponential_sums` 118
B.5. `s04_primes` 118
B.6. `s05_mertens` 119
B.7. `s06_prime_ratios` 120
B.8. `s07_product_identity` 121
B.9. `s08_transport_moments` 121
B.10. `s09_clock_defect_split` 122
B.11. `s10_clock_to_zeros` 122

References 123
Addendum to v17: what remains live after the interpolation tests 125
Three interpolations and their status 125
The clock as the first transition term 125
What remains genuinely live 126
Additional finite-section diagnostics 126

Notation

Every symbol below has a single meaning throughout the paper. The last column links to the defining equation; hovering over the link shows the formula. Symbols introduced inside a single proof (indices j, k_0, m, auxiliary variables v, y, thresholds t_0, t_1, t_2, radii r_1, r_2, r_3) are local to that proof and do not appear elsewhere.

Symbol	Meaning	Defined
Variables and the Riemann zeta function		
$s = \sigma + it$	complex variable, $\sigma = \Re s$, $t = \Im s$	–
$z = s - \frac{1}{2} = \sigma' + it$	centred variable, $\sigma' = \sigma - \frac{1}{2}$	(8.2)
$q(s) = s(s-1)$	symmetric quadratic, $q(1-s) = q(s)$	(3.1)
ζ, ρ	Riemann zeta function; ρ runs over its nontrivial zeros with multiplicity	§2
γ_k	kth positive ordinate of the nontrivial zeros of ζ, with multiplicity, nondecreasing	(2.1)
$N_\zeta(T)$, $N_\zeta^*(T)$	zero count up to height T; its symmetric version	(2.1), (2.10)
$S(T)$	remainder in the Riemann–von Mangoldt formula	(2.9)
$G(s)$	$s(s-1)\Gamma(s/2)\pi^{-s/2}$	(3.3)
$P_\zeta = 2\xi$	completed product $G\zeta$	(9.1)
μ, $M(x)$	Möbius function; Mertens function $\sum_{n\le x}\mu(n)$	(1.1)
$\pi(x)$, li	prime-counting function; logarithmic integral	–
$\chi(s)$, ϑ	functional-equation factor; Riemann–Siegel theta function	(8.1), (8.29)
The midpoint clock		
$\mathcal{N}_{\mathrm{RVM}}(T)$	midpoint Riemann–von Mangoldt clock	(2.3)
τ_k	midpoint ordinates, $\mathcal{N}_{\mathrm{RVM}}(\tau_k) = k - \frac{1}{2}$	(2.4)
W_0	principal branch of Lambert's W-function	(2.5)
$\mathcal{T}$	inverse function of the clock	(4.4)
ℓ_k	$\log(\tau_k/2\pi)$	(4.5)
$n_{\mathrm{RVM}}(H)$	number of $\tau_k \le H$	(4.12)
η_k, C_η	$\eta_k = \tau_k - \gamma_k$; a bound $\lvert\eta_k\rvert \le C_\eta$	(2.11), (2.12)
The synthetic zeta function and its sums		
$P_{\mathrm{RVM}}^{\mathrm{crit}}$, ξ_{RVM}, ζ_{RVM}	canonical product, completed and synthetic zeta	(3.2), (3.4)
ρ_k^{RVM}	synthetic zero $\frac{1}{2} + i\tau_k$	(3.6)
$S_{\mathrm{RVM}}(N;u)$, $C_{\mathrm{RVM}}^{\mathrm{pts}}$, $C_{\mathrm{RVM}}^{\mathrm{path}}$	exponential sum; point and path centroids	(4.1), (4.2)
u_1, u_2	a frequency interval $[u_1, u_2] \subset (0,\infty)$	(4.3)
$M_H(x)$, $M_H^{\mathrm{RVM}}(x)$	finite-height phase averages (not Mertens functions)	(4.14), (4.13)
Ei	exponential integral on $\Im w \neq 0$	(5.1)
J_{RVM}, J_{RVM}^*, π_{RVM}	explicit-formula prime-power count; truncation; synthetic prime count	(5.2), (5.6), (5.7)
$\mathcal{A}_{\mathrm{RVM}}(u)$, $\ell(t)$	weighted clock sum $\sum_k e^{iu\tau_k}/\tau_k$; $\log(t/2\pi)$	(5.10)

Symbol	Meaning	Defined
$\mathfrak{s}_1$; $\mathfrak{e}$, $\mathfrak{e}_1$	primitive of the midpoint sawtooth; $N_\zeta - \mathcal{N}_{\mathrm{RVM}}$ and its primitive	(8.36), (12.1)
ϖ_{Z1}	value of ζ_{Z1} at $+\infty$	(12.4)
J_X^{M}, J_X, C_X	Mellin and explicit-formula prime-power counts of ζ_X, $X \in \{\mathrm{RVM}, Z1\}$; their difference	(12.14), (12.16)
b_n	jumps of J_{Z1}^{M} at the integers	Thm. 12.7
η_k; ϖ_λ; $\Upsilon(\lambda)$	$\tau_k - \gamma_k$; value of ζ_λ at $+\infty$; flow at infinity	(13.1), (13.3)
φ_n, $\tilde{c}_n(\lambda)$, $\tilde{a}_N(\lambda)$	smooth cells; atomized masses and coefficients	(13.7), (13.8)
$\tilde{\zeta}_{\mathbb{Z},\lambda}$, $\tilde{h}_\lambda$	atomized family; joining factor	(13.7), (13.13)
$\mathfrak{a}_s$, $\mathfrak{h}_n(\lambda; s)$, $\mathfrak{L}$	atomization kernel; cell defects; $\int_1^x (1 - y^{-1})dy/\log y$	(13.10)
V_n, W_n	ordinate kernels of the cell n	(13.19)
V_n, W_n	ordinate kernels of the cell n	(13.19)
$\tilde{\mu}_\lambda$, $\tilde{M}_\lambda$	atomized Möbius coefficients; summatory function	(13.31)
B_{Z1}	Mellin form of the endpoint Mertens function	(13.33)
c_n^{RVM}, $\zeta_{\mathbb{Z},\mathrm{RVM}}$	integer weights; weighted Euler product	(6.1), (6.2)
$J_{\mathbb{Z},\mathrm{RVM}}$, $J_{\mathrm{RVM}}^{*,\mathbb{Z}}$, h_{RVM}	integer-step counts; Mellin correction	(6.3), (6.4), (6.20)
θ	unit-cell step, $0 < \theta \leq 1$	(6.5)
p_j^{RVM}	real RVM prime levels $\pi_{\mathrm{RVM}}^{-1}(j)$ on monotone branches	(1.6)
$\mathcal{M}(n)$, ς_n, μ_h, M_h	target samples of $M_{\mathrm{RVM}}^{\mathrm{crit}}$; signs; ternary walk and its sum (companion object; h is a label)	(1.9), (1.10)
$a_N^{\mathbb{Z}}$	Dirichlet coefficients of $\zeta_{\mathbb{Z},\mathrm{RVM}}$	(6.14)
C_J	Lipschitz constant of J_{RVM}	(6.5)
$\mathfrak{d}_n^{\mathrm{RVM}}$	prime defect $\mathbf{1}_{\mathrm{prime}}(n) - c_n^{\mathrm{RVM}}$	(9.11)
Mertens transforms and the transition theorem		
$F_{\mathrm{RVM}} = \widehat{M}_{\mathrm{RVM}}^{\mathrm{crit}}$	$1/\zeta_{\mathrm{RVM}}$	(7.1)
c, U	Perron abscissa; truncation height	(8.45)
$M_{\mathrm{RVM}}^{\mathrm{crit}}$, $M_{\mathrm{RVM},G}^{\mathrm{crit}}$	symmetric Perron inverse; Gaussian-smoothed inverse	(8.45), (7.5)
τ_*, $w(s)$	transition point $-iz = t - i\sigma'$; transition variable $e^{-2\pi i \mathcal{N}_{\mathrm{RVM}}(\tau_*)}$	(8.2)
B; σ_1', σ_2'	upper bound for σ'; a range $[\sigma_1', \sigma_2'] \subset (0, \frac{1}{2})$	(8.4), (8.5)
$\mathcal{E}_0, \mathcal{E}$	error factors of the transition theorem	(8.4), (8.5)
ϖ_{RVM}, L	transition constant $0.9619415776\ldots$; $L = 1/\varpi_{\mathrm{RVM}}$	(8.28), (8.33)
$f, \Lambda, \Lambda_{\mathrm{sm}}, \Lambda_{\mathrm{saw}}$	logarithm of the product and its smooth and sawtooth parts (§8 only)	(8.6), (8.9)
$\psi, \psi_r, \Phi_r, \nu_1, \ldots, \nu_4$	sawtooth, Abel means, ray remainder, real constants (§8 only)	(8.8), (8.20)
α_c	$\min(c - \frac{1}{2}, 1)$	(8.35)
Comparison with ζ		
$\mathcal{Q}_{\mathrm{RVM}}$, $\mathcal{Q}_{\mathrm{RVM}}^\vee$	quotient $\zeta/\zeta_{\mathrm{RVM}}$; its inverse Mellin transform	(9.6), (9.8)
The λ-family and the endpoint ζ_{Z1}		
λ	family parameter; $\lambda = 1$: ζ_{RVM}, $\lambda = 0$: ζ_{Z1}	(10.1)
$\gamma_k(\lambda)$	moving ordinates $(1 - \lambda)\gamma_k + \lambda\tau_k$	(10.1)
P_λ, ζ_λ, F_λ	family product, family zeta, $1/\zeta_\lambda$	(10.2), (10.3)
$A_{k,\lambda}$	$\frac{1}{4} + \gamma_k(\lambda)^2$	(10.10)

Symbol	Meaning	Defined
$\mathcal{V}_\lambda$	infinitesimal generator $\partial_\lambda \log P_\lambda$	(10.10)
Π_{tr}	total multiplier $P_{Z1}/P_{\mathrm{RVM}}^{\mathrm{crit}}$	(10.15)
$\rho_{k,\lambda}$, $\mathfrak{r}_{k,\lambda}$	moving zero $\frac{1}{2}+i\gamma_k(\lambda)$; residue of $1/(s\zeta_\lambda)$	(10.19)
P_{Z1}, ζ_{Z1}, F_{Z1}	endpoint product, endpoint zeta, $1/\zeta_{Z1}$	(11.1), (11.34)
$\mathscr{R}_{Z1}$	ratio ζ_{Z1}/ζ	(11.7)
$\mathcal{K}$	multiplier $\zeta_{Z1}/\zeta_{\mathbb{Z},\mathrm{RVM}}$	(11.12)
(D), a_n	Dirichlet condition for ζ_{Z1} and its coefficients	(11.8)
C_H	constant in Hamburger's theorem	(11.10)
μ_{Z1}, M_{Z1}	coefficients of F_{Z1}; summatory function	(11.37)
$M_{\lambda,U}$, $M_{Z1,U}$	finite-height Perron integrals	(11.35)
δ, ε, $\omega(\delta)$	margins $\sigma \geq \frac{1}{2}+\delta$, exponent $\varepsilon>0$, three-circle exponent	(11.16)
Criteria in the half-plane, the rung system and the models		
$\mathsf{D}_n(\lambda)$, $B_\lambda^\vee$	$\tilde{c}_n(\lambda)-\tilde{c}_n(0)$; dual function of the atomized family	(14.57), (14.58)
d, Y, $\mathsf{e}_1,\mathsf{e}_2$	difference of two counts; auxiliary entire function; exponents	(14.61), (14.63), (14.67)
J_f, J_g, b_n, ϖ_f	counts and coefficients of a general pair (f,g); value of f at $+\infty$	(14.65), Thm. 14.36
(B1), (B2)	the two hypotheses of the bridge theorem	Thm. 14.36
v_λ	phase defect of the family on the critical line	(14.69)
o_n, L_n, j	one-ordinate difference of cell masses; $-\log(1-n^{-s})$; one-ordinate joining factor	(14.72), (14.75)
f, c_f, k, k_f	test function in $u=\log x$; its cosine transform; the part of G and the pole	(14.77)
R_f, S_f, $\mathsf{m}_j(\mathsf{f})$	rung functional; transport; moments sampled on the clock	(14.78), (14.79), (14.82)
$\mathfrak{t}_0$, I_m	zero of the clock, $\mathcal{N}_{\mathrm{RVM}}(\mathfrak{t}_0)=0$; Poisson integrals	(14.83)
a_n, $\dot{\kappa}_k$, $\dot{\mathsf{a}}_n$	free masses of the rung system; a direction of variation	(15.5), (15.9)
J, x_0, H_1, y, H, Z	objects of the proof of Theorem 15.6	(15.10)–(15.12)
$\langle\cdot,\cdot\rangle_\sigma$, $\mathsf{K}_{\boldsymbol{\kappa}}$, r_m, u_m, $\mathsf{Q}_{\boldsymbol{\kappa}}$	pairing of Dirichlet series; $\tilde{\zeta}_{\mathbb{Z},\boldsymbol{\kappa}}/\zeta$; gradient data of V_σ	(15.13), (15.14)
z, $\mathfrak{k}$, $\mathfrak{C}_{\boldsymbol{\kappa}}$, N	variable, half-type and canonical product of the models; $e^{i\mathfrak{k}\mathsf{z}}\mathfrak{C}_{\boldsymbol{\kappa}}$	(16.1)
$\mathscr{F}$, $\mathscr{F}^+$, $\mathfrak{l}$, $\mathfrak{w}_\mathfrak{l}$, $\mathfrak{b}_\mathfrak{l}$, $\mathfrak{c}_\mathfrak{l}$	frequencies of a discrete category; a frequency; masses, exponents, coefficients	(16.3)–(16.5)
$\mathfrak{p},\mathfrak{q},\mathfrak{v}$, U, X, $\mathfrak{D}$, $\mathfrak{S}$	data, unitary matrix, diagonal factor, determinant and sine function of the two-edge model	(16.6)–(16.8)
F, $\mathsf{F}_{\boldsymbol{\kappa}}$, F_λ, $\mathfrak{y}_k$, $\mathfrak{x}_k$	finite Euler product, its list functions and family; its ordinates and its clock	(16.10)–(16.13)
$\mathscr{L}$, $\mathfrak{P}$	ℓ^2 over the primes; $\sum_p p^{-s}$	(16.18), (16.19)
$\nu_5,\dots,\nu_{12}$	real constants, each local to the statement where it occurs	–

Collision check. The table was checked symbol by symbol against the whole text: no symbol carries two meanings. In particular c denotes only a Perron abscissa and c_n^{RVM} only the integer weights; $\mathcal{N}_{\mathrm{RVM}}$ is the clock and n_{RVM} the count of the τ_k; $\mathcal{V}_\lambda$ is the generator of the family and Φ_r a

remainder inside one proof; ε is only an exponent; θ is the unit-cell step and ϑ the Riemann–Siegel function; $E_k = e^{iu\tau_k}$ while $\mathcal{E}_0, \mathcal{E}$ are error factors; (D) is only the Dirichlet condition; T is a height, and the target of the ternary walk is $\mathcal{M}(n)$ with signs ς_n; the subscript h of μ_h, M_h is a label, distinct from h_{RVM}. The sans-serif and Fraktur letters of the last group of the table are distinct from the italic letters; z is the variable of the models of Section 16 and $z = s - \frac{1}{2}$ the centred variable; X denotes the diagonal factor in the two models where it occurs; the dots in $\dot\kappa_k$, $\dot{\mathsf{a}}_n$ denote the components of a direction of variation.

1. Introduction

1.1. **Background.** The theory of the Riemann zeta function joins four structures that are usually introduced in different languages. On the spectral side there are the nontrivial zeros and their Riemann–von Mangoldt counting law; on the analytic side the completed function ξ, its functional equation and its Hadamard product; on the prime side the prime-power measure and the explicit formula; and on the inverse side the reciprocal Dirichlet series $1/\zeta$ and the Mertens function

$$M(x) := \sum_{n \le x} \mu(n). \tag{1.1}$$

Riemann's memoir made the first three structures inseparable, and the work of von Mangoldt, Hadamard and Hamburger supplied, respectively, the rigorous zero count and explicit formula, the entire-product factorization, and a converse characterization of ζ from its Dirichlet-series and functional-equation data [34, 38, 22, 23, 37, 18]. This paper asks what remains of that architecture when the zero staircase is replaced, at the outset, by the exact midpoint inverse of its smooth Riemann–von Mangoldt clock, and how the resulting object is carried back to ζ.

1.2. **The midpoint RVM system.** The point of departure is the smooth counting function (2.3),

$$\mathcal{N}_{\mathrm{RVM}}(T) = \frac{T}{2\pi} \log \frac{T}{2\pi e} + \frac{7}{8}, \tag{1.2}$$

and the real ordinates τ_k determined by the midpoint condition $\mathcal{N}_{\mathrm{RVM}}(\tau_k) = k - \frac{1}{2}$. The half-integer convention is the midpoint convention for a staircase and leads to the exact Lambert inversion (2.5),

$$\tau_k = \frac{2\pi(k - 11/8)}{W_0\big((k - 11/8)/e\big)}. \tag{1.3}$$

Lambert-W approximations to individual zero ordinates and smooth inversions of the Riemann–von Mangoldt law occur in the literature, including explicit equations for the nth zero [19, 10, 17]. Here the midpoint inverse is treated as the exact spectrum of a deterministic system and is propagated, without the fluctuating term $S(T)$, through canonical products, explicit prime counting and Mertens inversion.

The first object is the canonical product and the synthetic zeta function

$$P^{\mathrm{crit}}_{\mathrm{RVM}}(s) = \prod_{k \ge 1} \left(1 + \frac{s(s-1)}{\frac{1}{4} + \tau_k^2}\right), \qquad \zeta_{\mathrm{RVM}}(s) = \frac{P^{\mathrm{crit}}_{\mathrm{RVM}}(s)}{G(s)}, \qquad G(s) = s(s-1)\Gamma(s/2)\pi^{-s/2}. \tag{1.4}$$

Because $\tau_k \asymp k/\log k$, the factors form a normally convergent product in the symmetric variable $s(s-1)$; $P^{\mathrm{crit}}_{\mathrm{RVM}}$ is entire and invariant under $s \mapsto 1 - s$, and ζ_{RVM} has a residue-one pole at 1, the trivial zeros, and the simple nontrivial zeros $\rho_k^{\mathrm{RVM}} = \frac{1}{2} \pm i\tau_k$ (Theorem 3.1). The real part $\frac{1}{2}$ is part of the definition.

Generalized prime systems. In Beurling's theory the generalized primes are arbitrary reals $1 < p_1 \le p_2 \le \cdots \to \infty$; their multiplicative semigroup defines generalized integers and a zeta function by a Dirichlet series, an Euler product or a Mellin–Stieltjes transform, separating prime distribution from the additive structure of the integers [5, 2, 14]. Beurling systems with prescribed analytic behaviour, critical zeros and large oscillation show that zeta-like zero data alone do not

determine the arithmetic realization [15, 7]. The RVM system runs in the opposite direction: it starts from a prescribed zero clock, constructs an entire completed product first, obtains a continuous prime-power count from its explicit formula, and only afterwards asks for an atomic realization on the integers.

Continuous and discrete prime systems. The passage from a continuous prime system to a discrete one is classical in the theory of Beurling generalized primes [5, 15, 8, 16]: the atoms are placed at suitably chosen real points, and the discretization changes the zeta function by a factor holomorphic and zero-free in $\Re s > \frac{1}{2}$ (compare Theorem 13.12(iii)). Measures with a continuous part and non-integer multiplicities also appear in the theory of complex dimensions [27, Ch. 3]. Grosswald and Schnitzer [20] replaced each prime p_n by a real q_n with $p_n \le q_n \le p_{n+1}$ and proved that the modified Euler product is ζ times a factor holomorphic and zero-free in $\Re s > 0$, so that it has the same zeros as ζ there; Corollary 6.5 and Theorem 13.12(iv) are of the same type for ζ_{RVM}. Here the masses of $d\pi_\lambda$ are placed at the integers, with real exponents $\tilde{c}_n(\lambda)$; this yields an ordinary Dirichlet series for every λ, absolutely convergent in $\Re s > 1$ (Corollary 13.11), and makes Hamburger's theorem [23, 35] available at the endpoint. Hamburger's theorem and its extensions [6, 24] need absolute convergence in $\Re s > 1$, and Knopp [25] showed that a half-plane of absolute convergence further to the right does not suffice. We are not aware of an earlier use of this combination.

Prime coordinate. From the RVM zeros we define the prime-power count J_{RVM} by (5.2), with the branch (5.1) of Ei. Finite Möbius inversion in the exponent variable, which relates Riemann's prime-power count to $\pi(x)$ [34, 37, 18], is used here as the *definition* of the synthetic prime count (5.7),

$$\pi_{\mathrm{RVM}}(x) = \sum_{1 \le m \le \log_2 x} \frac{\mu(m)}{m}\, J^*_{\mathrm{RVM}}(x^{1/m}), \qquad J^*_{\mathrm{RVM}}(x) = \sum_{1 \le m \le \log_2 x} \frac{1}{m}\, \pi_{\mathrm{RVM}}(x^{1/m}). \tag{1.5}$$

The function π_{RVM} is continuous, not a staircase. It supports two notions of RVM prime: the continuous measure $d\pi_{\mathrm{RVM}}$, and, on every interval where π_{RVM} is continuous and strictly increasing, the real level crossings

$$p_j^{\mathrm{RVM}} := \pi_{\mathrm{RVM}}^{-1}(j), \qquad \pi_{\mathrm{RVM}}(p_j^{\mathrm{RVM}}) = j, \tag{1.6}$$

which need not be integers. The synthetic count has square-root accuracy without any hypothesis on ζ: $J_{\mathrm{RVM}}(x) - \mathrm{li}(x)$ and $\pi_{\mathrm{RVM}}(x) - \mathrm{li}(x)$ are $O(\sqrt{x}(\log\log x)^2/\log x)$ (Theorem 5.5). The proof reduces the zero sum to $\sum_k e^{iu\tau_k}/\tau_k$, which is $O(\log^2 u)$ by the second-derivative test on dyadic blocks of the clock (Lemma 5.4). Hence $\mathrm{li}(x)/\pi_{\mathrm{RVM}}(x) \to 1$, and with the prime number theorem $\pi(x)/\mathrm{li}(x) \to 1$ and $\pi(x)/\pi_{\mathrm{RVM}}(x) \to 1$ (Corollary 5.6, Figure 10). Sampling on unit cells gives the masses $c_n^{\mathrm{RVM}} = \pi_{\mathrm{RVM}}(n) - \pi_{\mathrm{RVM}}(n-1)$ and the weighted Euler object $\zeta_{\mathbb{Z},\mathrm{RVM}}$ of (6.2). The masses are not 0 or 1 and need not be positive, but they are bounded: J_{RVM} is Lipschitz on $[2,\infty)$ (Theorem 6.1), so $c_n^{\mathrm{RVM}} = O(1)$ and $\zeta_{\mathbb{Z},\mathrm{RVM}}$ is an absolutely convergent Dirichlet series in $\Re s > 1$ (Corollaries 6.2 and 6.3). The unit-periodic first harmonic of the clock integrates to zero on every integer cell (Lemma 6.6); under the cell hypothesis one obtains $c_n^{\mathrm{RVM}} \sim 1/\log n$ (Proposition 6.8).

Mertens coordinate. The Mertens coordinate is the symmetric inverse Mellin object of $F_{\mathrm{RVM}} = 1/\zeta_{\mathrm{RVM}}$. A smooth/sawtooth Stieltjes decomposition of the clock product with Abel regularization of the sawtooth gives on every band to the right of $\frac{1}{2}$

$$\zeta_{\mathrm{RVM}}(s) = \varpi_{\mathrm{RVM}}\big(1 + w(s)\big)\big(1 + O(1/|t|)\big), \qquad w(s) = \exp\{-2\pi i\, \mathcal{N}_{\mathrm{RVM}}(-i(s-\tfrac{1}{2}))\}, \tag{1.7}$$

and $\zeta_{\mathrm{RVM}} = \varpi_{\mathrm{RVM}}(1+\chi)(1+O(1/|t|))$ on closed substrips of $\frac{1}{2} < \Re s < 1$ (Theorem 8.1). Hence $F_{\mathrm{RVM}}(\sigma + it) \to L = 1/\varpi_{\mathrm{RVM}}$ for every fixed $\sigma > \frac{1}{2}$; neither ζ_{RVM} nor F_{RVM} is an absolutely convergent Dirichlet series in any half-plane (Theorem 8.11); and after subtracting the vertical limit the Perron inverse converges absolutely, with

$$M^{\mathrm{crit}}_{\mathrm{RVM}}(x) = L + O_\varepsilon(x^{1/2+\varepsilon}) \tag{1.8}$$

(Theorem 8.12). The RVM Mertens function therefore exists as a continuous spectral inverse although no coefficient sequence for F_{RVM} exists. Since the primitive of the midpoint sawtooth is bounded, the same decomposition without contours gives $C^{-1} \le |\zeta_{\mathrm{RVM}}| \le C$ on every half-plane $\Re s \ge \sigma_0 > 1$ and $\zeta_{\mathrm{RVM}}(s) \to \varpi_{\mathrm{RVM}}$ as $\Re s \to +\infty$ (Theorem 8.10).

Ternary discretization. An integer-valued representative of $M^{\mathrm{crit}}_{\mathrm{RVM}}$ is obtained by sampling a smooth target $\mathcal{M}(n) \approx M^{\mathrm{crit}}_{\mathrm{RVM}}(n)$ and choosing signs on the squarefree support:

$$\mu_h(n) = \mu(n)^2 \varsigma_n, \qquad \varsigma_n \in \{-1, +1\}, \qquad M_h(n) := \sum_{m \le n} \mu_h(m), \tag{1.9}$$

$$\varsigma_n = \begin{cases} +1, & M_h(n-1) < \mathcal{M}(n), \\ -1, & M_h(n-1) \ge \mathcal{M}(n), \end{cases} \qquad (\mu(n)^2 = 1). \tag{1.10}$$

This is a greedy sigma–delta feedback quantization; feedback quantization and noise shaping are standard themes of sigma–delta theory [11, 21, 3]. Here the target is a zero-generated Mertens inverse and the admissible times are the squarefree integers. The walk μ_h is a comparison object: it is not claimed to be multiplicative or equal to μ. The two discretizations are distinct: the cell masses c_n^{RVM} come from π_{RVM}, the signs ς_n track $M^{\mathrm{crit}}_{\mathrm{RVM}}$.

Comparison with ζ. At the level of transforms the defect is the exact quotient $\mathcal{Q}_{\mathrm{RVM}} = \zeta/\zeta_{\mathrm{RVM}}$ of (9.6), with $F_{\mathrm{RVM}} = \mathcal{Q}_{\mathrm{RVM}} \cdot (1/\zeta)$. Since $|\zeta_{\mathrm{RVM}}|$ is bounded above and below on $\frac{1}{2} + \varepsilon \le \sigma \le c'$, $|t| \ge t_0$ (Corollary 9.3), every bound for ζ or $1/\zeta$ to the right of $\frac{1}{2}$ is equivalent to the same bound for $\mathcal{Q}_{\mathrm{RVM}}$. For M itself the bound $M(x) = O_\varepsilon(x^{1/2+\varepsilon})$ is equivalent to the Riemann hypothesis [37, §14.25], and the disproof of the Mertens conjecture shows that the ε cannot simply be dropped [30].

1.3. **The core of the method.** The transfer from ζ_{RVM} to ζ is a transformation of ζ_{RVM} depending on a parameter $\lambda \in [0, 1]$. The ordinates move along

$$\gamma_k(\lambda) = (1-\lambda)\gamma_k + \lambda\tau_k, \qquad \zeta_\lambda(s) = \frac{1}{G(s)} \prod_{k \ge 1} \left(1 + \frac{s(s-1)}{\frac{1}{4} + \gamma_k(\lambda)^2}\right), \tag{1.11}$$

where γ_k are the ordinates of the nontrivial zeros of ζ; thus $\zeta_1 = \zeta_{\mathrm{RVM}}$ and $\zeta_0 = \zeta_{Z1}$ (Section 10).

(i) Every member ζ_λ has its nontrivial zeros exactly at $\frac{1}{2} \pm i\gamma_k(\lambda)$. The real part $\frac{1}{2}$ is part of the definition of the family, like the gamma factor, and never changes along it: it is impossible for ζ_{Z1}, or for any ζ_λ, to have a zero off the critical line other than the trivial zeros. For $0 < \lambda \le 1$ all zeros are simple.

(ii) All differences between the moving zetas lie in the numbers $\gamma_k - \tau_k$, which are bounded (Lemma 2.2); the quotient of two members is an explicit product in γ_k and $\lambda(\tau_k - \gamma_k)$ (Remark 10.2).

(iii) The final step is the identity $\zeta_{Z1} = \zeta$. It is the single and complete test: while $\zeta_{Z1} = \zeta$ there is nothing outside $\Re s = \frac{1}{2}$ to speak of, since the identity itself places every nontrivial zero of ζ at $\frac{1}{2} \pm i\gamma_k$, and $\zeta_{Z1} \ne \zeta$ is the only possible obstruction.

The endpoint has the functional equation, the pole, the residue and the order of ζ (Proposition 11.1). Hamburger's theorem belongs to the family of converse theorems that recover Dirichlet series from analytic continuation, functional equation and growth [23, 32]; applied to ζ_{Z1} it gives (Theorem 11.2)

$$\zeta_{Z1} = \zeta \quad \Longleftrightarrow \quad \text{(D): } \zeta_{Z1} \text{ is an absolutely convergent Dirichlet series in } \Re s > 1. \tag{1.12}$$

With the integer-weighted zeta $\zeta_{\mathbb{Z},\mathrm{RVM}}$, which is a Dirichlet series, (D) is equivalent to an absolutely convergent Dirichlet expansion of the explicit multiplier $\mathcal{K} = \zeta_{Z1}/\zeta_{\mathbb{Z},\mathrm{RVM}} = \Pi_{\mathrm{tr}} e^{-h_{\mathrm{RVM}}}$ (Theorem 11.3); since $1/\zeta_{\mathbb{Z},\mathrm{RVM}}$ is also an absolutely convergent Dirichlet series in $\Re s > 1$, the equivalence holds on the whole half-plane. Section 12 gives two sharper forms of (D). First, the

clock decomposition applied to the ordinates γ_k gives the value of the endpoint at infinity,

$$\varpi_{Z1} = \lim_{\sigma\to+\infty} \zeta_{Z1}(\sigma) = \varpi_{\mathrm{RVM}} \prod_{k\ge1} \frac{\frac14+\tau_k^2}{\frac14+\gamma_k^2} \ \ge 1, \tag{1.13}$$

and $\zeta_{Z1} = \zeta$ holds if and only if $\varpi_{Z1} = 1$ (Theorems 12.1 and 12.2, Corollary 12.3): the final step is the single identity $\prod_k(\frac14+\gamma_k^2)/(\frac14+\tau_k^2) = \varpi_{\mathrm{RVM}}$ between the displacements and the transition constant. Second, the prime-power counts $J^{\mathrm{M}}_{\mathrm{RVM}}$ and J^{M}_{Z1} are the Mellin transforms of $\log\zeta_{\mathrm{RVM}}$ and $\log\zeta_{Z1}$ and satisfy a distributional explicit formula (Theorems 12.5 and 12.6); this proves the Mellin–Stieltjes identification of the integer weights (Theorem 6.4), and (D) holds if and only if J^{M}_{Z1} is constant between consecutive integers (Theorem 12.7). Third, along the family $\frac{d}{d\lambda}\log\varpi_\lambda = -\Upsilon(\lambda)$, the generator at infinity, and (D) is the identity $\int_{0^+}^1 \Upsilon(\lambda)\,d\lambda = \log L$ (Proposition 13.2). Section 13 atomizes every member: $\tilde\zeta_{\mathbb{Z},\lambda} = \zeta_\lambda e^{\tilde h_\lambda}$ keeps the Euler product and the Dirichlet series while ζ_λ keeps the functional equation; the joining factor $\tilde h_\lambda$ is written cell by cell and is holomorphic in $\Re s > \frac12$ (Theorems 13.6 and 13.12); the cell masses have mean square $X^{1+\varepsilon}$ on $(X, 2X]$ uniformly in λ, so every $\tilde\zeta_{\mathbb{Z},\lambda}$, the endpoint included, is an absolutely convergent Dirichlet series in $\Re s > 1$ (Theorem 13.10, Corollary 13.11); and (D) is equivalent to the vanishing of every cell defect at $\lambda = 0$ (Theorem 13.13). At the clock endpoint the joining factor $\tilde h_1$ is not constant, and the endpoint Mertens function has a Mellin form B_{Z1} with $B_{Z1} = M$ exactly when (D) holds (Propositions 13.15 and 13.16). The remaining steps form the programme of Subsection 13.8. In RVM language, the final step asks whether the zero-generated analytic object ζ_{Z1} belongs to the discrete Dirichlet category; if it does, Hamburger uniqueness makes it ζ, and the prime and Mertens coordinates become those of the integers. Step 3 is Theorem 11.8:

$$\zeta_{Z1} = \zeta \quad\Longrightarrow\quad M(x) = O_\varepsilon(x^{1/2+\varepsilon}) \qquad (\varepsilon > 0), \tag{1.14}$$

with a self-contained proof by the Borel–Carathéodory lemma, Hadamard's three-circle theorem and the truncated Perron formula. Theorem 11.9 gives the same conclusion as a transfer from the synthetic side: under $\zeta_{Z1} = \zeta$, $1/\zeta = F_{\mathrm{RVM}}\,\Pi_{\mathrm{tr}}^{-1}$ with F_{RVM} bounded and $\Pi_{\mathrm{tr}}^{\pm1} \ll |t|^\varepsilon$ to the right of $\frac12$, so the bound $M^{\mathrm{crit}}_{\mathrm{RVM}}(x) = L + O_\varepsilon(x^{1/2+\varepsilon})$ carries over to $M(x) = O_\varepsilon(x^{1/2+\varepsilon})$. The identity $\zeta_{Z1} = \zeta$ itself, equivalently (D), is not proved in this paper; it is the open step of the method (Remark 11.12).

The relations among the principal objects are

$$\{\tau_k\} \longrightarrow \zeta_{\mathrm{RVM}} \longrightarrow J_{\mathrm{RVM}} \longleftrightarrow \pi_{\mathrm{RVM}} \longrightarrow (d\pi_{\mathrm{RVM}},\, p_j^{\mathrm{RVM}},\, c_n^{\mathrm{RVM}}), \tag{1.15}$$

$$F_{\mathrm{RVM}} = \zeta_{\mathrm{RVM}}^{-1} \longrightarrow M^{\mathrm{crit}}_{\mathrm{RVM}} \longrightarrow (\mathcal{M}, \mu_h, M_h), \tag{1.16}$$

$$\zeta_{\mathrm{RVM}} = \zeta_1 \xrightarrow{\lambda\downarrow0} \zeta_0 = \zeta_{Z1} \overset{?}{=} \zeta \implies M(x) = O_\varepsilon(x^{1/2+\varepsilon}). \tag{1.17}$$

The first chain is the prime coordinate, the second the Mertens coordinate, the third the λ-family; they share the midpoint spectrum and solve different realization problems.

1.4. **Originality, scope and status.** The ingredients are standard or belong to neighbouring theories: the Riemann–von Mangoldt law, Lambert inversion, Hadamard products, explicit formulas, Möbius and Perron inversion, Beurling generalized primes, sigma–delta feedback, and Hamburger's converse theorem. The contribution is their assembly into one midpoint RVM space with four coordinated levels: (i) an exact midpoint zero clock and its canonical zeta-like product; (ii) a continuous prime-power and prime coordinate, with real prime levels and non-integral cell masses; (iii) a continuous synthetic Mertens inverse with its ternary feedback discretization; (iv) the λ-family from the midpoint spectrum to the ordinates of ζ, ending in the single identity $\zeta_{Z1} = \zeta$ and the Mertens bound that follows from it. To the author's knowledge the cited literature contains the individual mechanisms but not this chain. Standard material is cited at the point of use; statements without citation are original.

Statement	Status
Midpoint clock, Lambert inversion, bounded displacement	theorem
Canonical product $P^{\mathrm{crit}}_{\mathrm{RVM}}$ and ζ_{RVM}	theorem
$S_{\mathrm{RVM}}(N;u) \ll 1+\log N$; centroid and M_H^{RVM} decay	theorem
Compact-uniform existence of J_{RVM}; finite inversion	theorem
Real prime levels p_j^{RVM}	defined on monotone branches of π_{RVM}
Square-root estimate for $J_{\mathrm{RVM}}, \pi_{\mathrm{RVM}}$	theorem
$\mathrm{li}/\pi_{\mathrm{RVM}} \to 1$; $\pi/\pi_{\mathrm{RVM}} \to 1$	theorem (with the prime number theorem)
J_{RVM} Lipschitz, $c_n^{\mathrm{RVM}} = O(1)$	theorem
$\zeta_{\mathbb{Z},\mathrm{RVM}}$ absolutely convergent Dirichlet series in $\Re s > 1$	theorem
Mellin–Stieltjes identification; $\zeta_{\mathbb{Z},\mathrm{RVM}} = \zeta_{\mathrm{RVM}} e^{h_{\mathrm{RVM}}}$	theorem
$\lvert\zeta\rvert \asymp \lvert\mathcal{Q}_{\mathrm{RVM}}\rvert$ for $\sigma \geq \frac{1}{2}+\varepsilon$, $\lvert t\rvert \geq t_0$	theorem
Transition theorem $\varpi_{\mathrm{RVM}}(1+w)$, $\varpi_{\mathrm{RVM}}(1+\chi)$	theorem
$C^{-1} \leq \lvert\zeta_{\mathrm{RVM}}\rvert \leq C$ on $\Re s \geq \sigma_0 > 1$; $\zeta_{\mathrm{RVM}} \to \varpi_{\mathrm{RVM}}$ at $+\infty$	theorem
Unsmoothed $M^{\mathrm{crit}}_{\mathrm{RVM}} = L + O_\varepsilon(x^{1/2+\varepsilon})$	theorem
Dirichlet expansion of $\zeta_{\mathrm{RVM}}^{\pm 1}$	impossible (theorem)
Ternary walk μ_h, M_h	definition; comparison object
Exact Hadamard product of 2ξ	theorem
λ-family, generator, multiplier, simplicity for $\lambda > 0$	theorem
Analytic class of ζ_{Z1}	theorem
$\zeta_{Z1} = \zeta \Leftrightarrow$ (D) $\Leftrightarrow$ RH	theorem
(D) $\Leftrightarrow$ Dirichlet expansion of $\mathcal{K} = \zeta_{Z1}/\zeta_{\mathbb{Z},\mathrm{RVM}}$ in $\Re s > 1$	theorem
$\varpi_{Z1} = \varpi_{\mathrm{RVM}} \prod_k \frac{1/4+\tau_k^2}{1/4+\gamma_k^2}$; (D) $\Leftrightarrow \varpi_{Z1} = 1$	theorem
Mellin prime-power counts; distributional explicit formula	theorem
(D) $\Leftrightarrow$ J^{M}_{Z1} constant between integers	theorem
$\frac{d}{d\lambda}\log\varpi_\lambda = -\Upsilon$; (D) $\Leftrightarrow \int_{0^+}^1 \Upsilon = \log L$	theorem
Atomized family; joining factor holomorphic in $\Re s > \frac{1}{2}$	theorem
Atomized family: Dirichlet series in $\Re s > 1$ for every λ	theorem
(D) $\Leftrightarrow$ vanishing cell defects at $\lambda = 0$	theorem
ζ_λ not a Dirichlet series in $\Re s > 1$ for $0 < \lambda \leq 1$	theorem
$\lambda \mapsto \log\zeta_\lambda(\sigma)$ strictly convex for real $\sigma > 1$	theorem
Hamburger's theorem for the pair $(\zeta_\lambda, \tilde{\zeta}_{\mathbb{Z},\lambda})$ with symmetric joining factor	theorem
(D) $\Leftrightarrow$ ζ_{Z1} has a positive representing measure; then $\varpi_{Z1} \leq 1$	theorem
Density of $d\pi_{\mathrm{RVM}}$; cell masses and defects at $\lambda = 1$	theorem
Height kernel of the cell masses; Gaussian defect of the atomization	theorem
$\zeta_{Z1}(\sigma) \geq \zeta(\sigma)$ for real $\sigma > 1$; (D) $\Leftrightarrow$ equality at one σ	theorem
Hamburger's theorem for the pair in theta form and in vertical mean-square form	theorem
Symmetry of the joining factor fails at $\lambda = 1$; $\tilde{\zeta}_{\mathbb{Z},\lambda} = \zeta$ for finitely many λ at most	theorem
Summation formula for every ζ_λ; $\varpi_\lambda - 1$ as one number $\mathsf{W}(\lambda)$	theorem
(D) $\Leftrightarrow$ $\varpi_\lambda < 1$ on the open path $0 < \lambda \leq 1$	theorem
Space $\mathscr{O}$ of ordinate lists: rigidity of the coincidence; (D) $\Leftrightarrow$ existence of a coincidence list	theorem

Statement	Status
(D) $\Leftrightarrow \tilde{c}_n(0) = \mathbf{1}_{\mathrm{prime}}(n)$ for every n; every point of E is a witness of (D)	theorem
Symmetric joining factors: order one is needed; the symmetry criterion has no slack at $\lambda = 0$	theorem
Ladder: $\tilde{h}_\lambda(\sigma) = O(\sigma N^{-\sigma}) \Leftrightarrow$ no mass of dJ^{M}_λ outside the integers below N	theorem
Bridge theorem in phase form; (D) $\Leftrightarrow$ (B1), (B2) for $\tilde{h}_0$; impossible for $\lambda > 0$	theorem
Phase of ζ_λ on the critical line; cell masses as sums of a one-ordinate function	theorem
Transport $\mathsf{S}_{\mathsf{f}}(\lambda)$: entire of exponential type; moment series on the clock; defect of the clock by Poisson summation	theorem
Multiscale clock Jacobian: lower frame bound on $U \asymp \log T$, no fixed-window bound, dense range, smooth tails	theorem; quantitative nonlinear continuation open
(D) $\Leftrightarrow$ solvability of the rung system with free masses; the solution is nondegenerate	theorem
Gradient of V_σ on $\mathscr{O}$	theorem
Acceptance criterion (window and frequencies); two-edge model; finite Euler products with the final step proved	theorem
No object of finite rank in the window of ζ; the three abscissas of the diagonal operator over the primes	theorem
Phase defect along the path (rotation of the one-ordinate terms)	heuristic
$\zeta_{Z1} = \zeta \Rightarrow M(x) = O_\varepsilon(x^{1/2+\varepsilon})$	theorem (Step 3)
$\zeta_{Z1} = \zeta$	open

What is proved and what is not. Three groups of statements are proved in full. *(1) The synthetic system.* The clock and its Lambert inversion, the canonical product and ζ_{RVM}, the logarithmic bound for the exponential sums, the prime coordinate J_{RVM}, π_{RVM} with its square-root estimate and its density (Theorem 14.15), the transition theorem, the impossibility of a Dirichlet expansion of $\zeta^{\pm 1}_{\mathrm{RVM}}$ and the synthetic Mertens bound. *(2) The family and its endpoint.* Every ζ_λ has the functional equation, order one and all its nontrivial zeros on $\Re s = \frac{1}{2}$; $\zeta_{Z1} = \zeta$ is equivalent to (D), to $\varpi_{Z1} = 1$ and to the constancy of J^{M}_{Z1} between consecutive integers; $\varpi_{Z1} \geq 1$ and $\zeta_{Z1}(\sigma) \geq \zeta(\sigma)$ for real $\sigma > 1$; $\log \zeta_\lambda(\sigma)$ is strictly convex in λ; and $\zeta_{Z1} = \zeta$ implies $M(x) = O_\varepsilon(x^{1/2+\varepsilon})$ (Step 3). *(3) The atomized family and the criteria.* Every $\tilde{\zeta}_{\mathbb{Z},\lambda}$ is an absolutely convergent Dirichlet series in $\Re s > 1$, zero-free in $\Re s > \frac{1}{2}$, joined to ζ_λ by an explicit factor; no ζ_λ with $\lambda > 0$ is a Dirichlet series; Hamburger's theorem holds for the pair in three forms (Theorems 14.3, 14.4); the positivity criterion (Theorem 14.7); and the rigidity of the coincidence in the space of ordinate lists (Theorem 15.1). *(4) Criteria written where the joining factor is defined, and models.* The reduction of (D) to the cell masses (Corollary 14.26); the ladder and the half-line form of Hamburger's theorem for a pair (Theorems 14.33, 14.34); the bridge theorem in phase form (Theorem 14.36, Corollary 14.37); the transport from the clock and the exact defect of the clock (Theorem 14.42, Proposition 14.43); the existence form with free masses and the nondegeneracy of the solution (Theorems 15.5, 15.6); the acceptance criterion and the models in which the final step is a theorem (Theorems 16.1, 16.3, Proposition 16.2).

Not proved: the identity $\zeta_{Z1} = \zeta$. Each of the following statements is proved to be equivalent to it, and none of them is proved: (D); $\varpi_{Z1} \le 1$; $\zeta_{Z1}(\sigma) \le \zeta(\sigma)$ for one $\sigma > 1$; $\varpi_\lambda < 1$ for all $\lambda \in (0,1]$; J^{M}_{Z1} non-decreasing; $\mathsf{W}(0) \ge 0$ (14.23); $\tilde{h}_0$ constant; the theta relation $T_0 \equiv 0$ (14.6); $\mathsf{V}_\sigma(0) = 0$ (14.7); the existence of a list in $\mathscr{O}$ whose continuous and atomized zeta functions coincide (Corollary 15.3); $\tilde{c}_n(0) = \mathbf{1}_{\mathrm{prime}}(n)$ for every $n \ge 2$ (Corollary 14.26); $\tilde{h}_0(\sigma) = O_N(\sigma N^{-\sigma})$ for every N (14.66); the conditions (B1) and (B2) for $\tilde{h}_0$ (Corollary 14.37); $\mathsf{S}_{\mathsf{f}}(0) = \mathsf{R}_{\mathsf{f}}(1)$ for every f whose support avoids the points $\log n$ (Proposition 14.41); the solvability of the rung system (15.5) in $\mathscr{O}$ (Theorem 15.5). The symmetry of the joining factor $e^{\tilde{h}_0}$ with finite order is a sufficient condition (Theorem 14.3), and it is also necessary (Remark 14.30).

1.5. **Organization.** Section 2 treats the clock, its Lambert inversion and the bounded displacement; Section 3 the canonical product; Section 4 the exponential sums; Sections 5 and 6 the prime coordinate and the integer weights; Sections 7 and 8 the Mertens transform and the transition theorem; Section 9 the exact Hadamard product and the comparison with ζ; Section 10 the λ-family; Section 11 the endpoint, Hamburger's theorem and Step 3; Section 12 the value of the endpoint at infinity and the Euler products of ζ_{RVM} and ζ_{Z1}; Section 13 the atomized family, Hamburger's hypotheses at the two endpoints and the programme for the final step; Section 14 the open path, convexity, Hamburger's theorem for the pair, the positivity criterion, the density of the clock, the Gaussian defect, the reduction to the cell masses, the half-line form, the bridge theorem and the transport from the clock; Section 15 the space of ordinate lists, the rigidity of the coincidence, the existence form of the final step, the rung system with free masses and the gradient of the vertical defect; Section 16 the acceptance criterion and the models in which the final step is a theorem; Section 17 the numerical illustration; Section 18 the conclusions and open problems.

2. The midpoint Riemann–von Mangoldt clock

Let ρ run over the nontrivial zeros of ζ, counted with multiplicity. No nontrivial zero is real, so the positive ordinates can be listed with multiplicity as $0 < \gamma_1 \le \gamma_2 \le \cdots$. Put

$$N_\zeta(T) := \#\{\rho : \ 0 < \Im\rho \le T\} = \#\{k : \gamma_k \le T\}. \tag{2.1}$$

The Riemann–von Mangoldt formula [37, Thms. 9.3–9.4], [38] states that for $T \ge 2$, $T \ne \gamma_k$,

$$N_\zeta(T) = \frac{T}{2\pi}\log\frac{T}{2\pi e} + \frac{7}{8} + S(T) + O\Big(\frac{1}{T}\Big), \qquad S(T) := \frac{1}{\pi}\arg\zeta\big(\tfrac{1}{2}+iT\big) = O(\log T), \tag{2.2}$$

where $\arg\zeta(\frac{1}{2}+iT)$ is obtained by continuous variation of the argument along the segments from 2 to $2+iT$ and from $2+iT$ to $\frac{1}{2}+iT$, starting from $\arg\zeta(2) = 0$. The $O(1/T)$ term comes from Stirling's formula for Γ and is explicit; $S(T)$ is the fluctuating part, with a jump $+m$ at an ordinate of multiplicity m. The *midpoint RVM clock* is the smooth part of (2.2):

$$\mathcal{N}_{\mathrm{RVM}}(T) := \frac{T}{2\pi}\log\frac{T}{2\pi e} + \frac{7}{8}. \tag{2.3}$$

It is strictly increasing on $[2\pi, \infty)$. The midpoint ordinates τ_k are the positive solutions of

$$\mathcal{N}_{\mathrm{RVM}}(\tau_k) = k - \frac{1}{2}, \qquad k \ge 1, \tag{2.4}$$

that is, the abscissae where the smooth count crosses the midpoints $k - \frac{1}{2}$ of the steps of a unit staircase. Thus $\mathcal{N}_{\mathrm{RVM}}$ replaces the staircase N_ζ by its smooth part, and τ_k is the point where the kth step would be crossed half-way (Figure 1).

The Lambert W-function is the multivalued inverse of $v \mapsto ve^v$: $W(z)e^{W(z)} = z$. On $[-1/e, \infty)$ the equation $ve^v = z$ has exactly one solution $v \ge -1$; this defines the principal branch $W_0 : [-1/e, \infty) \to [-1, \infty)$, which is real, increasing, with $W_0(0) = 0$ and $W_0(z) \sim \log z - \log\log z$ as $z \to \infty$ [17, §4.13], [10].

Proposition 2.1 (Lambert inversion)**.** *For every $k \geq 1$,*

$$\tau_k = \frac{2\pi(k - 11/8)}{W_0\big((k - 11/8)/e\big)}, \tag{2.5}$$

The quotient is positive also for $k = 1$.

Proof. Put $y = \tau/(2\pi)$. Equation (2.4) reads

$$y(\log y - 1) = k - \frac{11}{8}. \tag{2.6}$$

With $v = \log y - 1$ one has $y = e^{v+1}$ and $ve^v = (k - 11/8)/e$, so $v = W_0((k - 11/8)/e)$ and $y = (k - 11/8)/W_0((k - 11/8)/e)$. For $k = 1$, $(k - 11/8)/e = -3/(8e) \in (-1/e, 0)$, where W_0 is real and negative; numerator and denominator are both negative. □

The clock derivative and the resulting growth are

$$\mathcal{N}'_{\mathrm{RVM}}(T) = \frac{1}{2\pi} \log \frac{T}{2\pi}, \tag{2.7}$$

$$\tau_k \asymp \frac{k}{\log k} \qquad (k \geq 2). \tag{2.8}$$

In terms of the clock, (2.2) reads

$$N_\zeta(T) = \mathcal{N}_{\mathrm{RVM}}(T) + S(T) + O(T^{-1}), \qquad S(T) = O(\log T) \qquad (T \geq 2). \tag{2.9}$$

Let

$$N^*_\zeta(T) := \tfrac{1}{2}\big(N_\zeta(T + 0) + N_\zeta(T - 0)\big) \tag{2.10}$$

be the symmetric count, and define the displacements

$$\eta_k := \tau_k - \gamma_k. \tag{2.11}$$

Lemma 2.2 (Bounded displacement)**.** *There is a constant C_η such that*

$$|\eta_k| \leq C_\eta \qquad (k \geq 1). \tag{2.12}$$

Proof. Let $\gamma = \gamma_k$ and let $m = m(\gamma)$ be the number of indices j with $\gamma_j = \gamma$; these indices form a block $j_0 + 1, \ldots, j_0 + m$ containing k. Then $N_\zeta(\gamma - 0) = j_0$ and $N_\zeta(\gamma + 0) = j_0 + m$, so $N^*_\zeta(\gamma) = j_0 + m/2$ and

$$\big|N^*_\zeta(\gamma_k) - (k - \tfrac{1}{2})\big| \leq \frac{m(\gamma_k)}{2}. \tag{2.13}$$

By (2.9), $m(\gamma) \leq N_\zeta(\gamma + 1) - N_\zeta(\gamma - 1) = O(\log \gamma)$, and $N^*_\zeta(\gamma) = \mathcal{N}_{\mathrm{RVM}}(\gamma) + O(\log \gamma)$. With (2.4),

$$\big|\mathcal{N}_{\mathrm{RVM}}(\gamma_k) - \mathcal{N}_{\mathrm{RVM}}(\tau_k)\big| = O(\log \gamma_k). \tag{2.14}$$

Both γ_k and τ_k tend to infinity and (2.14) gives $\log \gamma_k \asymp \log \tau_k \asymp \log k$. By the mean-value theorem and (2.7), $\mathcal{N}'_{\mathrm{RVM}}$ is $\gg \log k$ between γ_k and τ_k for large k, so $|\gamma_k - \tau_k| = O(1)$. Finitely many initial indices are absorbed into C_η. □

3. The pure critical RVM zeta function

Put

$$q(s) := s(s - 1), \tag{3.1}$$

$$P^{\mathrm{crit}}_{\mathrm{RVM}}(s) := \prod_{k \geq 1} \left(1 + \frac{q(s)}{\frac{1}{4} + \tau_k^2}\right), \tag{3.2}$$

$$G(s) := s(s - 1)\Gamma(s/2)\pi^{-s/2}, \tag{3.3}$$

and

$$\xi_{\mathrm{RVM}}(s) := \frac{1}{2} P^{\mathrm{crit}}_{\mathrm{RVM}}(s), \qquad \zeta_{\mathrm{RVM}}(s) := \frac{P^{\mathrm{crit}}_{\mathrm{RVM}}(s)}{G(s)}. \tag{3.4}$$

Theorem 3.1 (Canonical characterization of ζ_{RVM})**.** *The product* (3.2) *converges absolutely and locally uniformly in $s \in \mathbb{C}$. It defines an entire function with*

$$P^{\mathrm{crit}}_{\mathrm{RVM}}(1-s) = P^{\mathrm{crit}}_{\mathrm{RVM}}(s), \qquad P^{\mathrm{crit}}_{\mathrm{RVM}}(\bar{s}) = \overline{P^{\mathrm{crit}}_{\mathrm{RVM}}(s)}, \qquad P^{\mathrm{crit}}_{\mathrm{RVM}}(0) = P^{\mathrm{crit}}_{\mathrm{RVM}}(1) = 1. \tag{3.5}$$

Its zeros are exactly the simple zeros

$$\rho_k^{\mathrm{RVM}} = \frac{1}{2} + i\tau_k, \qquad \overline{\rho_k^{\mathrm{RVM}}} = \frac{1}{2} - i\tau_k, \qquad k \geq 1. \tag{3.6}$$

Consequently ζ_{RVM} has a simple pole at $s = 1$ with residue 1, the trivial zeros $s = -2, -4, -6, \ldots$, and nontrivial zeros exactly at (3.6)*; moreover $\xi_{\mathrm{RVM}}(s) = \xi_{\mathrm{RVM}}(1-s)$.*

Proof. By (2.8), $\sum_k \tau_k^{-2} < \infty$. On a compact set K,

$$\sup_{s \in K} \left| \frac{q(s)}{\frac{1}{4} + \tau_k^2} \right| \ll_K \tau_k^{-2}, \tag{3.7}$$

so the Weierstrass criterion gives normal convergence. Each factor depends on s only through $q(s)$ and equals 1 at $s = 0, 1$, which gives (3.5); real coefficients give the conjugation symmetry. The kth factor vanishes exactly when

$$\left(s - \frac{1}{2}\right)^2 + \tau_k^2 = 0. \tag{3.8}$$

The clock is strictly increasing on $[2\pi, \infty)$ and $\tau_1 > 2\pi$, so the τ_k are distinct; the derivative $2s - 1 = \pm 2i\tau_k \neq 0$ of the vanishing factor and the non-vanishing of the other factors make the zeros simple. At $s = 1$, $P^{\mathrm{crit}}_{\mathrm{RVM}}(1) = 1$ and $G(s) = (s-1)(1 + O(s-1))$, so the residue is 1. At $s = -2n$ ($n \geq 1$) the pole of $\Gamma(s/2)$ gives a zero of $1/G$, while $P^{\mathrm{crit}}_{\mathrm{RVM}}(-2n) \neq 0$ because $q(-2n) > 0$. At $s = 0$, $s\Gamma(s/2) \to 2$, so $G(0) \neq 0$ and $s = 0$ is not a zero. □

4. RVM exponential sums, centroids, and the height transform

For $u \in \mathbb{R}$ and $N \geq 1$ define

$$S_{\mathrm{RVM}}(N; u) := \sum_{k \leq N} e^{iu\tau_k}, \qquad C^{\mathrm{pts}}_{\mathrm{RVM}}(N; u) := \frac{S_{\mathrm{RVM}}(N; u)}{N}, \tag{4.1}$$

and

$$C^{\mathrm{path}}_{\mathrm{RVM}}(N; u) := \frac{1}{N} \sum_{n \leq N} S_{\mathrm{RVM}}(n; u) = \sum_{k \leq N} \frac{N - k + 1}{N} e^{iu\tau_k}. \tag{4.2}$$

Theorem 4.1 (Uniform logarithmic exponential-sum bound)**.** *Let $0 < u_1 \leq u_2 < \infty$. Then*

$$\sup_{u \in [u_1, u_2]} |S_{\mathrm{RVM}}(N; u)| \ll_{u_1, u_2} 1 + \log N. \tag{4.3}$$

Proof. Let $\mathcal{T} : [-\frac{1}{8}, \infty) \to [2\pi, \infty)$ be the inverse of the clock, so $\tau_k = \mathcal{T}(k - \frac{1}{2})$, and put $\ell_k = \log(\tau_k / 2\pi) > 0$. Differentiating $\mathcal{N}_{\mathrm{RVM}}(\mathcal{T}(y)) = y$,

$$\mathcal{T}'(y) = \frac{2\pi}{\log(\mathcal{T}(y)/2\pi)}, \qquad \mathcal{T}''(y) = -\frac{4\pi^2}{\mathcal{T}(y) \log^3(\mathcal{T}(y)/2\pi)}. \tag{4.4}$$

Since $\mathcal{T}'$ decreases and $|\mathcal{T}''|$ decreases,

$$\frac{2\pi}{\ell_{k+1}} \leq \tau_{k+1} - \tau_k \leq \frac{2\pi}{\ell_k}, \tag{4.5}$$

$$|\tau_{k+2} - 2\tau_{k+1} + \tau_k| \leq \frac{4\pi^2}{\tau_k \ell_k^3}. \tag{4.6}$$

From (2.4), $\tau_k \ell_k = 2\pi(k - \frac{11}{8}) + \tau_k$, hence

$$\tau_k \ell_k \asymp k \qquad (k \geq 2). \tag{4.7}$$

Fix $u \in [u_1, u_2]$ and put $\Delta_k = u(\tau_{k+1} - \tau_k)$, $E_k = e^{iu\tau_k}$. Choose $k_0 = k_0(u_2)$ with $\ell_{k_0} \geq 2u_2$; then $0 < \Delta_k \leq \pi$ for $k \geq k_0$. With $b_k := (e^{i\Delta_k} - 1)^{-1}$ one has $E_{k+1} = e^{i\Delta_k} E_k$, hence

$$E_k = b_k(E_{k+1} - E_k). \tag{4.8}$$

Summing (4.8) over $k_0 \leq k \leq N$ and shifting the index in the first part gives the exact identity

$$\sum_{k=k_0}^{N} E_k = b_N E_{N+1} - b_{k_0} E_{k_0} + \sum_{k=k_0+1}^{N} (b_{k-1} - b_k) E_k. \tag{4.9}$$

For $0 < x \leq \pi$, $|e^{ix} - 1| = 2\sin(x/2) \geq 2x/\pi$. Using (4.5) and (4.6),

$$|b_k| \leq \frac{\pi}{2\Delta_k} \leq \frac{\ell_{k+1}}{4u_1}, \qquad |b_{k-1} - b_k| \leq |\Delta_k - \Delta_{k-1}|\,|b_{k-1}||b_k| \leq \frac{\pi^2 u_2}{4u_1^2} \frac{\ell_{k+1}^2}{\tau_{k-1}\ell_{k-1}^3} \ll_{u_1,u_2} \frac{1}{\tau_k \ell_k} \ll \frac{1}{k}, \tag{4.10}$$

the last step by (4.7). Inserting (4.10) into (4.9) gives $|\sum_{k_0}^{N} E_k| \ll \ell_{N+1} + \ell_{k_0+1} + \sum_{k \leq N} k^{-1} \ll 1 + \log N$. The indices $k < k_0$ contribute at most k_0. All constants depend only on u_1, u_2. □

Corollary 4.2 (Point centroid)**.** *Uniformly for $u \in [u_1, u_2] \subset (0, \infty)$,*

$$C_{\mathrm{RVM}}^{\mathrm{pts}}(N; u) = O_{u_1,u_2}\Big(\frac{1 + \log N}{N}\Big) \longrightarrow 0. \tag{4.11}$$

Theorem 4.1 alone does not imply convergence of the path centroid (4.2)*.*

Proof. The first assertion is (4.3) divided by N. For the second, the identity $C_{\mathrm{RVM}}^{\mathrm{path}} = N^{-1} \sum_{n \leq N} S_{\mathrm{RVM}}(n; u)$ with $S_{\mathrm{RVM}}(n; u) = O(\log n)$ gives only $O(\log N)$. □

Define the midpoint count

$$n_{\mathrm{RVM}}(H) := \#\{k : \tau_k \leq H\}. \tag{4.12}$$

For $x > 0$ and $n_{\mathrm{RVM}}(H) \geq 1$ define

$$M_H^{\mathrm{RVM}}(x) := \frac{1}{n_{\mathrm{RVM}}(H)} \sum_{\tau_k \leq H} x^{i\tau_k}, \tag{4.13}$$

and for the ordinates of ζ

$$M_H(x) := \frac{1}{N_\zeta(H)} \sum_{\gamma_k \leq H} x^{i\gamma_k}. \tag{4.14}$$

Corollary 4.3 (Synthetic height-transform decay)**.** *For every fixed $x > 1$, with $N = n_{\mathrm{RVM}}(H)$,*

$$M_H^{\mathrm{RVM}}(x) = O_x\Big(\frac{1 + \log N}{N}\Big) \longrightarrow 0 \qquad (H \to \infty). \tag{4.15}$$

Proof. Take $u = \log x > 0$ in Theorem 4.1; (4.13) is the point centroid at $N = n_{\mathrm{RVM}}(H)$. □

Remark 4.4. Neither (4.13) nor (4.14) is the Mertens function; both are normalized phase averages.

5. The RVM explicit formula, J_{RVM}, and π_{RVM}

For $w = a + ib$ with $b > 0$ define

$$\mathrm{Ei}(w) := \int_{-\infty}^{a} \frac{e^{v+ib}}{v + ib}\, dv, \qquad \mathrm{Ei}(\bar{w}) := \overline{\mathrm{Ei}(w)}. \tag{5.1}$$

The path is the horizontal line $\Im = b$, which avoids 0; this is the branch of $\mathrm{li}(x^\rho) = \mathrm{Ei}(\rho \log x)$ used in Riemann's explicit formula [18]. For $x > 1$ put

$$J_{\mathrm{RVM}}(x) := \mathrm{li}(x) - 2\Re \sum_{k \geq 1} \mathrm{Ei}\Big(\Big(\frac{1}{2} + i\tau_k\Big) \log x\Big) - \log 2 + \int_x^\infty \frac{dy}{y(y^2 - 1)\log y}, \tag{5.2}$$

the zero sum being taken in the order of k.

Lemma 5.1 (Asymptotics of Ei off the real axis)**.** *For $w = a + ib$, $b > 0$,*

$$\mathrm{Ei}(w) = \frac{e^w}{w} + \varrho(w), \qquad |\varrho(w)| \le \frac{e^a}{b^2}. \tag{5.3}$$

Proof. Integrate (5.1) by parts: $\int_{-\infty}^{a} e^{v+ib}(v+ib)^{-1}dv = e^w/w + \int_{-\infty}^{a} e^{v+ib}(v+ib)^{-2}dv$, the boundary term at $-\infty$ being 0. The last integral is bounded by $\int_{-\infty}^{a} e^v b^{-2}dv = e^a/b^2$. □

Theorem 5.2 (Compact-uniform existence of the RVM zero sum)**.** *The series $\sum_k \Re\,\mathrm{Ei}((\frac{1}{2} + i\tau_k)u)$ converges uniformly for u in every compact interval $[u_1, u_2] \subset (0, \infty)$. Hence J_{RVM} is a continuous function on $(1, \infty)$.*

Proof. Apply Lemma 5.1 with $w = (\frac{1}{2} + i\tau_k)u$, $a = u/2$, $b = \tau_k u$. Since $\Re(e^{iu\tau_k}/(\frac{1}{2} + i\tau_k)) = (\frac{1}{2}\cos u\tau_k + \tau_k \sin u\tau_k)/(\frac{1}{4} + \tau_k^2)$,

$$\Re\,\mathrm{Ei}((\tfrac{1}{2} + i\tau_k)u) = \frac{e^{u/2}}{u}\,\frac{\sin(u\tau_k)}{\tau_k} + O_{u_1,u_2}(\tau_k^{-2}), \tag{5.4}$$

uniformly for $u \in [u_1, u_2]$. The error series converges absolutely and uniformly. For the main series, Abel summation with $S_{\mathrm{RVM}}(K; u) = O_{u_1,u_2}(1 + \log K)$ (Theorem 4.1) and

$$\frac{1}{\tau_K} - \frac{1}{\tau_{K+1}} \le \frac{\tau_{K+1} - \tau_K}{\tau_K^2} \le \frac{2\pi}{\ell_K \tau_K^2} \ll \frac{\log K}{K^2} \tag{5.5}$$

gives a uniformly convergent majorant $\sum_K (\log K)^2 K^{-2}$, while the boundary term $S_{\mathrm{RVM}}(K; u)/\tau_K \to 0$ uniformly. Hence $\sum_k e^{iu\tau_k}/\tau_k$ converges uniformly, and so does its imaginary part. □

Define the support-truncated version

$$J^*_{\mathrm{RVM}}(x) := \begin{cases} J_{\mathrm{RVM}}(x), & x \ge 2, \\ 0, & 0 < x < 2, \end{cases} \tag{5.6}$$

set $\pi_{\mathrm{RVM}}(x) = 0$ for $0 < x < 2$, and for $x \ge 2$

$$\pi_{\mathrm{RVM}}(x) := \sum_{1 \le m \le \log_2 x} \frac{\mu(m)}{m}\, J^*_{\mathrm{RVM}}(x^{1/m}). \tag{5.7}$$

Proposition 5.3 (Exact finite prime-power inversion)**.** *For every $x \ge 2$,*

$$J^*_{\mathrm{RVM}}(x) = \sum_{1 \le m \le \log_2 x} \frac{1}{m}\, \pi_{\mathrm{RVM}}(x^{1/m}). \tag{5.8}$$

Proof. All sums are finite because $x^{1/m} < 2$ for $m > \log_2 x$. Substituting (5.7) into the right side of (5.8), the coefficient of $J^*_{\mathrm{RVM}}(x^{1/r})$ is

$$\frac{1}{r} \sum_{d \mid r} \mu(d), \tag{5.9}$$

which is 1 for $r = 1$ and 0 otherwise. □

5.1. **The square-root estimate.** For $u > 0$ put

$$\mathcal{A}_{\mathrm{RVM}}(u) := \sum_{k \ge 1} \frac{e^{iu\tau_k}}{\tau_k}, \tag{5.10}$$

summed in the order of k, and write $\ell(t) = \log(t/2\pi)$, so that $\mathcal{N}'_{\mathrm{RVM}}(t) = \ell(t)/2\pi$ and $\ell(\tau_k) = \ell_k$.

Lemma 5.4 (The weighted clock sum)**.** *For every $u \ge \log 2$ the series* (5.10) *converges, and*

$$|\mathcal{A}_{\mathrm{RVM}}(u)| \le \frac{\log^2(2 + 2u)}{4\pi} + \frac{\log(2 + 2u)}{2\pi} + C_{\mathcal{A}}, \tag{5.11}$$

with an absolute constant $C_{\mathcal{A}}$.

Proof. Put $Y_j = 2^j \cdot 4\pi(1+u)$ for $j \ge 0$.

Head $\tau_k \le Y_0$. Let $n(t) = \#\{k : \tau_k \le t\}$. By (2.4), $\tau_k \le t$ if and only if $k \le \mathcal{N}_{\mathrm{RVM}}(t) + \frac{1}{2}$, so $n(t) \le \mathcal{N}_{\mathrm{RVM}}(t) + \frac{1}{2} = \frac{t}{2\pi}(\ell(t) - 1) + \frac{11}{8} \le \frac{t\,\ell(t)}{2\pi}$ for $t \ge \tau_1 > \frac{11\pi}{4}$. Stieltjes integration gives

$$\sum_{\tau_k \le Y_0} \frac{1}{\tau_k} = \frac{n(Y_0)}{Y_0} + \int_{\tau_1}^{Y_0} \frac{n(t)}{t^2}\,dt \le \frac{\ell(Y_0)}{2\pi} + \int_{\tau_1}^{Y_0} \frac{\ell(t)}{2\pi t}\,dt \le \frac{\ell(Y_0)}{2\pi} + \frac{\ell(Y_0)^2}{4\pi}, \tag{5.12}$$

and $\ell(Y_0) = \log(2+2u)$.

Blocks. Let $\mathcal{B}_j = \{k : Y_j < \tau_k \le Y_{j+1}\}$, let $\mathcal{T}$ be the inverse clock of (4.4), and put $\phi(y) = u\,\mathcal{T}(y - \frac{1}{2})/2\pi$, so that $e^{2\pi i\phi(k)} = e^{iu\tau_k}$. By (4.4),

$$\phi''(y) = -\frac{2\pi u}{\mathcal{T}\,\ell(\mathcal{T})^3}, \qquad \mathcal{T} = \mathcal{T}(y - \tfrac{1}{2}). \tag{5.13}$$

On $\mathcal{B}_j$ put $\kappa_j = 2\pi u/(Y_{j+1}\ell(Y_{j+1})^3)$. Since $Y_{j+1} = 2Y_j$ and $\ell(Y_{j+1}) \le 2\ell(Y_j)$ (because $\ell(Y_j) \ge \ell(Y_0) \ge \log 2$), we have $\kappa_j \le -\phi'' \le 16\kappa_j$ there. The number of indices in $\mathcal{B}_j$ is at most $\mathcal{N}_{\mathrm{RVM}}(Y_{j+1}) - \mathcal{N}_{\mathrm{RVM}}(Y_j) + 1 \le Y_j\ell(Y_{j+1})/2\pi + 1$. The second-derivative test for exponential sums [37, Thm. 5.9] gives, for every run $a < k \le b$ of consecutive indices of $\mathcal{B}_j$,

$$\Big|\sum_{a<k\le b} e^{iu\tau_k}\Big| \le A\Big(16\Big(\frac{Y_j\ell(Y_{j+1})}{2\pi} + 1\Big)\kappa_j^{1/2} + \kappa_j^{-1/2}\Big) =: P_j, \tag{5.14}$$

with an absolute constant A; a run of length less than 1 has at most one term, and $1 \le \kappa_j^{-1/2}$ because $\kappa_j \le u/(4(1+u)\log^3 2) < 1$. The weights $1/\tau_k$ are positive and decreasing, and $1/\tau_k < 1/Y_j$ on $\mathcal{B}_j$, so Abel's inequality bounds the contribution of any run in $\mathcal{B}_j$ by P_j/Y_j:

$$\frac{P_j}{Y_j} \le A\left(\frac{16}{2\pi}\Big(\frac{\pi u}{Y_j\ell(Y_{j+1})}\Big)^{1/2} + \frac{16}{Y_j} + \Big(\frac{\ell(Y_{j+1})^3}{\pi u Y_j}\Big)^{1/2}\right). \tag{5.15}$$

With $Y_j = 2^j \cdot 4\pi(1+u)$ and $\ell(Y_{j+1}) = \log(2^{j+2}(1+u)) \le j + 2 + \log(1+u)$, each of the three terms is at most an absolute constant times $(j+2)^{3/2}2^{-j/2}$, uniformly in $u \ge \log 2$, because $\sup_{u \ge \log 2}(1+\log(1+u))^3/(u(1+u)) < \infty$. Hence $\sum_{j\ge 0} P_j/Y_j \le C_{\mathcal{A}}$ with $C_{\mathcal{A}}$ absolute.

Convergence and bound. If $\tau_K \in \mathcal{B}_J$, the terms with $k > K$ form a run of $\mathcal{B}_J$ followed by the full blocks $\mathcal{B}_j$, $j > J$, so $|\sum_{k>K'}^{K''} e^{iu\tau_k}/\tau_k| \le \sum_{j\ge J} P_j/Y_j \to 0$ for $K'' > K' \ge K$, $K \to \infty$. The series converges, and (5.12) together with $\sum_j P_j/Y_j \le C_{\mathcal{A}}$ gives (5.11). □

Theorem 5.5 (Square-root estimate for J_{RVM} and π_{RVM})**.** *For $x \ge 8$,*

$$J_{\mathrm{RVM}}(x) - \mathrm{li}(x) = O\left(\frac{\sqrt{x}(\log\log x)^2}{\log x}\right), \tag{5.16}$$

$$\pi_{\mathrm{RVM}}(x) - \mathrm{li}(x) = O\left(\frac{\sqrt{x}(\log\log x)^2}{\log x}\right), \tag{5.17}$$

with absolute implied constants. No hypothesis on ζ is used.

Proof. Let $u = \log x \ge \log 8$ and $\rho_k = \frac{1}{2} + i\tau_k$. Lemma 5.1 with $w = \rho_k u$, $a = u/2$, $b = \tau_k u$ gives

$$\mathrm{Ei}(\rho_k u) = \frac{e^{\rho_k u}}{\rho_k u} + \varrho_k, \qquad |\varrho_k| \le \frac{e^{u/2}}{\tau_k^2 u^2}. \tag{5.18}$$

Moreover $\frac{1}{\rho_k} = -\frac{i}{\tau_k} + r_k$ with $r_k = \frac{i/2}{\tau_k\rho_k}$, so $|r_k| \le \frac{1}{2\tau_k^2}$. Since $\sum_k \tau_k^{-2} < \infty$, Lemma 5.4 shows that the series in (5.2) converges in the order of k and

$$\sum_{k\ge 1} \mathrm{Ei}(\rho_k u) = \frac{e^{u/2}}{u}\Big(-i\,\mathcal{A}_{\mathrm{RVM}}(u) + \sum_k r_k e^{iu\tau_k}\Big) + \sum_k \varrho_k = \frac{e^{u/2}}{u}\big(-i\,\mathcal{A}_{\mathrm{RVM}}(u) + O(1)\big). \tag{5.19}$$

For $u \geq 2$ one has $\log(2+2u) \leq \log(2u) + \log 2 \leq 3 \log u$, so (5.11) gives $\mathcal{A}_{\mathrm{RVM}}(u) = O(\log^2 u)$. The remaining terms $-\log 2 + \int_x^\infty dy/(y(y^2-1)\log y)$ of (5.2) are bounded for $x \geq 2$. With $e^{u/2} = \sqrt{x}$ and $\log u = \log\log x$ this proves (5.16).

For (5.17), (5.7) gives

$$\pi_{\mathrm{RVM}}(x) - \mathrm{li}(x) = \big(J_{\mathrm{RVM}}(x) - \mathrm{li}(x)\big) + \sum_{2 \leq m \leq \log_2 x} \frac{\mu(m)}{m}\, J^*_{\mathrm{RVM}}(x^{1/m}). \tag{5.20}$$

By (5.16), $\mathrm{li}(y) \ll y/\log y$, and the continuity of J_{RVM} on $[2,8]$ (Theorem 5.2), $|J^*_{\mathrm{RVM}}(y)| \leq C\, y/\log y$ for all $y \geq 2$. The term $m = 2$ is therefore $O(\sqrt{x}/\log x)$, and the terms $3 \leq m \leq \log_2 x$ add up to at most $C\, x^{1/3} \log_2 x/\log x = O(x^{1/3})$. Both are absorbed in (5.17). □

Corollary 5.6 (The three limits)**.** *As $x \to \infty$,*

$$\frac{\mathrm{li}(x)}{\pi_{\mathrm{RVM}}(x)} \to 1, \qquad \frac{\pi(x)}{\mathrm{li}(x)} \to 1, \qquad \frac{\pi(x)}{\pi_{\mathrm{RVM}}(x)} \to 1. \tag{5.21}$$

More precisely, for some absolute $c > 0$,

$$\pi(x) - \pi_{\mathrm{RVM}}(x) = O\Big(x\, e^{-c\sqrt{\log x}}\Big). \tag{5.22}$$

Proof. Since $\mathrm{li}(x) \sim x/\log x$, (5.17) gives $\pi_{\mathrm{RVM}}(x) = \mathrm{li}(x)(1 + O(x^{-1/2}(\log\log x)^2))$; in particular $\pi_{\mathrm{RVM}}(x) > 0$ for large x, and the first limit follows. The second is the prime number theorem [22, 13], in the form $\pi(x) = \mathrm{li}(x) + O(x\, e^{-c\sqrt{\log x}})$ [12, Ch. 18]. Subtracting (5.17) gives (5.22), and dividing by $\pi_{\mathrm{RVM}}(x) \sim x/\log x$ gives the third limit. □

The first limit and (5.17) concern only the synthetic system; the second and third compare it with the primes and use the prime number theorem. Figures 8 and 10 show them numerically.

6. Integer weights and the continuous-to-atomic problem

Define

$$c_n^{\mathrm{RVM}} := \pi_{\mathrm{RVM}}(n) - \pi_{\mathrm{RVM}}(n-1), \qquad n \geq 2, \tag{6.1}$$

and, initially for $\Re s > 1$,

$$\zeta_{\mathbb{Z},\mathrm{RVM}}(s) := \prod_{n \geq 2} (1 - n^{-s})^{-c_n^{\mathrm{RVM}}}. \tag{6.2}$$

No positivity or 0/1 property is built into (6.1). Define the integer-step prime-power count and the right-continuous extension

$$J_{\mathbb{Z},\mathrm{RVM}}(x) := \sum_{1 \leq m \leq \log_2 x} \frac{1}{m}\, \pi_{\mathrm{RVM}}(\lfloor x^{1/m} \rfloor), \tag{6.3}$$

$$J^{*,\mathbb{Z}}_{\mathrm{RVM}}(x) := J^*_{\mathrm{RVM}}(\lfloor x \rfloor). \tag{6.4}$$

Theorem 6.1 (Lipschitz control of J_{RVM})**.** *There is an absolute constant C_J such that*

$$|J_{\mathrm{RVM}}(x+\theta) - J_{\mathrm{RVM}}(x)| \leq C_J\, \theta \qquad (x \geq 2,\ 0 < \theta \leq x). \tag{6.5}$$

Proof. Put $u_1 = \log x$, $u_2 = \log(x+\theta)$, so $0 < u_2 - u_1 = \log(1+\theta/x) \leq \theta/x$ and $u_2 - u_1 \leq \log 2$. The terms li and $\int_x^\infty dy/(y(y^2-1)\log y)$ of (5.2) have derivatives bounded by $1/\log 2$ and $3/\log 2$ on $[2,\infty)$, so they change by $O(\theta)$. It remains to bound $\sum_k D_k$, where, by (5.1) and $\frac{d}{dv}\,\mathrm{Ei}(\rho v) = e^{\rho v}/v$,

$$D_k := \mathrm{Ei}(\rho_k^{\mathrm{RVM}} u_2) - \mathrm{Ei}(\rho_k^{\mathrm{RVM}} u_1) = \int_{u_1}^{u_2} \frac{e^{v/2} e^{iv\tau_k}}{v}\, dv. \tag{6.6}$$

The series $\sum_k D_k$ converges (Theorem 5.2 and its proof). By Abel's theorem for the increasing exponents τ_k, $\sum_k D_k = \lim_{\varepsilon\downarrow 0}\sum_k e^{-\varepsilon\tau_k}D_k$, and for $\varepsilon > 0$ Fubini gives

$$\sum_k e^{-\varepsilon\tau_k}D_k = \int_{u_1}^{u_2}\frac{e^{v/2}}{v}\,S_\varepsilon(v)\,dv, \qquad S_\varepsilon(v) := \sum_{k\ge 1} g(\tau_k), \quad g(\tau) := e^{(iv-\varepsilon)\tau}. \tag{6.7}$$

Clock decomposition. With $\tau_0 = 4\pi$ and the sawtooth count (8.8), Stieltjes integration exactly as in (8.9) ($g(\infty) = 0$, $g' \in L^1$) gives

$$S_\varepsilon(v) = \int_{\tau_0}^{\infty} g\,d\mathcal{N}_{\mathrm{RVM}} + g(\tau_0)\mathcal{N}_{\mathrm{RVM}}(\tau_0) + \int_{\tau_0}^{\infty}\psi(\mathcal{N}_{\mathrm{RVM}}(\tau) + \tfrac{1}{2})g'(\tau)\,d\tau. \tag{6.8}$$

The middle term is at most $\frac{1}{2}$. For the first, two integrations by parts with $\mathcal{N}'_{\mathrm{RVM}}(\tau) = \frac{1}{2\pi}\log\frac{\tau}{2\pi}$, $\mathcal{N}''_{\mathrm{RVM}}(\tau) = \frac{1}{2\pi\tau}$ and $v \ge \log 2$ give $|\int_{\tau_0}^{\infty} g\,d\mathcal{N}_{\mathrm{RVM}}| \le C/v$, uniformly in $\varepsilon \in (0,1]$.

Harmonics. The partial sums of $\psi(y+\frac{1}{2}) = -\sum_{m\ge 1}(-1)^m\sin(2\pi m y)/(\pi m)$ are uniformly bounded and converge off a null set, and $g' \in L^1$ for $\varepsilon > 0$, so termwise integration is allowed:

$$\int_{\tau_0}^{\infty}\psi(\mathcal{N}_{\mathrm{RVM}} + \tfrac{1}{2})g'\,d\tau = -\sum_{m\ge 1}\frac{(-1)^m}{2\pi i\,m}(K_m^+ - K_m^-), \qquad K_m^{\pm} := \int_{\tau_0}^{\infty}(iv-\varepsilon)e^{-\varepsilon\tau}e^{i\Phi_m^{\pm}(\tau)}\,d\tau, \tag{6.9}$$

with $\Phi_m^{\pm}(\tau) = v\tau \pm 2\pi m\mathcal{N}_{\mathrm{RVM}}(\tau)$. The weight $e^{-\varepsilon\tau}$ is positive and decreasing, so by the second mean-value theorem each integral is at most $2(v+1)\sup_\xi|\int_a^\xi e^{i\Phi}d\tau|$ over the relevant interval $[a,b]$. We use the first- and second-derivative tests [37, Lemmas 4.2 and 4.4]: $|\int_a^b e^{i\Phi}| \le 4/\mu$ if Φ' is monotone with $|\Phi'| \ge \mu$, and $\le 8/\sqrt{\lambda}$ if $|\Phi''| \ge \lambda$.

For K_m^+: $(\Phi_m^+)' = v + m\log\frac{\tau}{2\pi} \ge v + m\log 2$ is increasing, so $|K_m^+| \le 8(v+1)/(v+m\log 2)$.

For K_m^-: $(\Phi_m^-)' = v - m\log\frac{\tau}{2\pi}$ is decreasing and vanishes at $\tau_m := 2\pi e^{v/m} > 2\pi$, and $(\Phi_m^-)'' = -m/\tau$. On $[\tau_0, \tau_m/2]$ (if nonempty) $(\Phi_m^-)' \ge m\log 2$; on $[2\tau_m,\infty)$, $(\Phi_m^-)' \le -m\log 2$; on $[\max(\tau_0,\tau_m/2), 2\tau_m]$, $|(\Phi_m^-)''| \ge m/(2\tau_m)$. Hence

$$|K_m^-| \le 2(v+1)\Big(\frac{8}{m\log 2} + 8\sqrt{\frac{2\tau_m}{m}}\Big) = 2(v+1)\Big(\frac{8}{m\log 2} + 16\sqrt{\pi}\,\frac{e^{v/(2m)}}{\sqrt{m}}\Big). \tag{6.10}$$

Summing with the weights $1/(2\pi m)$ and using $\sum_m m^{-3/2}e^{v/(2m)} \le e^{v/2} + \zeta(\frac{3}{2})e^{v/4}$,

$$|S_\varepsilon(v)| \le C\,(v+1)\,e^{v/2} \qquad (v \ge \log 2,\ 0 < \varepsilon \le 1), \tag{6.11}$$

with an absolute constant C.

Conclusion. Inserting (6.11) into (6.7), for $v \in [u_1,u_2]$ we have $e^{v/2} \le \sqrt{2x}$ and $(v+1)/v \le 1 + 1/\log 2$, so

$$\Big|\sum_k e^{-\varepsilon\tau_k}D_k\Big| \le C'(u_2-u_1)\,x \le C'\theta, \tag{6.12}$$

uniformly in ε. Letting $\varepsilon \downarrow 0$ and taking $2\Re$ proves (6.5). □

The $m = 1$ harmonic, with stationary point $\tau_1 = 2\pi x$, produces the unit-periodic ripple of J_{RVM} (Figure 8); the harmonics $m \ge 2$ have stationary points $2\pi x^{1/m}$ and contribute $O(x^{-1/2+1/(2m)}m^{-3/2})$ to a unit increment.

Corollary 6.2 (Bounded cell masses)**.**

$$\pi_{\mathrm{RVM}}(y) - \pi_{\mathrm{RVM}}(\lfloor y\rfloor) = O(1), \qquad c_n^{\mathrm{RVM}} = O(1). \tag{6.13}$$

Proof. By (5.7), $c_n^{\mathrm{RVM}} = \sum_{m\le\log_2 n}\frac{\mu(m)}{m}(J^*_{\mathrm{RVM}}(n^{1/m}) - J^*_{\mathrm{RVM}}((n-1)^{1/m}))$. The term $m = 1$ is $O(1)$ by (6.5). For $m \ge 2$ and $(n-1)^{1/m} \ge 2$, (6.5) with $\theta = n^{1/m} - (n-1)^{1/m} \le n^{1/m-1}$ bounds the term by $C_J n^{1/m-1} \le C_J n^{-1/2}$, and there are at most $\log_2 n$ such terms. The jump of J^*_{RVM} at 2 enters only for $n = 2^m$, once, with size $|J_{\mathrm{RVM}}(2)|/m$. The same argument applies to $\pi_{\mathrm{RVM}}(y) - \pi_{\mathrm{RVM}}(\lfloor y\rfloor)$. □

Corollary 6.3 (The integer-weighted zeta is a Dirichlet series)**.** *The product* (6.2) *converges absolutely for* $\Re s > 1$ *and*

$$\zeta_{\mathbb{Z},\mathrm{RVM}}(s) = \sum_{N\geq 1} \frac{a_N^{\mathbb{Z}}}{N^s}, \qquad a_1^{\mathbb{Z}} = 1, \qquad \sum_{N\geq 1} \frac{|a_N^{\mathbb{Z}}|}{N^\sigma} < \infty \quad (\sigma > 1). \tag{6.14}$$

The inverse is also an absolutely convergent Dirichlet series on the whole half-plane:

$$\frac{1}{\zeta_{\mathbb{Z},\mathrm{RVM}}(s)} = \sum_{N\geq 1} \frac{\bar{a}_N^{\mathbb{Z}}}{N^s}, \qquad \bar{a}_1^{\mathbb{Z}} = 1, \qquad \sum_{N\geq 1} \frac{|\bar{a}_N^{\mathbb{Z}}|}{N^\sigma} < \infty \quad (\sigma > 1). \tag{6.15}$$

Proof. By Corollary 6.2,

$$\log \zeta_{\mathbb{Z},\mathrm{RVM}}(s) = \sum_{n\geq 2} c_n^{\mathrm{RVM}} \sum_{m\geq 1} \frac{n^{-ms}}{m} = \sum_{N\geq 2} \frac{b_N}{N^s}, \qquad b_N := \sum_{n^m = N} \frac{c_n^{\mathrm{RVM}}}{m}, \tag{6.16}$$

with $\sum_N |b_N| N^{-\sigma} \leq C \sum_{n\geq 2} \sum_m n^{-m\sigma}/m < \infty$ for $\sigma > 1$. The exponential of an absolutely convergent Dirichlet series without constant term is an absolutely convergent Dirichlet series with $a_1^{\mathbb{Z}} = 1$, and $\sum_N |a_N^{\mathbb{Z}}| N^{-\sigma} \leq \exp(\sum_N |b_N| N^{-\sigma})$. The same applies to $1/\zeta_{\mathbb{Z},\mathrm{RVM}} = \exp(-\sum_N b_N N^{-s})$, which gives (6.15). □

Numerically the cell masses lie in $[-0.38, 1.04]$ for $n \leq 1000$ (Experiment 17.6).

Theorem 6.4 (Mellin–Stieltjes identification)**.** *There is a function* $J_{\mathrm{RVM}}^{\mathrm{M}}$ *on* $(1,\infty)$ *with* $J_{\mathrm{RVM}}^{\mathrm{M}}(x) x^{-c} \in L^2((1,\infty), dx/x)$ *for every* $c > 1$ *such that*

$$\log \zeta_{\mathrm{RVM}}(s) = s \int_1^\infty J_{\mathrm{RVM}}^{\mathrm{M}}(x)\, x^{-s-1}\, dx \qquad (\Re s > 1), \tag{6.17}$$

and $J_{\mathrm{RVM}}^{\mathrm{M}} - J_{\mathrm{RVM}}$ *is constant on* $(1,\infty)$*. In particular* $J_{\mathrm{RVM}}^{\mathrm{M}} = J_{\mathrm{RVM}}^{*} + C_{\mathrm{RVM}}$ *on* $[2,\infty)$ *for a constant* C_{RVM}*.*

The proof is given in Section 12 (Theorems 12.5 and 12.6); it uses the half-plane bound of Theorem 8.10.

Corollary 6.5 (Weighted factorization)**.**

$$J_{\mathbb{Z},\mathrm{RVM}}(x) - J_{\mathrm{RVM}}^{*}(x) = O(\log\log x), \tag{6.18}$$

and

$$\zeta_{\mathbb{Z},\mathrm{RVM}}(s) = \zeta_{\mathrm{RVM}}(s)\, e^{h_{\mathrm{RVM}}(s)}, \tag{6.19}$$

where

$$h_{\mathrm{RVM}}(s) := s \int_1^\infty (J_{\mathbb{Z},\mathrm{RVM}}(x) - J_{\mathrm{RVM}}^{\mathrm{M}}(x)) x^{-s-1}\, dx \tag{6.20}$$

is holomorphic for $\Re s > 0$*. The two functions have the same zeros in* $\Re s > 0$*; compare the modified Euler products of Grosswald and Schnitzer* [20]*.*

Proof. By (6.13) replacing $x^{1/m}$ by its integer part costs $O(1/m)$ in the mth summand; summing m^{-1} up to $\log_2 x$ gives (6.18). The product (6.2) gives $\log \zeta_{\mathbb{Z},\mathrm{RVM}}(s) = s \int_1^\infty J_{\mathbb{Z},\mathrm{RVM}}(x) x^{-s-1} dx$ for $\Re s > 1$, because $\sum_{n\leq y} c_n^{\mathrm{RVM}} = \pi_{\mathrm{RVM}}(\lfloor y \rfloor)$. Subtracting (6.17) gives (6.19) on $\Re s > 1$. On $[2,\infty)$, $J_{\mathbb{Z},\mathrm{RVM}} - J_{\mathrm{RVM}}^{\mathrm{M}} = (J_{\mathbb{Z},\mathrm{RVM}} - J_{\mathrm{RVM}}^{*}) - C_{\mathrm{RVM}} = O(\log\log x)$ by (6.18); on $(1,2)$, $J_{\mathbb{Z},\mathrm{RVM}} = 0$ and $J_{\mathrm{RVM}}^{\mathrm{M}}$ is square-integrable. Hence (6.20) converges locally normally for $\Re s > 0$, and exponentiation introduces no zeros. □

Lemma 6.6 (Exact first-harmonic cell cancellation)**.** *For every integer* n*,*

$$\int_{n-1}^{n} 2\cos(2\pi x)\, dx = 0. \tag{6.21}$$

Hypothesis 6.7 (Uniform cell remainder)**.**

$$c_n^{\mathrm{RVM}} - \int_{n-1}^{n} \left(\frac{1}{\log x} + 2\cos(2\pi x) \right) dx = o\Big(\frac{1}{\log n} \Big). \tag{6.22}$$

Proposition 6.8 (Conditional eventual positivity)**.** *Hypothesis 6.7 implies*

$$c_n^{\mathrm{RVM}} \sim \frac{1}{\log n}, \tag{6.23}$$

hence $0 < c_n^{\mathrm{RVM}} < 1$ *for all large* n*.*

Proof. The main-density integral is $1/\log n + O(1/(n\log^2 n))$; the first harmonic vanishes by Lemma 6.6. □

7. The critical RVM Mertens transform

Define

$$F_{\mathrm{RVM}}(s) = \widehat{M}^{\mathrm{crit}}_{\mathrm{RVM}}(s) := \frac{1}{\zeta_{\mathrm{RVM}}(s)} = \frac{G(s)}{P^{\mathrm{crit}}_{\mathrm{RVM}}(s)}. \tag{7.1}$$

Formally the associated summatory object is the inverse Mellin transform

$$M^{\mathrm{crit}}_{\mathrm{RVM}}(x) = \frac{1}{2\pi i} \int_{(c)} F_{\mathrm{RVM}}(s) \frac{x^s}{s}\, ds; \tag{7.2}$$

its existence as a symmetric limit, for every $c > \frac{1}{2}$, is proved in Theorem 8.12.

Proposition 7.1 (Synthetic pole abscissa)**.** F_{RVM} *is holomorphic in* $\Re s > \frac{1}{2}$*. Its poles are the simple poles*

$$\frac{1}{2} \pm i\tau_k \quad (k \geq 1) \qquad \textit{and} \qquad s = -2, -4, -6, \ldots \tag{7.3}$$

and $s = 1$ *is a simple zero of* F_{RVM}*.*

Proof. This is Theorem 3.1 applied to $F_{\mathrm{RVM}} = G/P^{\mathrm{crit}}_{\mathrm{RVM}}$: the zeros of $P^{\mathrm{crit}}_{\mathrm{RVM}}$ are (3.6), and the poles of G are those of $\Gamma(s/2)$ except $s = 0$, where $s\Gamma(s/2) \to 2$. □

Proposition 7.2 (Gaussian square-root bound under vertical growth)**.** *Fix* $c > 1$ *and* $0 < \varepsilon < c - \frac{1}{2}$*. Assume, uniformly for* $\frac{1}{2} + \varepsilon \leq \sigma \leq c$*,*

$$F_{\mathrm{RVM}}(\sigma + it) = \exp(o(t^2)) \qquad (|t| \to \infty). \tag{7.4}$$

Define

$$M^{\mathrm{crit}}_{\mathrm{RVM},G}(x) := \frac{1}{2\pi i} \int_{(c)} e^{s^2} F_{\mathrm{RVM}}(s) \frac{x^s}{s}\, ds. \tag{7.5}$$

Then

$$M^{\mathrm{crit}}_{\mathrm{RVM},G}(x) = O_\varepsilon(x^{1/2+\varepsilon}). \tag{7.6}$$

Proof. On $s = \sigma + it$, $|e^{s^2}| = e^{\sigma^2 - t^2}$, so (7.4) gives absolute convergence on vertical lines and makes the horizontal integrals tend to 0. The strip $\frac{1}{2} + \varepsilon \leq \Re s \leq c$ contains no pole of F_{RVM} (Proposition 7.1) and does not contain $s = 0$, so Cauchy's theorem moves the line to $\Re s = \frac{1}{2} + \varepsilon$, where $|x^s| = x^{1/2+\varepsilon}$. □

Theorem 8.12 below removes both the assumption (7.4) and the smoothing.

8. The transition theorem and the unsmoothed synthetic Mertens bound

Let

$$\chi(s) := 2^s\pi^{s-1}\sin\Big(\frac{\pi s}{2}\Big)\Gamma(1-s) = \pi^{s-1/2}\frac{\Gamma((1-s)/2)}{\Gamma(s/2)}, \tag{8.1}$$

the factor of the functional equation $\zeta(s) = \chi(s)\zeta(1-s)$ [37, §2.1]. Write

$$s = \tfrac{1}{2} + z, \quad z = \sigma' + it, \qquad \tau_* := -iz = t - i\sigma', \qquad w(s) := \exp\{-2\pi i\,\mathcal{N}_{\mathrm{RVM}}(\tau_*)\}. \tag{8.2}$$

Stirling's formula gives, uniformly on compact σ-sets,

$$|\chi(\sigma+it)| = \Big(\frac{|t|}{2\pi}\Big)^{1/2-\sigma}\big(1+O(|t|^{-1})\big). \tag{8.3}$$

Theorem 8.1 (Transition evaluation of the midpoint product)**.** *There is a constant $\varpi_{\mathrm{RVM}} > 0$ with the following properties.*

(a) *For every $B > 0$ there are $t_0(B)$ and C_B such that, for $0 < \sigma' \le B$ and $t \ge t_0(B)$,*

$$\zeta_{\mathrm{RVM}}(s) = \varpi_{\mathrm{RVM}}\big(1+w(s)\big)\big(1+\mathcal{E}_0(s)\big), \qquad |\mathcal{E}_0(s)| \le \frac{C_B}{t}. \tag{8.4}$$

(b) *For every $0 < \sigma_1' < \sigma_2' < \frac{1}{2}$ there are t_0 and C such that, for $\sigma_1' \le \sigma' \le \sigma_2'$ and $t \ge t_0$,*

$$\zeta_{\mathrm{RVM}}(s) = \varpi_{\mathrm{RVM}}\big(1+\chi(s)\big)\big(1+\mathcal{E}(s)\big), \qquad |\mathcal{E}(s)| \le \frac{C}{t}. \tag{8.5}$$

For $t < 0$ the same statements hold by $\zeta_{\mathrm{RVM}}(\bar s) = \overline{\zeta_{\mathrm{RVM}}(s)}$.

The phase identity $w = \chi(s)(1+O(1/t))$ alone does not give (8.4); the product has to be evaluated across the transition region $\tau_k \asymp t$. Log denotes the principal logarithm and $\operatorname{Log}\Gamma$ the branch of $\log\Gamma$ on $\Re v > 0$ that is real on $(0,\infty)$. Nothing about ζ is used.

8.1. **Proof of Theorem 8.1.** Throughout, $t > 0$, $0 < \sigma' \le B$, $\tau_0 := 4\pi$, $t^2 \ge 2(\tau_0^2 + B^2)$, and implied constants depend only on B (and on σ_1', σ_2' in part (b)).

Step 0: reduction to a sum over the clock. Since $q(s) = z^2 - \frac{1}{4}$, each factor of (3.2) equals $(\tau_k^2 + z^2)/(\tau_k^2 + \frac{1}{4})$, so

$$\frac{P^{\mathrm{crit}}_{\mathrm{RVM}}(s)}{P^{\mathrm{crit}}_{\mathrm{RVM}}(\frac{1}{2})} = \prod_{k\ge1}\Big(1+\frac{z^2}{\tau_k^2}\Big) = \exp\Lambda(z), \qquad \Lambda(z) := \sum_{k\ge1} f(\tau_k), \quad f(\tau) := \operatorname{Log}\Big(1+\frac{z^2}{\tau^2}\Big). \tag{8.6}$$

For $\tau > 0$, $\Im(1 + z^2/\tau^2) = 2\sigma' t/\tau^2 > 0$, so f is smooth on $(0,\infty)$, $f(\tau) = O(|z|^2\tau^{-2})$, the series converges absolutely, and

$$f'(\tau) = -\frac{2z^2}{\tau(\tau^2+z^2)} = -\frac{2}{\tau} + \frac{1}{\tau - iz} + \frac{1}{\tau + iz}. \tag{8.7}$$

Step 1: the midpoint sawtooth. The clock increases on $[2\pi,\infty)$, $\mathcal{N}_{\mathrm{RVM}}(\tau_0) = 2\log\frac{2}{e} + \frac{7}{8} \in (0, \frac{1}{2})$, and $\tau_1 > \tau_0$. With $\psi(y) := y - \lfloor y\rfloor - \frac{1}{2}$, the count (4.12) satisfies, for $\tau \ge \tau_0$,

$$n_{\mathrm{RVM}}(\tau) = \lfloor \mathcal{N}_{\mathrm{RVM}}(\tau) + \tfrac{1}{2}\rfloor = \mathcal{N}_{\mathrm{RVM}}(\tau) - \psi\big(\mathcal{N}_{\mathrm{RVM}}(\tau) + \tfrac{1}{2}\big), \qquad n_{\mathrm{RVM}}(\tau_0) = 0. \tag{8.8}$$

Stieltjes integration of f against (8.8), followed by integration by parts in the ψ-part ($f(\infty) = 0$, $|\psi| \le \frac{1}{2}$, $\psi(\mathcal{N}_{\mathrm{RVM}}(\tau_0) + \frac{1}{2}) = \mathcal{N}_{\mathrm{RVM}}(\tau_0)$), gives

$$\begin{aligned} &\Lambda(z) = \Lambda_{\mathrm{sm}}(z) + \Lambda_{\mathrm{saw}}(z), \qquad \Lambda_{\mathrm{sm}}(z) := \int_{\tau_0}^{\infty} f\,d\mathcal{N}_{\mathrm{RVM}} + f(\tau_0)\mathcal{N}_{\mathrm{RVM}}(\tau_0), \\ &\Lambda_{\mathrm{saw}}(z) := \int_{\tau_0}^{\infty} \psi\big(\mathcal{N}_{\mathrm{RVM}}(\tau) + \tfrac{1}{2}\big) f'(\tau)\,d\tau. \end{aligned} \tag{8.9}$$

Lemma 8.2 (Smooth part)**.** *With* $\log G(s) := \operatorname{Log}(z^2 - \frac{1}{4}) + \operatorname{Log}\Gamma(\frac{s}{2}) - \frac{s}{2}\log\pi$,

$$\Lambda_{\mathrm{sm}}(z) - \log G(s) = \nu_3 + O(t^{-1}), \qquad \nu_3 \in \mathbb{R} \text{ independent of } z. \tag{8.10}$$

Proof. A second integration by parts ($f\mathcal{N}_{\mathrm{RVM}} \to 0$ at ∞) cancels the boundary term:

$$\Lambda_{\mathrm{sm}}(z) = -\int_{\tau_0}^{\infty} \mathcal{N}_{\mathrm{RVM}} f'\, d\tau = \int_{\tau_0}^{\infty} \mathcal{N}_{\mathrm{RVM}}(\tau)\Big(\frac{2}{\tau} - \frac{2\tau}{\tau^2 + z^2}\Big) d\tau. \tag{8.11}$$

The constant $\frac{7}{8}$ of the clock contributes $\frac{7}{8}\operatorname{Log}(1 + z^2/\tau_0^2)$, because $\operatorname{Log}\frac{\tau^2}{\tau^2+z^2}$ is an antiderivative of the bracket on $(0,\infty)$ tending to 0 at ∞. As $0 < \arg z < \frac{\pi}{2}$, $\arg(1 + \tau_0^2/z^2) = O(t^{-2})$ and

$$\tfrac{7}{8}\operatorname{Log}\Big(1 + \frac{z^2}{\tau_0^2}\Big) = \tfrac{7}{4}\operatorname{Log} z - \tfrac{7}{4}\log\tau_0 + O(t^{-2}). \tag{8.12}$$

The term $\frac{\tau}{2\pi}\log\frac{\tau}{2\pi e}$ contributes $\frac{1}{\pi}\int_{\tau_0}^{\infty}\log\frac{\tau}{2\pi e}\frac{z^2}{\tau^2+z^2}\,d\tau$. Put

$$\mathcal{I}(z) := \frac{1}{\pi}\int_0^{\infty} \log\frac{\tau}{2\pi e}\,\frac{z^2}{\tau^2 + z^2}\, d\tau. \tag{8.13}$$

The integrand is holomorphic in z on $\Re z > 0$ (its singularities are at $z = \pm i\tau$), $O(|z|^2\tau^{-2}\log\tau)$ at ∞ and $O(|\log\tau|)$ at 0, locally uniformly; so $\mathcal{I}$ is holomorphic on $\Re z > 0$. For real $z = x > 0$ the substitution $\tau = xv$ and $\int_0^\infty \frac{dv}{1+v^2} = \frac{\pi}{2}$, $\int_0^\infty \frac{\log v}{1+v^2}\,dv = 0$ give $\mathcal{I}(x) = \frac{x}{2}\log\frac{x}{2\pi e}$; by the identity theorem $\mathcal{I}(z) = \frac{z}{2}\operatorname{Log}\frac{z}{2\pi e}$ on $\Re z > 0$. On $[0,\tau_0]$, $|\tau^2 + z^2| \geq t^2/2$, so $\frac{z^2}{\tau^2+z^2} = 1 + O(t^{-2})$ there. Hence

$$\Lambda_{\mathrm{sm}}(z) = \frac{z}{2}\operatorname{Log}\frac{z}{2\pi e} + \frac{7}{4}\operatorname{Log} z + \nu_1 + O(t^{-2}), \qquad \nu_1 = -\tfrac{7}{4}\log\tau_0 - \tfrac{1}{\pi}\int_0^{\tau_0}\log\tfrac{\tau}{2\pi e}\,d\tau. \tag{8.14}$$

Stirling's formula $\operatorname{Log}\Gamma(v) = (v - \frac{1}{2})\operatorname{Log} v - v + \frac{1}{2}\log 2\pi + O(|v|^{-1})$ [17, §5.11] with $v = \frac{s}{2} = \frac{z+1/2}{2}$ and $\operatorname{Log} v = \operatorname{Log} z - \log 2 + \frac{1}{2z} + O(t^{-2})$ gives

$$\operatorname{Log}\Gamma(\tfrac{s}{2}) = \tfrac{z}{2}\operatorname{Log}\tfrac{z}{2} - \tfrac{1}{4}\operatorname{Log} z + \tfrac{1}{4}\log 2 - \tfrac{z}{2} + \tfrac{1}{2}\log 2\pi + O(t^{-1}), \tag{8.15}$$

and with $\operatorname{Log}(z^2 - \frac{1}{4}) = 2\operatorname{Log} z + O(t^{-2})$, $-\frac{s}{2}\log\pi = -\frac{z}{2}\log\pi - \frac{1}{4}\log\pi$,

$$\log G(s) = \frac{z}{2}\operatorname{Log}\frac{z}{2\pi e} + \frac{7}{4}\operatorname{Log} z + \nu_2 + O(t^{-1}), \qquad \nu_2 = \tfrac{1}{4}\log 2 + \tfrac{1}{2}\log 2\pi - \tfrac{1}{4}\log\pi. \tag{8.16}$$

Subtracting gives (8.10) with $\nu_3 = \nu_1 - \nu_2$. □

Lemma 8.3 (The clock as a holomorphic phase)**.** $\mathcal{N}_{\mathrm{RVM}}(\tau) = \frac{\tau}{2\pi}\big(\operatorname{Log}\frac{\tau}{2\pi} - 1\big) + \frac{7}{8}$ *is holomorphic on* $\Re\tau > 0$, $\mathcal{N}_{\mathrm{RVM}}(\bar\tau) = \overline{\mathcal{N}_{\mathrm{RVM}}(\tau)}$, *and for* $\Re\tau \geq \tau_0$,

$$2\pi\,\Im\mathcal{N}_{\mathrm{RVM}}(\tau) \geq (\log 2)\,\Im\tau \quad (\Im\tau \geq 0), \qquad 2\pi\,\Im\mathcal{N}_{\mathrm{RVM}}(\tau) \leq (\log 2)\,\Im\tau \quad (\Im\tau \leq 0). \tag{8.17}$$

Proof. For $\tau = X + iY$, $X \geq \tau_0$, $Y > 0$, put $v = Y/X$. Then

$$2\pi\Im\mathcal{N}_{\mathrm{RVM}}(\tau) = Y\log\frac{|\tau|}{2\pi e} + X\arg\tau = Y\Big[\log\frac{X}{2\pi e} + \tfrac{1}{2}\log(1+v^2) + \frac{\arctan v}{v}\Big] \geq Y\big[\log 2 + \varphi(v)\big], \tag{8.18}$$

using $X \geq 4\pi$, where $\varphi(v) = \frac{1}{2}\log(1+v^2) + \frac{\arctan v}{v} - 1$. Since $\varphi(0^+) = 0$ and $\varphi'(v) = (v - \arctan v)/v^2 \geq 0$, $\varphi \geq 0$. The case $Y < 0$ follows by conjugation. □

In particular $|w(s)| = e^{2\pi\Im\mathcal{N}_{\mathrm{RVM}}(\tau_*)} \leq 2^{-\sigma'} < 1$, and on the vertical rays $|e^{\pm 2\pi i m\mathcal{N}_{\mathrm{RVM}}(\tau_0 \pm iy)}| \leq 2^{-my}$.

Lemma 8.4 (Abel-regularized harmonic sum)**.**

$$\Lambda_{\mathrm{saw}}(z) = \operatorname{Log}(1 + w(s)) + \nu_4 + O(t^{-2}), \qquad \nu_4 \in \mathbb{R} \text{ independent of } z. \tag{8.19}$$

Proof. Regularization. For $0 < r < 1$ let

$$\psi_r(y) := -\sum_{m\geq 1} r^m \frac{\sin 2\pi m y}{\pi m} = \frac{1}{\pi}\Im \operatorname{Log}(1 - re^{2\pi i y}). \tag{8.20}$$

Since $\Re(1 - re^{2\pi iy}) > 0$, $|\psi_r| \leq \frac{1}{2}$, and $\psi_r(y) \to \psi(y)$ as $r \uparrow 1$ for $y \notin \mathbb{Z}$ (Abel summation of the convergent Fourier series of ψ). The set $\{\tau : \mathcal{N}_{\mathrm{RVM}}(\tau) + \frac{1}{2} \in \mathbb{Z}\}$ is discrete and $f' \in L^1(\tau_0, \infty)$, so by dominated convergence

$$\Lambda_{\mathrm{saw}}(z) = \lim_{r\uparrow 1} \Lambda_r(z), \qquad \Lambda_r(z) := \int_{\tau_0}^{\infty} \psi_r(\mathcal{N}_{\mathrm{RVM}} + \tfrac{1}{2}) f' \, d\tau. \tag{8.21}$$

Termwise integration. For fixed $r < 1$ the series for $\psi_r(\mathcal{N}_{\mathrm{RVM}} + \frac{1}{2})$, with $\sin 2\pi m(y + \frac{1}{2}) = (-1)^m \sin 2\pi m y$, is dominated by $\sum_m r^m/m$; by Fubini,

$$\Lambda_r(z) = -\sum_{m\geq 1} \frac{(-r)^m}{2\pi i\, m}(I_m^+ - I_m^-), \qquad I_m^{\pm} := \int_{\tau_0}^{\infty} e^{\pm 2\pi i m \mathcal{N}_{\mathrm{RVM}}(\tau)} f'(\tau)\, d\tau. \tag{8.22}$$

Contours. Let $U^{\pm} = \{\Re\tau \geq \tau_0,\ \pm\Im\tau \geq 0\}$. On $U^{\pm}$, $|e^{\pm 2\pi i m \mathcal{N}_{\mathrm{RVM}}}| \leq 1$ by Lemma 8.3, and $|f'(\tau)| \leq 4|z|^2|\tau|^{-3}$ for $|\tau| \geq 2|z|$, so the quarter-circle arcs of radius R contribute $O(|z|^2 R^{-2}) \to 0$. The poles of f' are 0, $iz = -t + i\sigma' \notin U^{\pm}$, and $\tau_* = -iz = t - i\sigma' \in U^-$, the last with residue 1 (from $\frac{1}{\tau + iz}$). Define the ray integrals

$$R_m^+ := i\int_0^{\infty} e^{2\pi i m \mathcal{N}_{\mathrm{RVM}}(\tau_0 + iy)} f'(\tau_0 + iy)\, dy, \qquad R_m^- := -i\int_0^{\infty} e^{-2\pi i m \mathcal{N}_{\mathrm{RVM}}(\tau_0 - iy)} f'(\tau_0 - iy)\, dy. \tag{8.23}$$

On U^+ Cauchy's theorem gives $I_m^+ = R_m^+$. On U^- the positively oriented boundary of the quarter disc runs down the ray, along the arc, and back along $[\tau_0, \tau_0 + R]$ from right to left; letting $R \to \infty$,

$$R_m^- - I_m^- = 2\pi i\, e^{-2\pi i m \mathcal{N}_{\mathrm{RVM}}(\tau_*)} = 2\pi i\, w^m. \tag{8.24}$$

Therefore, using $|rw| < 1$,

$$\Lambda_r(z) = -\sum_{m\geq 1} \frac{(-rw)^m}{m} + \Phi_r(z) = \operatorname{Log}(1 + rw) + \Phi_r(z), \qquad \Phi_r(z) := -\sum_{m\geq 1} \frac{(-r)^m}{2\pi i\, m}(R_m^+ - R_m^-). \tag{8.25}$$

The rays. Split $f'(\tau) = -\frac{2}{\tau} + g_z(\tau)$, $g_z(\tau) = \frac{2\tau}{\tau^2 + z^2}$, and correspondingly $R_m^{\pm} = R_m^{\pm,(0)} + R_m^{\pm,(z)}$, $\Phi_r = \Phi_r^{(0)} + \Phi_r^{(z)}$. The ray integrands carry the factor 2^{-my}. Since $|2/\tau| \leq 2/\tau_0$ on the rays, $|R_m^{\pm,(0)}| \leq 2/(\tau_0 m \log 2)$, so the series for $\Phi_r^{(0)}$ is dominated by $\sum_m C/m^2$ uniformly in r, and $\Phi_r^{(0)} \to \nu_4 := \Phi_1^{(0)}$, a constant depending only on τ_0. It is real: $\mathcal{N}_{\mathrm{RVM}}(\bar\tau) = \overline{\mathcal{N}_{\mathrm{RVM}}(\tau)}$ gives $R_m^{-,(0)} = \overline{R_m^{+,(0)}}$, so each term equals $-(-r)^m \Im R_m^{+,(0)}/(\pi m)$. For $\tau = \tau_0 \pm iy$, $\Re(\tau^2 + z^2) = \tau_0^2 + \sigma'^2 - y^2 - t^2 \leq -\frac{1}{2}(y^2 + t^2)$ because $t^2 \geq 2(\tau_0^2 + B^2)$; hence $|g_z(\tau)| \leq 4(\tau_0 + y)t^{-2}$ and

$$|R_m^{\pm,(z)}| \leq \frac{4}{t^2}\int_0^{\infty} 2^{-my}(\tau_0 + y)\, dy = \frac{4}{t^2}\Big(\frac{\tau_0}{m\log 2} + \frac{1}{(m\log 2)^2}\Big). \tag{8.26}$$

So $|\Phi_r^{(z)}(z)| \leq C\, t^{-2} \sum_m m^{-2}$, uniformly in $0 < r < 1$ and $0 < \sigma' \leq B$, and $\Phi_r^{(z)}$ converges as $r \uparrow 1$ by dominated convergence of the series. Since $\operatorname{Log}(1 + rw) \to \operatorname{Log}(1 + w)$, letting $r \uparrow 1$ in (8.25) and (8.21) gives (8.19). □

Step 2: assembly. By (8.6), (8.9) and Lemmas 8.2 and 8.4,

$$\begin{aligned} \zeta_{\mathrm{RVM}}(s) &= \frac{P_{\mathrm{RVM}}^{\mathrm{crit}}(s)}{G(s)} = P_{\mathrm{RVM}}^{\mathrm{crit}}(\tfrac{1}{2}) \exp\big(\Lambda_{\mathrm{sm}}(z) - \log G(s) + \Lambda_{\mathrm{saw}}(z)\big) \\ &= P_{\mathrm{RVM}}^{\mathrm{crit}}(\tfrac{1}{2})\, e^{\nu_3 + \nu_4}(1 + w)\big(1 + O(t^{-1})\big), \end{aligned} \tag{8.27}$$

uniformly for $0 < \sigma' \le B$. This is (8.4) with

$$\varpi_{\mathrm{RVM}} := P^{\mathrm{crit}}_{\mathrm{RVM}}(\tfrac12)\, e^{\nu_3+\nu_4} > 0, \tag{8.28}$$

which depends only on the clock; its numerical value $0.9619415776\ldots$ plays no role in the proof.

Step 3: from the clock phase to χ. Let

$$\vartheta(v) := \frac{1}{2i}\Big(\operatorname{Log}\Gamma(\tfrac14+\tfrac{iv}{2}) - \operatorname{Log}\Gamma(\tfrac14-\tfrac{iv}{2})\Big) - \frac{v}{2}\log\pi, \tag{8.29}$$

holomorphic on $|\Im v| < \frac12$. The identity $\chi(\frac12+iv) = e^{-2i\vartheta(v)}$ holds for real v, hence on the strip, and $s = \frac12 + i\tau_*$ with $|\Im\tau_*| = \sigma' < \frac12$. Stirling's formula in the strip gives $\vartheta(v) = \frac{v}{2}\operatorname{Log}\frac{v}{2\pi} - \frac{v}{2} - \frac{\pi}{8} + O(|v|^{-1})$, i.e.

$$\mathcal{N}_{\mathrm{RVM}}(v) = \frac{\vartheta(v)}{\pi} + 1 + O(|v|^{-1}). \tag{8.30}$$

Therefore

$$w = e^{-2i\vartheta(\tau_*) - 2\pi i + O(1/t)} = \chi(s)\big(1 + O(t^{-1})\big). \tag{8.31}$$

For $\sigma' \ge \sigma'_1$, (8.3) gives $|\chi(s)| \le \frac12$ when $t \ge t_0(\sigma'_1)$, so $1 + w = (1+\chi)(1 + (w-\chi)/(1+\chi)) = (1+\chi)(1+O(t^{-1}))$, which with (8.4) gives (8.5). This completes the proof of Theorem 8.1. □

Lemma 8.5 (Extension to the critical line)**.** *The function $\mathcal{E}_0 = \zeta_{\mathrm{RVM}}/(\varpi_{\mathrm{RVM}}(1+w)) - 1$ extends holomorphically to $\{|\sigma'| < \frac12,\ t > t_0\}$, and $|\mathcal{E}_0| \le C/t$ for $0 \le \sigma' \le \frac14$.*

Proof. By Lemma 8.3, $\Im\mathcal{N}_{\mathrm{RVM}}(\tau_*) \neq 0$ for $\sigma' \neq 0$, so $1+w$ vanishes in the strip only on $\sigma' = 0$, exactly at $t = \tau_k$; these zeros are simple because $\frac{d}{dt}w = -\log(\tau_*/2\pi)\, w \neq 0$. The zeros of ζ_{RVM} in the strip are the same simple points. Hence $\mathcal{E}_0$ has only removable singularities, and the bound extends to $\sigma' = 0$ by continuity. □

Corollary 8.6 (Derivative at the zeros)**.** *For k with $\tau_k > t_0$,*

$$\zeta'_{\mathrm{RVM}}(\rho_k^{\mathrm{RVM}}) = \varpi_{\mathrm{RVM}} \log\frac{\tau_k}{2\pi}\Big(1 + O\Big(\frac{1}{\tau_k}\Big)\Big). \tag{8.32}$$

Proof. By Lemma 8.5, $\zeta_{\mathrm{RVM}} = \varpi_{\mathrm{RVM}}(1+w)(1+\mathcal{E}_0)$ holds holomorphically near $\sigma' = 0$. At $s = \rho_k^{\mathrm{RVM}}$, $1 + w = 0$, so $\zeta'_{\mathrm{RVM}}(\rho_k^{\mathrm{RVM}}) = \varpi_{\mathrm{RVM}} w'(\rho_k^{\mathrm{RVM}})(1 + \mathcal{E}_0(\rho_k^{\mathrm{RVM}}))$. From (8.2), $w'(s) = -2\pi i\, \mathcal{N}'_{\mathrm{RVM}}(\tau_*) \cdot (-i)\, w(s) = -\log(\tau_*/2\pi)\, w(s)$, and at the zero $w = -1$, $\tau_* = \tau_k$. Finally $|\mathcal{E}_0| \le C/\tau_k$. □

Remark 8.7 (Structure of the proof). The delicate points are handled as follows: the smooth part is computed with principal branches by the identity theorem (Lemma 8.2); the clock estimate is quantified (Lemma 8.3); the sawtooth series is replaced by its Abel means, termwise integration is justified for $r < 1$, and $r \uparrow 1$ is taken after the contour argument uniformly in r and z (Lemma 8.4); the orientation of the lower contour and the sign of the residue are written out in (8.24); the z-dependence of the rays is isolated in g_z and is $O(t^{-2})$, the rest is the explicit real constant ν_4; Stirling is used uniformly for $0 < \sigma' \le B$, and the only place where $\sigma' < \frac12$ enters is Step 3.

Corollary 8.8 (Vertical constant limit)**.** *For every fixed $\sigma > \frac12$,*

$$\zeta_{\mathrm{RVM}}(\sigma+it) \longrightarrow \varpi_{\mathrm{RVM}}, \qquad F_{\mathrm{RVM}}(\sigma+it) \longrightarrow L := \frac{1}{\varpi_{\mathrm{RVM}}} \qquad (|t| \to \infty). \tag{8.33}$$

More precisely, with $\sigma' = \sigma - \frac12$,

$$|w(\sigma+it)| = O_\sigma(|t|^{-\sigma'}), \tag{8.34}$$

and, with $\alpha_c := \min(c - \frac12, 1)$ for $c > \frac12$,

$$F_{\mathrm{RVM}}(c+it) - L = O_c(|t|^{-\alpha_c}). \tag{8.35}$$

Proof. Expanding the clock at $\tau_* = t - i\sigma'$ gives $\Im\mathcal{N}_{\mathrm{RVM}}(\tau_*) = -\frac{\sigma'}{2\pi}\log\frac{t}{2\pi} + O_\sigma(t^{-1})$, hence $|w| = e^{2\pi\Im\mathcal{N}_{\mathrm{RVM}}(\tau_*)} = O_\sigma(t^{-\sigma'})$. Inverting (8.4) for large t gives $F_{\mathrm{RVM}} = L(1 + O(|w|) + O(t^{-1}))$. □

8.2. **Bounds on the whole half-plane.** The constants of Theorem 8.1 depend on the width B of the band. The following estimate holds on the whole half-plane and does not use contours. Put

$$\mathfrak{s}_1(\tau) := \int_{\tau_0}^{\tau} \psi(\mathcal{N}_{\mathrm{RVM}}(v) + \tfrac{1}{2})\, dv \qquad (\tau \geq \tau_0). \tag{8.36}$$

Lemma 8.9 (Bounded primitive of the midpoint sawtooth)**.** $|\mathfrak{s}_1(\tau)| \leq \dfrac{\pi}{4\log 2}$ *for all* $\tau \geq \tau_0$.

Proof. Substitute $y = \mathcal{N}_{\mathrm{RVM}}(v)$, $dv = \mathcal{T}'(y)\, dy$ with $\mathcal{T}'(y) = 2\pi/\log(\mathcal{T}(y)/2\pi)$ from (4.4). The weight $\mathcal{T}'$ is positive and decreasing, with $\mathcal{T}' \leq 2\pi/\log 2$ on $y \geq \mathcal{N}_{\mathrm{RVM}}(\tau_0)$ because $\log(\tau_0/2\pi) = \log 2$. The primitive of $y \mapsto \psi(y + \frac{1}{2})$ is periodic with values in $[-\frac{1}{8}, 0]$, so every integral of $\psi(y + \frac{1}{2})$ over an interval has modulus at most $\frac{1}{8}$. The second mean-value theorem gives $|\mathfrak{s}_1(\tau)| \leq \frac{2\pi}{\log 2} \cdot \frac{1}{8}$. □

Theorem 8.10 (Two-sided bound and limit at $+\infty$)**.** *There is an absolute constant* R_0 *such that, for* $s = \frac{1}{2} + z$, $z = \sigma' + it$, $\sigma' > \frac{1}{2}$ *and* $|z| \geq R_0$,

$$\zeta_{\mathrm{RVM}}(s) = \varpi_{\mathrm{RVM}}\, e^{\theta_{\mathrm{RVM}}(s)}, \qquad |\theta_{\mathrm{RVM}}(s)| \leq \frac{\pi^2}{2\log 2} \cdot \frac{1}{\sigma'} + \frac{C}{|z|}. \tag{8.37}$$

Consequently, for every $\sigma_0 > 1$ *there is* $C(\sigma_0)$ *with*

$$C(\sigma_0)^{-1} \leq |\zeta_{\mathrm{RVM}}(s)| \leq C(\sigma_0) \qquad (\Re s \geq \sigma_0), \tag{8.38}$$

and $\zeta_{\mathrm{RVM}}(s) \to \varpi_{\mathrm{RVM}}$ *uniformly in* t *as* $\Re s \to +\infty$. *Moreover*

$$\nu_4 = -2 \int_{\tau_0}^{\infty} \frac{\mathfrak{s}_1(\tau)}{\tau^2}\, d\tau. \tag{8.39}$$

Proof. Smooth part. For $\Re z > 0$ the computation in the proof of Lemma 8.2 is exact:

$$\Lambda_{\mathrm{sm}}(z) = \frac{z}{2} \operatorname{Log} \frac{z}{2\pi e} + \frac{7}{8} \operatorname{Log}\Big(1 + \frac{z^2}{\tau_0^2}\Big) - \frac{1}{\pi} \int_0^{\tau_0} \log \frac{\tau}{2\pi e}\, \frac{z^2}{\tau^2 + z^2}\, d\tau. \tag{8.40}$$

For $|z| \geq 2\tau_0$ one has $|\tau^2 + z^2| \geq \frac{3}{4}|z|^2$ on $[0, \tau_0]$, and Stirling's formula is uniform in $|\arg(s/2)| < \frac{\pi}{2}$; the expansions (8.14) and (8.16) therefore hold with $O(|z|^{-1})$ in place of $O(t^{-1})$, uniformly on $\Re z > \frac{1}{2}$, and $\Lambda_{\mathrm{sm}}(z) - \log G(s) = \nu_3 + O(|z|^{-1})$.

Sawtooth part. Integrating (8.9) by parts with (8.36) ($\mathfrak{s}_1(\tau_0) = 0$, $f'(\tau) = O(|z|^2\tau^{-3})$),

$$\Lambda_{\mathrm{saw}}(z) = -\int_{\tau_0}^{\infty} \mathfrak{s}_1 f''\, d\tau, \qquad f''(\tau) = \frac{2}{\tau^2} - \frac{1}{(\tau - iz)^2} - \frac{1}{(\tau + iz)^2}. \tag{8.41}$$

Since $|\tau \mp iz|^2 = (\tau \pm t)^2 + \sigma'^2$ and $\int_{\mathbb{R}} \frac{d\tau}{(\tau - a)^2 + \sigma'^2} = \frac{\pi}{\sigma'}$, Lemma 8.9 gives

$$\Big|\Lambda_{\mathrm{saw}}(z) + 2 \int_{\tau_0}^{\infty} \frac{\mathfrak{s}_1(\tau)}{\tau^2}\, d\tau\Big| \leq \frac{\pi}{4\log 2} \cdot \frac{2\pi}{\sigma'}. \tag{8.42}$$

Assembly. By (8.27), $\log \zeta_{\mathrm{RVM}}(s) = \log \varpi_\infty + \theta_{\mathrm{RVM}}(s)$ with $\varpi_\infty := P_{\mathrm{RVM}}^{\mathrm{crit}}(\frac{1}{2}) \exp(\nu_3 - 2\int_{\tau_0}^{\infty} \mathfrak{s}_1 \tau^{-2} d\tau) > 0$ and θ_{RVM} bounded as in (8.37). For fixed σ' and $t \to \infty$, Corollary 8.8 gives $|\log \varpi_{\mathrm{RVM}} - \log \varpi_\infty| \leq \pi^2/(2\sigma' \log 2)$; letting $\sigma' \to \infty$, $\varpi_\infty = \varpi_{\mathrm{RVM}}$, which is (8.39) by (8.28). On the compact set $\{\Re s \geq \sigma_0,\ |z| \leq R_0\}$ the function ζ_{RVM} is continuous and has no zeros or poles, which gives (8.38). □

Theorem 8.10 extends the two-sided bound (9.9) of Corollary 9.3 from bands to the whole half-plane $\Re s \geq \sigma_0 > 1$, and identifies ϖ_{RVM} also as the limit of $\zeta_{\mathrm{RVM}}(\sigma)$ as $\sigma \to +\infty$.

Theorem 8.11 (No absolutely convergent Dirichlet expansion)**.** *Neither* ζ_{RVM} *nor* $F_{\mathrm{RVM}} = 1/\zeta_{\mathrm{RVM}}$ *is represented by a Dirichlet series* $\sum_n a_n n^{-s}$ *converging absolutely in some half-plane* $\Re s > \sigma_0$.

Proof. Suppose $F_{\mathrm{RVM}}(s) = \sum_{n\ge1} a_n n^{-s}$ absolutely for $\Re s > \sigma_0$. Fix $\sigma > \max(\sigma_0, \frac12)$ and put $\tilde a_1 = a_1 - L$, $\tilde a_n = a_n$ $(n \ge 2)$, so $F_{\mathrm{RVM}}(\sigma+it) - L = \sum_n \tilde a_n n^{-\sigma-it}$ with $\sum_n |\tilde a_n| n^{-\sigma} < \infty$. Then

$$\frac{1}{2T}\int_{-T}^{T} \big|F_{\mathrm{RVM}}(\sigma+it) - L\big|^2 dt = \sum_{m,n} \tilde a_m \overline{\tilde a_n} (mn)^{-\sigma} \, \frac{1}{2T}\int_{-T}^{T} \Big(\frac{n}{m}\Big)^{it} dt, \tag{8.43}$$

the interchange being justified by absolute convergence of the double series, uniformly in t. The inner average is 1 for $m = n$ and is bounded by $\min\big(1, 1/(T|\log(n/m)|)\big) \to 0$ for $m \ne n$; each term is bounded by $|\tilde a_m \tilde a_n|(mn)^{-\sigma}$, which is summable, so dominated convergence gives

$$\lim_{T\to\infty} \frac{1}{2T}\int_{-T}^{T} \big|F_{\mathrm{RVM}}(\sigma+it) - L\big|^2 dt = \sum_{n\ge1} |\tilde a_n|^2 n^{-2\sigma}. \tag{8.44}$$

By (8.33) the integrand tends to 0 and is bounded on the line (continuous, with a finite limit), so the left side of (8.44) is 0. Hence every $\tilde a_n = 0$, $F_{\mathrm{RVM}} \equiv L$ in the half-plane and, by analytic continuation, everywhere, which contradicts $F_{\mathrm{RVM}}(1) = 0$. The same argument applied to $\zeta_{\mathrm{RVM}} - \varpi_{\mathrm{RVM}}$ forces an absolutely convergent Dirichlet series for ζ_{RVM} to be the constant ϖ_{RVM}, which contradicts its pole at $s = 1$. □

Theorem 8.12 (Unsmoothed synthetic Mertens bound)**.** *For every $x > 1$ and every $c > \frac12$ the symmetric Perron integral*

$$M^{\mathrm{crit}}_{\mathrm{RVM}}(x) := \lim_{U\to\infty} \frac{1}{2\pi i}\int_{c-iU}^{c+iU} F_{\mathrm{RVM}}(s)\frac{x^s}{s}\, ds \tag{8.45}$$

exists, is independent of $c > \frac12$, and equals the absolutely convergent expression

$$M^{\mathrm{crit}}_{\mathrm{RVM}}(x) = L + \frac{1}{2\pi i}\int_{(c)} (F_{\mathrm{RVM}}(s) - L)\frac{x^s}{s}\, ds. \tag{8.46}$$

For every $\varepsilon \in (0, \frac12)$,

$$M^{\mathrm{crit}}_{\mathrm{RVM}}(x) = L + O_\varepsilon(x^{1/2+\varepsilon}). \tag{8.47}$$

Proof. Fix $c > \frac12$. By (8.35) and continuity of F_{RVM} on $\Re s = c$ (no poles there by Proposition 7.1), the integral in (8.46) converges absolutely. The Perron kernel gives, for $x > 1$ and $c > 0$ [12, Ch. 17],

$$\lim_{U\to\infty} \frac{1}{2\pi i}\int_{c-iU}^{c+iU} \frac{x^s}{s}\, ds = 1. \tag{8.48}$$

Splitting $F_{\mathrm{RVM}} = L + (F_{\mathrm{RVM}} - L)$ proves existence of (8.45) and (8.46). If $\frac12 < c < c'$, the integrand of (8.45) is holomorphic on the closed strip $c \le \Re s \le c'$ (Proposition 7.1); by Theorem 8.1(a) with $B = c' - \frac12$, F_{RVM} is bounded on the horizontal segments $\Im s = \pm U$, $U \ge t_0(B)$, so these contribute $O(x^{c'}U^{-1}) \to 0$, and Cauchy's theorem gives independence of c. Take $c = \frac12 + \varepsilon$ in (8.46):

$$|M^{\mathrm{crit}}_{\mathrm{RVM}}(x) - L| \le \frac{x^{1/2+\varepsilon}}{2\pi}\int_{-\infty}^{\infty} \frac{|F_{\mathrm{RVM}}(\frac12+\varepsilon+it) - L|}{|\frac12+\varepsilon+it|}\, dt, \tag{8.49}$$

and the integral is finite by (8.35), the integrand being $O(|t|^{-1-\alpha_c})$. □

Remark 8.13 (Scope). Theorem 8.12 concerns $1/\zeta_{\mathrm{RVM}}$. The transfer to M goes through the endpoint of the λ-family and is Theorem 11.8.

9. The exact Hadamard product and the comparison with ζ

Put

$$P_\zeta(s) := G(s)\zeta(s) = 2\xi(s), \qquad \xi(s) = \tfrac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s). \tag{9.1}$$

Theorem 9.1 (Exact Hadamard product)**.** *Let ρ run over the nontrivial zeros of ζ with $\Im\rho > 0$, with multiplicity, ordered by $\Im\rho$. The product*

$$P_\zeta(s) = \prod_{\Im\rho>0} \Big(1 + \frac{q(s)}{\rho(1-\rho)}\Big) \tag{9.2}$$

converges absolutely and locally uniformly in $s \in \mathbb{C}$, and the identity holds with no exponential or constant factor.

Proof. ξ is entire of order one with $\xi(0) = \frac{1}{2}$, so Hadamard's factorization [22], [37, §2.12], [12, Ch. 12] gives

$$2\xi(s) = e^{Bs}\prod_\rho \Big(1 - \frac{s}{\rho}\Big)e^{s/\rho}, \tag{9.3}$$

the product running over all nontrivial zeros and converging absolutely because $\sum_\rho |\rho|^{-2} < \infty$. The functional equation $\xi(s) = \xi(1-s)$ makes $\rho \mapsto 1-\rho$ an involution of the zero multiset; no nontrivial zero is real, since $(1-2^{1-\sigma})\zeta(\sigma) = \sum_{n\ge1}(-1)^{n-1}n^{-\sigma} > 0$ for $0 < \sigma < 1$; so this involution exchanges $\Im\rho > 0$ and $\Im\rho < 0$. Grouping ρ with $1-\rho$, which absolute convergence permits,

$$\Big(1 - \frac{s}{\rho}\Big)\Big(1 - \frac{s}{1-\rho}\Big)e^{s/\rho + s/(1-\rho)} = \Big(1 + \frac{q(s)}{\rho(1-\rho)}\Big)e^{s/(\rho(1-\rho))}. \tag{9.4}$$

Since $|\rho(1-\rho)| \ge (\Im\rho)^2$ and $\sum_{\Im\rho>0}(\Im\rho)^{-2} < \infty$, the numbers $1/(\rho(1-\rho))$ are absolutely summable, so the exponentials collect into a single factor:

$$2\xi(s) = e^{B's}\,\Pi_\zeta(s), \qquad \Pi_\zeta(s) := \prod_{\Im\rho>0}\Big(1 + \frac{q(s)}{\rho(1-\rho)}\Big), \qquad B' = B + \sum_{\Im\rho>0}\frac{1}{\rho(1-\rho)}. \tag{9.5}$$

Π_ζ depends on s only through $q(s)$, hence $\Pi_\zeta(1-s) = \Pi_\zeta(s)$. From $\xi(1-s) = \xi(s)$ we get $e^{B's} = e^{B'(1-s)}$ on the open set where $\Pi_\zeta \neq 0$, so $B' = 0$. □

Define the exact quotient

$$\mathcal{Q}_{\mathrm{RVM}}(s) := \frac{\zeta(s)}{\zeta_{\mathrm{RVM}}(s)} = \frac{P_\zeta(s)}{P^{\mathrm{crit}}_{\mathrm{RVM}}(s)}. \tag{9.6}$$

Proposition 9.2 (Exact transform relation)**.** *For $\Re s > 1$, and by meromorphic continuation,*

$$F_{\mathrm{RVM}}(s) = \mathcal{Q}_{\mathrm{RVM}}(s)\,\frac{1}{\zeta(s)}. \tag{9.7}$$

If the inverse Mellin transform $\mathcal{Q}^\vee_{\mathrm{RVM}}$ of $\mathcal{Q}_{\mathrm{RVM}}$ and the inverse defining $M^{\mathrm{crit}}_{\mathrm{RVM}}$ exist in a common distribution space with Mellin convolution, then

$$M^{\mathrm{crit}}_{\mathrm{RVM}} = \mathcal{Q}^\vee_{\mathrm{RVM}} *_{\mathrm{Mel}} M. \tag{9.8}$$

Proof. (9.7) is the identity $1/\zeta_{\mathrm{RVM}} = (\zeta/\zeta_{\mathrm{RVM}})(1/\zeta)$; (9.8) is the Mellin convolution theorem under the stated assumptions. □

Corollary 9.3 (All growth of ζ to the right of $\frac{1}{2}$ lies in $\mathcal{Q}_{\mathrm{RVM}}$)**.** *Let $0 < \varepsilon < c' - \frac{1}{2}$. There are t_0 and $0 < c_- \le c_+$ such that*

$$c_- \le |\zeta_{\mathrm{RVM}}(\sigma + it)| \le c_+ \qquad (\tfrac{1}{2} + \varepsilon \le \sigma \le c',\ |t| \ge t_0), \tag{9.9}$$

and therefore, in the same region,

$$c_-|\mathcal{Q}_{\mathrm{RVM}}(s)| \le |\zeta(s)| \le c_+|\mathcal{Q}_{\mathrm{RVM}}(s)|, \qquad \frac{1}{c_+|\mathcal{Q}_{\mathrm{RVM}}(s)|} \le \Big|\frac{1}{\zeta(s)}\Big| \le \frac{1}{c_-|\mathcal{Q}_{\mathrm{RVM}}(s)|}. \tag{9.10}$$

Consequently:

(i) *the Lindelöf hypothesis holds if and only if* $\mathcal{Q}_{\mathrm{RVM}}(\sigma+it) = O_{\sigma,\varepsilon}(|t|^{\varepsilon})$ *for every* $\sigma > \frac{1}{2}$*;*

(ii) *every bound for* ζ *or* $1/\zeta$ *on* $\frac{1}{2}+\varepsilon \le \sigma \le c'$, $|t| \ge t_0$*, is equivalent, up to the constants* $c_\pm$*, to the same bound for* $\mathcal{Q}_{\mathrm{RVM}}$ *or* $1/\mathcal{Q}_{\mathrm{RVM}}$*.*

Proof. By Theorem 8.1(a) with $B = c' - \frac{1}{2}$, $\zeta_{\mathrm{RVM}} = \varpi_{\mathrm{RVM}}(1+w)(1+\mathcal{E}_0)$ with $|\mathcal{E}_0| \le C_B/|t|$ and, by (8.34), $|w| \le C(|t|/2\pi)^{-\varepsilon}$; for $|t| \ge t_0$ this gives $\frac{1}{2} \le |1+w| \le 2$ and $\frac{1}{2} \le |1+\mathcal{E}_0| \le 2$, hence (9.9) with $c_- = \varpi_{\mathrm{RVM}}/4$, $c_+ = 4\varpi_{\mathrm{RVM}}$. Then (9.10) follows from $\zeta = \mathcal{Q}_{\mathrm{RVM}}\zeta_{\mathrm{RVM}}$. For (i): the Lindelöf hypothesis is equivalent to $\zeta(\sigma+it) = O(|t|^{\varepsilon})$ for every $\sigma > \frac{1}{2}$ [37, Ch. XIII], and (9.10) transfers this to $\mathcal{Q}_{\mathrm{RVM}}$. (ii) is (9.10). □

On the prime side define

$$\mathfrak{d}_n^{\mathrm{RVM}} := \mathbf{1}_{\{n \text{ prime}\}} - c_n^{\mathrm{RVM}}. \tag{9.11}$$

Since $\pi_{\mathrm{RVM}}(1) = 0$, for every integer $N \ge 2$

$$\sum_{2\le n\le N} \mathfrak{d}_n^{\mathrm{RVM}} = \pi(N) - \pi_{\mathrm{RVM}}(N). \tag{9.12}$$

10. The λ-family from ζ_{RVM} to ζ_{Z1}

For $\lambda \in [0,1]$ define the moving ordinates, the family product and the family zeta function

$$\gamma_k(\lambda) := (1-\lambda)\gamma_k + \lambda\tau_k = \gamma_k + \lambda\eta_k, \tag{10.1}$$

$$P_\lambda(s) := \prod_{k\ge1} \left(1 + \frac{q(s)}{\frac{1}{4}+\gamma_k(\lambda)^2}\right), \tag{10.2}$$

$$\zeta_\lambda(s) := \frac{P_\lambda(s)}{G(s)}, \qquad F_\lambda(s) := \frac{1}{\zeta_\lambda(s)}, \tag{10.3}$$

and the endpoint

$$P_{Z1} := P_0, \qquad \zeta_{Z1} := \zeta_0, \qquad \zeta_1 = \zeta_{\mathrm{RVM}}. \tag{10.4}$$

Theorem 10.1 (The λ-family)**.** *The product* (10.2) *converges absolutely and locally uniformly jointly on* $[0,1]\times\mathbb{C}$*. For every* λ*,* P_λ *is entire of order exactly one,* $P_\lambda(1-s) = P_\lambda(s)$*,* $P_\lambda(\bar{s}) = \overline{P_\lambda(s)}$*,* $P_\lambda(0) = P_\lambda(1) = 1$*, and its zeros, with multiplicity, are exactly*

$$\frac{1}{2} \pm i\gamma_k(\lambda), \qquad k \ge 1. \tag{10.5}$$

For $0 < \lambda \le 1$ *the ordinates* $\gamma_k(\lambda)$ *are strictly increasing in* k *and all zeros of* P_λ *are simple; at* $\lambda = 0$ *the multiplicities are those of the ordinate list* (γ_k)*. For every* $j \ge 0$*,* $P_\lambda^{(j)} \to P_{\lambda_0}^{(j)}$ *locally uniformly as* $\lambda \to \lambda_0$*. In particular no member* ζ_λ *of the family,* $\zeta_{Z1} = \zeta_0$ *included, can have a zero off the critical line other than* $-2, -4, \dots$*.*

Proof. By Lemma 2.2, $|\gamma_k(\lambda) - \gamma_k| \le C_\eta$, and $\gamma_k \asymp k/\log k$; so $\gamma_k(\lambda) \ge \frac{1}{2}\gamma_k$ for $k \ge k_0$, uniformly in λ. On a compact $K \subset \mathbb{C}$,

$$\sum_{k\ge1} \sup_{s\in K,\ 0\le\lambda\le1} \left|\frac{q(s)}{\frac{1}{4}+\gamma_k(\lambda)^2}\right| < \infty, \tag{10.6}$$

which gives joint normal convergence and, by Cauchy's formula, convergence of all s-derivatives. For $|s| \le R_0$, uniformly in λ,

$$\log \max_{|s|\le R_0} |P_\lambda(s)| \le \sum_{k\ge 1} \log\Big(1 + \frac{R_0^2 + R_0}{\frac{1}{4} + \gamma_k(\lambda)^2}\Big) = O(R_0 \log R_0), \tag{10.7}$$

so the order is at most one; the zero count is $\asymp T \log T$, so the exponent of convergence is one and the order is exactly one. Symmetries, normalization and divisor follow factor by factor as in Theorem 3.1. For $0 < \lambda \le 1$,

$$\gamma_{k+1}(\lambda) - \gamma_k(\lambda) = (1-\lambda)(\gamma_{k+1} - \gamma_k) + \lambda(\tau_{k+1} - \tau_k) \ge \lambda(\tau_{k+1} - \tau_k) > 0, \tag{10.8}$$

so distinct factors vanish at distinct points and each zero is simple; at $\lambda = 0$ equal ordinates give equal factors. □

Remark 10.2 (All differences lie in $\gamma_k - \tau_k$). For $\lambda, \lambda' \in [0,1]$ and s off the divisors,

$$\frac{\zeta_\lambda(s)}{\zeta_{\lambda'}(s)} = \prod_{k\ge 1} \frac{(\frac{1}{4} + (\gamma_k + \lambda'\eta_k)^2)\,(\frac{1}{4} + (\gamma_k + \lambda\eta_k)^2 + q(s))}{(\frac{1}{4} + (\gamma_k + \lambda\eta_k)^2)\,(\frac{1}{4} + (\gamma_k + \lambda'\eta_k)^2 + q(s))}. \tag{10.9}$$

The gamma factor cancels; the members differ only through $\eta_k = \tau_k - \gamma_k$, and all of them coincide if every η_k vanishes.

Theorem 10.3 (Infinitesimal generator)**.** *Put $A_{k,\lambda} := \frac{1}{4} + \gamma_k(\lambda)^2$. Off the divisor of P_λ,*

$$\mathcal{V}_\lambda(s) := \partial_\lambda \log P_\lambda(s) = -2q(s) \sum_{k\ge 1} \frac{\gamma_k(\lambda)\,\eta_k}{A_{k,\lambda}\big(A_{k,\lambda} + q(s)\big)}, \tag{10.10}$$

the series converging locally uniformly on compact sets separated from the divisor. Since G does not depend on λ,

$$\partial_\lambda \log \zeta_\lambda = \mathcal{V}_\lambda, \qquad \partial_\lambda \log F_\lambda = -\mathcal{V}_\lambda. \tag{10.11}$$

Proof. Since $\partial_\lambda A_{k,\lambda} = 2\gamma_k(\lambda)\eta_k$,

$$\partial_\lambda \log\Big(1 + \frac{q}{A_{k,\lambda}}\Big) = -\frac{2q\,\gamma_k(\lambda)\eta_k}{A_{k,\lambda}(A_{k,\lambda} + q)}. \tag{10.12}$$

On compact sets separated from the divisor the summand is $O_K(|\eta_k|\gamma_k^{-3})$, and $\sum_k \gamma_k^{-3} < \infty$, so termwise differentiation is justified. □

Proposition 10.4 (Integrated multiplier)**.** *On every component avoiding the moving divisor,*

$$\zeta_\lambda(s) = \zeta_{Z1}(s) \exp\Big(\int_0^\lambda \mathcal{V}_{\lambda'}(s)\,d\lambda'\Big), \tag{10.13}$$

$$F_\lambda(s) = F_{Z1}(s) \exp\Big(-\int_0^\lambda \mathcal{V}_{\lambda'}(s)\,d\lambda'\Big), \tag{10.14}$$

and the total multiplier is

$$\Pi_{\mathrm{tr}}(s) := \exp\Big(-\int_0^1 \mathcal{V}_{\lambda'}(s)\,d\lambda'\Big) = \frac{P_{Z1}(s)}{P_{\mathrm{RVM}}^{\mathrm{crit}}(s)}, \qquad \zeta_{Z1} = \zeta_{\mathrm{RVM}}\,\Pi_{\mathrm{tr}}. \tag{10.15}$$

The identities extend meromorphically to $\mathbb{C}$ through the product form.

Proof. (10.13) and (10.14) are the fundamental theorem of calculus applied to (10.11). For (10.15), integrate each mode with $r = \gamma_k(\lambda')$, $dr = \eta_k\,d\lambda'$:

$$\int_0^1 \partial_{\lambda'} \log\Big(1 + \frac{q}{A_{k,\lambda'}}\Big) d\lambda' \equiv \log \frac{1 + q/(\frac{1}{4} + \tau_k^2)}{1 + q/(\frac{1}{4} + \gamma_k^2)} \pmod{2\pi i}. \tag{10.16}$$

Normal convergence permits summation, and exponentiation removes the branch ambiguity. □

Corollary 10.5 (Expansion at the endpoint)**.** *Let $K \subset \mathbb{C}$ be compact and disjoint from the zeros of P_{Z1}. Uniformly for $s \in K$, as $\lambda \downarrow 0$,*

$$\zeta_\lambda(s) = \zeta_{Z1}(s)(1 + \lambda \mathcal{V}_0(s) + O_K(\lambda^2)), \tag{10.17}$$

$$F_\lambda(s) = F_{Z1}(s)(1 - \lambda \mathcal{V}_0(s) + O_K(\lambda^2)). \tag{10.18}$$

Proof. By Theorem 10.1 the zeros of P_λ converge to those of P_{Z1}, so K avoids the moving divisor for small λ. The series for $\mathcal{V}_\lambda$ and its λ-derivative converge normally on K (the differentiated summands are $O_K(\gamma_k^{-3})$), so $\mathcal{V}_\lambda = \mathcal{V}_0 + O_K(\lambda)$, and (10.13) gives $\log(\zeta_\lambda/\zeta_{Z1}) = \lambda \mathcal{V}_0 + O_K(\lambda^2)$. □

Proposition 10.6 (Residues along the family)**.** *Suppose γ_k is a simple ordinate (it occurs once in the list). Put $\rho_{k,\lambda} = \frac{1}{2} + i\gamma_k(\lambda)$ and, for $0 < \lambda \le 1$,*

$$\mathfrak{r}_{k,\lambda} := \operatorname*{Res}_{s=\rho_{k,\lambda}} \frac{1}{s\,\zeta_\lambda(s)} = \frac{G(\rho_{k,\lambda})}{\rho_{k,\lambda} P_\lambda'(\rho_{k,\lambda})}. \tag{10.19}$$

Then

$$\mathfrak{r}_{k,\lambda} \longrightarrow \frac{G(\frac{1}{2} + i\gamma_k)}{(\frac{1}{2} + i\gamma_k) P_{Z1}'(\frac{1}{2} + i\gamma_k)} \qquad (\lambda \downarrow 0). \tag{10.20}$$

Proof. Theorem 10.1 gives convergence of the zero locations and local uniform convergence of P_λ'; the limit denominator is nonzero because the endpoint zero is simple, and G is continuous. □

Remark 10.7 (Multiplicity). For $0 < \lambda \le 1$ every zero of ζ_λ is simple. At $\lambda = 0$ a repeated ordinate of ζ appears in ζ_{Z1} with the same multiplicity, and the residues of Proposition 10.6 are replaced by the full principal part. Multiplicity plays no role in Step 3: Theorem 11.8 uses only that ζ does not vanish in $\Re s > \frac{1}{2}$.

11. The endpoint ζ_{Z1}, Hamburger's theorem, and Step 3

11.1. The analytic class of the endpoint.

Proposition 11.1 (Analytic class of ζ_{Z1})**.** *The endpoint*

$$P_{Z1}(s) = \prod_{k \ge 1} \left(1 + \frac{q(s)}{\frac{1}{4} + \gamma_k^2}\right), \qquad \zeta_{Z1}(s) = \frac{P_{Z1}(s)}{G(s)}, \tag{11.1}$$

has the following properties.

(i) *P_{Z1} is entire of order exactly one, $P_{Z1}(1 - s) = P_{Z1}(s)$, $P_{Z1}(0) = P_{Z1}(1) = 1$;*

(ii) *the nontrivial zeros of ζ_{Z1} are exactly*

$$\frac{1}{2} \pm i\gamma_k, \qquad k \ge 1, \tag{11.2}$$

with the multiplicities of the ordinate list; its other zeros are $s = -2, -4, \ldots$. It is impossible for ζ_{Z1} to vanish at any point off the critical line other than $-2, -4, \ldots$: every factor of P_{Z1} vanishes only at $\frac{1}{2} \pm i\gamma_k$;

(iii) *ζ_{Z1} has a simple pole at $s = 1$ with residue 1;*

(iv) *ζ_{Z1} satisfies the functional equation of ζ:*

$$\pi^{-s/2}\Gamma(s/2)\zeta_{Z1}(s) = \pi^{-(1-s)/2}\Gamma((1-s)/2)\zeta_{Z1}(1-s); \tag{11.3}$$

(v) *$(s-1)\zeta_{Z1}(s)$ is entire of order exactly one.*

Proof. (i) and the divisor in (ii) are the case $\lambda = 0$ of Theorem 10.1; the trivial zeros and the absence of other zeros come from $1/G$ as in Theorem 3.1. For (iii), $P_{Z1}(1) = 1$ and

$$G(s) = (s-1) + O\big((s-1)^2\big) \qquad (s \to 1). \tag{11.4}$$

For (iv),

$$\pi^{-s/2}\Gamma(s/2)\zeta_{Z1}(s) = \frac{P_{Z1}(s)}{q(s)}, \tag{11.5}$$

and both P_{Z1} and q are invariant under $s \mapsto 1-s$. For (v),

$$(s-1)\zeta_{Z1}(s) = P_{Z1}(s)\,\frac{\pi^{s/2}}{s\,\Gamma(s/2)}, \tag{11.6}$$

where $s\Gamma(s/2) \to 2$ at $s=0$ and $1/\Gamma$ is entire of order one; the product has order at most one, and the zero count $\asymp T\log T$ gives order at least one. □

Define the ratio

$$\mathscr{R}_{Z1}(s) := \frac{\zeta_{Z1}(s)}{\zeta(s)} = \frac{P_{Z1}(s)}{P_\zeta(s)}, \tag{11.7}$$

a meromorphic function; $\mathscr{R}_{Z1} \equiv 1$ is the identity $\zeta_{Z1} = \zeta$.

11.2. **Hamburger's theorem at the endpoint.**

Theorem 11.2 (Hamburger closure)**.** *Consider the condition*

$$\textit{(D)} \qquad \zeta_{Z1}(s) = \sum_{n\ge 1}\frac{a_n}{n^s} \qquad \textit{absolutely for } \Re s > 1. \tag{11.8}$$

Then

$$\text{(D)} \iff \mathscr{R}_{Z1} \equiv 1 \iff \zeta_{Z1} = \zeta \iff \text{RH}. \tag{11.9}$$

Proof. Assume (D). Then $\zeta_{Z1}(1-s) = \sum_n a_n n^{-(1-s)}$ converges absolutely for $\Re s < 0$, $(s-1)\zeta_{Z1}$ is entire of finite order (Proposition 11.1(v)), and (11.3) holds. These are the hypotheses of Hamburger's theorem [23], [37, §2.13], which gives

$$\zeta_{Z1} = C_H\,\zeta \tag{11.10}$$

for a constant C_H. Both functions have residue 1 at $s=1$, so $C_H = 1$. If $\zeta_{Z1} = \zeta$, the nontrivial zeros of ζ are those of ζ_{Z1}, namely (11.2), which is RH. If RH holds, every ρ with $\Im\rho > 0$ is $\frac12 + i\gamma_k$, so $\rho(1-\rho) = \frac14 + \gamma_k^2$, and Theorem 9.1 gives

$$P_\zeta(s) = \prod_{k\ge 1}\Big(1 + \frac{q(s)}{\frac14 + \gamma_k^2}\Big) = P_{Z1}(s) \tag{11.11}$$

exactly, hence $\zeta = \zeta_{Z1}$. Finally $\zeta = \sum_n n^{-s}$ converges absolutely for $\Re s > 1$, so $\zeta_{Z1} = \zeta$ gives (D). □

11.3. **Condition (D) and the integer-weighted zeta.** By Corollary 6.3, $\zeta_{\mathbb{Z},\mathrm{RVM}}$ is an absolutely convergent Dirichlet series in $\Re s > 1$ with first coefficient 1, unconditionally. Define the multiplier

$$\mathcal{K}(s) := \frac{\zeta_{Z1}(s)}{\zeta_{\mathbb{Z},\mathrm{RVM}}(s)}. \tag{11.12}$$

Theorem 11.3 (Reduction of (D) to the multiplier)**.** (i) *If $\mathcal{K}$ has an expansion $\mathcal{K}(s) = \sum_{n\ge1} k_n n^{-s}$ that converges absolutely in $\Re s > 1$, then* (11.8) *holds.*

(ii) *If* (11.8) *holds, then $\mathcal{K}(s) = \sum_{n\ge1} k_n n^{-s}$ absolutely on the whole half-plane $\Re s > 1$.*

Thus (D) holds if and only if $\mathcal{K}$ is an absolutely convergent Dirichlet series in $\Re s > 1$. Hence, by Theorem 11.2, an absolutely convergent Dirichlet expansion of $\mathcal{K}$ in $\Re s > 1$ gives $\zeta_{Z1} = \zeta$, RH, and, by Theorem 11.8, $M(x) = O_\varepsilon(x^{1/2+\varepsilon})$.

Proof. (i) $\zeta_{Z1} = \zeta_{\mathbb{Z},\mathrm{RVM}}\,\mathcal{K}$ is the Dirichlet convolution of two absolutely convergent series in $\Re s > 1$, and $\sum_N |(a^{\mathbb{Z}} * k)_N| N^{-\sigma} \le \sum_N |a^{\mathbb{Z}}_N| N^{-\sigma} \sum_n |k_n| n^{-\sigma} < \infty$. (ii) By (6.15), $1/\zeta_{\mathbb{Z},\mathrm{RVM}}$ is an absolutely convergent Dirichlet series in $\Re s > 1$, and so is its product $\mathcal{K}$ with (11.8). □

By (10.15) and (6.19) (Theorem 6.4),

$$\mathcal{K}(s) = \Pi_{\mathrm{tr}}(s)\, e^{-h_{\mathrm{RVM}}(s)} = \frac{P_{Z1}(s)}{P^{\mathrm{crit}}_{\mathrm{RVM}}(s)}\, e^{-h_{\mathrm{RVM}}(s)}, \tag{11.13}$$

an explicit function of the displacements $\gamma_k - \tau_k$ and of the cell masses c_n^{RVM}. Condition (D) is thereby located in a single object built from the two constructions of this paper: the λ-family and the integer weights.

Remark 11.4 (Continuous-to-atomic meaning of (D))*.* The prime-power count J^{M}_{Z1} of ζ_{Z1} (Theorem 12.5) is a continuous object built from the zeros, and (D) makes its measure the atomic measure $\sum_{p,m} m^{-1}\delta_{p^m}$. The final step of the method is therefore the passage of ζ_{Z1} into the discrete Dirichlet category. Theorem 12.7 states it exactly: (D) holds if and only if J^{M}_{Z1} is constant between consecutive integers, and Corollary 12.3 reduces it to the single identity (12.12) between the displacements $\gamma_k - \tau_k$ and ϖ_{RVM}.

11.4. **Step 3: the Mertens bound from $\zeta_{Z1} = \zeta$.**

Lemma 11.5 (Vertical growth of ζ)**.** *For $\frac{1}{4} \le \sigma \le 4$ and $|t| \ge 2$,*

$$|\zeta(\sigma + it)| \le 15|t|. \tag{11.14}$$

Proof. For $\sigma > 0$ [37, (2.1.5)],

$$\zeta(s) = \frac{s}{s-1} - s\int_1^\infty \{y\} y^{-s-1}\, dy. \tag{11.15}$$

For $|t| \ge 2$ and $\sigma \le 4$ one has $|s| \le 3|t|$, $|s/(s-1)| \le |s|/|t| \le 3$, and the integral is at most $1/\sigma \le 4$; hence $|\zeta(s)| \le 3 + 12|t| \le 15|t|$. □

Lemma 11.6 (Logarithm of ζ off the critical line)**.** *Assume $\zeta(s) \ne 0$ for $\Re s > \frac{1}{2}$. Let $0 < \delta \le \frac{1}{4}$ and*

$$\omega(\delta) := \frac{\log(3 - 2\delta)}{\log(3 - \delta)} < 1. \tag{11.16}$$

Then, with the branch of $\log\zeta$ equal to $\sum_{n\ge 2}\Lambda(n)n^{-s}/\log n$ on $\Re s > 1$,

$$|\log\zeta(\sigma + it)| \le \Big(\frac{48}{\delta}\log|t|\Big)^{\omega(\delta)} \qquad (\sigma \ge \tfrac{1}{2} + \delta,\ |t| \ge 4). \tag{11.17}$$

Consequently, for every $\varepsilon > 0$ there is $t_2 = t_2(\delta, \varepsilon)$ with

$$\Big|\frac{1}{\zeta(\sigma + it)}\Big| \le |t|^{\varepsilon} \qquad (\sigma \ge \tfrac{1}{2} + \delta,\ |t| \ge t_2). \tag{11.18}$$

Proof. Fix t with $|t| \ge 4$ and put $s_0 = 2 + it$, $R = \frac{3}{2} - \frac{\delta}{4}$, $r_3 = \frac{3}{2} - \frac{\delta}{2}$. The closed disc $|s - s_0| \le R$ lies in $\frac{1}{2} + \frac{\delta}{4} \le \Re s \le \frac{7}{2}$, $|\Im s - t| \le \frac{3}{2}$; it avoids $s = 1$ and, by hypothesis, every zero of ζ. Hence $\log\zeta$ has a holomorphic branch on a neighbourhood of it, equal to the Dirichlet series on its intersection with $\Re s > 1$, which is convex. On the disc $|\Im s| \ge |t|/2 \ge 2$ and $|\Im s| \le 2|t|$, so Lemma 11.5 gives $\Re\log\zeta = \log|\zeta| \le \log(30|t|)$ there, while $|\log\zeta(s_0)| \le \log\zeta(2)$. The Borel–Carathéodory lemma [36, §5.5], applied with radii $r_3 < R$,

$$\max_{|s-s_0|\le r_3} |\log\zeta(s)| \le \frac{2r_3}{R - r_3} \max_{|s-s_0|=R} \Re\log\zeta(s) + \frac{R + r_3}{R - r_3}|\log\zeta(s_0)|, \tag{11.19}$$

together with $R - r_3 = \delta/4$, $2r_3 \le 3$, $R + r_3 \le 3$ and $\log 30 + \log\zeta(2) < 4 \le 3\log|t|$, gives

$$\max_{|s-s_0|\le r_3} |\log\zeta(s)| \le \frac{12}{\delta}\big(\log(30|t|) + \log\zeta(2)\big) \le \frac{48}{\delta}\log|t|. \tag{11.20}$$

Now let $\frac12+\delta\le\sigma<\frac32$ and set $r_1=\frac12$, $r_2=2-\sigma\in(\frac12,\frac32-\delta]$. On $|s-s_0|=r_1$ one has $\Re s\ge\frac32$, so $|\log\zeta(s)|\le\log\zeta(\frac32)<1$. Hadamard's three-circle theorem [36, §5.3] gives, with $M_j=\max_{|s-s_0|=r_j}|\log\zeta|$,

$$M_2\le M_1^{1-\omega}M_3^{\omega},\qquad \omega=\frac{\log(r_2/r_1)}{\log(r_3/r_1)}\le\frac{\log(3-2\delta)}{\log(3-\delta)}=\omega(\delta). \tag{11.21}$$

Since $M_1<1$ and $M_3\le\frac{48}{\delta}\log|t|$ with $\frac{48}{\delta}\log|t|>1$, (11.21) gives (11.17) at $s=\sigma+it$, which lies on the circle of radius r_2. For $\sigma\ge\frac32$, $|\log\zeta|\le\log\zeta(\frac32)<1$. Finally $|1/\zeta|=\exp(-\Re\log\zeta)\le\exp|\log\zeta|$, and $(\frac{48}{\delta}\log|t|)^{\omega(\delta)}\le\varepsilon\log|t|$ for $|t|\ge t_2(\delta,\varepsilon)$ because $\omega(\delta)<1$. □

Lemma 11.7 (Perron kernel)**.** *For $y>0$, $y\neq1$, $c>0$, $T>0$* [12, Ch. 17],

$$\left|\frac{1}{2\pi i}\int_{c-iT}^{c+iT}\frac{y^s}{s}\,ds-\mathbf{1}_{\{y>1\}}\right|\le y^c\min\Big(1,\frac{1}{T|\log y|}\Big). \tag{11.22}$$

Theorem 11.8 (Step 3)**.** *If $\zeta_{Z1}=\zeta$, then for every $\varepsilon>0$*

$$M(x)=O_\varepsilon(x^{1/2+\varepsilon})\qquad(x\ge1). \tag{11.23}$$

Proof. Non-vanishing. By Proposition 11.1(ii), ζ_{Z1} has no zeros in $\Re s>\frac12$; since $\zeta=\zeta_{Z1}$, neither has ζ, and Lemma 11.6 applies.

Truncation. Let $x=N+\frac12$ with $N\ge3$ an integer, $c=1+1/\log x$ and $T=x$. Applying (11.22) with $y=x/n$ to each term of $\sum_n\mu(n)n^{-s}=1/\zeta(s)$, which converges absolutely on $\Re s=c$, gives

$$M(x)=\frac{1}{2\pi i}\int_{c-iT}^{c+iT}\frac{x^s}{s\,\zeta(s)}\,ds+O\Big(\sum_{n\ge1}\Big(\frac{x}{n}\Big)^c\min\Big(1,\frac{1}{T|\log(x/n)|}\Big)\Big). \tag{11.24}$$

For $n\le x/2$ or $n\ge2x$, $|\log(x/n)|\ge\log2$, and these terms contribute at most $x^c\zeta(c)/(T\log2)\le e\,x(1+\log x)/(T\log2)$. For $x/2<n<2x$, $(x/n)^c\le4$ and $|\log(x/n)|\ge|n-x|/(2x)$; as $|n-x|\in\{j-\frac12:j\ge1\}$, each value taken at most twice, these terms contribute at most $8\sum_{1\le j\le2x}\min(1,4x/(T(2j-1)))\ll1+(x/T)\log x$. Hence

$$M(x)=\frac{1}{2\pi i}\int_{c-iT}^{c+iT}\frac{x^s}{s\,\zeta(s)}\,ds+O(\log x). \tag{11.25}$$

Contour shift. Fix $0<\delta\le\frac14$. The integrand is holomorphic on the closed rectangle with vertices $c\pm iT$, $\frac12+\delta\pm iT$: ζ has no zeros there, $s=1$ is a zero of $1/\zeta$, and $s=0$ lies outside. By Cauchy's theorem

$$\int_{c-iT}^{c+iT}=\int_{\frac12+\delta-iT}^{\frac12+\delta+iT}+\int_{\frac12+\delta+iT}^{c+iT}-\int_{\frac12+\delta-iT}^{c-iT}. \tag{11.26}$$

For $x\ge t_2(\delta,\varepsilon)$, (11.18) on $|t|=T=x$ bounds the horizontal integrals by

$$\frac{1}{2\pi}\int_{\frac12+\delta}^{c}\frac{x^\sigma T^\varepsilon}{T}\,d\sigma\le x^cT^{\varepsilon-1}=e\,x^\varepsilon. \tag{11.27}$$

On the left side, $1/\zeta$ is continuous on the compact segment $\{\frac12+\delta+it:|t|\le t_2\}$, bounded by a constant $C(\delta,\varepsilon)$, and (11.18) holds for $|t|\ge t_2$; hence

$$\frac{1}{2\pi}\int_{-T}^{T}\frac{x^{1/2+\delta}}{|\frac12+\delta+it|\,|\zeta(\frac12+\delta+it)|}\,dt\le x^{1/2+\delta}\Big(C'(\delta,\varepsilon)+\int_{t_2}^{T}t^{\varepsilon-1}dt\Big)\ll_{\delta,\varepsilon}x^{1/2+\delta+\varepsilon}. \tag{11.28}$$

Conclusion. (11.25)–(11.28) give $M(x)\ll_{\delta,\varepsilon}x^{1/2+\delta+\varepsilon}$ for $x=N+\frac12$; taking $\delta=\varepsilon=\varepsilon_0/2$ (with $\varepsilon_0\le\frac12$) gives $O_{\varepsilon_0}(x^{1/2+\varepsilon_0})$. For arbitrary $x\ge3$, $M(x)=M(\lfloor x\rfloor+\frac12)$ and $\lfloor x\rfloor+\frac12\le2x$; for $1\le x<3$, $|M(x)|\le2$. □

Theorem 11.9 (Transfer of the synthetic Mertens bound to M)**.** *Assume $\zeta_{Z1}=\zeta$ and let $0<\delta\le\frac14$, $\varepsilon>0$. Then:*

(i) *on $\Re s>\frac12$,*

$$\frac{1}{\zeta(s)}=F_{\mathrm{RVM}}(s)\,\Pi_{\mathrm{tr}}(s)^{-1},\qquad \Pi_{\mathrm{tr}}=\frac{P_{Z1}}{P^{\mathrm{crit}}_{\mathrm{RVM}}}=\frac{\zeta}{\zeta_{\mathrm{RVM}}},\tag{11.29}$$

where Π_{tr} is holomorphic and zero-free in $\Re s>\frac12$ and $\Pi_{\mathrm{tr}}(1)=1$;

(ii) *there is $t_3=t_3(\delta,\varepsilon)$ such that, for $\frac12+\delta\le\sigma\le2$ and $|t|\ge t_3$,*

$$|F_{\mathrm{RVM}}(\sigma+it)|\le 2L,\qquad |\Pi_{\mathrm{tr}}(\sigma+it)^{\pm1}|\le|t|^{\varepsilon};\tag{11.30}$$

(iii) *the synthetic bound of Theorem 8.12, $M^{\mathrm{crit}}_{\mathrm{RVM}}(x)=L+O_\varepsilon(x^{1/2+\varepsilon})$, carries over to the Mertens function:*

$$M(x)=O_\varepsilon\big(x^{1/2+\varepsilon}\big).\tag{11.31}$$

Proof. (i) By (10.15), $\zeta_{Z1}=\zeta_{\mathrm{RVM}}\Pi_{\mathrm{tr}}$; with $\zeta_{Z1}=\zeta$ this is (11.29). In $\Re s>\frac12$, $\zeta=\zeta_{Z1}$ has no zeros (Proposition 11.1(ii)) and ζ_{RVM} has none (Theorem 3.1); both have a simple pole of residue 1 at $s=1$, so Π_{tr} is holomorphic there with $\Pi_{\mathrm{tr}}(1)=1$, and it is zero-free in $\Re s>\frac12$.

(ii) By Theorem 8.1(a) with $B=\frac32$, for $\delta\le\sigma'\le\frac32$ and $|t|\ge t_0$ we have $\zeta_{\mathrm{RVM}}=\varpi_{\mathrm{RVM}}(1+w)(1+\mathcal{E}_0)$ with $|w|\le C|t|^{-\delta}\le\frac12$ and $|\mathcal{E}_0|\le\frac12$ for $|t|$ large; hence $|F_{\mathrm{RVM}}|\le2L$ and

$$\log\zeta_{\mathrm{RVM}}=\log\varpi_{\mathrm{RVM}}+\mathrm{Log}(1+w)+\mathrm{Log}(1+\mathcal{E}_0)=O(1).\tag{11.32}$$

By Lemma 11.6, which applies because $\zeta=\zeta_{Z1}$ has no zeros in $\Re s>\frac12$, $|\log\zeta(\sigma+it)|\le(\frac{48}{\delta}\log|t|)^{\omega(\delta)}$ with $\omega(\delta)<1$. Therefore

$$|\log\Pi_{\mathrm{tr}}(\sigma+it)|=|\log\zeta-\log\zeta_{\mathrm{RVM}}|\le\Big(\frac{48}{\delta}\log|t|\Big)^{\omega(\delta)}+O(1)\le\varepsilon\log|t|\tag{11.33}$$

for $|t|\ge t_3(\delta,\varepsilon)$, which gives (11.30).

(iii) The proof of Theorem 8.12 moves the Perron line of $F_{\mathrm{RVM}}x^s/s$ to $\Re s=\frac12+\delta$ using only the holomorphy of F_{RVM} in $\Re s>\frac12$ and the vertical control of F_{RVM}. By (11.29) the integrand for M is the same integrand multiplied by Π_{tr}^{-1}, which is holomorphic in $\Re s>\frac12$ by (i) and at most $|t|^\varepsilon$ by (ii). Hence $|1/\zeta(\sigma+it)|\le2L|t|^\varepsilon$ for $\frac12+\delta\le\sigma\le2$, $|t|\ge t_3$, and the truncated Perron formula, contour shift and estimates (11.25)–(11.28) of Theorem 11.8, applied with this bound, give $M(x)\ll_{\delta,\varepsilon}x^{1/2+\delta+\varepsilon}$. Choosing $\delta=\varepsilon=\varepsilon_0/2$ gives (11.31). □

Remark 11.10 (The transfer in Mellin form). Formally, (11.29) reads $M=(\Pi_{\mathrm{tr}}^{-1})^\vee*_{\mathrm{Mel}}M^{\mathrm{crit}}_{\mathrm{RVM}}$: the Mertens function is the synthetic Mertens function convolved with the inverse Mellin transform of the inverse transport multiplier. The synthetic factor F_{RVM} contributes the exponent $\frac12$ and is bounded (Theorem 8.1); the multiplier contributes at most $|t|^\varepsilon$. All the information of the λ-family enters through $\Pi_{\mathrm{tr}}=P_{Z1}/P^{\mathrm{crit}}_{\mathrm{RVM}}$, that is, through the displacements $\gamma_k-\tau_k$.

Corollary 11.11 (Step 3 through (D))**.** *Condition* (11.8) *implies $M(x)=O_\varepsilon(x^{1/2+\varepsilon})$ for every $\varepsilon>0$.*

Proof. Theorem 11.2 and Theorem 11.8. □

Remark 11.12 (Status of Step 3). The implication $\zeta_{Z1}=\zeta\Rightarrow M(x)=O_\varepsilon(x^{1/2+\varepsilon})$ is proved completely above, and it requires nothing beyond the identity: no separate statement about the zeros of ζ enters, because $\zeta=\zeta_{Z1}$ already carries the divisor (11.2). The identity $\zeta_{Z1}=\zeta$ itself, equivalently (D), equivalently an absolutely convergent Dirichlet expansion of the multiplier $\mathcal{K}$ of (11.12) in $\Re s>1$ (Theorem 11.3), is not proved in this paper. It is the open step of the method.

11.5. **Reciprocal endpoint and finite-height Perron integrals.** Put

$$F_{Z1}(s) := \frac{1}{\zeta_{Z1}(s)} = F_{\mathrm{RVM}}(s)\,\Pi_{\mathrm{tr}}(s)^{-1}, \tag{11.34}$$

the second identity by (10.15). For $c > 1$ and finite U define

$$M_{\lambda,U}(x) := \frac{1}{2\pi i}\int_{c-iU}^{c+iU} F_\lambda(s)\frac{x^s}{s}\,ds, \qquad M_{Z1,U} := M_{0,U}. \tag{11.35}$$

Proposition 11.13 (Finite-height Perron integrals along the family)**.** *For fixed $c > 1$, $U < \infty$ and $x > 0$,*

$$M_{\lambda,U}(x) \longrightarrow M_{Z1,U}(x) \qquad (\lambda \downarrow 0). \tag{11.36}$$

Proof. The segment $\{c + it : |t| \le U\}$ is compact and disjoint from every divisor (10.5), and $P_\lambda \to P_{Z1}$ uniformly on it, so $F_\lambda \to F_{Z1}$ uniformly there. □

Theorem 11.14 (Reciprocal-coefficient criterion)**.** *Assume that in $\Re s > 1$*

$$F_{Z1}(s) = \sum_{n\ge1} \frac{\mu_{Z1}(n)}{n^s} \tag{11.37}$$

absolutely, and let $M_{Z1}(x) = \sum_{n\le x} \mu_{Z1}(n)$, so that

$$\mu_{Z1}(n) = M_{Z1}(n) - M_{Z1}(n-1). \tag{11.38}$$

Then the following are equivalent: (i) $M_{Z1}(n) = M(n)$ for every $n \ge 1$; (ii) $\mu_{Z1} = \mu$; (iii) $F_{Z1} = 1/\zeta$ in $\Re s > 1$; (iv) $\zeta_{Z1} = \zeta$; (v) RH.

Proof. (i)⇔(ii) by (11.38); (ii)⇔(iii) by uniqueness of absolutely convergent Dirichlet coefficients; (iii)⇔(iv) by analytic continuation, both functions being nonzero in $\Re s > 1$; (iv)⇔(v) by Theorem 11.2. □

Remark 11.15 (A route that does not close Step 3). The following approach does not work. One may try to expand both factors of $F_{Z1} = F_{\mathrm{RVM}}\,\Pi_{\mathrm{tr}}^{-1}$ in (11.34) as absolutely convergent Dirichlet series and identify their Dirichlet convolution with μ. By Theorem 8.11 the expansion of F_{RVM} does not exist in any half-plane, so there are no coefficients to convolve. The endpoint criterion concerns ζ_{Z1} itself, as in Theorem 11.2.

12. The endpoint at infinity and the Euler products

This section applies the clock decomposition of Theorem 8.10 to the ordinates γ_k of the endpoint, and constructs the prime-power counts of ζ_{RVM} and ζ_{Z1} as Mellin transforms of their logarithms. No hypothesis is used.

12.1. **The value of the endpoint at infinity.** With the zero count N_ζ of (2.2) put

$$\mathfrak{e}(\tau) := N_\zeta(\tau) - \mathcal{N}_{\mathrm{RVM}}(\tau), \qquad \mathfrak{e}_1(\tau) := \int_{\tau_0}^{\tau} \mathfrak{e}(v)\,dv \qquad (\tau \ge \tau_0 = 4\pi). \tag{12.1}$$

By (2.2), $\mathfrak{e}(\tau) = S(\tau) + O(1/\tau)$, and Littlewood's bound $\int_0^T S(t)\,dt = O(\log T)$ [37, Thm. 9.9] gives an absolute constant $C_{\mathfrak{e}}$ with

$$|\mathfrak{e}_1(\tau)| \le C_{\mathfrak{e}} \log\tau \qquad (\tau \ge \tau_0). \tag{12.2}$$

For the clock the corresponding primitive is $-\mathfrak{s}_1$, bounded by Lemma 8.9.

Theorem 12.1 (Clock decomposition of the endpoint)**.** *Let $s = \frac{1}{2} + z$, $z = \sigma' + it$, $\sigma' > 0$. Then*

$$\log \zeta_{Z1}(s) = \log \varpi_{Z1} - \int_{\tau_0}^{\infty} \mathfrak{e}_1(\tau)\Big(\frac{1}{(\tau - iz)^2} + \frac{1}{(\tau + iz)^2}\Big)d\tau + O\Big(\frac{1}{|z|}\Big) \qquad (|z| \geq R_0), \tag{12.3}$$

where

$$\varpi_{Z1} := P_{Z1}(\tfrac{1}{2}) \exp\Big(\nu_3 + 2\int_{\tau_0}^{\infty} \frac{\mathfrak{e}_1(\tau)}{\tau^2}\, d\tau\Big) > 0. \tag{12.4}$$

Consequently:

(i) *for $\sigma' > 0$ and $|z| \geq R_0$,*

$$|\log \zeta_{Z1}(s) - \log \varpi_{Z1}| \leq \frac{2(\pi+2)C_{\mathfrak{e}}}{\sigma'}\Big(1 + \log(2|t| + 2\sigma' + \tau_0)\Big) + \frac{C}{|z|}; \tag{12.5}$$

(ii) *$\zeta_{Z1}(\sigma) \to \varpi_{Z1}$ as $\sigma \to +\infty$;*

(iii) *the value at infinity is the ordered product*

$$\varpi_{Z1} = \varpi_{\mathrm{RVM}} \prod_{k\geq 1} \frac{\frac{1}{4} + \tau_k^2}{\frac{1}{4} + \gamma_k^2} := \varpi_{\mathrm{RVM}} \lim_{K\to\infty} \prod_{k \leq K} \frac{\frac{1}{4} + \tau_k^2}{\frac{1}{4} + \gamma_k^2}. \tag{12.6}$$

Proof. As in (8.6), $\log P_{Z1}(s) - \log P_{Z1}(\frac{1}{2}) = \sum_k f(\gamma_k)$ with $f(\tau) = \mathrm{Log}(1 + z^2/\tau^2)$. Since $\gamma_1 > \tau_0$, $N_\zeta(\tau_0) = 0$ and $\mathfrak{e}(\tau_0) = -\mathcal{N}_{\mathrm{RVM}}(\tau_0)$. Stieltjes integration against $N_\zeta = \mathcal{N}_{\mathrm{RVM}} + \mathfrak{e}$ and two integrations by parts ($f\mathfrak{e} \to 0$ and $\mathfrak{e}_1 f' \to 0$ at ∞, $\mathfrak{e}_1(\tau_0) = 0$) give

$$\sum_k f(\gamma_k) = \Lambda_{\mathrm{sm}}(z) + \int_{\tau_0}^{\infty} \mathfrak{e}_1 f''\, d\tau, \tag{12.7}$$

with the same Λ_{sm} as in (8.9). The proof of Theorem 8.10 gives $\Lambda_{\mathrm{sm}} - \log G = \nu_3 + O(|z|^{-1})$, and inserting f'' from (8.41) gives (12.3).

For (i), $|\tau \mp iz|^2 = (\tau \pm t)^2 + \sigma'^2$. For real a, splitting at $X = 2|a| + 2\sigma' + \tau_0$ and using $(\tau - a)^2 \geq \tau^2/4$ for $\tau \geq X$,

$$\int_{\tau_0}^{\infty} \frac{\log \tau}{(\tau - a)^2 + \sigma'^2}\, d\tau \leq \frac{\pi}{\sigma'} \log X + 4\,\frac{1 + \log X}{X} \leq \frac{\pi + 2}{\sigma'}(1 + \log X). \tag{12.8}$$

With (12.2) this gives (12.5). For (ii) take $t = 0$: the right side of (12.5) tends to 0 as $\sigma' \to \infty$.

For (iii), Theorem 8.10 gives $\varpi_{\mathrm{RVM}} = P^{\mathrm{crit}}_{\mathrm{RVM}}(\frac{1}{2}) \exp(\nu_3 - 2\int \mathfrak{s}_1 \tau^{-2} d\tau)$, so

$$\frac{\varpi_{Z1}}{\varpi_{\mathrm{RVM}}} = \frac{P_{Z1}(\frac{1}{2})}{P^{\mathrm{crit}}_{\mathrm{RVM}}(\frac{1}{2})} \exp\Big(2\int_{\tau_0}^{\infty} \frac{\mathfrak{e}_1(\tau) + \mathfrak{s}_1(\tau)}{\tau^2}\, d\tau\Big). \tag{12.9}$$

Since $\mathfrak{e} + \psi(\mathcal{N}_{\mathrm{RVM}} + \frac{1}{2}) = N_\zeta - n_{\mathrm{RVM}}$ and $(\mathfrak{e}_1 + \mathfrak{s}_1)(X)/X \to 0$,

$$\int_{\tau_0}^{\infty} \frac{\mathfrak{e}_1 + \mathfrak{s}_1}{\tau^2}\, d\tau = \lim_{X\to\infty} \int_{\tau_0}^{X} \frac{N_\zeta(\tau) - n_{\mathrm{RVM}}(\tau)}{\tau}\, d\tau = \lim_{K\to\infty} \sum_{k\leq K} \log \frac{\tau_k}{\gamma_k}. \tag{12.10}$$

In the last step, $\int_{\tau_0}^{X} (N_\zeta - n_{\mathrm{RVM}})\tau^{-1} d\tau = \sum_{\gamma_k \leq X} \log \frac{X}{\gamma_k} - \sum_{\tau_k \leq X} \log \frac{X}{\tau_k}$; the indices counted by only one of the two sums are $O(\log X)$ in number and, by Lemma 2.2, have ordinates in $[X - C_\eta, X]$, so they contribute $O(\log X/X)$. Finally $P_{Z1}(\frac{1}{2})/P^{\mathrm{crit}}_{\mathrm{RVM}}(\frac{1}{2}) = \prod_k \frac{\gamma_k^2}{\frac{1}{4}+\gamma_k^2} \cdot \frac{\frac{1}{4}+\tau_k^2}{\tau_k^2}$, an absolutely convergent product, and multiplying by $\prod_{k\leq K} \tau_k^2/\gamma_k^2$ gives (12.6). □

The ordinates enter ϖ_{Z1} only through the displacements $\gamma_k - \tau_k$, in agreement with Remark 10.2.

Theorem 12.2 (The value at infinity decides the endpoint identity)**.** *$\varpi_{Z1} \geq 1$, and $\varpi_{Z1} = 1$ if and only if $\zeta_{Z1} = \zeta$.*

Proof. Write $\zeta_{Z1} = \zeta\,\mathscr{R}_{Z1}$ with $\mathscr{R}_{Z1} = P_{Z1}/P_\zeta$ as in (11.7), and compare the products factor by factor, with the zeros $\rho_k = \beta_k + i\gamma_k$ listed with the ordinates (Theorem 9.1). Factors with $\rho_k(1-\rho_k) = \frac14 + \gamma_k^2$ cancel. The remaining factors of P_ζ come in pairs $\{\rho, 1-\bar\rho\}$ with the same ordinate γ; with $\delta = \Re\rho - \frac12$ and $W = (s-\frac12)^2 + \gamma^2$, the corresponding factor of $\mathscr{R}_{Z1}$ is

$$\frac{W^2}{(W-\delta^2)^2 + 4\delta^2\gamma^2}\cdot c_\gamma, \qquad c_\gamma = \frac{|\rho(1-\rho)|^2}{(\frac14+\gamma^2)^2} = 1 + \frac{2\delta^2(\gamma^2-\frac14)+\delta^4}{(\frac14+\gamma^2)^2} > 1. \tag{12.11}$$

For real $\sigma > 1$, $W \ge \gamma^2 \ge \gamma_1^2$, and the logarithm of the first factor is at most $14\delta^2/W \le 14\delta^2/\gamma^2$ in modulus; since $\delta \le \frac12$ and $\sum\gamma^{-2} < \infty$, dominated convergence and $\zeta(\sigma)\to 1$ give $\varpi_{Z1} = \lim_{\sigma\to\infty}\zeta_{Z1}(\sigma) = \prod c_\gamma \ge 1$, with equality exactly when there are no such pairs, that is, when $\mathscr{R}_{Z1} \equiv 1$. □

Corollary 12.3 (The endpoint identity as an identity of the displacements)**.** *The following are equivalent: (D); $\zeta_{Z1} = \zeta$; $\varpi_{Z1} = 1$;*

$$\prod_{k\ge1} \frac{\frac14+\gamma_k^2}{\frac14+\tau_k^2} = \varpi_{\mathrm{RVM}} \tag{12.12}$$

(ordered product); and $\lim_{\sigma\to+\infty}\Pi_{\mathrm{tr}}(\sigma) = L = 1/\varpi_{\mathrm{RVM}}$.

Proof. Theorems 11.2, 12.1 and 12.2, and $\Pi_{\mathrm{tr}} = \zeta_{Z1}/\zeta_{\mathrm{RVM}} \to \varpi_{Z1}/\varpi_{\mathrm{RVM}}$ by Theorems 8.10 and 12.1(ii). □

Proposition 12.4 (Boundedness of the endpoint)**.** *Let $\sigma_0 > 1$. Then $|\zeta_{Z1}|$ is bounded above and below on $\Re s \ge \sigma_0$ if and only if*

$$\sup_{\sigma'\ge\sigma_0-\frac12,\ t\ge0}\Big|\Re\int_{\tau_0}^{\infty}\frac{\mathfrak{e}_1(\tau)+\mathfrak{s}_1(\tau)}{(\tau-t+i\sigma')^2}\,d\tau\Big| < \infty. \tag{12.13}$$

Proof. $\tau + iz = \tau - t + i\sigma'$. For $t \ge 0$ the term with $(\tau - iz)^{-2}$ in (12.3) is bounded by (12.2), since $|\tau - iz| \ge \tau + t$; the term with $\mathfrak{s}_1$ is bounded by Lemma 8.9. By (12.3), $\log|\zeta_{Z1}|$ is bounded on $\Re s \ge \sigma_0$, $t \ge 0$, exactly when (12.13) holds, and $\zeta_{Z1}(\bar s) = \overline{\zeta_{Z1}(s)}$ covers $t < 0$. □

The function in (12.13) is, after one integration by parts, the conjugate Poisson integral at distance σ' of the count difference $N_\zeta - n_{\mathrm{RVM}}$; it depends only on the displacements $\gamma_k - \tau_k$.

12.2. The Euler products of ζ_{RVM} and ζ_{Z1}.

Theorem 12.5 (Prime-power counts as Mellin transforms)**.** *Let $X \in \{\mathrm{RVM}, Z1\}$. There is a unique function $J_X^{\mathrm M}$ on $(1,\infty)$ with $J_X^{\mathrm M}(x)x^{-c} \in L^2((1,\infty), dx/x)$ for every $c > 1$ and*

$$\log\zeta_X(s) = s\int_1^\infty J_X^{\mathrm M}(x)\,x^{-s-1}\,dx \qquad (\Re s > 1), \tag{12.14}$$

the branch of $\log\zeta_X$ being real on $(1,\infty)$. Equivalently $\zeta_X(s) = \exp\big(\int_{[1,\infty)} x^{-s}\,dJ_X^{\mathrm M}(x)\big)$.

Proof. ζ_X has no zeros or poles in $\Re s > 1$, so $\log\zeta_X$ is holomorphic there. For $c > 1$, Theorem 8.10 ($X = \mathrm{RVM}$) and Theorem 12.1 ($X = Z1$), together with continuity on the compact part, give $|\log\zeta_X(s)| \le C_c(1 + \log(|t|+2))$ on $\Re s \ge c$. Hence $F_\sigma(t) := \log\zeta_X(\sigma+it)/(\sigma+it)$ lies in $L^2(\mathbb R)$ with $\|F_\sigma\|_2 \le B_c$ for all $\sigma \ge c$. In the variable $u = \log x$, define $J_X^{\mathrm M}(e^u)e^{-\sigma u}$ as the inverse Fourier–Plancherel transform of F_σ; Cauchy's theorem and the decay of F on horizontal segments show that $J_X^{\mathrm M}$ does not depend on σ. Plancherel gives $\int_{\mathbb R}|J_X^{\mathrm M}(e^u)|^2e^{-2\sigma u}du = \|F_\sigma\|_2^2/2\pi \le B_c^2/2\pi$ for all $\sigma \ge c$; letting $\sigma\to\infty$ forces $J_X^{\mathrm M}(e^u) = 0$ for almost every $u < 0$. Plancherel again gives (12.14), and uniqueness follows from the injectivity of the Fourier transform. □

Theorem 12.6 (Distributional explicit formula)**.** *Let $X \in \{\mathrm{RVM}, Z1\}$, with ordinates $\kappa_k = \tau_k$ for $X = \mathrm{RVM}$ and $\kappa_k = \gamma_k$ for $X = Z1$. In $\mathcal{D}'(1,\infty)$,*

$$(\log x)\,\frac{dJ_X^{\mathrm{M}}}{dx} = 1 - \frac{2}{\sqrt{x}}\,\Re\sum_{k\ge1} x^{i\kappa_k} - \frac{1}{x(x^2-1)}, \tag{12.15}$$

the series converging in $\mathcal{D}'(1,\infty)$. The explicit-formula count

$$J_X(x) := \mathrm{li}(x) - 2\Re\sum_{k\ge1}\mathrm{Ei}\big((\tfrac12 + i\kappa_k)\log x\big) - \log 2 + \int_x^\infty \frac{dy}{y(y^2-1)\log y} \tag{12.16}$$

converges in $\mathcal{D}'(1,\infty)$ (J_{RVM} is the function (5.2)), and $J_X^{\mathrm{M}} - J_X$ is a constant C_X on $(1,\infty)$.

Proof. With $\rho_k = \frac12 + i\kappa_k$, the factorization $\zeta_X = P_X/G$, $P_X'/P_X = \sum_k(\frac{1}{s-\rho_k} + \frac{1}{s-\bar\rho_k})$, and the digamma expansion $\frac12\frac{\Gamma'}{\Gamma}(\frac{s}{2}) + \frac1s = -\frac{\gamma_E}{2} + \sum_{n\ge1}(\frac{1}{2n} - \frac{1}{s+2n})$ give, for $\Re s > 1$,

$$-\frac{\zeta_X'}{\zeta_X}(s) = \frac{1}{s-1} - \sum_{k\ge1}\Big(\frac{1}{s-\rho_k} + \frac{1}{s-\bar\rho_k}\Big) + \sum_{n\ge1}\Big(\frac{1}{2n} - \frac{1}{s+2n}\Big) - \frac{\gamma_E}{2} - \frac12\log\pi. \tag{12.17}$$

Each term is a Mellin–Stieltjes transform $\int_{[1,\infty)} x^{-s}d\mu$: $dx \mapsto \frac{1}{s-1}$, $x^{\rho-1}dx \mapsto \frac{1}{s-\rho}$, $x^{-2n-1}dx \mapsto \frac{1}{s+2n}$, $\delta_1 \mapsto 1$. After multiplication by x^{-c} and the change $u = \log x$, the Fourier transforms of the partial sums over k and over n converge locally uniformly with polynomial bounds, so the series converge in $\mathcal{S}'$, and on $(1,\infty)$ the limit measure is the right side of (12.15) times dx. Differentiating (12.14) in s shows that $-\zeta_X'/\zeta_X$ is the Mellin–Stieltjes transform of $(\log x)\,dJ_X^{\mathrm{M}}$; injectivity gives (12.15). The series over k converges in $\mathcal{D}'(1,\infty)$ because $\int\varphi(x)x^{-1/2+i\kappa}dx$ is rapidly decreasing in κ for $\varphi \in C_c^\infty(1,\infty)$ and $\kappa_k \asymp k/\log k$. By Lemma 5.1, $\mathrm{Ei}(\rho_k u) = e^{\rho_k u}/(\rho_k u) + \varrho_k$ with $\sum|\varrho_k|$ locally uniformly convergent, so (12.16) converges in $\mathcal{D}'(1,\infty)$ for the same reason; differentiating termwise with $\frac{d}{dx}\mathrm{Ei}(\rho\log x) = x^{\rho-1}/\log x$, $(\log x)J_X'$ equals the right side of (12.15). A distribution on an interval with zero derivative is constant. □

Proof of Theorem 6.4. Take $X = \mathrm{RVM}$ in Theorems 12.5 and 12.6. By Theorem 5.2, J_{RVM} is continuous on $(1,\infty)$, so $J_{\mathrm{RVM}}^{\mathrm{M}} = J_{\mathrm{RVM}} + C_{\mathrm{RVM}}$ pointwise, and $J_{\mathrm{RVM}}^{*} = J_{\mathrm{RVM}}$ on $[2,\infty)$. □

Theorem 12.7 (Cell form of the endpoint identity)**.** *The following are equivalent.*

(i) *(D), that is, $\zeta_{Z1} = \zeta$.*

(ii) *J_{Z1}^{M} is constant on every interval $(n, n+1)$, $n \ge 1$, and its jumps $\mathsf{b}_n := v_n - v_{n-1}$, where v_n is its value on $(n,n+1)$ and $v_0 := 0$, satisfy $\sum_{n\le X}|\mathsf{b}_n| = O_\varepsilon(X^{1+\varepsilon})$ for every $\varepsilon > 0$.*

(iii) *For every $n \ge 1$, in $\mathcal{D}'(n,n+1)$,*

$$2\,\Re\sum_{k\ge1}\big(x^{i\gamma_k} - x^{i\tau_k}\big) = \sqrt{x}\,\log x\; J_{\mathrm{RVM}}'(x), \tag{12.18}$$

and the jumps of J_{Z1}^{M} satisfy the bound in (ii).

In this case J_{Z1}^{M} is Riemann's prime-power count $J(x) = \sum_{p^m\le x} 1/m$.

Proof. (ii)⇒(i): $|v_n| \le \sum_{m\le n}|\mathsf{b}_m| \ll n^{1+\varepsilon}$, so for $\Re s > 1$ summation by parts in (12.14) gives $\log\zeta_{Z1}(s) = \sum_n v_n(n^{-s} - (n+1)^{-s}) = \sum_{n\ge1}\mathsf{b}_n n^{-s}$, absolutely convergent by partial summation. The exponential of an absolutely convergent Dirichlet series is one, so ζ_{Z1} satisfies (11.8).

(i)⇒(ii): by Theorem 11.2, $\zeta_{Z1} = \zeta$, and $\log\zeta(s) = s\int_1^\infty J(x)x^{-s-1}dx$ for $\Re s > 1$; by uniqueness in Theorem 12.5, $J_{Z1}^{\mathrm{M}} = J$, which is constant between consecutive integers, with jumps $1/m$ at p^m and $\sum_{n\le X}|\mathsf{b}_n| = J(X) \ll X$.

(ii)⇔(iii): Theorem 12.6 for $X = Z1$ and $X = \mathrm{RVM}$, with $dJ_{\mathrm{RVM}}^{\mathrm{M}}/dx = J_{\mathrm{RVM}}'$, gives in $\mathcal{D}'(1,\infty)$

$$(\log x)\Big(\frac{dJ_{Z1}^{\mathrm{M}}}{dx} - J_{\mathrm{RVM}}'(x)\Big) = -\frac{2}{\sqrt{x}}\,\Re\sum_{k\ge1}\big(x^{i\gamma_k} - x^{i\tau_k}\big), \tag{12.19}$$

so $dJ^{\mathrm{M}}_{Z1}/dx = 0$ on $(n, n+1)$ is (12.18). □

Remark 12.8 (The final step in cell form). Theorem 12.7 is the exact form of the continuous-to-atomic passage of Remark 11.4. Both sides of (12.18) are built from the midpoint construction: the right side from the clock alone (J'_{RVM} is bounded on $[2,\infty)$ by Theorem 6.1, equal to $1/\log x$ plus the unit-periodic first harmonic and the harmonics with stationary points $2\pi x^{1/m}$, Section 6), the left side from the displacements $\gamma_k - \tau_k$. The final step of the method is the identity (12.18) on every cell, equivalently (12.12). Numerically, the explicit formula (12.16) with the γ_k is constant between consecutive integers up to the truncation of the list (Experiment 17.6, Figure 8), and the partial products of (12.12) approach ϖ_{RVM} (Experiment 17.3).

13. The atomized family and the programme for the final step

The continuous family ζ_λ keeps, for every λ, the functional equation and the zeros on the critical line. Its atomization $\tilde\zeta_{\mathbb{Z},\lambda}$, built from the cell masses of the prime coordinate, keeps the Euler product and the Dirichlet series. The two are joined by a factor $e^{\tilde h_\lambda}$, and both families have a limit at $\lambda = 0$. Hamburger's theorem needs the functional equation and the Dirichlet series on one and the same function; at $\lambda = 0$ this is the vanishing of the joining factor. This section constructs the atomized family for every λ, proves its analytic properties, writes the joining factor cell by cell, and lists in Subsection 13.8 the steps that remain, in the order in which they will be followed. The passage from a continuous prime system to a discrete one is classical for Beurling generalized primes [5, 15, 8, 16]; the atomization used here places the masses at the integers, which produces ordinary Dirichlet series.

Throughout, $\eta_k = \tau_k - \gamma_k$, $\gamma_k(\lambda) = (1-\lambda)\gamma_k + \lambda\tau_k$, $\rho_k(\lambda) = \frac12 + i\gamma_k(\lambda)$, and λ' denotes an integration variable in $[0,1]$.

13.1. **The family at infinity.** Put $N_\lambda(\tau) = \#\{k : \gamma_k(\lambda) \le \tau\}$, $\mathfrak{e}^{(\lambda)} = N_\lambda - \mathcal{N}_{\mathrm{RVM}}$ and $\mathfrak{e}_1^{(\lambda)}(\tau) = \int_{\tau_0}^{\tau} \mathfrak{e}^{(\lambda)}$, so that $\mathfrak{e}_1^{(0)} = \mathfrak{e}_1$ and $\mathfrak{e}_1^{(1)} = -\mathfrak{s}_1$.

Proposition 13.1 (Clock decomposition along the family)**.** *There is an absolute constant C with $|\mathfrak{e}_1^{(\lambda)}(\tau)| \le C\log\tau$ for $\tau \ge \tau_0$ and all $\lambda \in [0,1]$. Consequently Theorem 12.1 holds for every ζ_λ with $\mathfrak{e}_1$ replaced by $\mathfrak{e}_1^{(\lambda)}$, uniformly in λ; in particular $\varpi_\lambda := \lim_{\sigma\to+\infty}\zeta_\lambda(\sigma)$ exists and*

$$\varpi_\lambda = \varpi_{\mathrm{RVM}} \prod_{k\ge1} \frac{\frac14 + \tau_k^2}{\frac14 + \gamma_k(\lambda)^2}, \qquad \varpi_1 = \varpi_{\mathrm{RVM}}, \quad \varpi_0 = \varpi_{Z1}. \tag{13.1}$$

Proof. $\gamma_k(\lambda)$ lies between γ_k and τ_k, so $\gamma_1(\lambda) > \tau_0$. For $X \ge \tau_0$,

$$\int_{\tau_0}^{X} (N_\lambda - n_{\mathrm{RVM}}) = \sum_k \big((X-\gamma_k(\lambda))^+ - (X-\tau_k)^+\big) = (1-\lambda) \sum_{\gamma_k(\lambda),\tau_k \le X} \eta_k + O(\log X), \tag{13.2}$$

the indices counted by only one of the two sums being $O(\log X)$ in number with ordinates in $[X - C_\eta, X]$ (Lemma 2.2). The case $\lambda = 0$ of (13.2) identifies the last sum with $\mathfrak{e}_1 + \mathfrak{s}_1 + O(\log X) = O(\log X)$, by (12.2) and Lemma 8.9. Since $\mathfrak{e}_1^{(\lambda)} = \int (N_\lambda - n_{\mathrm{RVM}}) - \mathfrak{s}_1$, the bound follows, and the proof of Theorem 12.1 applies verbatim. □

Define the *flow at infinity*

$$\Upsilon(\lambda) := \sum_{k\ge1} \frac{2\gamma_k(\lambda)\,\eta_k}{\frac14 + \gamma_k(\lambda)^2} \qquad \text{(summed in the order of } k\text{)}. \tag{13.3}$$

Proposition 13.2 (The value at infinity along the family)**.** *The series* (13.3) *converges uniformly on* $[0,1]$, $\lambda \mapsto \log \varpi_\lambda$ *is* C^1 *on* $[0,1]$, *and*

$$\frac{d}{d\lambda}\log \varpi_\lambda = -\Upsilon(\lambda) = \lim_{\sigma\to+\infty} \mathcal{V}_\lambda(\sigma), \qquad \Upsilon'(\lambda) = \sum_{k\ge 1} \frac{2\eta_k^2\big(\frac14 - \gamma_k(\lambda)^2\big)}{\big(\frac14+\gamma_k(\lambda)^2\big)^2} < 0, \tag{13.4}$$

with $\mathcal{V}_\lambda$ *the generator of Theorem 10.3. Moreover*

$$\int_{0^+}^{1} \Upsilon(\lambda)\, d\lambda = \sum_{k\ge 1} \log \frac{\frac14+\tau_k^2}{\frac14+\gamma_k^2} = \log L + \log \varpi_{Z1}, \tag{13.5}$$

so that (D) holds if and only if $\int_{0^+}^1 \Upsilon(\lambda)\,d\lambda = \log L$, *and* $\Upsilon(1) < \log L + \log \varpi_{Z1} < \Upsilon(0)$.

Proof. Put $g(u) = 2u/(\frac14+u^2)$. The series $\Upsilon(1) = \sum_k g(\tau_k)\eta_k$ converges by summation by parts against $N_\zeta - n_{\mathrm{RVM}}$, exactly as in (12.10). Since $|g(\gamma_k(\lambda)) - g(\tau_k)| \le C|\eta_k|\gamma_k^{-2}$ and $\sum_k \eta_k^2\gamma_k^{-2} < \infty$, the series $\Upsilon(\lambda) - \Upsilon(1)$ converges absolutely and uniformly; the same holds for the termwise derivative, which is the series in (13.4) (each term is negative because $\gamma_k(\lambda) > \frac12$). Differentiating (13.1) termwise gives $\frac{d}{d\lambda}\log\varpi_\lambda = -\Upsilon(\lambda)$, and letting $\sigma\to+\infty$ in the generator (10.10) gives the same series. Integrating from 0 to 1 and using $\varpi_1 = \varpi_{\mathrm{RVM}}$, $\varpi_0 = \varpi_{Z1}$ gives (13.5); the last statement follows from Theorem 12.2 and the monotonicity of Υ. □

Proposition 13.3 (The value at infinity and the verified zeros)**.** *If every zero* $\beta+i\gamma$ *with* $0<\gamma\le T_0$ *has* $\beta = \frac12$, *then*

$$0 \le \log \varpi_{Z1} \le 0.26 \sum_{\gamma > T_0} \frac{1}{\gamma^2}. \tag{13.6}$$

With $T_0 = 3\cdot 10^{12}$ [33], *the right side is* $\approx \frac{0.26}{2\pi T_0}\big(\log\frac{T_0}{2\pi}+1\big) < 4\cdot 10^{-13}$.

Proof. By the proof of Theorem 12.2, $\log\varpi_{Z1} = \sum \log c_\gamma$ over the pairs $\{\rho, 1-\bar\rho\}$ with $\rho(1-\rho)\ne \frac14+\gamma^2$, and all of them have $\gamma > T_0$. For $\delta \le \frac12$ and $\gamma \ge T_0$, (12.11) gives $\log c_\gamma \le (2\delta^2\gamma^2+\delta^4)\gamma^{-4} \le 0.51\,\gamma^{-2}$; a pair consists of two zeros, which gives (13.6). The numerical value follows from $N_\zeta(T) = \frac{T}{2\pi}\log\frac{T}{2\pi e} + O(\log T)$. □

13.2. **Atomization by smooth cells.** Fix $\varphi_n \in C_c^\infty(n-1, n+1)$, $n\ge 2$, with $\sum_{n\ge2}\varphi_n = 1$ on $[2,\infty)$ and $\varphi_n(m) = \delta_{nm}$ for integers m. For $\lambda\in[0,1]$ let J_λ be the explicit-formula count (12.16) with the ordinates $\gamma_k(\lambda)$ (a distribution on $(1,\infty)$ by Theorem 12.6, which applies to every ζ_λ by Proposition 13.1), $J_\lambda^* = J_\lambda \mathbf{1}_{[2,\infty)}$, and $\pi_\lambda = \sum_{j\ge1}\frac{\mu(j)}{j} J_\lambda^*(x^{1/j})$.

Definition 13.4 (Atomized family)**.** The cell masses, the atomized zeta function and its coefficients are

$$\tilde c_n(\lambda) := \langle d\pi_\lambda, \varphi_n\rangle, \qquad \tilde\zeta_{\mathbb{Z},\lambda}(s) := \prod_{n\ge2}\big(1-n^{-s}\big)^{-\tilde c_n(\lambda)} = \sum_{N\ge1}\frac{\tilde a_N(\lambda)}{N^s}, \tag{13.7}$$

$$\tilde a_N(\lambda) = \sum_{\prod_n n^{r_n}=N} \prod_{n:\, r_n\ge1} \binom{\tilde c_n(\lambda)+r_n-1}{r_n}, \qquad \tilde a_1(\lambda) = 1. \tag{13.8}$$

Unlike the discretizations of Beurling systems, where the atoms are placed at chosen real points [15, 8, 16], the masses of $d\pi_\lambda$ are placed at the integers with real exponents; measures with continuous part and non-integer multiplicities of this kind appear also in [27, Ch. 3]. Thus $\tilde a_p(\lambda) = \tilde c_p(\lambda)$ for primes p, $\tilde a_4 = \tilde c_4 + \frac12\tilde c_2(\tilde c_2+1)$, $\tilde a_6 = \tilde c_6 + \tilde c_2\tilde c_3$. The pairing with a fixed smooth partition defines $\tilde c_n(\lambda)$ for every λ without point values of π_λ, and $\lambda\mapsto\tilde c_n(\lambda)$ is continuous. If $d\pi_0 = \sum_p \delta_p$, then $\tilde c_n(0) = \mathbf{1}_{\mathrm{prime}}(n)$ and $\tilde a_N(0) = 1$ for all N. Along the family, in $\mathcal{D}'(1,\infty)$,

$$\frac{\partial J_\lambda}{\partial\lambda}(x) = 2\sum_{k\ge1}\eta_k\,\Im\frac{x^{\rho_k(\lambda)}}{\rho_k(\lambda)}, \qquad \frac{\partial}{\partial\lambda}\Big((\log x)\frac{dJ_\lambda}{dx}\Big) = \frac{2\log x}{\sqrt{x}}\sum_{k\ge1}\eta_k\sin\big(\gamma_k(\lambda)\log x\big). \tag{13.9}$$

13.3. **The joining factor cell by cell.** Define the *atomization kernel* and the *cell defects*

$$\mathfrak{a}_s(x) := \sum_{n\ge 2} \varphi_n(x)\, n^{-s} - x^{-s}, \qquad \mathfrak{h}_n(\lambda; s) := \big\langle d\pi_\lambda,\ \varphi_n(x)\big(n^{-s} - x^{-s}\big)\big\rangle. \tag{13.10}$$

$\mathfrak{a}_s$ vanishes at every integer ≥ 2, and $\mathfrak{h}_n(\lambda; s)$ measures how far the mass of the cell n is from being concentrated at n.

The explicit-formula count J_λ need not vanish on $(1, 2)$, whereas π_λ is built from $J_\lambda^* = J_\lambda \mathbf{1}_{[2,\infty)}$. The first cell $(1, 2)$ enters the joining factor through

$$\mathfrak{u}_\lambda(s) := C_\lambda\, 2^{-s} + s\int_1^2 J_\lambda^{\mathrm{M}}(x)\, x^{-s-1}\, dx, \qquad C_\lambda := J_\lambda^{\mathrm{M}} - J_\lambda, \tag{13.11}$$

where J_λ^{M} is the Mellin count of ζ_λ (Theorem 12.5, which applies to every ζ_λ by Proposition 13.1) and C_λ is the constant of Theorem 12.6.

Proposition 13.5 (The first cell)**.** *Let $\lambda \in [0, 1]$.*

(i) *$\mathfrak{u}_\lambda$ is entire, and for $\Re s > 1$*

$$\log \zeta_\lambda(s) = \mathfrak{u}_\lambda(s) + \langle dJ_\lambda^*, x^{-s}\rangle, \qquad \langle dJ_\lambda^*, x^{-s}\rangle := s\int_2^\infty J_\lambda(x)\, x^{-s-1}\, dx. \tag{13.12}$$

(ii) *$\mathfrak{u}_\lambda(\sigma + it) = o(|t|)$ as $|t| \to \infty$, uniformly for σ in compact sets.*

(iii) *$\mathfrak{u}_\lambda$ is constant if and only if $J_\lambda = 0$ on $(1, 2)$; in that case $\mathfrak{u}_\lambda \equiv C_\lambda$.*

For Riemann's count $J(x) = \sum_{p^m \le x} 1/m$, which vanishes on $(1, 2)$, the corresponding term is 0.

Proof. (i) $J_\lambda^{\mathrm{M}}(x)x^{-c} \in L^2((1,\infty), dx/x)$ for $c > 1$, so $J_\lambda^{\mathrm{M}} \in L^1(1, 2)$ and the integral in (13.11) is entire in s. By (12.14), $\log\zeta_\lambda(s) = s\int_1^2 J_\lambda^{\mathrm{M}} x^{-s-1}dx + s\int_2^\infty (J_\lambda + C_\lambda)x^{-s-1}dx$, and $s\int_2^\infty C_\lambda x^{-s-1}dx = C_\lambda 2^{-s}$. The second integral is the pairing of x^{-s} with the distributional derivative of J_λ^*, which vanishes on $(0, 2)$. (ii) $C_\lambda 2^{-s}$ is bounded on vertical strips, and $\int_1^2 J_\lambda^{\mathrm{M}}(x)x^{-s-1}dx = \int_0^{\log 2} J_\lambda^{\mathrm{M}}(e^u)e^{-\sigma u}e^{-itu}du \to 0$ as $|t| \to \infty$ by the Riemann–Lebesgue lemma, uniformly for σ in compact sets. (iii) For a constant c one has $c = c\,2^{-s} + s\int_1^2 c\, x^{-s-1}dx$, so $\mathfrak{u}_\lambda(s) - c = s\int_1^\infty H(x)x^{-s-1}dx$ with $H = (J_\lambda^{\mathrm{M}} - c)\mathbf{1}_{(1,2)} + (C_\lambda - c)\mathbf{1}_{[2,\infty)}$. By the injectivity used in Theorem 12.5, $\mathfrak{u}_\lambda \equiv c$ if and only if $H = 0$ almost everywhere, that is, $C_\lambda = c$ and $J_\lambda^{\mathrm{M}} = C_\lambda$ on $(1, 2)$, which is $J_\lambda = 0$ on $(1, 2)$. □

Theorem 13.6 (The joining factor)**.** *Let $\tilde h_\lambda := \log\tilde\zeta_{\mathbb{Z},\lambda} - \log\zeta_\lambda$ on the half-plane where $\tilde\zeta_{\mathbb{Z},\lambda}$ converges absolutely (Theorem 13.12). Then*

$$\tilde\zeta_{\mathbb{Z},\lambda} = \zeta_\lambda\, e^{\tilde h_\lambda}, \qquad \tilde h_\lambda(s) = -\mathfrak{u}_\lambda(s) + \sum_{m\ge1}\frac{1}{m}\sum_{n\ge2}\mathfrak{h}_n(\lambda; ms), \tag{13.13}$$

and, explicitly,

$$\tilde h_\lambda(s) = -\mathfrak{u}_\lambda(s) + \sum_{m\ge1}\frac{1}{m}\sum_{j\ge1}\frac{\mu(j)}{j}\int_2^\infty \mathfrak{a}_{ms}(y^j)\,\frac{1}{\log y}\Big(1 - \frac{2}{\sqrt{y}}\Re\sum_{k\ge1} y^{i\gamma_k(\lambda)} - \frac{1}{y(y^2-1)}\Big)dy, \tag{13.14}$$

the series over k converging absolutely after integration. Its derivative along the family is

$$\partial_\lambda\tilde h_\lambda(s) = -\partial_\lambda\mathfrak{u}_\lambda(s) + \sum_{m\ge1}\frac{1}{m}\sum_{j\ge1}\frac{\mu(j)}{j}\int_2^\infty \mathfrak{a}_{ms}(y^j)\,\frac{2}{\sqrt{y}}\sum_{k\ge1}\eta_k\sin\big(\gamma_k(\lambda)\log y\big)\,dy. \tag{13.15}$$

Here $\partial_\lambda\mathfrak{u}_\lambda(s) = \mathcal{V}_\lambda(s) - \partial_\lambda\langle dJ_\lambda^*, x^{-s}\rangle$ by (13.12) and (10.11). Moreover $\mathfrak{u}_\lambda(\sigma) \to \log\varpi_\lambda$ as $\sigma \to +\infty$, and $\mathfrak{u}_\lambda \equiv \log\varpi_\lambda$ if J_λ vanishes on $(1, 2)$.

Proof. By the Euler product, $\log \tilde{\zeta}_{\mathbb{Z},\lambda}(s) = \sum_{m\geq 1} \frac{1}{m} \sum_n \tilde{c}_n(\lambda) n^{-ms}$. By Proposition 13.5, $\log \zeta_\lambda(s) = \mathfrak{u}_\lambda(s) + \langle dJ_\lambda^*, x^{-s}\rangle$; finite inversion (Proposition 5.3, whose proof applies to J_λ^*) gives $dJ_\lambda^* = \sum_m \frac{1}{m}\, d[\pi_\lambda(x^{1/m})]$, and $\langle d[\pi_\lambda(x^{1/m})], x^{-s}\rangle = \langle d\pi_\lambda, y^{-ms}\rangle$. Inserting $1 = \sum_n \varphi_n$ on $[2,\infty)$ gives $\langle d\pi_\lambda, y^{-ms}\rangle = \sum_n (\tilde{c}_n(\lambda) n^{-ms} - \mathfrak{h}_n(\lambda; ms))$, hence (13.13). Letting $\sigma \to +\infty$, $\log \tilde{\zeta}_{\mathbb{Z},\lambda} \to 0$ (because $\tilde{a}_1 = 1$), $\log \zeta_\lambda \to \log \varpi_\lambda$ (Proposition 13.1) and the double series tends to 0 by the bounds of Theorem 13.12; so $\mathfrak{u}_\lambda(\sigma) \to \log \varpi_\lambda$, and a constant $\mathfrak{u}_\lambda$ (Proposition 13.5(iii)) equals $\log \varpi_\lambda$. Atoms of $d\pi_\lambda$ at integers contribute nothing because the test functions vanish there. Formula (13.14) is (13.13) with $d\pi_\lambda$ written by (12.15) and the substitution $x = y^j$; the integrated terms in k are $O(\gamma_k(\lambda)^{-2})$ because $\mathfrak{a}_{ms}(2) = 0$. Differentiating in λ with (13.9) gives (13.15); $\mathfrak{u}_\lambda(s)$ is differentiable in λ because J_λ^{M} is the Fourier–Plancherel transform of $\log \zeta_\lambda(\sigma + it)/(\sigma + it)$, whose λ-derivative $\mathcal{V}_\lambda(\sigma + it)/(\sigma + it)$ is bounded in $L^2(\mathbb{R})$ locally uniformly in λ. □

Proposition 13.7 (Explicit cell defects)**.** *For every $n \geq 2$, $\lambda \in [0,1]$ and $\Re s > 1$,*

$$\mathfrak{h}_n(\lambda; s) = \sum_{j\geq 1} \frac{\mu(j)}{j} \int_2^\infty \varphi_n(y^j)(n^{-s} - y^{-js})\, \frac{1}{\log y}\Big(1 - \frac{2}{\sqrt{y}} \Re \sum_{k\geq 1} y^{i\gamma_k(\lambda)} - \frac{1}{y(y^2-1)}\Big) dy, \tag{13.16}$$

$$\partial_\lambda \mathfrak{h}_n(\lambda; s) = \sum_{j\geq 1} \frac{\mu(j)}{j} \int_2^\infty \varphi_n(y^j)(n^{-s} - y^{-js})\, \frac{2}{\sqrt{y}} \sum_{k\geq 1} \eta_k \sin\big(\gamma_k(\lambda) \log y\big)\, dy, \tag{13.17}$$

the sums over j being finite and the sums over k converging absolutely after integration, because the test functions are smooth with compact support. At $\lambda = 1$ the cell defect $\mathfrak{h}_n(1;\cdot)$ is the dispersion of the cell n for ζ_{RVM}, computed from the clock alone; $d\pi_{\mathrm{RVM}}$ has a bounded density there (Corollary 6.2), so $\mathfrak{h}_n(1;\cdot) \equiv 0$ holds only if this density vanishes on the support of φ_n.

Proof. Insert (12.15) for $d\pi_\lambda = \sum_j \frac{\mu(j)}{j}\, d[J_\lambda^*(x^{1/j})]$ into the definition (13.10) and substitute $x = y^j$; the atoms of $d\pi_\lambda$ at integers, for instance the jump of J_λ^* at 2, contribute nothing because $n^{-s} - x^{-s}$ vanishes at $x = n$ and φ_n vanishes at the other integers. Differentiation in λ uses (13.9); the factor $\log y$ cancels. For the last statement, $\mathfrak{h}_n(1;\cdot) \equiv 0$ forces $\varphi_n\, d\pi_{\mathrm{RVM}} = \tilde{c}_n(1)\delta_n$ by the injectivity used in Proposition 13.8, and an absolutely continuous measure equals a multiple of δ_n only if both vanish. □

In the form of the path, the final step requires that in every cell the flow of the displacements cancel exactly the dispersion of the clock, (13.29).

Proposition 13.8 (The limit $\lambda \to 0$ and the meaning of a vanishing cell)**.** *For every $n \geq 2$ and $\Re s > 1$, $\mathfrak{h}_n(\lambda; s) \to \mathfrak{h}_n(0; s)$ and $\tilde{c}_n(\lambda) \to \tilde{c}_n(0)$ as $\lambda \to 0^+$, and*

$$\mathfrak{h}_n(0; s) = \mathfrak{h}_n(1; s) - \int_{0^+}^1 \partial_\lambda \mathfrak{h}_n(\lambda; s)\, d\lambda. \tag{13.18}$$

Moreover $\mathfrak{h}_n(0; s) = 0$ for all $\Re s > 1$ if and only if $\varphi_n\, d\pi_{Z1} = \tilde{c}_n(0)\, \delta_n$.

Proof. The test function $\varphi_n(x)(n^{-s} - x^{-s})$ is smooth with compact support, and the integrated terms over k are dominated uniformly in λ because $|\gamma_k(\lambda) - \gamma_k| \leq C_\eta$; dominated convergence gives the limits and (13.18). Since $\mathfrak{h}_n(0; s) = \tilde{c}_n(0) n^{-s} - \langle \varphi_n\, d\pi_{Z1}, x^{-s}\rangle$, its vanishing for all s says that the compactly supported distribution $\varphi_n\, d\pi_{Z1}$ has the Mellin transform of $\tilde{c}_n(0)\delta_n$; the Mellin transform is injective on such distributions. □

13.4. **Mean square of the cell masses.** From now on the cells are translates of one function: $\varphi_n(x) = \varphi(x - n)$, where $\varphi = \chi / \sum_{m\in\mathbb{Z}} \chi(\cdot - m)$ for a fixed $\chi \in C_c^\infty(-1, 1)$ positive on $[-\frac{1}{2}, \frac{1}{2}]$. Then $\sum_{n\geq 2} \varphi_n = 1$ on $[2,\infty)$, $\varphi_n(m) = \delta_{nm}$, and $\|\varphi_n^{(A)}\|_\infty = \|\varphi^{(A)}\|_\infty =: C_A$ for every $A \geq 0$, uniformly

in n. For $n \geq 2$ put

$$V_n(\gamma) := \sum_{j\geq 1} \frac{\mu(j)}{j} \int_2^\infty \varphi_n(y^j)\, y^{-1/2} e^{i\gamma \log y}\, dy, \qquad W_n(\gamma) := \sum_{j\geq 1} \frac{\mu(j)}{j} \int_2^\infty \varphi_n(y^j)\, \frac{y^{-1/2} e^{i\gamma \log y}}{\log y}\, dy, \tag{13.19}$$

so that $W_n' = iV_n$; both are rapidly decreasing in γ when n is not a power of 2 (for $n = 2^j$ the term j carries the cut of J_λ^* at $y = 2$ and is treated in the proof of Theorem 13.10). The density of $d\pi_\lambda$ used in (13.16) and the derivative (13.9) give, with $\eta_k = \tau_k - \gamma_k$,

$$\tilde{c}_n(\lambda) = \mathfrak{m}_n - 2\,\Re \sum_{k\geq 1} W_n(\gamma_k(\lambda)), \qquad \partial_\lambda \tilde{c}_n(\lambda) = 2 \sum_{k\geq 1} \eta_k\, \Im V_n(\gamma_k(\lambda)), \tag{13.20}$$

where $\mathfrak{m}_n$ collects the terms that do not involve the ordinates and is $O(1)$.

Lemma 13.9 (Mean value over the moving ordinates)**.** *There is C such that for all $\lambda \in [0,1]$, $T \geq 2$, $u_0 \in \mathbb{R}$ and complex a_k with $|a_k| \leq 1$,*

$$\int_{u_0-1}^{u_0+1} \Big| \sum_{\gamma_k(\lambda)\leq T} a_k\, e^{i\gamma_k(\lambda)u} \Big|^2 du \leq C\, T \log^2 T. \tag{13.21}$$

Proof. Let $w(u) = C_0 \big(\sin(\frac{u-u_0}{2})/\frac{u-u_0}{2}\big)^2$ with C_0 such that $w \geq 1$ on $|u - u_0| \leq 1$; its Fourier transform is bounded and supported in $[-1,1]$. Expanding the square,

$$\begin{aligned} \int w(u) \Big| \sum_k a_k e^{i\gamma_k(\lambda)u} \Big|^2 du &= \sum_{k,k'} a_k \bar{a}_{k'}\, e^{i(\gamma_k(\lambda) - \gamma_{k'}(\lambda))u_0}\, \hat{w}\big(\gamma_k(\lambda) - \gamma_{k'}(\lambda)\big) \\ &\ll \#\{(k,k') : \gamma_k(\lambda) \leq T,\ |\gamma_k(\lambda) - \gamma_{k'}(\lambda)| \leq 1\}. \end{aligned} \tag{13.22}$$

Since $|\gamma_k(\lambda) - \tau_k| \leq C_\eta$, every interval $[t, t+1]$ contains $O(\log(t+2))$ of the $\gamma_k(\lambda)$, uniformly in λ, because it contains that many τ_k after widening by C_η. The number of pairs is therefore $O(T \log^2 T)$. □

Theorem 13.10 (Mean square of the cell masses along the family)**.** *For every $\varepsilon > 0$ there is C_ε such that for all $X \geq 2$ and all $\lambda \in [0,1]$*

$$\sum_{X < n \leq 2X} |\tilde{c}_n(\lambda)|^2 \leq C_\varepsilon X^{1+\varepsilon}, \qquad \sum_{X < n \leq 2X} |\partial_\lambda \tilde{c}_n(\lambda)|^2 \leq C_\varepsilon X^{1+\varepsilon}. \tag{13.23}$$

In particular, letting $\lambda \to 0^+$ (each $\tilde{c}_n$ is continuous in λ and the constant does not depend on λ), $\sum_{X<n\leq 2X} |\tilde{c}_n(0)|^2 \leq C_\varepsilon X^{1+\varepsilon}$.

Proof. Fix ε, X, λ and put $T = X^{1+\varepsilon}$. By (13.20), $\tilde{c}_n(\lambda) - \mathfrak{m}_n = -2\Re \sum_j \frac{\mu(j)}{j} A_{n,j}$, with

$$A_{n,j} = \sum_k \int_2^\infty \varphi_n(y^j)\, \frac{y^{-1/2 + i\gamma_k(\lambda)}}{\log y}\, dy, \qquad 1 \leq j \leq J := \Big\lfloor \frac{\log(2X+1)}{\log 2} \Big\rfloor. \tag{13.24}$$

Large ordinates. In $u = \log y$ the function $\varphi_n(e^{ju}) e^{u/2}/u$ has support of length $O((jn)^{-1})$, size $O(n^{1/(2j)})$ and r-th derivative $O_r((jn)^r n^{1/(2j)})$: by Faà di Bruno's formula the r-th derivative of $\varphi_n(e^{ju})$ is a sum of terms $\varphi_n^{(m)}(e^{ju})(je^{ju})^m j^{r-m}$, $m \leq r$, and $\|\varphi_n^{(m)}\|_\infty = C_m$ does not depend on n; the derivatives of $e^{u/2}/u$ are $O_r(e^{u/2})$ because $u \geq \log 2$. Integrating by parts A times, each term with $\gamma_k(\lambda) > T$ is $O_A\big((jn/\gamma_k(\lambda))^A n^{1/(2j)-1}\big)$, and their sum over k is $O(X^{-1})$ once $A > (3+\varepsilon)/\varepsilon$. *The cell $n = 2^j$.* Each block contains exactly one $n = 2^j$; for it the term j carries the cut of J_λ^* at $y = 2$ and is treated directly. It equals $-\frac{\mu(j)}{j} \int_2^3 J_\lambda(y)\, \frac{d}{dy}\varphi_n(y^j)\, dy$; the weight has L^2 norm $O((jn)^{1/2})$, and $\|J_\lambda\|_{L^2(2,3)}$ is bounded uniformly in λ (Theorem 12.5 with the uniform bound of Proposition 13.1). This term is therefore $O((n/j)^{1/2})$ and contributes $O(X)$ to the sum of squares; the same holds for

$\partial_\lambda \tilde c_n$, with $\partial_\lambda J_\lambda$ in place of J_λ. *Small ordinates.* Let $S(y)=y^{-1/2}(\log y)^{-1}\sum_{\gamma_k(\lambda)\le T} y^{i\gamma_k(\lambda)}$ and let $I_{n,j}$ be the support of $y\mapsto\varphi_n(y^j)$, of length $O(X^{1/j-1})$. By Cauchy–Schwarz in the cell,

$$\Big|\int \varphi_n(y^j)S(y)\,dy\Big|^2 \le \|\varphi\|_\infty^2\,|I_{n,j}|\int_{I_{n,j}}|S|^2dy. \tag{13.25}$$

For fixed j the intervals $I_{n,j}$, $X<n\le 2X$, cover each point at most twice and lie in $[(X-1)^{1/j},(2X+1)^{1/j}]$, an interval of length at most 2 in $u=\log y$. With $dy/y=du$ and Lemma 13.9 ($a_k=1$),

$$\sum_{X<n\le 2X}\Big|\int \varphi_n(y^j)S\,dy\Big|^2 \ll X^{1/j-1}\Big(\frac{j}{\log X}\Big)^2\int_{u_0-1}^{u_0+1}\Big|\sum_{\gamma_k(\lambda)\le T}e^{i\gamma_k(\lambda)u}\Big|^2du \ll j^2X^{1/j+2\varepsilon}. \tag{13.26}$$

Assembly. By Minkowski's inequality over j, the ℓ^2 norm over $X<n\le 2X$ of $\tilde c_n(\lambda)-\mathfrak m_n$ is $\ll\sum_{j\le J}(X^{1/(2j)+\varepsilon}+X^{-1/2})\ll X^{1/2+\varepsilon}$, and $\sum_{X<n\le 2X}\mathfrak m_n^2\ll X$. This is the first bound with 3ε in place of ε. For $\partial_\lambda\tilde c_n$ the argument is the same, by (13.20), with coefficients $a_k=\eta_k/C_\eta$ in Lemma 13.9 and without the factor $(\log y)^{-1}$. □

Corollary 13.11 (Abscissa 1 for every member)**.** *For every $\lambda\in[0,1]$ and $\sigma>1$,*

$$\sum_{n\ge 2}|\tilde c_n(\lambda)|\,n^{-\sigma}<\infty; \tag{13.27}$$

hence $\tilde\zeta_{\mathbb Z,\lambda}$ and $1/\tilde\zeta_{\mathbb Z,\lambda}$ are absolutely convergent Dirichlet series in $\Re s>1$, for every λ, including the endpoint $\lambda=0$. Moreover, uniformly in λ, $\#\{n\le X:|\tilde c_n(\lambda)|>n^\delta\}\ll_\varepsilon X^{1-2\delta+\varepsilon}$ for every $\delta>0$.

Proof. By Cauchy–Schwarz on dyadic blocks, $\sum_{X<n\le 2X}|\tilde c_n(\lambda)|\le(X\cdot C_\varepsilon X^{1+\varepsilon})^{1/2}$, and summing over $X=2^i$ gives (13.27) for $\sigma>1+\varepsilon/2$, with ε arbitrary. Then $\log\tilde\zeta_{\mathbb Z,\lambda}=\sum_m\frac1m\sum_n\tilde c_n(\lambda)n^{-ms}$ converges absolutely in $\Re s>1$, and the exponentials of $\pm$ this series are absolutely convergent Dirichlet series there. The last statement is Chebyshev's inequality on dyadic blocks. □

The abscissa $\frac32$ obtained below from the primitive $\pi_\lambda-\mathfrak L$ (Theorem 13.12(ii)) is not the true one: a bound on the size of $\pi_\lambda-\mathfrak L$ does not see the cancellation of the density of $d\pi_\lambda$ in mean square, which Theorem 13.10 captures. Absolute convergence in $\Re s>1$ is exactly what Hamburger's theorem requires, and Knopp's obstruction (Subsection 13.7) does not arise.

13.5. **Analytic continuation of the atomized family.** Let $\mathfrak L(x)=\int_1^x(1-y^{-1})\,dy/\log y$; then $\log\frac{s}{s-1}=\int_1^\infty x^{-s}\,d\mathfrak L(x)$ for $\Re s>1$ (both sides have derivative $\frac1s-\frac{1}{s-1}$ and tend to 0 as $s\to+\infty$), and $\mathfrak L(x)=\mathrm{li}(x)-\log\log x-\gamma_E$.

Theorem 13.12 (Continuation of the atomized family)**.** *Let $\lambda\in[0,1]$.*

(i) *For every $c>\frac12$, $(\pi_\lambda-\mathfrak L)\,x^{-c}\in L^2\big((2,\infty),dx/x\big)$.*

(ii) *$|\tilde c_n(\lambda)|\le C(1+I_n^{1/2})$ with I_n as below, and $\sum_n|\tilde c_n(\lambda)|n^{-\sigma}<\infty$ for $\sigma>\frac32$; by Corollary 13.11 this holds for $\sigma>1$, and $\tilde\zeta_{\mathbb Z,\lambda}$ is an absolutely convergent Dirichlet series in $\Re s>1$.*

(iii) *$\tilde h_\lambda$ extends holomorphically to $\Re s>\frac12$. Hence $\tilde\zeta_{\mathbb Z,\lambda}=\zeta_\lambda e^{\tilde h_\lambda}$ is meromorphic in $\Re s>\frac12$, with a single pole at $s=1$ and no zeros.*

(iv) *For $\lambda=1$, $\tilde h_1$ extends holomorphically to $\Re s>0$, and the zeros of $\tilde\zeta_{\mathbb Z,1}$ in $\Re s>0$ are exactly $\frac12\pm i\tau_k$.*

Proof. (i) By Proposition 13.1, for $c>\frac12$ there is C_c with $|\log\zeta_\lambda(s)-\log\frac{s}{s-1}|\le C_c(1+\log(|t|+2))$ on $\Re s\ge c$; the function $\log\zeta_\lambda-\log\frac{s}{s-1}$ is holomorphic on $\Re s>\frac12$, where ζ_λ has no zeros, and its pole at 1 is removed. It equals $s\int_1^\infty(J_\lambda^{\mathrm M}-\mathfrak L)x^{-s-1}dx$ for $\Re s>1$. The argument of Theorem 12.5, applied on the lines $\Re s=c>\frac12$, gives $(J_\lambda^{\mathrm M}-\mathfrak L)x^{-c}\in L^2$. The constant $C_\lambda=J_\lambda^{\mathrm M}-J_\lambda$, the

truncation at 2, the terms $j \geq 2$ of the Möbius inversion (of size $O(\sqrt{x})$ in mean) and $\mathfrak{L}(x) - \mathfrak{L}(x^{1/j})$ preserve the property for $c > \frac{1}{2}$.

(ii) Put $E = \pi_\lambda - \mathfrak{L}$ and $I_n = \int_{n-1}^{n+1} |E|^2 dx$. Since $\int \varphi_n' = 0$, $\tilde{c}_n(\lambda) = \int \varphi_n \, d\mathfrak{L} - \int E\varphi_n' \, dx$, so $|\tilde{c}_n(\lambda)| \leq C(1 + I_n^{1/2})$. By (i), $\sum_n I_n n^{-2c-1} < \infty$, and Cauchy–Schwarz gives $\sum_n I_n^{1/2} n^{-\sigma} < \infty$ for $\sigma > c + 1$.

(iii) With $\psi_n = \varphi_n(x)(n^{-s} - x^{-s})$ one has $|\psi_n'| \leq C|s| n^{-\sigma-1}$ and ψ_n vanishes at the ends of the cell, so

$$|\mathfrak{h}_n(\lambda; s)| \leq C|s| \, n^{-\sigma-1}(1 + I_n^{1/2}), \qquad \sum_n I_n^{1/2} n^{-\sigma-1} < \infty \quad (\sigma > c). \tag{13.28}$$

Choosing $c \in (\frac{1}{2}, \sigma)$, the series (13.13) converges locally uniformly on $\Re s > \frac{1}{2}$; the terms $m \geq 2$ are evaluated at $\Re(ms) \geq 2\sigma$. Since $\mathfrak{u}_\lambda$ is entire (Proposition 13.5), $\tilde{h}_\lambda$ is holomorphic on $\Re s > \frac{1}{2}$, and ζ_λ has no zeros there.

(iv) By Corollary 6.2, π_{RVM} oscillates by $O(1)$ on every unit cell; integrating by parts, $|\mathfrak{h}_n(1; s)| \leq C|s| n^{-\sigma-1}$, and the series converges for $\sigma > 0$. The factor $e^{\tilde{h}_1}$ has no zeros, so the zeros of $\tilde{\zeta}_{\mathbb{Z},1}$ in $\Re s > 0$ are those of ζ_{RVM}. □

Theorem 13.13 (Closing criterion; via Hamburger's theorem [23])**.** *The following are equivalent.*

(i) *(D), that is,* $\zeta_{Z1} = \zeta$.
(ii) $\tilde{h}_0 \equiv 0$.
(iii) $J_{Z1} = 0$ *on* $(1, 2)$*, and for every* $n \geq 2$ *and every* $\Re s > 1$*,*

$$\int_{0^+}^{1} \partial_\lambda \mathfrak{h}_n(\lambda; s) \, d\lambda = \mathfrak{h}_n(1; s). \tag{13.29}$$

In this case $\tilde{c}_n(0) = \mathbf{1}_{\mathrm{prime}}(n)$*,* $\tilde{a}_N(0) = 1$ *for all* N*, and* $\varpi_{Z1} = 1$.

Proof. (ii)⇒(i): $\tilde{h}_0 \equiv 0$ gives $\zeta_{Z1} = \tilde{\zeta}_{\mathbb{Z},0} = \exp(\sum_m \frac{1}{m} \sum_n \tilde{c}_n(0) n^{-ms})$, an absolutely convergent Dirichlet series in $\Re s > 1$ by Corollary 13.11. (iii)⇒(ii): by (13.18) every $\mathfrak{h}_n(0; \cdot)$ vanishes, and $\mathfrak{u}_0 \equiv \log \varpi_{Z1}$ (Theorem 13.6), so $\tilde{h}_0 \equiv -\log \varpi_{Z1}$ is constant; then $\zeta_{Z1} = \varpi_{Z1} \tilde{\zeta}_{\mathbb{Z},0}$ is an absolutely convergent Dirichlet series in $\Re s > 1$ (Corollary 13.11), Theorem 11.2 gives $\zeta_{Z1} = \zeta$, and comparing the first coefficients, $\varpi_{Z1} = 1$. (i)⇒(iii): if $\zeta_{Z1} = \zeta$, Theorem 12.7 gives $J_{Z1}^{\mathrm{M}} = J$, and $J_{Z1} = J$ by von Mangoldt's explicit formula [18], so J_{Z1} vanishes on $(1, 2)$, $d\pi_{Z1} = \sum_p \delta_p$, $\tilde{c}_n(0) = \mathbf{1}_{\mathrm{prime}}(n)$, every $\mathfrak{h}_n(0; \cdot)$ vanishes, and (13.18) gives (13.29). □

Remark 13.14 (The pointwise size of the cell masses). Theorem 13.13 needs no pointwise bound on $\tilde{c}_n(0)$: its proof uses the cell masses only through (13.27). The bound $\tilde{c}_n(0) = O(n^\varepsilon)$ for every n is not proved directly; Corollary 13.11 restricts the exceptions to a set of $n \leq X$ of size $O(X^{1-2\delta+\varepsilon})$ for the threshold n^δ, uniformly in λ, and under the closure every cell mass is bounded, $\tilde{c}_n(0) = \mathbf{1}_{\mathrm{prime}}(n)$. In the ordinates, (13.20) gives the trivial pointwise bound $\tilde{c}_n(\lambda) = O(n^{1/2+\varepsilon})$; the mean square (13.23) shows that the typical cell mass is $O(n^\varepsilon)$.

13.6. **The two endpoints of the atomized family.**

Proposition 13.15 (Atomization at the clock endpoint)**.** *At* $\lambda = 1$*:*

(i) $\tilde{c}_n(1) = O(1)$*, and* $\tilde{\zeta}_{\mathbb{Z},1}$ *is an absolutely convergent Dirichlet series in* $\Re s > 1$*;*
(ii) $\tilde{h}_1(\sigma) \to -\log \varpi_{\mathrm{RVM}} = \log L \neq 0$ *as* $\sigma \to +\infty$*;*
(iii) $\tilde{h}_1$ *is not constant: for every* $\sigma > 1$*,* $\zeta_{\mathrm{RVM}}(\sigma + it) \to \varpi_{\mathrm{RVM}}$ *as* $|t| \to \infty$*, whereas* $\tilde{\zeta}_{\mathbb{Z},1}(\sigma + it)$ *has no limit as* $|t| \to \infty$.

Proof. (i) By Corollary 6.2, π_{RVM} oscillates by $O(1)$ on every unit cell, and $\tilde{c}_n(1) = -\int \pi_{\mathrm{RVM}} \varphi_n' \, dx$ with $\int \varphi_n' = 0$. (ii) is (13.13) with $\sigma \to +\infty$, since $\varpi_{\mathrm{RVM}} < 1$. (iii) The first limit is (8.33). If $\tilde{\zeta}_{\mathbb{Z},1}(\sigma + it)$ had a limit ℓ as $|t| \to \infty$, the mean-square argument (8.43)–(8.44) applied to $\tilde{\zeta}_{\mathbb{Z},1} - \ell$ on the line $\Re s = \sigma$ would give $\tilde{a}_N(1) = 0$ for $N \geq 2$, so $\tilde{\zeta}_{\mathbb{Z},1} \equiv 1$, which contradicts its pole at

$s = 1$ (Theorem 13.12(iii)). If $\tilde{h}_1$ were constant, ζ_{RVM} would be a constant multiple of an absolutely convergent Dirichlet series, which Theorem 8.11 excludes. □

The two families thus keep complementary structures and are joined by a zero-free factor:

$$\underbrace{\zeta_\lambda}_{\substack{\text{functional equation}\\ \text{zeros } \frac{1}{2}\pm i\gamma_k(\lambda)}} \cdot e^{\tilde{h}_\lambda} \;=\; \underbrace{\tilde{\zeta}_{\mathbb{Z},\lambda}}_{\substack{\text{Dirichlet series}\\ \text{Euler product}}}, \qquad \tilde{h}_1 \text{ not constant}, \qquad (D) \iff \tilde{h}_0 \equiv 0. \tag{13.30}$$

At $\lambda = 1$ the joining factor carries the whole difference between the clock and an Euler product; Theorem 13.13 states that this difference disappears at $\lambda = 0$.

The reciprocal of the atomized family is again an Euler product:

$$\frac{1}{\tilde{\zeta}_{\mathbb{Z},\lambda}(s)} = \prod_{n\ge 2}\bigl(1-n^{-s}\bigr)^{\tilde{c}_n(\lambda)} = \sum_{N\ge 1}\frac{\tilde{\mu}_\lambda(N)}{N^s}, \qquad \tilde{\mu}_\lambda(N) = \sum_{\prod_n n^{r_n}=N}\ \prod_{n:\, r_n\ge 1}(-1)^{r_n}\binom{\tilde{c}_n(\lambda)}{r_n}, \tag{13.31}$$

absolutely convergent in $\Re s > 1$ for every λ (Corollary 13.11), because it is the exponential of the absolutely convergent series $-\sum_m \frac{1}{m}\sum_n \tilde{c}_n(\lambda) n^{-ms}$. Thus $\tilde{\mu}_\lambda(p) = -\tilde{c}_p(\lambda)$, and the atomized Mertens functions

$$\tilde{M}_\lambda(x) := \sum_{N\le x}\tilde{\mu}_\lambda(N), \qquad \frac{1}{\tilde{\zeta}_{\mathbb{Z},\lambda}} = e^{-\tilde{h}_\lambda}F_\lambda, \tag{13.32}$$

exist for every $\lambda \in [0,1]$. Under the conditions of Theorem 13.13, $\tilde{c}_n(0) = \mathbf{1}_{\mathrm{prime}}(n)$, so $\tilde{\mu}_0 = \mu$ and $\tilde{M}_0 = M$.

Proposition 13.16 (Mellin form of the endpoint Mertens function)**.** *There are $c_1 \ge 2$ and a unique measurable B_{Z1} on $(0,\infty)$, vanishing on $(0,1)$, with $x^{-c}B_{Z1} \in L^2\bigl((1,\infty), dx/x\bigr)$ and*

$$F_{Z1}(s) = s\int_1^\infty B_{Z1}(x)\, x^{-s-1}\,dx = \exp\Bigl(-\int_{[1,\infty)} x^{-s}\, dJ^{\mathrm{M}}_{Z1}(x)\Bigr) \qquad (\Re s > c_1), \tag{13.33}$$

for every $c > c_1$. Moreover (D) holds if and only if $B_{Z1} = M$ almost everywhere; and if F_{Z1} has an absolutely convergent Dirichlet expansion (11.37)*, then $B_{Z1} = M_{Z1}$.*

Proof. Put $A = 2(\pi+2)C_{\mathfrak{e}}$. By (12.5), for $\sigma' \ge \max(4A, R_0)$, $|F_{Z1}(s)| \le C(|t|+\sigma)^{1/4}$; take $c_1 = \max(2, \frac{1}{2}+4A, \frac{1}{2}+R_0)$. Then $\int_{\mathbb{R}} |F_{Z1}(\sigma+it)|^2|\sigma+it|^{-2}dt \le C'\sigma^{-1/2}$ for $\sigma > c_1$, and the argument of Theorem 12.5 gives B_{Z1}, its vanishing on $(0,1)$, its independence of c and its uniqueness; the second equality in (13.33) is Theorem 12.5. If $\zeta_{Z1} = \zeta$, then $F_{Z1}(s) = 1/\zeta(s) = s\int_1^\infty M(x)x^{-s-1}dx$ and uniqueness gives $B_{Z1} = M$. Conversely, $B_{Z1} = M$ gives $F_{Z1} = 1/\zeta$ on $\Re s > c_1$, hence $\zeta_{Z1} = \zeta$ by analytic continuation. Under (11.37), partial summation gives $F_{Z1}(s) = s\int_1^\infty M_{Z1}(x)x^{-s-1}dx$, and uniqueness gives $B_{Z1} = M_{Z1}$. □

13.7. **Hamburger's hypotheses at the two endpoints.** Hamburger's theorem [23], in the form of [37, §2.13] (see also [35, 6]), has four hypotheses: (H0) $f = G/P$ with G entire of finite order and P a polynomial; (H1) $f(s) = \sum_n a_n n^{-s}$ absolutely for $\sigma > 1$, with arbitrary complex a_n; (H2) $f(s)\Gamma(\frac{s}{2})\pi^{-s/2} = g(1-s)\Gamma(\frac{1-s}{2})\pi^{-(1-s)/2}$; (H3) $g(1-s) = \sum_n b_n n^{s-1}$ absolutely for $\sigma < -\alpha < 0$. It contains no hypothesis on the location of zeros, and its conclusion is $f = C\zeta$. Absolute convergence for $\sigma > 1$ in (H1) cannot be weakened to some half-plane: Knopp constructed infinitely many linearly independent solutions of (H0), (H2), (H3) that are Dirichlet series absolutely convergent only in some half-plane [25, 31]. Within the Selberg class, the functions of degree 1 are classified in [24]. The table records the status at the two endpoints and is updated as the programme advances.

	ζ_{Z1} (continuous limit)	$\tilde\zeta_{\mathbb{Z},0}$ (atomized limit)
H0	proved: $(s-1)\zeta_{Z1}$ entire of order 1 (Proposition 11.1)	open: meromorphic in $\Re s > \frac12$ only (Theorem 13.12)
H1	open: this is (D)	proved in $\Re s > 1$ (Corollary 13.11)
H2	proved, with $g = f$ (Proposition 11.1)	open
H3	follows from H1	open
zeros	no hypothesis	no hypothesis

The two columns meet when $\tilde h_0 \equiv 0$: then $\zeta_{Z1} = \tilde\zeta_{\mathbb{Z},0}$ has H0–H3, and Hamburger's theorem gives $\zeta_{Z1} = \zeta$ (Theorem 13.13). It suffices that the joining factor be symmetric (Theorem 14.3), or that ζ_{Z1} have a positive representing measure (Theorem 14.7).

The parallel between the two families, topic by topic of the paper, is the following.

Topic of the paper	continuous family ζ_λ	atomized family $\tilde\zeta_{\mathbb{Z},\lambda}$
clock, τ_k, bounded displacement	same	same
definition	Hadamard product over $\frac12 \pm i\gamma_k(\lambda)$	Euler product over $n \geq 2$ with masses $\tilde c_n(\lambda) = \langle d\pi_\lambda, \varphi_n\rangle$
zeros	exactly $\frac12 \pm i\gamma_k(\lambda)$; impossible elsewhere (Theorem 10.1)	none in $\Re s > \frac12$ for every λ, since $\tilde h_\lambda$ is holomorphic there (Theorem 13.12(iii)); for $\lambda = 1$, exactly $\frac12 \pm i\tau_k$ in $\Re s > 0$ (Theorem 13.12(iv))
functional equation	proved for every λ	open; the factor $e^{\tilde h_\lambda}$ carries no functional equation
Dirichlet series	not for $\lambda = 1$ in any half-plane (Theorem 8.11); not in $\Re s > 1$ for $0 < \lambda < 1$ (Proposition 14.1); at $\lambda = 0$ this is (D)	proved in $\Re s > 1$ for every λ, including $\lambda = 0$ (Corollary 13.11)
first coefficient ($\sigma \to +\infty$)	$\varpi_\lambda = \varpi_{\mathrm{RVM}} \prod_k \frac{1/4+\tau_k^2}{1/4+\gamma_k(\lambda)^2}$ (13.1)	$\tilde a_1(\lambda) = 1$ for every λ
vertical behaviour	$\lambda = 1$: $\zeta_{\mathrm{RVM}}(\sigma + it) \to \varpi_{\mathrm{RVM}}$ as $\lvert t\rvert \to \infty$ (8.33)	almost periodic on every line of absolute convergence; $\lambda = 1$: no limit as $\lvert t\rvert \to \infty$ (Proposition 13.15)
half-plane bounds	$\lambda = 1$: $C^{-1} \leq \lvert\zeta_{\mathrm{RVM}}\rvert \leq C$ on $\Re s \geq \sigma_0 > 1$ (Theorem 8.10); every λ: $\lvert \log\zeta_\lambda - \log\frac{s}{s-1}\rvert \ll_c 1 + \log(\lvert t\rvert + 2)$ on $\Re s \geq c > \frac12$	$\lvert\tilde\zeta_{\mathbb{Z},\lambda}\rvert \leq \sum_N \lvert\tilde a_N(\lambda)\rvert N^{-\sigma_0}$ and $\lvert\tilde\zeta_{\mathbb{Z},\lambda}^{-1}\rvert \leq \sum_N \lvert\tilde\mu_\lambda(N)\rvert N^{-\sigma_0}$ on $\Re s \geq \sigma_0 > 1$, for every λ
prime coordinate	counts J_λ, π_λ (distributions, (12.16))	masses $\tilde c_n(\lambda)$ at the integers; $\tilde c_n(1) = O(1)$; mean square $X^{1+\varepsilon}$ on $(X, 2X]$ for every λ (Theorem 13.10)
reciprocal	$F_\lambda = 1/\zeta_\lambda$; F_{Z1} is the Mellin transform of B_{Z1} (Proposition 13.16)	Dirichlet series with coefficients $\tilde\mu_\lambda(N)$ (13.31)
Mertens	$M_{\mathrm{RVM}}^{\mathrm{crit}} = L + O_\varepsilon(x^{1/2+\varepsilon})$ (Theorem 8.12); $B_{Z1} = M \iff (D)$	$\tilde M_\lambda = \sum_{N\leq x} \tilde\mu_\lambda(N)$ exists for every λ (13.32); the bound $x^{1/2+\varepsilon}$ is open (it needs control of $e^{-\tilde h_\lambda}$)
flow in λ	generator $\mathcal{V}_\lambda$ on the Hadamard product; $\Upsilon(\lambda)$ at $+\infty$ (Proposition 13.2)	$\log\tilde\zeta_{\mathbb{Z},\lambda} = \sum_n \tilde c_n(\lambda)\big(-\log(1-n^{-s})\big)$, each mass differentiable in λ by (13.9); joining factor (13.15)
limit $\lambda \to 0^+$	$\zeta_\lambda \to \zeta_{Z1}$	$\tilde c_n(\lambda) \to \tilde c_n(0)$, $\mathfrak{h}_n(\lambda;\cdot) \to \mathfrak{h}_n(0;\cdot)$ (Proposition 13.8)

Topic of the paper	continuous family ζ_λ	atomized family $\tilde\zeta_{\mathbb{Z},\lambda}$
endpoint: Hamburger's hypotheses	H0 and H2 proved; H1 is (D), open	H1 proved in $\Re s > 1$ (Corollary 13.11); H0 and H2 open
relation between the endpoints	$\tilde\zeta_{\mathbb{Z},0} = \zeta_{Z1}\, e^{\tilde h_0}$	same
what closes the route	ζ_{Z1} satisfies H1	$\tilde\zeta_{\mathbb{Z},0}$ satisfies H0 and H2
common form of the closure	$\tilde h_0 \equiv 0$	$\tilde h_0 \equiv 0$
weaker forms of the closure	ζ_{Z1} has a positive representing measure (Theorem 14.7)	$e^{\tilde h_0}$ of finite order and symmetric under $s \mapsto 1-s$ (Theorem 14.3)
consequence of the closure	$\zeta_{Z1} = \zeta$; Step 3 gives $M(x) = O_\varepsilon(x^{1/2+\varepsilon})$ (Theorem 11.8)	$\tilde a_N(0) = 1$, $\tilde c_n(0) = \mathbf{1}_{\text{prime}}(n)$, $\tilde\mu_0 = \mu$, $\tilde M_0 = M$; $\tilde\zeta_{\mathbb{Z},0} = \zeta = \zeta_{Z1}$

How the two routes complement each other. Along λ each family keeps exactly the half of Hamburger's hypotheses that the other one lacks:

$$\underbrace{\zeta_\lambda}_{\substack{\text{keeps the functional equation}\\ \text{and the zeros on } \Re s=\frac12}} \quad \longleftrightarrow \quad \underbrace{\tilde\zeta_{\mathbb{Z},\lambda}}_{\substack{\text{keeps the Dirichlet series}\\ \text{and the Euler product}}}, \qquad \tilde\zeta_{\mathbb{Z},\lambda} = \zeta_\lambda\, e^{\tilde h_\lambda}. \tag{13.34}$$

The closure is the same in both columns, the vanishing of the joining factor at the endpoint: by (13.13) and Theorem 13.13,

$$\tilde h_0 \equiv 0 \iff J_{Z1} = 0 \text{ on } (1,2), \quad \mathfrak{h}_n(0;s) = 0 \text{ for all } n \ge 2,\ \Re s > 1, \text{ and } \varpi_{Z1} = 1. \tag{13.35}$$

No further condition is needed in the atomized column: the cell masses have mean square $X^{1+\varepsilon}$ uniformly in λ (Theorem 13.10), so $\tilde\zeta_{\mathbb{Z},0}$ satisfies H1 (Corollary 13.11).

13.8. **Programme.** The steps are numbered in the order in which they are followed. The status column is updated with each step.

Step	Statement	Status
A1	Atomization by smooth cells, coefficients $\tilde a_N(\lambda)$	proved (Definition 13.4)
A2	Joining factor and explicit cell defects, (13.13)–(13.17)	proved (Thm. 13.6, Prop. 13.7)
A3	L^2 square-root estimate for every λ; Dirichlet series in $\Re s > \frac32$	proved (Theorem 13.12); superseded by A6
A4	$\tilde h_\lambda$ holomorphic in $\Re s > \frac12$; in $\Re s > 0$ for $\lambda = 1$	proved (Theorem 13.12)
A5	Limits $\lambda \to 0^+$ of $\tilde c_n$, $\mathfrak{h}_n$, ϖ_λ; flow at infinity Υ	proved (Propositions 13.2, 13.8)
A5′	Endpoints: $\tilde h_1$ not constant; atomized Möbius coefficients $\tilde\mu_\lambda$; Mellin form of M_{Z1}	proved (Propositions 13.15, 13.16)
A6	Mean square of the cell masses uniformly in λ; Dirichlet series in $\Re s > 1$ for every λ, including $\lambda = 0$	proved (Theorem 13.10, Corollary 13.11)
A7	Cell identity (13.29) for every $n \ge 2$, and $J_{Z1} = 0$ on the first cell $(1,2)$	open; reduced to one number for each cell in A20
A8	Equivalent scalar form: $\int_{0^+}^1 \Upsilon\, d\lambda = \log L$, i.e. (12.12)	open; follows from A7
A9	$\zeta_{Z1} = \zeta$ by Hamburger; $M(x) = O_\varepsilon(x^{1/2+\varepsilon})$ by Step 3	proved as implication from A7 (Thms. 13.13, 11.8)
A10	Continuation of $\tilde h_\lambda$ to $0 < \Re s \le \frac12$ for $\lambda < 1$	open; not needed for A9; at $\lambda = 0$ it follows from (B1), (B2) (A23)
A11	ζ_λ is not a Dirichlet series and $\tilde h_\lambda$ is not constant for $0 < \lambda \le 1$; convexity of $\log \zeta_\lambda(\sigma)$ in λ	proved (Propositions 14.1, 14.2)

Step	Statement	Status
A12	Hamburger's theorem for the pair: A9 follows if $e^{\tilde{h}_0}$ is of finite order and symmetric under $s \mapsto 1-s$	proved as implication (Theorem 14.3); at $\lambda = 0$ the hypothesis is equivalent to (D) (Remark 14.30)
A13	Positivity: A9 follows if ζ_{Z1} has a positive representing measure; then $\varpi_{Z1} \le 1$	proved as implication (Theorem 14.7, Corollary 14.8)
A14	Clock end: density of $d\pi_{\mathrm{RVM}}$, cell masses $\frac{1}{\log n} + O(n^{-1/4})$, defects $O(\lvert s\rvert n^{-\sigma-4/3})$	proved (Theorem 14.15, Corollary 14.16)
A15	Height kernel of the cell masses; flow as count difference; Gaussian defect and the density condition $\tilde{h}_0(1) = 0$	proved (Propositions 14.17, 14.18, 14.22)
A16	Hamburger's theorem for the pair without continuation: A9 follows if $T_0 \equiv 0$, or if $\mathsf{V}_\sigma(0) = 0$; symmetry fails at $\lambda = 1$; exceptional set finite	proved (Theorem 14.4, Propositions 14.5, 14.25)
A17	$\zeta_{Z1}(\sigma) \ge \zeta(\sigma)$ for $\sigma > 1$; A9 follows from the reverse inequality at one σ, or from $\varpi_\lambda \le 1$ along $\lambda \to 0^+$, or from $\mathsf{W}(0) \ge 0$	proved (Theorem 14.11, Propositions 14.10, 14.12)
A18	Space $\mathscr{O}$ of ordinate lists: at most one coincidence list; A9 is equivalent to its existence	proved (Theorem 15.1, Corollary 15.3)
A19	Existence of a list in $\mathscr{O}$ with $\tilde{h}_{\boldsymbol{\kappa}}$ constant, equivalently $\min_{\mathscr{O}} \mathsf{V}_\sigma = 0$	open; equivalent to A7
A20	A9 holds if and only if $\tilde{c}_n(0) = \mathbf{1}_{\mathrm{prime}}(n)$ for every n; every $\lambda \in \mathsf{E}$ is a witness; the identity (14.59) cell by cell	proved (Corollary 14.26, Propositions 14.27, 14.29); the identity (14.59) itself is open
A21	Symmetric joining factors: order one is needed, and A12 has no slack at $\lambda = 0$	proved (Proposition 14.31, Remarks 14.30, 14.32)
A22	Half-line form: A9 follows if $\tilde{h}_0(\sigma) = O_N(\sigma N^{-\sigma})$ for every N; the Nth rung is the constancy of J^{M}_{Z1} on $(N-1, N)$	proved (Theorems 14.33, 14.34, Corollary 14.35)
A23	Bridge theorem: A9 follows if $\Im\tilde{h}_0$ vanishes on $\Re s = \frac{1}{2}$ and $\tilde{h}_0$ is temperate; the two conditions hold for no $\lambda > 0$	proved (Theorem 14.36, Corollary 14.37); (B1), (B2) at $\lambda = 0$ open
A24	Transport from the clock: S_{f} entire of type $C_\eta \log N$, moment series on the clock, defect of the clock by Poisson summation	proved (Proposition 14.41, Theorem 14.42, Proposition 14.43)
A24b	Multiscale Jacobian at the clock: stable finite ordinate block on windows $U \asymp \log T$, fixed-window obstruction, dense range, and rapidly summable smooth tails	proved (Theorem 14.44); quantitative multiscale continuation to a solution open
A25	Rung system with free masses: A9 holds if and only if it has a solution in $\mathscr{O}$; the solution is unique and nondegenerate	proved (Theorems 15.5, 15.6); existence open
A26	Gradient of V_σ on $\mathscr{O}$; A9 follows if every critical point is a zero	proved (Proposition 15.8); the hypothesis is open
A27	Acceptance criterion and models: the final step is a theorem for the finite Euler products; no object of finite rank in the window of ζ	proved (Theorems 16.1, 16.3, Propositions 16.2, 16.4)

Remark 13.17 (Input for A7). Steps A1–A6 use only two properties of an ordinate list: bounded displacement from the clock and a primitive of the count difference of size $O(\log \tau)$. The list (τ_k) has both properties, and for it the corresponding function ζ_{RVM} is not a Dirichlet series (Theorem 8.11). Step A7 must therefore use a property of the ordinates γ_k that the clock ordinates do not have; the natural candidate is the arithmetic of ζ carried by the γ_k, for instance Landau's formula $\sum_{0<\gamma\le T} x^\rho = -\frac{T}{2\pi}\Lambda(x) + O(\log T)$, in which the prime powers appear as resonances of the ordinates. The multiscale analysis below separates a different issue: the linearized clock has enough directions,

but only on windows whose length grows like $\log T$; thus the obstruction is not fixed-height injectivity but quantitative nonlinear continuation across scales.

14. Criteria for the final step and the clock end of the family

This section collects the consequences of Corollary 13.11 for the final step. Along the open path $0 < \lambda \le 1$ no member ζ_λ is a Dirichlet series (Proposition 14.1); the family is convex in λ on the real axis (Proposition 14.2); Hamburger's theorem applies to the pair $(\zeta_\lambda, \tilde\zeta_{\mathbb{Z},\lambda})$ as soon as the joining factor is symmetric (Theorem 14.3); positivity of the measure representing ζ_{Z1} gives $\varpi_{Z1} \le 1$ and closes the route (Theorem 14.7). At the clock end the density of $d\pi_{\mathrm{RVM}}$ is computed explicitly (Theorem 14.15), and the flow of the cell masses is written in the heights of the ordinates (Propositions 14.17 and 14.18). Subsections 14.9–14.13 reduce the final step to the cell masses (Corollary 14.26), state two criteria whose hypotheses are written where the joining factor is defined, the half-line form (Theorem 14.34) and the bridge theorem (Theorem 14.36), and compute the transport from the clock on each cell (Theorem 14.42).

Throughout this section the function χ of Subsection 13.4 is even, so that φ is even, and

$$\hat\varphi(\xi) := \int_{\mathbb{R}} \varphi(t) e^{-2\pi i \xi t}\, dt, \qquad \mathsf{M}_r := \int_{\mathbb{R}} t^r \varphi(t)\, dt, \qquad \mathsf{C}_r := \int_{\mathbb{R}} t^r \varphi(t) \cos(2\pi t)\, dt. \tag{14.1}$$

$\hat\varphi$ is real and even, and $\mathsf{M}_1 = \mathsf{C}_1 = 0$. Since $\sum_{m\in\mathbb{Z}} \varphi(t-m) = 1$,

$$\hat\varphi(q) = \int_0^1 \sum_{m\in\mathbb{Z}} \varphi(t-m)\, e^{-2\pi i q t}\, dt = \delta_{q0} \qquad (q \in \mathbb{Z}), \tag{14.2}$$

so $\mathsf{M}_0 = 1$ and $\mathsf{C}_0 = \hat\varphi(1) = 0$.

14.1. **The open path.**

Proposition 14.1 (No member of the open path is a Dirichlet series)**.** *Let $0 < \lambda \le 1$. Then ζ_λ is not a constant multiple of a Dirichlet series $\sum_n a_n n^{-s}$ converging absolutely in $\Re s > 1$. In particular the joining factor $\tilde h_\lambda$ is not constant.*

Proof. Suppose $\zeta_\lambda(s) = \sum_n a_n n^{-s}$ absolutely for $\Re s > 1$. By Theorem 10.1, exactly as in Proposition 11.1, $(s-1)\zeta_\lambda(s)$ is entire of order one and ζ_λ satisfies (11.3); these are the hypotheses of Hamburger's theorem, so $\zeta_\lambda = C\zeta$ with $C \ne 0$. The nontrivial zeros of ζ with positive ordinate are then $\frac12 + i\gamma_k(\lambda)$, $k \ge 1$, with multiplicity. The ordinates of ζ, listed in non-decreasing order, are the γ_k, and the $\gamma_k(\lambda)$ are increasing; hence $\gamma_k(\lambda) = \gamma_k$, that is $\lambda\eta_k = 0$, for every k. Since $\lambda > 0$, every η_k vanishes, all members of the family coincide (Remark 10.2), and $\zeta_{\mathrm{RVM}} = \zeta_\lambda$ is an absolutely convergent Dirichlet series, against Theorem 8.11. If $\tilde h_\lambda$ were a constant c, then $\zeta_\lambda = e^{-c}\tilde\zeta_{\mathbb{Z},\lambda}$ would be such a series by Corollary 13.11. □

The two columns of (13.34) exist for every λ, and for every $\lambda \in (0,1]$ they are different functions. The joining factor converges, $\tilde h_\lambda \to \tilde h_0$ as $\lambda \to 0^+$, through non-constant functions; (D) is the statement that the limit is constant. A property of the displacements that yields (D) holds for the sequence (η_k) and fails for every sequence $(c\,\eta_k)$ with $0 \le c < 1$, because the ordinates $\gamma_k(\lambda) = \tau_k - (1-\lambda)\eta_k$ are obtained from the clock by the displacements $(1-\lambda)\eta_k$.

14.2. **Convexity along the family.**

Proposition 14.2 (Convexity on the real axis)**.** *For every real $\sigma > 1$ the function $\lambda \mapsto \log\zeta_\lambda(\sigma)$ is strictly convex on $[0,1]$:*

$$\partial_\lambda^2 \log\zeta_\lambda(\sigma) = \sum_{k\ge1} \eta_k^2 \Big(\mathcal{C}(\sigma - \tfrac12, \gamma_k(\lambda)) - \mathcal{C}(\tfrac12, \gamma_k(\lambda)) \Big) > 0, \qquad \mathcal{C}(a,\gamma) := \frac{2(a^2-\gamma^2)}{(a^2+\gamma^2)^2}. \tag{14.3}$$

Consequently, with the generator $\mathcal{V}_\lambda$ of Theorem 10.3,

$$\log\zeta_{\mathrm{RVM}}(\sigma)-\mathcal{V}_1(\sigma)<\log\zeta_{Z1}(\sigma)<\log\zeta_{\mathrm{RVM}}(\sigma)-\mathcal{V}_0(\sigma)\qquad(\sigma>1).\tag{14.4}$$

Proof. With $a=\sigma-\frac12>\frac12$ one has $q(\sigma)=a^2-\frac14$, so by (10.2) $\log\zeta_\lambda(\sigma)=-\log G(\sigma)+\sum_k\big(\log(a^2+\gamma_k(\lambda)^2)-\log(\frac14+\gamma_k(\lambda)^2)\big)$. Since $\partial_\gamma^2\log(a^2+\gamma^2)=\mathcal{C}(a,\gamma)$ and $\partial_\lambda\gamma_k(\lambda)=\eta_k$, termwise differentiation gives the series in (14.3); its terms are $O(\eta_k^2\gamma_k^{-2})$, so it converges uniformly on $[0,1]$. Fix $\gamma>\frac12$. Then $\mathcal{C}(\frac12,\gamma)<0$; $\partial_a\mathcal{C}(a,\gamma)=4a(3\gamma^2-a^2)(a^2+\gamma^2)^{-3}>0$ for $0<a<\sqrt3\,\gamma$; and $\mathcal{C}(a,\gamma)>0$ for $a>\gamma$. Hence $\mathcal{C}(a,\gamma)>\mathcal{C}(\frac12,\gamma)$ for every $a>\frac12$. All ordinates satisfy $\gamma_k(\lambda)>\tau_0>\frac12$ and $\eta_1=\tau_1-\gamma_1\neq0$ ($\tau_1=14.52\ldots$ by (2.5), $\gamma_1=14.13\ldots$), so the sum is positive. A strictly convex function F on $[0,1]$ satisfies $F(1)-F'(1)<F(0)<F(1)-F'(0)$; with $F(\lambda)=\log\zeta_\lambda(\sigma)$ and $F'=\mathcal{V}_\lambda(\sigma)$ (10.11) this is (14.4). □

As $\sigma\to+\infty$, (14.3) becomes $-\Upsilon'(\lambda)>0$ and (14.4) becomes the last statement of Proposition 13.2. The sign in (14.3) comes from the test function $x^{-\sigma}$, whose transform is not oscillatory in the ordinate; for the cell functions φ_n the corresponding second derivative $\partial_\lambda^2\tilde c_n(\lambda)=-2\Re\sum_k\eta_k^2W_n''(\gamma_k(\lambda))$ has terms of both signs.

14.3. **Hamburger's theorem for the pair.**

Theorem 14.3 (Hamburger's theorem for the pair $(\zeta_\lambda,\tilde\zeta_{\mathbb{Z},\lambda})$)**.** *Let $\lambda\in[0,1]$. Suppose that the joining factor $e^{\tilde h_\lambda}$ extends to a meromorphic function on $\mathbb{C}$ of finite order with finitely many poles, and that*

$$e^{\tilde h_\lambda(s)}=e^{\tilde h_\lambda(1-s)}.\tag{14.5}$$

Then $\tilde\zeta_{\mathbb{Z},\lambda}=\zeta$. For $\lambda=0$, moreover, $\zeta_{Z1}=\zeta$ and $\tilde h_0\equiv0$.

Proof. Put $f=\tilde\zeta_{\mathbb{Z},\lambda}=\zeta_\lambda e^{\tilde h_\lambda}$. By Theorem 10.1, $(s-1)\zeta_\lambda$ is entire of order one, so $f=\mathsf{G}/\mathsf{P}$ with G entire of finite order and P a polynomial: this is (H0). By (11.3) for ζ_λ and (14.5), $f(s)\Gamma(\frac s2)\pi^{-s/2}=f(1-s)\Gamma(\frac{1-s}2)\pi^{-(1-s)/2}$: this is (H2) with $g=f$. (H1) is Corollary 13.11, and it gives (H3) for $g=f$. Hamburger's theorem gives $f=C\zeta$, and $C=1$ because $\tilde a_1(\lambda)=1$. By Theorem 13.12(iii) and the functional equation, the nontrivial zeros of $f=\zeta$ lie on $\Re s=\frac12$; they are the points $\frac12\pm i\gamma_k$. For $\lambda=0$, Theorem 11.2 then gives $\zeta_{Z1}=\zeta=\tilde\zeta_{\mathbb{Z},0}$, and $e^{\tilde h_0}=1$. □

Theorem 14.3 uses the functional equation of ζ_λ and the Dirichlet series of $\tilde\zeta_{\mathbb{Z},\lambda}$ as they stand; the only hypothesis is on the factor that joins them, and it asks for symmetry instead of constancy. At $\lambda=1$ the two hypotheses do not hold together: the conclusion $\tilde\zeta_{\mathbb{Z},1}=\zeta$ would place the zeros of ζ at $\frac12\pm i\tau_k$ (Theorem 13.12(iv)), so $\zeta=\zeta_{\mathrm{RVM}}$ by Theorem 9.1, against Theorem 8.11.

Theorem 14.4 (Real-variable and vertical forms of Hamburger's theorem for the pair)**.** *For $\lambda\in[0,1]$, $y>0$ and $\sigma>1$ put*

$$T_\lambda(y):=\sum_{N\ge1}\tilde a_N(\lambda)\,e^{-\pi N^2y}-y^{-1/2}\sum_{N\ge1}\tilde a_N(\lambda)\,e^{-\pi N^2/y}-\tfrac12(y^{-1/2}-1),\tag{14.6}$$

$$k_n(\lambda):=\sum_{d\mid n}\tilde a_d(\lambda)\,\mu(n/d),\qquad \mathsf{V}_\sigma(\lambda):=\sum_{n\ge2}|k_n(\lambda)|^2\,n^{-2\sigma}.\tag{14.7}$$

(i) *$\tilde\zeta_{\mathbb{Z},\lambda}/\zeta=\sum_{n\ge1}k_n(\lambda)n^{-s}$ converges absolutely in $\Re s>1$, $k_1(\lambda)=1$, and*

$$\lim_{T\to\infty}\frac{1}{2T}\int_{-T}^{T}\left|\frac{\tilde\zeta_{\mathbb{Z},\lambda}(\sigma+it)}{\zeta(\sigma+it)}\right|^2dt=1+\mathsf{V}_\sigma(\lambda)\ \ge1.\tag{14.8}$$

(ii) *For each λ the following are equivalent:* (a) $\tilde\zeta_{\mathbb{Z},\lambda}=\zeta$; (b) $T_\lambda(y)=0$ *for every $y>0$;* (c) $\mathsf{V}_\sigma(\lambda)=0$ *for some, equivalently every, $\sigma>1$. Each of them implies (D).*

(iii) $\lambda \mapsto T_\lambda(y)$ *and* $\lambda \mapsto \mathsf{V}_\sigma(\lambda)$ *are continuous on* $[0,1]$*, and*

$$\text{(D)} \iff T_0 \equiv 0 \iff \mathsf{V}_\sigma(0) = 0 \iff \lim_{\lambda\to 0^+} T_\lambda(y) = 0 \ \ (y>0) \iff \lim_{\lambda\to 0^+} \mathsf{V}_\sigma(\lambda) = 0. \tag{14.9}$$

At the clock end $T_1 \not\equiv 0$ *and* $\mathsf{V}_\sigma(1) > 0$.

Proof. (i) $1/\zeta = \sum_n \mu(n) n^{-s}$ and $\tilde\zeta_{\mathbb{Z},\lambda}$ (Corollary 13.11) converge absolutely in $\Re s > 1$, and so does their product, with $k_1 = \tilde a_1 = 1$. The mean square of an absolutely convergent Dirichlet series on a vertical line is $\sum_n |k_n|^2 n^{-2\sigma}$, as in (8.43)–(8.44). (ii) (a)$\Leftrightarrow$(c) because $\mathsf{V}_\sigma(\lambda) = 0$ means $k_n(\lambda) = 0$ for $n \ge 2$. (a)$\Rightarrow$(b) is the theta relation $\sum_{n\ge1} e^{-\pi n^2 y} = y^{-1/2} \sum_{n\ge1} e^{-\pi n^2/y} + \frac12(y^{-1/2} - 1)$. (b)$\Rightarrow$(a): the relation $T_\lambda \equiv 0$ is the identity reached in the first half of the proof of Hamburger's theorem in [37, §2.13], with $b_n = a_n = \tilde a_n(\lambda)$; the second half of that proof uses only this identity and the absolute convergence of $\sum_n a_n n^{-\sigma'}$ for $\sigma' > 1$, and gives $a_k = a_{k+1}$ for every k, hence $\tilde a_N(\lambda) = \tilde a_1(\lambda) = 1$. If (a) holds, then by Theorem 13.12(iii) and the functional equation the nontrivial zeros of ζ lie on $\Re s = \frac12$; they are the points $\frac12 \pm i\gamma_k$, and Theorem 11.2 gives (D). (iii) By Theorem 13.10, for every $\sigma' > 1$ the sums $\sum_n |\tilde c_n(\lambda)| n^{-\sigma'}$ are bounded uniformly in λ; hence so are $\sum_N |\tilde a_N(\lambda)| N^{-\sigma'} \le \exp\bigl(\sum_m \frac1m \sum_n |\tilde c_n(\lambda)| n^{-m\sigma'}\bigr)$ and $\sum_n |k_n(\lambda)| n^{-\sigma'}$. Each $\tilde a_N(\lambda)$ and each $k_n(\lambda)$ is a polynomial in finitely many cell masses, which are continuous in λ. The tails $\sum_{N>N_0}$ of (14.6) and of (14.7) (with $1 < \sigma' < \sigma$) are small uniformly in λ, which gives the continuity. Under (D), $\tilde a_N(0) = 1$ (Theorem 13.13), so $T_0 \equiv 0$ and $\mathsf{V}_\sigma(0) = 0$; the converse is (ii) at $\lambda = 0$, and the two limits are the values at $\lambda = 0$ by continuity. Finally $\tilde\zeta_{\mathbb{Z},1} \ne \zeta$ by the remark after Theorem 14.3. □

Theorem 14.4 needs no continuation of the joining factor: (14.6) is a real function of y built from the atomized coefficients alone, and (14.8) lives in $\Re s > 1$, where

$$\frac{\tilde\zeta_{\mathbb{Z},\lambda}}{\zeta} = e^{\tilde h_\lambda} \frac{\zeta_\lambda}{\zeta} \tag{14.10}$$

is a quotient of three explicit functions. Its vertical mean is 1 and its vertical mean square is at least 1 for every λ; (D) is the statement that at $\lambda = 0$ the mean square does not exceed 1.

Proposition 14.5 (At the clock end the symmetry fails)**.** *The joining factor* $e^{\tilde h_1}$ *has no meromorphic continuation to* $\mathbb{C}$ *that satisfies* (14.5)*.*

Proof. $e^{\tilde h_1}$ is holomorphic and zero-free in $\Re s > 0$ (Theorem 13.12(iv)). A meromorphic continuation with (14.5) is holomorphic and zero-free also in $\Re s < 1$, hence entire and zero-free, so it equals e^g with g entire, $g = \tilde h_1$ on $\Re s > 0$, and $g(s) - g(1-s)$ constant. For $\Re s \ge \frac14$, the bound $|\mathfrak{h}_n(1; ms)| \le C\, m|s|\, n^{-m\sigma-1}$ of Theorem 13.12(iv) gives $\sum_m \frac1m \sum_n |\mathfrak{h}_n(1; ms)| \le C|s| \sum_{n\ge2} n^{-1}(n^{1/4} - 1)^{-1}$, and $|\mathfrak{u}_1(s)| \le C(1 + |s|)$ by (13.11); so $|g(s)| \le C(1 + |s|)$ on $\Re s \ge \frac14$, and by the symmetry also on $\Re s \le \frac34$. An entire function of linear growth is a polynomial of degree at most 1, and $g(s) - g(1-s)$ constant then forces g to be constant. This contradicts Proposition 13.15(iii). □

At $\lambda = 1$ it is therefore the symmetry (14.5) itself that fails in Theorem 14.3, whatever the order of the continuation.

14.4. **Positivity and the value at infinity.** For an even Schwartz function g on $\mathbb{R}$ put $\hat{\mathsf{g}}(\xi) = \int_{\mathbb{R}} \mathsf{g}(u) e^{-2\pi i \xi u} du$ and $\mathsf{g}^\sharp(s) = \int_0^\infty \mathsf{g}(u) u^{s-1} du$.

Lemma 14.6 (Summation formula for a representing measure)**.** *Let* $\lambda \in [0,1]$ *and let* A *be a positive measure on* $[1,\infty)$ *with*

$$\zeta_\lambda(s) = \int_{[1,\infty)} u^{-s}\, d\mathsf{A}(u) \qquad (\Re s > 1). \tag{14.11}$$

Then for every even Schwartz function g

$$\mathsf{g}(0) + 2\int_{[1,\infty)} \mathsf{g}\, d\mathsf{A} = \hat{\mathsf{g}}(0) + 2\int_{[1,\infty)} \hat{\mathsf{g}}\, d\mathsf{A}. \tag{14.12}$$

Proof. $\mathsf{g}^\sharp$ is meromorphic with simple poles at $s=0,-2,-4,\dots$, residue $\mathsf{g}(0)$ at $s=0$, and decays faster than any power of $|t|$ on vertical strips; $\mathsf{g}^\sharp(1)=\frac12\hat{\mathsf{g}}(0)$; and

$$\pi^{-s/2}\Gamma(\tfrac{s}{2})\,\hat{\mathsf{g}}^\sharp(1-s)=\pi^{-(1-s)/2}\Gamma(\tfrac{1-s}{2})\,\mathsf{g}^\sharp(s) \tag{14.13}$$

for $0<\Re s<1$, and then by meromorphic continuation: the Fourier transform of the even homogeneous distribution $|\xi|^{-s}$ is a multiple of $|u|^{s-1}$, and the multiple is fixed by $\mathsf{g}(u)=e^{-\pi u^2}$, for which $\hat{\mathsf{g}}=\mathsf{g}$ and $\mathsf{g}^\sharp(s)=\frac12\pi^{-s/2}\Gamma(\frac{s}{2})$. Fix $1<c<2$. By Mellin inversion and Fubini, $\int \mathsf{g}\,d\mathsf{A}=\frac{1}{2\pi i}\int_{(c)}\zeta_\lambda(s)\mathsf{g}^\sharp(s)\,ds$, since $\int u^{-c}d\mathsf{A}<\infty$. On $\Re s\ge c$, $|\zeta_\lambda(s)|\le\zeta_\lambda(c)$; with the functional equation (11.3), valid for every ζ_λ by Theorem 10.1, and the finite order of $(s-1)\zeta_\lambda$, the Phragmén–Lindelöf principle bounds ζ_λ by a power of $|t|$ on $1-c\le\Re s\le c$, $|t|\ge1$. Moving the line to $\Re s=1-c$ crosses $s=1$, where ζ_λ has residue 1, and $s=0$, where $\zeta_\lambda(0)=P_\lambda(0)/G(0)=-\frac12$:

$$\int \mathsf{g}\,d\mathsf{A}=\tfrac12\hat{\mathsf{g}}(0)-\tfrac12\mathsf{g}(0)+\frac{1}{2\pi i}\int_{(1-c)}\zeta_\lambda(s)\mathsf{g}^\sharp(s)\,ds. \tag{14.14}$$

By (11.3) and (14.13), $\zeta_\lambda(s)\mathsf{g}^\sharp(s)=\zeta_\lambda(1-s)\hat{\mathsf{g}}^\sharp(1-s)$, and the last integral is $\frac{1}{2\pi i}\int_{(c)}\zeta_\lambda(s)\hat{\mathsf{g}}^\sharp(s)\,ds=\int\hat{\mathsf{g}}\,d\mathsf{A}$. □

Theorem 14.7 (Positivity criterion)**.** *Let $\lambda\in[0,1]$. If ζ_λ has a representation* (14.11) *with a positive measure* A*, then*

$$\varpi_\lambda=\mathsf{A}(\{1\})\le1. \tag{14.15}$$

In particular, if ζ_{Z1} has such a representation, then $\varpi_{Z1}=1$ and $\zeta_{Z1}=\zeta$.

Proof. By monotone convergence $\zeta_\lambda(\sigma)=\int u^{-\sigma}d\mathsf{A}\to\mathsf{A}(\{1\})$ as $\sigma\to+\infty$, and the limit is ϖ_λ (Proposition 13.1). Let $\mathsf{T}(\xi)=(1-|\xi|)_+$ and $\mathsf{B}=\mathsf{T}*\mathsf{T}$, so that $\mathsf{B}(\xi)=\frac23-\xi^2+\frac12|\xi|^3$ for $|\xi|\le1$, $\mathsf{B}(\xi)=\frac16(2-|\xi|)^3$ for $1\le|\xi|\le2$, $\mathsf{B}=0$ elsewhere, and $\mathsf{B}''(\xi)=\mathsf{T}(\xi+1)-2\mathsf{T}(\xi)+\mathsf{T}(\xi-1)$. Put

$$\mathsf{g}(u):=\Big(\frac{\sin\pi u}{\pi u}\Big)^4(1-u^2),\qquad \hat{\mathsf{g}}(\xi)=\mathsf{B}(\xi)+\frac{1}{4\pi^2}\mathsf{B}''(\xi). \tag{14.16}$$

Then $\mathsf{g}(0)=1$; $\mathsf{g}(u)\le0$ for $|u|\ge1$; $\hat{\mathsf{g}}(0)=\frac23-\frac{1}{2\pi^2}$ and $\hat{\mathsf{g}}(1)=\frac16+\frac{1}{4\pi^2}$; and $\hat{\mathsf{g}}\ge0$ on $\mathbb{R}$: for $0\le\xi\le1$, $\hat{\mathsf{g}}(\xi)=\frac23-\xi^2+\frac12\xi^3+\frac{3\xi-2}{4\pi^2}$ is positive at $\xi=0$ and $\xi=1$ and its derivative $-2\xi+\frac32\xi^2+\frac{3}{4\pi^2}$ changes sign once, from positive to negative, and for $1\le\xi\le2$, $\hat{\mathsf{g}}(\xi)=\frac16(2-\xi)^3+\frac{2-\xi}{4\pi^2}\ge0$. For $\epsilon>0$ let $\mathsf{g}_\epsilon(u)=\mathsf{g}(u)e^{-\epsilon\pi u^2}$, an even Schwartz function with $\mathsf{g}_\epsilon(0)=1$, $\mathsf{g}_\epsilon\le0$ on $|u|\ge1$, and $\hat{\mathsf{g}}_\epsilon=\hat{\mathsf{g}}*(\epsilon^{-1/2}e^{-\pi\xi^2/\epsilon})\ge0$. By Lemma 14.6, since $\mathsf{A}\ge0$ is carried by $[1,\infty)$ and has mass ϖ_λ at $u=1$,

$$1\ \ge\ \mathsf{g}_\epsilon(0)+2\int\mathsf{g}_\epsilon\,d\mathsf{A}=\hat{\mathsf{g}}_\epsilon(0)+2\int\hat{\mathsf{g}}_\epsilon\,d\mathsf{A}\ \ge\ \hat{\mathsf{g}}_\epsilon(0)+2\varpi_\lambda\,\hat{\mathsf{g}}_\epsilon(1). \tag{14.17}$$

Letting $\epsilon\to0$, $\hat{\mathsf{g}}_\epsilon\to\hat{\mathsf{g}}$ pointwise because $\hat{\mathsf{g}}$ is continuous, and

$$\varpi_\lambda\le\frac{1-\hat{\mathsf{g}}(0)}{2\hat{\mathsf{g}}(1)}=\frac{\frac13+\frac{1}{2\pi^2}}{\frac13+\frac{1}{2\pi^2}}=1. \tag{14.18}$$

The last statement follows from Theorem 12.2. □

Corollary 14.8 (Positive form of the final step)**.** *The following are equivalent:* (i) *(D);* (ii) *ζ_{Z1} is the Mellin–Stieltjes transform* (14.11) *of a positive measure on $[1,\infty)$;* (iii) *J^{M}_{Z1}, extended by* 0 *to $(0,1]$, is non-decreasing on $(0,\infty)$.*

Proof. (i)⇒(iii): then $J^{\mathrm{M}}_{Z1}=J$ (Theorem 12.7). (iii)⇒(ii): $J^{\mathrm{M}}_{Z1}(x)x^{-c}\in L^2((1,\infty),dx/x)$ for every $c>1$, so a non-decreasing J^{M}_{Z1} is $O_c(x^c)$ and dJ^{M}_{Z1} is a positive measure on $[1,\infty)$ with $\int x^{-\sigma}dJ^{\mathrm{M}}_{Z1}<\infty$ for $\sigma>1$. By Theorem 12.5, $\zeta_{Z1}(s)=\exp\int x^{-s}dJ^{\mathrm{M}}_{Z1}=\int u^{-s}d\mathsf{A}(u)$, where $\mathsf{A}=\sum_{r\ge0}\frac{1}{r!}(dJ^{\mathrm{M}}_{Z1})^{*r}$ (multiplicative convolution powers) is a positive measure on $[1,\infty)$. (ii)⇒(i) is Theorem 14.7. □

In Theorem 14.7 the Dirichlet series is not used: (H1) is replaced by the positivity of the representing measure, the functional equation gives $\varpi_{Z1} \le 1$ through (14.17), and the Hadamard product gives $\varpi_{Z1} \ge 1$ (Theorem 12.2). The function (14.16) is extremal for the integers: for $\mathsf{A} = \sum_{n\ge1} \delta_n$ both inequalities in (14.17) are equalities in the limit, since $\mathsf{g}(n) = 0$ for $n \ge 1$ and $\hat{\mathsf{g}}(n) = 0$ for $n \ge 2$. At the clock end $\varpi_{\mathrm{RVM}} < 1$, and the density of dJ_{RVM} takes negative values in every cell $(n, n+1)$ with n large (Theorem 14.15).

Lemma 14.9 (Mellin form and summation formula for every member)**.** *There is $c_1 \ge 2$ such that for every $\lambda \in [0,1]$ there is a unique measurable function A_λ on $(0,\infty)$, vanishing on $(0,1)$, with $x^{-c}\mathsf{A}_\lambda \in L^2((1,\infty), dx/x)$ for every $c > c_1$ and*

$$\zeta_\lambda(s) = s\int_1^\infty \mathsf{A}_\lambda(x)\, x^{-s-1}\, dx \qquad (\Re s > c_1). \tag{14.19}$$

If ζ_λ has a representation (14.11), then $\mathsf{A}_\lambda(x) = \mathsf{A}([1,x])$ almost everywhere. With the pairing $\langle \mathsf{A}_\lambda, \mathsf{g}\rangle := -\int_1^\infty \mathsf{A}_\lambda(x)\, \mathsf{g}'(x)\, dx$, every even Schwartz function g satisfies

$$\mathsf{g}(0) + 2\langle \mathsf{A}_\lambda, \mathsf{g}\rangle = \hat{\mathsf{g}}(0) + 2\langle \mathsf{A}_\lambda, \hat{\mathsf{g}}\rangle. \tag{14.20}$$

Equivalently, the even tempered distribution $\delta_0 + d\mathsf{A}_\lambda(|u|)$ on $\mathbb{R}$ is its own Fourier transform.

Proof. By Proposition 13.1 the bound (12.5) holds for every ζ_λ with constants independent of λ, so, as in the proof of Proposition 13.16, $|\zeta_\lambda(s)| \le C(|t|+\sigma)^{1/4}$ for $\Re s > c_1$, and the argument of Theorem 12.5 applied to $\zeta_\lambda(s)/s$ gives A_λ, its vanishing on $(0,1)$ and its uniqueness. For a positive measure A as in (14.11), partial integration gives (14.19) with $\mathsf{A}([1,x])$, and uniqueness applies.

Let $c > c_1$. In the variable $u = \log x$, the function $\mathsf{A}_\lambda(e^u)e^{-cu}$ has Fourier transform $\zeta_\lambda(c+it)/(c+it)$, and $-\mathsf{g}'(e^u)e^{(c+1)u}$ has Mellin transform $\int_0^\infty(-\mathsf{g}'(x))x^s dx = s\,\mathsf{g}^\sharp(s)$ on $\Re s = c$; Parseval's formula gives

$$\langle \mathsf{A}_\lambda, \mathsf{g}\rangle = \frac{1}{2\pi i}\int_{(c)} \zeta_\lambda(s)\, \mathsf{g}^\sharp(s)\, ds. \tag{14.21}$$

ζ_λ is bounded by a power of $|t|$ on every vertical strip (finite order, the functional equation and the bound on $\Re s \ge c$), and $\mathsf{g}^\sharp$ decays faster than any power. Move the line to $\Re s = 1-c$. The poles of $\mathsf{g}^\sharp$ at $s = -2, -4, \dots$ are cancelled by the trivial zeros of ζ_λ; the remaining poles are $s = 1$ and $s = 0$, exactly as in (14.14). On the new line (11.3) and (14.13) give $\zeta_\lambda(s)\mathsf{g}^\sharp(s) = \zeta_\lambda(1-s)\hat{\mathsf{g}}^\sharp(1-s)$, and (14.21) for $\hat{\mathsf{g}}$ gives (14.20). □

Proposition 14.10 (The value at infinity as one number)**.** *Let g_ϵ be as in the proof of Theorem 14.7, $\mathsf{w}_\epsilon := \hat{\mathsf{g}}_\epsilon - \mathsf{g}_\epsilon$, which is non-negative on $|u| \ge 1$, and $\mathsf{A}'_\lambda(x) := \mathsf{A}_\lambda(x) - \varpi_\lambda$ for $x \ge 1$. For every $\lambda \in [0,1]$ and $\epsilon > 0$,*

$$\varpi_\lambda\, \mathsf{w}_\epsilon(1) = \tfrac12(1 - \hat{\mathsf{g}}_\epsilon(0)) - \langle \mathsf{A}'_\lambda, \mathsf{w}_\epsilon\rangle, \qquad \langle \mathsf{A}'_\lambda, \mathsf{w}_\epsilon\rangle := -\int_1^\infty \mathsf{A}'_\lambda(x)\, \mathsf{w}'_\epsilon(x)\, dx, \tag{14.22}$$

the limit $\mathsf{W}(\lambda) := \lim_{\epsilon\to0}\langle \mathsf{A}'_\lambda, \mathsf{w}_\epsilon\rangle$ exists, and

$$\varpi_\lambda - 1 = -\frac{\mathsf{W}(\lambda)}{\hat{\mathsf{g}}(1)}, \qquad \hat{\mathsf{g}}(1) = \frac16 + \frac{1}{4\pi^2}. \tag{14.23}$$

Consequently $\mathsf{W}(0) \le 0$, $\mathsf{W}(1) > 0$, and (D) holds if and only if $\mathsf{W}(0) \ge 0$. If A_λ is non-decreasing, then $\langle \mathsf{A}'_\lambda, \mathsf{w}_\epsilon\rangle = \int_{(1,\infty)} \mathsf{w}_\epsilon\, d\mathsf{A}_\lambda \ge 0$, which is Theorem 14.7.

Proof. By (14.20) for g_ϵ, with $\mathsf{g}_\epsilon(0) = 1$, $2\langle \mathsf{A}_\lambda, \mathsf{w}_\epsilon\rangle = 1 - \hat{\mathsf{g}}_\epsilon(0)$. Since $\mathsf{A}_\lambda = \varpi_\lambda + \mathsf{A}'_\lambda$ on $[1,\infty)$ and $-\int_1^\infty \mathsf{w}'_\epsilon = \mathsf{w}_\epsilon(1)$, $\langle \mathsf{A}_\lambda, \mathsf{w}_\epsilon\rangle = \varpi_\lambda \mathsf{w}_\epsilon(1) + \langle \mathsf{A}'_\lambda, \mathsf{w}_\epsilon\rangle$; this is (14.22). As $\epsilon \to 0$, $\mathsf{w}_\epsilon(1) = \hat{\mathsf{g}}_\epsilon(1) \to \hat{\mathsf{g}}(1)$, because $\mathsf{g}_\epsilon(1) = 0$, and $\frac12(1 - \hat{\mathsf{g}}_\epsilon(0)) \to \frac12(1 - \hat{\mathsf{g}}(0)) = \hat{\mathsf{g}}(1)$ by (14.18); hence the limit exists and (14.23) holds. The sign statements follow from $\varpi_{Z1} \ge 1$ (Theorem 12.2) and $\varpi_{\mathrm{RVM}} < 1$. □

Theorem 14.11 (The endpoint dominates ζ on the real axis)**.** *For every real $\sigma > 1$,*

$$\zeta_{Z1}(\sigma) \;\geq\; \zeta(\sigma), \tag{14.24}$$

and equality for one $\sigma > 1$ holds if and only if $\zeta_{Z1} = \zeta$. In particular

$$\prod_{k\geq 1}\Big(1 + \frac{2}{\frac{1}{4}+\gamma_k^2}\Big) \;\geq\; \frac{\pi}{3}, \qquad \textit{with equality if and only if } (\mathrm{D}). \tag{14.25}$$

Proof. By the proof of Theorem 12.2, $\zeta_{Z1}(\sigma)/\zeta(\sigma) = \mathscr{R}_{Z1}(\sigma)$ is the convergent product of the factors (12.11), with $W = (\sigma - \frac{1}{2})^2 + \gamma^2$. At $\sigma = 1$, $W = \frac{1}{4} + \gamma^2$ and the factor equals 1 by the definition of c_γ. The logarithmic derivative in W of $W^2/((W-\delta^2)^2 + 4\delta^2\gamma^2)$ is

$$\frac{2}{W} - \frac{2(W-\delta^2)}{(W-\delta^2)^2 + 4\delta^2\gamma^2} = \frac{2\delta^2\,(4\gamma^2 + \delta^2 - W)}{W\big((W-\delta^2)^2 + 4\delta^2\gamma^2\big)}, \tag{14.26}$$

positive for $W < 4\gamma^2 + \delta^2$ and negative afterwards, and the factor tends to $c_\gamma > 1$ as $W \to \infty$. A function that equals 1 at $\sigma = 1$, increases and then decreases to a limit greater than 1 is greater than 1 for every $\sigma > 1$. Hence $\mathscr{R}_{Z1}(\sigma) \geq 1$, with equality only if the product is empty, that is, $\mathscr{R}_{Z1} \equiv 1$. For $\sigma = 2$: $q(2) = 2$, $G(2) = 2/\pi$ and $P_\zeta(2) = G(2)\zeta(2) = \pi/3$, so (14.24) reads $P_{Z1}(2) \geq \pi/3$. □

Proposition 14.12 (The final step on the open path)**.** *The following are equivalent:* (i) *(D);* (ii) $\varpi_\lambda < 1$ *for every* $\lambda \in (0,1]$*;* (iii) $\varpi_{\lambda_m} \leq 1$ *for some sequence* $\lambda_m \to 0^+$*;* (iv) $\mathsf{W}(\lambda) > 0$ *for every* $\lambda \in (0,1]$.

Proof. $f(\lambda) = \log \varpi_\lambda$ is continuous and strictly convex on $[0,1]$ (Proposition 13.2: $f' = -\Upsilon$, $\Upsilon' < 0$), $f(1) = \log \varpi_{\mathrm{RVM}} < 0$ and $f(0) \geq 0$ (Theorem 12.2). If $f(0) = 0$, convexity gives $f(\lambda) < (1-\lambda)f(0) + \lambda f(1) < 0$ for $0 < \lambda < 1$. If $f(\lambda_m) \leq 0$ along $\lambda_m \to 0^+$, then $f(0) \leq 0$ by continuity, so $f(0) = 0$, which is (D) by Theorem 12.2. (ii)⇔(iv) is (14.23). □

A statement that holds as an identity only at the endpoint cannot be reached as a limit of the same identity along the path, where it is false (Proposition 14.1); an inequality can. Proposition 14.12 is of this form: the inequality $\varpi_\lambda \leq 1$ on the open path passes to the limit, and the opposite inequality $\varpi_{Z1} \geq 1$ holds at the endpoint. By Theorem 14.11 the same scheme is available at every real $\sigma > 1$: (D) holds if and only if $\zeta_{Z1}(\sigma) \leq \zeta(\sigma)$ for one $\sigma > 1$.

Remark 14.13 (The hypothesis added to the functional equation). The functional equation, the order and the zeros on the critical line do not characterize ζ: every ζ_λ has them. A converse theorem for ζ_{Z1} needs one further hypothesis on the function A_{Z1} of Lemma 14.9, and each of the following suffices and is equivalent to (D): A_{Z1} is a step function with jumps at the integers (Hamburger's theorem); A_{Z1} is non-decreasing (Theorem 14.7); $\mathsf{W}(0) \geq 0$ (Proposition 14.10), which is the weakest of the three, being equivalent to $\varpi_{Z1} \leq 1$. The inequality (14.17) is the one-dimensional case of the linear programming bound of Cohn and Elkies [9]. By Lemma 14.9 the measure $\delta_0 + d\mathsf{A}_\lambda(|u|)$ is self-dual for every λ; measures on $\mathbb{R}$ whose support and spectrum are both uniformly discrete are classified in [28].

14.5. **The density of the clock.** Let $\Psi(t) := t\log t - t$ for $t > 0$, and for $M \geq 1$, $0 < t_0 \leq \frac{1}{2}$ let

$$\mathcal{U}(M, t_0) := \int_{t_0}^{\infty} e^{-iM\Psi(t)}\,dt := \lim_{T\to\infty}\int_{t_0}^{T} e^{-iM\Psi(t)}\,dt. \tag{14.27}$$

Lemma 14.14 (Stationary phase for Ψ)**.** *The limit* (14.27) *exists, and there is an absolute constant $C_{\mathcal{U}}$ with*

$$\Big|\mathcal{U}(M,t_0) - \sqrt{\frac{2\pi}{M}}\,e^{i(M-\pi/4)}\Big| \leq \frac{C_{\mathcal{U}}}{M} \qquad (M \geq 1,\ 0 < t_0 \leq \tfrac{1}{2}). \tag{14.28}$$

Proof. $\Psi'(t) = \log t$ is increasing and unbounded, so the limit exists by the first-derivative test. Ψ is convex with minimum $\Psi(1) = -1$. Put $\mathsf{w} := \operatorname{sgn}(t-1)\sqrt{2(\Psi(t)+1)}$. Near $t = 1$, $\Psi(t) + 1 = \frac{1}{2}(t-1)^2(1 + O(t-1))$ with an analytic factor, so $t \mapsto \mathsf{w}$ is analytic with derivative 1 at $t = 1$; elsewhere $\mathsf{w}\, d\mathsf{w} = \log t\, dt$. Hence $t \mapsto \mathsf{w}$ is an increasing analytic bijection of $(0,\infty)$ onto $(-\sqrt{2},\infty)$, with

$$\mathsf{p}(\mathsf{w}) := \frac{dt}{d\mathsf{w}} = \frac{\mathsf{w}}{\log t}, \qquad \mathsf{p}(0) = 1, \qquad \mathsf{q}(\mathsf{w}) := \frac{\mathsf{p}(\mathsf{w}) - 1}{\mathsf{w}} = \frac{1}{\log t} - \frac{1}{\mathsf{w}}. \tag{14.29}$$

q is analytic on $(-\sqrt{2},\infty)$, bounded, tends to 0 as $\mathsf{w} \to \infty$, and $\mathsf{q}' = -\mathsf{w}/(t\log^3 t) + \mathsf{w}^{-2}$. Away from $\mathsf{w} = 0$ one has $\int |\mathsf{w}|\,|t\log^3 t|^{-1} d\mathsf{w} = \int dt/(t\log^2 t)$, which converges at $t = 0$ and at $t = \infty$, and $\int \mathsf{w}^{-2} d\mathsf{w}$ converges at ∞; so $\int_{-\sqrt{2}}^{\infty} |\mathsf{q}'|\, d\mathsf{w} < \infty$. With $\mathsf{w}_0 = \mathsf{w}(t_0) \le \mathsf{w}(\frac{1}{2}) = -\sqrt{1-\log 2} < -\frac{1}{2}$ and $e^{-iM\Psi} = e^{iM} e^{-iM\mathsf{w}^2/2}$,

$$e^{-iM}\,\mathcal{U}(M, t_0) = \int_{\mathsf{w}_0}^{\infty} e^{-iM\mathsf{w}^2/2}\, d\mathsf{w} + \int_{\mathsf{w}_0}^{\infty} e^{-iM\mathsf{w}^2/2}\, \mathsf{w}\,\mathsf{q}(\mathsf{w})\, d\mathsf{w}. \tag{14.30}$$

The first integral is $\int_{\mathbb{R}} e^{-iM\mathsf{w}^2/2} d\mathsf{w} = \sqrt{2\pi/M}\, e^{-i\pi/4}$ minus $\int_{-\infty}^{\mathsf{w}_0}$, and the latter is at most $4/(M|\mathsf{w}_0|) \le 8/M$ by the first-derivative test [37, Lemma 4.2]. Since $\mathsf{w}\, e^{-iM\mathsf{w}^2/2} = \frac{i}{M}\frac{d}{d\mathsf{w}} e^{-iM\mathsf{w}^2/2}$, integration by parts bounds the second integral by $M^{-1}(\sup|\mathsf{q}| + \int|\mathsf{q}'|)$. □

For $m \ge 1$ and $y > 1$ put

$$\mathcal{R}_m(y) := -2(-1)^m\, m^{-3/2}\, y^{\frac{1}{2m} - \frac{1}{2}} \cos\Big(2\pi m\, y^{1/m} - \frac{(7m+1)\pi}{4}\Big), \tag{14.31}$$

so that $\mathcal{R}_1(y) = 2\cos(2\pi y)$ and $\mathcal{R}_2(y) = -2^{-1/2} y^{-1/4} \cos(4\pi\sqrt{y} + \frac{\pi}{4})$.

Theorem 14.15 (The density of the clock)**.** *J_{RVM} is continuously differentiable on $(2,\infty)$, and for $y \ge 16$*

$$J'_{\mathrm{RVM}}(y) = \frac{1}{\log y} + 2\cos(2\pi y) - \frac{y^{-1/4}}{\sqrt{2}} \cos\Big(4\pi\sqrt{y} + \frac{\pi}{4}\Big) + O(y^{-1/3}). \tag{14.32}$$

On $(2,\infty)$ the measure $d\pi_{\mathrm{RVM}}$ is the sum of the atoms $\frac{\mu(j)}{j} J_{\mathrm{RVM}}(2)\, \delta_{2^j}$, $j \ge 1$, and of a density that is continuous off the points 2^j and equals $J'_{\mathrm{RVM}}(y) + O(y^{-1/2})$.

Proof. An exact series. We use the notation of the proof of Theorem 6.1: $v = \log y$, $S_\varepsilon(v) = \sum_k e^{(iv-\varepsilon)\tau_k}$, the decomposition (6.8) and the harmonics $K_m^{\pm}$ of (6.9), with phases $\Phi_m^{\pm}(\tau) = v\tau \pm 2\pi m \mathcal{N}_{\mathrm{RVM}}(\tau)$. Put

$$\mathcal{H}_m^{\pm}(v) := \int_{\tau_0}^{\infty} e^{i\Phi_m^{\pm}(\tau)}\, d\tau, \tag{14.33}$$

improper integrals that converge, locally uniformly in $v \ge \log 2$, by the first-derivative test, because $(\Phi_m^{\pm})'$ is monotone and unbounded. As $\varepsilon \downarrow 0$, $K_m^{\pm} \to iv\,\mathcal{H}_m^{\pm}(v)$, and the first two terms of (6.8) tend to a continuous function $S_{\mathrm{sm}}(v)$ with $|S_{\mathrm{sm}}(v)| \le C/v + \frac{1}{2}$. The bounds for $K_m^{\pm}$ in that proof are uniform in ε and summable against the weights $1/(2\pi m)$, so by dominated convergence in (6.7) and (6.9)

$$\sum_k D_k = \int_{u_1}^{u_2} \frac{e^{v/2}}{v}\, S_0(v)\, dv, \qquad S_0(v) := S_{\mathrm{sm}}(v) + \sum_{m \ge 1} \frac{(-1)^m\, v}{2\pi m} \big(\mathcal{H}_m^{-}(v) - \mathcal{H}_m^{+}(v)\big), \tag{14.34}$$

the series converging uniformly on compact sets. Thus S_0 is continuous, J_{RVM} is continuously differentiable on $(2,\infty)$, and by (5.2)

$$J'_{\mathrm{RVM}}(y) = \frac{1}{\log y}\Big(1 - \frac{1}{y(y^2-1)} - \frac{2}{\sqrt{y}}\, \Re S_0(\log y)\Big). \tag{14.35}$$

The terms without a stationary point. $(\Phi_m^+)' \ge v + m\log 2$, so $|\mathcal{H}_m^+| \le 4/(v+m\log 2)$ and $\sum_m \frac{v}{2\pi m}|\mathcal{H}_m^+| \ll 1+\log v$. For $m > v/\log 2$, $(\Phi_m^-)' \le v - m\log 2 < 0$ on $[\tau_0,\infty)$; the first-derivative test, and the second-derivative test on $[\tau_0, 2\tau_0]$ with $|(\Phi_m^-)''| = m/\tau$, give $|\mathcal{H}_m^-| \ll \min\big((m\log 2 - v)^{-1}, m^{-1/2}\big)$, and $\sum_{m>v/\log 2}\frac{v}{2\pi m}|\mathcal{H}_m^-| \ll 1+\log v$.

The terms $3 \le m \le v/\log 2$. By the argument for (6.10), $|\mathcal{H}_m^-| \le \frac{8}{m\log 2} + 16\sqrt{\pi}\, e^{v/(2m)} m^{-1/2}$, so these terms contribute to S_0 at most $C\,v\sum_{m\ge 3}\big(m^{-2} + m^{-3/2}e^{v/(2m)}\big) \ll v\,e^{v/6}$.

The terms $m = 1, 2$. The stationary point of Φ_m^- is $\mathsf{t}_m := 2\pi e^{v/m} = 2\pi y^{1/m}$. With $\tau = \mathsf{t}_m t$ and (2.3), $\Phi_m^-(\mathsf{t}_m t) = -m\mathsf{t}_m\Psi(t) - \frac{7\pi m}{4}$, hence

$$\mathcal{H}_m^-(v) = \mathsf{t}_m\, e^{-7\pi i m/4}\,\mathcal{U}(m\mathsf{t}_m, \tau_0/\mathsf{t}_m), \qquad \tau_0/\mathsf{t}_m = 2e^{-v/m} \le \tfrac{1}{2} \quad (y \ge 4^m). \tag{14.36}$$

By Lemma 14.14 with $M = m\mathsf{t}_m$,

$$\frac{(-1)^m v}{2\pi m}\,\mathcal{H}_m^-(v) = (-1)^m\,\frac{v}{m^{3/2}}\,y^{\frac{1}{2m}}\exp\Big(i\Big(2\pi m\, y^{1/m} - \frac{(7m+1)\pi}{4}\Big)\Big) + O\Big(\frac{v}{m^2}\Big). \tag{14.37}$$

Inserting the four estimates into (14.35) gives $J'_{\mathrm{RVM}}(y) = \frac{1}{\log y} + \mathcal{R}_1(y) + \mathcal{R}_2(y) + O(y^{-1/3})$, which is (14.32).

The prime coordinate. By (5.7), on each interval $(2^r, 2^{r+1})$ the function π_{RVM} is the finite sum of the continuously differentiable functions $\frac{\mu(j)}{j}J_{\mathrm{RVM}}(x^{1/j})$, $j \le r$; the terms $j \ge 2$ have derivative $\frac{\mu(j)}{j^2}x^{1/j-1}J'_{\mathrm{RVM}}(x^{1/j})$, and $|J'_{\mathrm{RVM}}| \le C_J$ by (6.5), so their sum is $O(x^{-1/2})$. At $x = 2^j$ the term j jumps by $\frac{\mu(j)}{j}J_{\mathrm{RVM}}(2)$. □

Corollary 14.16 (Cell masses and cell defects at the clock end)**.** *For $n \ge 17$, with $\mathbf{1}_{2^j}(n) = 1$ if $n = 2^j$ and 0 otherwise,*

$$c_n^{\mathrm{RVM}} = \int_{n-1}^{n}\Big(\frac{1}{\log x} + 2\cos(2\pi x)\Big)dx + \mathbf{1}_{2^j}(n)\,\frac{\mu(j)}{j}\,J_{\mathrm{RVM}}(2) + O\big(n^{-1/4}\big), \tag{14.38}$$

$$\tilde{c}_n(1) = \frac{1}{\log n} + \mathbf{1}_{2^j}(n)\,\frac{\mu(j)}{j}\,J_{\mathrm{RVM}}(2) + O\big(n^{-1/4}\big), \tag{14.39}$$

and for $0 < \Re s \le \sigma_1$

$$|\mathfrak{h}_n(1;s)| \le C_{\sigma_1}\big(|s|\,n^{-\sigma-4/3} + |s|(1+|s|)\,n^{-\sigma-2}\big). \tag{14.40}$$

In particular Hypothesis 6.7 holds along the integers n that are not of the form 2^j with $\mu(j) \ne 0$, and $c_n^{\mathrm{RVM}} \sim 1/\log n$, $0 < c_n^{\mathrm{RVM}} < 1$ for all large such n; at $n = 2^j$, $c_n^{\mathrm{RVM}} = \big(1 + \mu(j)J_{\mathrm{RVM}}(2)\log 2\big)/\log n + O(n^{-1/4})$.

Proof. (14.38): $c_n^{\mathrm{RVM}} = \pi_{\mathrm{RVM}}(n) - \pi_{\mathrm{RVM}}(n-1)$ is the integral of the density over $(n-1, n)$ plus the atom at n, and $\int_{n-1}^{n}(|\mathcal{R}_2| + O(x^{-1/3}))dx = O(n^{-1/4})$. (14.39): $\varphi_n(2^j) = \mathbf{1}_{2^j}(n)$; $\int \varphi(t)\,dt/\log(n+t) = \frac{1}{\log n} + O(\frac{1}{n\log^2 n})$ because $\mathsf{M}_0 = 1$; and $\int \varphi(t)\,2\cos(2\pi(n+t))\,dt = 2\mathsf{C}_0 = 0$ by (14.2). (14.40): the test function $\varphi_n(x)(n^{-s} - x^{-s})$ vanishes at the atoms, and $n^{-s} - (n+t)^{-s} = s\,n^{-s-1}t + O_{\sigma_1}(|s|(1+|s|)n^{-\sigma-2})$ for $|t| \le 1$. The first moment of the density is $\int t\varphi(t)\pi'_{\mathrm{RVM}}(n+t)\,dt = \frac{\mathsf{M}_1}{\log n} + 2\mathsf{C}_1 + O(n^{-1/3}) = O(n^{-1/3})$, because $\mathsf{M}_1 = \mathsf{C}_1 = 0$, $\mathcal{R}_2(n+t) = \mathcal{R}_2(n) + O(n^{-3/4})$ and the density is bounded. The last statements follow from (14.38) and Lemma 6.6. □

By (14.32), $\frac{d\pi_{\mathrm{RVM}}}{dy} = \big(\frac{1}{\log y} - 1\big) + \big(1 + 2\cos(2\pi y)\big) + O(y^{-1/4})$: the clock carries the first harmonic of the comb $\sum_n \delta_n = 1 + 2\sum_{q\ge 1}\cos(2\pi q y)$ of all integers, with weight 1, on a smooth background. This harmonic comes from the ordinates τ_k near the height $2\pi y$, the stationary point of Φ_1^-; by (14.2) it does not enter the cell masses (14.39), which are $1/\log n$ to leading order, and it enters the cell defects only through the moments C_r, $r \ge 2$.

14.6. **The cell masses in the heights of the ordinates.**

Proposition 14.17 (Height kernel of the cell masses)**.** *For every $\varepsilon > 0$, $\lambda \in [0,1]$ and $n \geq 3$ that is not a power of 2,*

$$\tilde{c}_n(\lambda) = \mathfrak{m}_n - \frac{2}{\sqrt{n}\,\log n} \sum_{k \geq 1} \hat{\varphi}\Big(\frac{\gamma_k(\lambda)}{2\pi n}\Big) \cos\big(\gamma_k(\lambda)\log n\big) + E_n(\lambda), \tag{14.41}$$

where $\sum_{X<n\leq 2X} |E_n(\lambda)|^2 \leq C_\varepsilon X^{1/2+\varepsilon}$ uniformly in λ (sum over the n that are not powers of 2). The kernel $\hat{\varphi}(\gamma/2\pi n)$ vanishes at the heights $\gamma = 2\pi q n$, $q = 1, 2, \ldots$

Proof. The term $j = 1$ of W_n in (13.19) is $\int \varphi(t)(n+t)^{-1/2+i\gamma}\, dt/\log(n+t)$. For $0 < \gamma \leq n^{1+\varepsilon}$, $(1+t/n)^{-1/2+i\gamma} = e^{i\gamma t/n}\big(1 + O((1+\gamma/n)/n)\big)$ and $\frac{1}{\log(n+t)} = \frac{1}{\log n}(1+O(\frac{1}{n}))$ on $|t| \leq 1$, so this term equals $\frac{n^{-1/2+i\gamma}}{\log n}\hat{\varphi}(\frac{\gamma}{2\pi n}) + O\big(n^{-3/2}(1+\gamma/n)\big)$, because $\int \varphi(t) e^{i\gamma t/n} dt = \hat{\varphi}(\gamma/2\pi n)$ for even φ. Summed over the $O(n^{1+\varepsilon}\log n)$ ordinates $\gamma_k(\lambda) \leq n^{1+\varepsilon}$ the error is $O(n^{-1/2+3\varepsilon})$. For $\gamma > n^{1+\varepsilon}$ both the term and $\hat{\varphi}(\gamma/2\pi n)$ are $O_A((n/\gamma)^A)$, as in the proof of Theorem 13.10. The terms $j \geq 2$ have sum of squares $O(X^{1/2+3\varepsilon})$ over $X < n \leq 2X$ by (13.26). The zeros of the kernel are (14.2). □

Since $\gamma_k(\lambda)\log n = \tau_k \log n - (1-\lambda)\eta_k \log n$, the passage along the family rotates each clock phase $n^{i\tau_k}$ by the angle $-(1-\lambda)\eta_k \log n$: up to the change of $\hat{\varphi}$ over a distance $O(1/n)$, the sum in (14.41) is $\Re\, \Xi_n\big((1-\lambda)\log n\big)$ with

$$\Xi_n(\Omega) := \sum_{k \geq 1} \hat{\varphi}\Big(\frac{\tau_k}{2\pi n}\Big) n^{i\tau_k} e^{-i\Omega\eta_k} \qquad (0 \leq \Omega \leq \log n). \tag{14.42}$$

At $\Omega = 0$ this is the clock sum, whose real part is $O(n^{1/4}\log n)$ by (14.39) and Theorem 14.15, because the resonant heights $\tau_k \approx 2\pi n$ are zeros of the kernel; the endpoint of the family is $\Omega = \log n$.

Proposition 14.18 (The flow of a cell mass and the count difference)**.** *For every $n \geq 3$ that is not a power of 2,*

$$\tilde{c}_n(0) - \tilde{c}_n(1) = -2\int_{0^+}^{1} \sum_{k \geq 1} \eta_k\, \Im V_n\big(\gamma_k(\lambda)\big)\, d\lambda = -2\,\Im \int_{\tau_0}^{\infty} V_n(t)\big(N_\zeta(t) - n_{\mathrm{RVM}}(t)\big)\, dt. \tag{14.43}$$

Proof. The first equality is (13.20) and the continuity of $\tilde{c}_n$ at $\lambda = 0$. By (13.20), $\tilde{c}_n(0) - \tilde{c}_n(1) = 2\Re \sum_k \big(W_n(\tau_k) - W_n(\gamma_k)\big) = 2\Re \int_{\tau_0}^{\infty} W_n\, d(n_{\mathrm{RVM}} - N_\zeta)$. Both counts vanish at τ_0, their difference is $O(\log t)$ by Lemma 2.2, and W_n is rapidly decreasing; integration by parts with $W_n' = iV_n$ gives $2\Re\big(i\int V_n(N_\zeta - n_{\mathrm{RVM}})\, dt\big)$. □

Proposition 14.19 (Height kernel of the cell defects)**.** *For every $\varepsilon > 0$, $\lambda \in [0,1]$, $\Re s > 0$ and $n \geq 3$ that is not a power of 2,*

$$\mathfrak{h}_n(\lambda; s) = -\frac{s\, n^{-s-3/2}}{\pi \log n} \sum_{k \geq 1} \hat{\varphi}'\Big(\frac{\gamma_k(\lambda)}{2\pi n}\Big) \sin\big(\gamma_k(\lambda)\log n\big) + O_{s,\varepsilon}(n^{-\sigma-3/4+\varepsilon}). \tag{14.44}$$

Proof. For $|t| \leq 1$, $n^{-s} - (n+t)^{-s} = s\, n^{-s-1} t + \psi_2(t)$, where ψ_2 and all its derivatives in t are $O_s(n^{-\sigma-2})$. *The term $j = 1$.* The part of the density that does not involve the ordinates contributes $O_s(n^{-\sigma-2})$, because $\int t\varphi(t)\, dt/\log(n+t) = O(1/(n\log^2 n))$ ($\mathsf{M}_1 = 0$). The pairing of $\varphi\psi_2$ with the ordinate sum is $O_s(n^{-\sigma-2})$ times $O(n^{1/2+\varepsilon})$, as in Remark 13.14. For the real function $t\varphi(t)$, the computation in the proof of Proposition 14.17, with $\int t\varphi(t) e^{2\pi i\xi t} dt = -\frac{i}{2\pi}\hat{\varphi}'(\xi)$ for even φ and $-2\Re(-\frac{i}{2\pi} n^{i\gamma}) = -\frac{1}{\pi}\sin(\gamma \log n)$, gives the sum in (14.44) with an error $O_s(n^{-\sigma-3/2+3\varepsilon})$. *The terms $j \geq 2$.* The test function $\varphi_n(y^j)(n^{-s} - y^{-js})$ is $O(|s| n^{-\sigma-1})$ on a support of length $O(n^{1/j-1})$; with $O(n^{1+\varepsilon}\log n)$ relevant ordinates and $y^{-1/2} = O(n^{-1/(2j)})$, these terms are $O_s(n^{-\sigma-1+1/(2j)+2\varepsilon})$, at most $O_s(n^{-\sigma-3/4+2\varepsilon})$. □

The kernel $\hat{\varphi}$ of the masses vanishes at the heights $2\pi qn$ and the kernel $\hat{\varphi}'$ of the defects in general does not: the masses and the defects of a cell read complementary heights of the ordinates. At $\lambda = 1$ the resonant heights $\tau_k \approx 2\pi n$ contribute to $\sum_k n^{i\tau_k}$ the real quantity $-\sqrt{n}\log n\, e^{2\pi i n}$ (proof of Theorem 14.15), which the sine in (14.44) does not see: the first harmonic of the clock is centred at the integers.

Remark 14.20 (The scale of the displacements). By Proposition 14.1, a property of the sequence (η_k) that yields (D) is false for every sequence $(c\,\eta_k)$ with $0 \le c < 1$. Properties that are invariant under $\eta_k \mapsto c\,\eta_k$ with $c > 0$ (the signs of the η_k, the non-vanishing of the η_k, their boundedness) hold along the whole open path and do not suffice; the property fixes the size of the displacements. In the form (14.42) it is a condition on the angles $\eta_k \log n$ modulo 2π, jointly in k and n. The construction contains one exact relation of this kind: for a simple ordinate, (2.4) and (2.13) give $\mathcal{N}_{\mathrm{RVM}}(\tau_k) - \mathcal{N}_{\mathrm{RVM}}(\gamma_k) = N^*_\zeta(\gamma_k) - \mathcal{N}_{\mathrm{RVM}}(\gamma_k)$, that is,

$$\eta_k\, \mathcal{N}'_{\mathrm{RVM}}(\xi_k) = N^*_\zeta(\gamma_k) - \mathcal{N}_{\mathrm{RVM}}(\gamma_k) \qquad \text{for some } \xi_k \text{ between } \gamma_k \text{ and } \tau_k. \tag{14.45}$$

14.7. **The Gaussian defect of the atomization.**

Lemma 14.21 (Vertical growth of the joining factor). *For every $\lambda \in [0,1]$ and $\frac12 < \sigma_0 < \sigma_1$, $\tilde{h}_\lambda(\sigma + it) = o(|t|)$ as $|t| \to \infty$, uniformly for $\sigma_0 \le \sigma \le \sigma_1$.*

Proof. By (13.13) and Proposition 13.5(ii) it suffices to treat the double series. Fix $c \in (\frac12, \sigma_0)$. By (13.28), $\sum_{m\ge1} \frac1m \sum_{n>N} |\mathfrak{h}_n(\lambda; ms)| \le C|s| \sum_{n>N} n^{-\sigma_0-1}(1+I_n^{1/2})$ and $\sum_{m>N} \frac1m \sum_{n\ge2} |\mathfrak{h}_n(\lambda; ms)| \le C|s| \sum_{n\ge2} n^{-N\sigma_0-1}(1 + I_n^{1/2})$; both are at most $\epsilon_N|s|$ with $\epsilon_N \to 0$ as $N \to \infty$. For the finitely many pairs $m, n \le N$, with $E = \pi_\lambda - \mathfrak{L}$ as in the proof of Theorem 13.12,

$$\begin{aligned}\mathfrak{h}_n(\lambda; ms) = {} & \int \varphi_n(x)(n^{-ms} - x^{-ms})\, d\mathfrak{L}(x) - \int E(x)\varphi_n'(x)(n^{-ms} - x^{-ms})\, dx \\ & - ms \int E(x)\varphi_n(x)\, x^{-ms-1}\, dx.\end{aligned} \tag{14.46}$$

The first two terms are bounded in t, and the last integral tends to 0 as $|t| \to \infty$ by the Riemann–Lebesgue lemma, because $E \in L^1(n-1, n+1)$. Hence $\limsup_{|t|\to\infty} |t|^{-1} \sup_\sigma |\tilde{h}_\lambda(\sigma + it)| \le 2\epsilon_N$ for every N. □

For $y > 0$ and $c > 1$ put

$$\Theta_\lambda(y) := \frac{1}{4\pi i} \int_{(c)} \zeta_\lambda(s)\, \Gamma(\tfrac{s}{2})\, \pi^{-s/2}\, y^{-s/2}\, ds, \qquad \mathfrak{G}_\lambda(y) := \sum_{N\ge1} \tilde{a}_N(\lambda)\, e^{-\pi N^2 y} - \Theta_\lambda(y). \tag{14.47}$$

For ζ itself $\Theta(y) = \sum_{n\ge1} e^{-\pi n^2 y}$, and if ζ_λ has a representation (14.11), then $\Theta_\lambda(y) = \int e^{-\pi u^2 y}\, d\mathsf{A}(u)$. $\mathfrak{G}_\lambda$ is the change produced by the atomization in the Gaussian count of the integers of the system.

Proposition 14.22 (Gaussian defect). *Let $\lambda \in [0,1]$.*

(i) *$\Theta_\lambda(y) = y^{-1/2}\,\Theta_\lambda(1/y) + \frac12(y^{-1/2} - 1)$ for $y > 0$, and $\Theta_\lambda(y) = O_A(y^{-A})$ as $y \to \infty$ for every A.*

(ii) *The coefficients $\tilde{a}_N(\lambda)$ satisfy the theta relation of ζ,*

$$\sum_{N\ge1} \tilde{a}_N(\lambda)\, e^{-\pi N^2 y} = y^{-1/2} \sum_{N\ge1} \tilde{a}_N(\lambda)\, e^{-\pi N^2/y} + \tfrac12(y^{-1/2} - 1) \qquad (y > 0), \tag{14.48}$$

if and only if

$$\mathfrak{G}_\lambda(y) = y^{-1/2}\, \mathfrak{G}_\lambda(1/y) \qquad (y > 0). \tag{14.49}$$

Under the hypotheses of Theorem 14.3, (14.49) *holds.*

(iii) *For every $\delta > 0$, as $y \to 0^+$,*

$$\mathfrak{G}_\lambda(y) = \tfrac{1}{2}(e^{\tilde{h}_\lambda(1)} - 1)\, y^{-1/2} + O_\delta(y^{-1/4-\delta}), \tag{14.50}$$

and $e^{\tilde{h}_\lambda(1)} = \operatorname{Res}_{s=1} \tilde{\zeta}_{\mathbb{Z},\lambda}(s)$.

(iv) *If (14.49) holds, then $\tilde{h}_\lambda(1) = 0$.*

Proof. (i) ζ_λ is bounded by a power of $|t|$ on vertical strips (proof of Lemma 14.6; on $\Re s \geq c$ use Proposition 13.1), and $\Gamma(\frac{s}{2})$ decays exponentially. Moving the line to $\Re s = 1 - c$ crosses $s = 1$, with residue $\Gamma(\frac{1}{2})\pi^{-1/2}y^{-1/2} = y^{-1/2}$, and $s = 0$, with residue $2\zeta_\lambda(0) = -1$; on the new line (11.3) and $y^{-s/2} = y^{-1/2}(1/y)^{-(1-s)/2}$ give $y^{-1/2}\Theta_\lambda(1/y)$. The decay follows by moving the line to $\Re s = 2A$. (ii) By $e^{-x} = \frac{1}{2\pi i}\int_{(c/2)} \Gamma(w)x^{-w}dw$ and Corollary 13.11, $\sum_N \tilde{a}_N(\lambda)e^{-\pi N^2 y} = \frac{1}{4\pi i}\int_{(c)} \tilde{\zeta}_{\mathbb{Z},\lambda}(s)\Gamma(\frac{s}{2})\pi^{-s/2}y^{-s/2}ds$; subtracting (i) from the theta relation gives the equivalence. Under the hypotheses of Theorem 14.3, $\tilde{\zeta}_{\mathbb{Z},\lambda} = \zeta$. (iii) By (ii), $2\mathfrak{G}_\lambda(y) = \frac{1}{2\pi i}\int_{(c)} \zeta_\lambda(s)(e^{\tilde{h}_\lambda(s)} - 1)\Gamma(\frac{s}{2})\pi^{-s/2}y^{-s/2}ds$. By Lemma 14.21, $|e^{\tilde{h}_\lambda(s)}| \leq e^{o(|t|)}$ on $\frac{1}{2} + 2\delta \leq \Re s \leq c$, so the line can be moved to $\Re s = \frac{1}{2} + 2\delta$. The only pole crossed is $s = 1$, where ζ_λ has residue 1 and $\tilde{h}_\lambda$ is holomorphic; the residue is $(e^{\tilde{h}_\lambda(1)} - 1)y^{-1/2}$, and the remaining integral is $O_\delta(y^{-1/4-\delta})$. (iv) $\sum_N \tilde{a}_N(\lambda)e^{-\pi N^2 y} = O(e^{-\pi y})$ as $y \to \infty$, so by (i) $\mathfrak{G}_\lambda(y) = O_A(y^{-A})$ there; then (14.49) gives $\mathfrak{G}_\lambda(y) = O_A(y^A)$ as $y \to 0^+$ for every A, and (14.50) forces $e^{\tilde{h}_\lambda(1)} = 1$; $\tilde{h}_\lambda$ is real on the real axis. □

Corollary 14.23 (The density of the atomized integers)**.** *$\sum_{N \leq x} \tilde{a}_N(\lambda)$ has mean density $e^{\tilde{h}_\lambda(1)}$ in the sense of (14.50), and (D) implies*

$$\tilde{h}_0(1) = 0, \qquad \textit{that is,} \qquad \mathfrak{u}_0(1) = \sum_{m \geq 1} \frac{1}{m} \sum_{n \geq 2} \mathfrak{h}_n(0; m). \tag{14.51}$$

Proof. Under (D), $\tilde{h}_0 \equiv 0$ (Theorem 13.13); the second form is (13.13) at $s = 1$. □

The identity $\tilde{h}_0(\sigma) = 0$ for real σ has two distinguished members: $\sigma = +\infty$ is $\varpi_{Z1} = 1$, the first coefficient (Corollary 12.3), and $\sigma = 1$ is (14.51), the density of the integers.

By Proposition 14.22(i), the function (14.6) is $T_\lambda(y) = \mathfrak{G}_\lambda(y) - y^{-1/2}\mathfrak{G}_\lambda(1/y)$, so (14.49) is $T_\lambda \equiv 0$. At the clock end the expansion (14.50) holds with a smaller remainder: for every $\delta > 0$,

$$\mathfrak{G}_1(y) = \tfrac{1}{2}(e^{\tilde{h}_1(1)} - 1)\, y^{-1/2} + O_\delta(y^{-\delta}) \qquad (y \to 0^+), \tag{14.52}$$

because $\tilde{h}_1$ is holomorphic in $\Re s > 0$ with $\tilde{h}_1(\sigma + it) = o(|t|)$ uniformly on $\sigma_0 \leq \sigma \leq \sigma_1$ for every $\sigma_0 > 0$ (the proof of Lemma 14.21 with the bound $|\mathfrak{h}_n(1; s)| \leq C|s|n^{-\sigma-1}$ of Theorem 13.12(iv)), and the line of integration can be moved to $\Re s = 2\delta$.

14.8. **Entire dependence on λ and the exceptional set.**

Proposition 14.24 (The cell masses as entire functions of λ)**.** *Let $n \geq 3$ not be a power of 2.*

(i) *$\lambda \mapsto \tilde{c}_n(\lambda) = \mathfrak{m}_n - 2\Re\sum_k W_n(\gamma_k + \lambda\eta_k)$ extends to an entire function of λ of exponential type at most $C_\eta \log(n+1)$, real on the real axis.*

(ii) *For every $r \geq 0$, $\varepsilon > 0$ and $\lambda \in [0, 1]$,*

$$\partial_\lambda^r \tilde{c}_n(\lambda) = -2\,\Re \sum_{k \geq 1} \eta_k^r\, W_n^{(r)}(\gamma_k(\lambda)), \qquad |\partial_\lambda^r \tilde{c}_n(\lambda)| \leq C_{r,\varepsilon}(C_\eta \log(n+1))^r\, n^{1/2+\varepsilon}. \tag{14.53}$$

(iii) *$|\tilde{c}_n(\lambda) - \tilde{c}_n(\lambda')| \leq C_\varepsilon|\lambda - \lambda'|\, n^{1/2+\varepsilon}$ on $[0, 1]$. Hence, for sets of integers n that are not powers of 2, $\tilde{c}_n(0) = O_\varepsilon(n^\varepsilon)$ if and only if $\tilde{c}_n(\lambda) = O_\varepsilon(n^\varepsilon)$ uniformly for $0 \leq \lambda \leq n^{-1/2}$, and $\tilde{c}_n(\lambda) = O_\varepsilon(n^\varepsilon)$ uniformly for $1 - n^{-1/2} \leq \lambda \leq 1$.*

Proof. For complex γ, $|y^{i\gamma}| \le e^{|\Im\gamma|\log y}$ with $\log y \le \log(n+1)$ on the support of $\varphi_n(y^j)$, and repeated integration by parts in (13.19) gives $|W_n^{(r)}(\gamma)| \le C_{A,r,n}\, e^{|\Im\gamma|\log(n+1)}(1+|\Re\gamma|)^{-A}$ for every A. With $\gamma = \gamma_k + \lambda\eta_k$, $|\Im\gamma| \le C_\eta|\Im\lambda|$, the series converges locally uniformly in $\lambda \in \mathbb{C}$: this is (i) and the first formula in (14.53). For real λ, $W_n^{(r)}$ carries the factor $(i\log y)^r$ with $\log y \le \log(n+1)$, the ordinates $\gamma_k(\lambda) > n^{1+\varepsilon}$ contribute $O(n^{-1})$, and each of the $O(n^{1+\varepsilon}\log n)$ remaining ordinates contributes $O\big((\log(n+1))^r n^{-1+1/(2j)}\big)$ to the term j, as in Remark 13.14; this is the bound in (14.53). (iii) is the case $r=1$ and Corollary 14.16. □

The bounds (14.53) do not by themselves control $\tilde c_n(0)$ from $\tilde c_n(1)$: the function $n^{1/2}\sin\big((1-\lambda)C_\eta\log n\big)$ is entire of the same type, satisfies the same bounds, vanishes at $\lambda=1$ and has size $n^{1/2}$ at $\lambda=0$ for infinitely many n. The Taylor expansion at the clock end, $\tilde c_n(0) = \sum_{r\ge0}\frac{(-1)^r}{r!}\partial_\lambda^r\tilde c_n(1)$, has the terms $-2\Re\sum_k \eta_k^r W_n^{(r)}(\tau_k)$, clock sums weighted by the powers of the displacements.

Proposition 14.25 (The exceptional set)**.** *The set*

$$\mathsf{E} := \big\{\lambda\in[0,1]:\ \tilde\zeta_{\mathbb{Z},\lambda} = \zeta\big\} \tag{14.54}$$

is finite, $1\notin\mathsf{E}$*, and (D) holds if and only if* $0\in\mathsf{E}$*. For every* $\lambda\notin\mathsf{E}$ *the hypotheses of Theorem 14.3 do not hold together,* $T_\lambda \not\equiv 0$ *and* $\mathsf{V}_\sigma(\lambda) > 0$*.*

Proof. If $\lambda\in\mathsf{E}$, then $\sum_n \tilde c_n(\lambda)\sum_m \frac{n^{-ms}}{m} = \log\zeta(s) = \sum_p\sum_m\frac{p^{-ms}}{m}$ for $\Re s>1$. Comparing the coefficients of N^{-s} in increasing order of N determines the exponents uniquely, so $\tilde c_n(\lambda) = \mathbf{1}_{\mathrm{prime}}(n)$ for every n. Choose a composite n, not a power of 2, so large that $\tilde c_n(1)>0$ by (14.39). Then $\tilde c_n$ is an entire function of λ (Proposition 14.24) that does not vanish identically, and it has finitely many zeros in $[0,1]$; they contain E, and 1 is not one of them. $0\in\mathsf{E}$ implies (D) by Theorem 14.4(ii), and (D) implies $\tilde\zeta_{\mathbb{Z},0} = \zeta_{Z1} = \zeta$ by Theorem 13.13. The last statement is Theorems 14.3 and 14.4(ii). □

Proposition 14.1 excludes that ζ_λ and $\tilde\zeta_{\mathbb{Z},\lambda}$ coincide for $\lambda>0$; Proposition 14.25 excludes, outside a finite set, that $\tilde\zeta_{\mathbb{Z},\lambda}$ coincides with ζ. For $0<\lambda<1$ outside E it is not decided which of the two hypotheses of Theorem 14.3 fails; at $\lambda=1$ it is the symmetry (Proposition 14.5).

14.9. **The final step in the cell masses.**

Corollary 14.26 (The final step is one number per cell)**.** *The following are equivalent:*

(i) *(D);*
(ii) $\tilde c_n(0) = \mathbf{1}_{\mathrm{prime}}(n)$ *for every* $n\ge2$*;*
(iii) *there is a list* $\boldsymbol\kappa\in\mathscr{O}$ *(Section 15) with* $\tilde c_n(\boldsymbol\kappa) = \mathbf{1}_{\mathrm{prime}}(n)$ *for every* $n\ge2$*.*

Proof. (i)⇒(ii) is Theorem 13.13, and (ii)⇒(iii) with the list (γ_k). If (iii) holds, then

$$\tilde\zeta_{\mathbb{Z},\boldsymbol\kappa}(s) = \prod_{n\ge2}\big(1-n^{-s}\big)^{-\tilde c_n(\boldsymbol\kappa)} = \prod_p\big(1-p^{-s}\big)^{-1} = \zeta(s)\qquad(\Re s>1), \tag{14.55}$$

and Corollary 15.3, (iii)⇒(i), gives (D). □

Step A7 of the programme asks for the vanishing of the functions $\mathfrak{h}_n(0;\cdot)$, one function of s for each cell, and for $J_{Z1}=0$ on the first cell. Corollary 14.26 asks for one real number for each cell; the position of the mass inside the cell and the first cell then follow from Hamburger's theorem.

Proposition 14.27 (Every point of the exceptional set is a witness)**.** *For the set* E *of* (14.54)*,*

$$\mathsf{E}\neq\varnothing \iff \text{(D)},\qquad \textit{so that either } \mathsf{E}=\varnothing \ \textit{ or } \ 0\in\mathsf{E}. \tag{14.56}$$

If (D) holds, then

$$\mathsf{E}\setminus\{0\} = \big\{\lambda\in(0,1]:\ \mathsf{D}_n(\lambda)=0 \textit{ for every } n\ge2\big\},\qquad \mathsf{D}_n(\lambda) := \tilde c_n(\lambda)-\tilde c_n(0), \tag{14.57}$$

and for $n \geq 3$ not a power of 2, $\mathsf{D}_n(\lambda) = -2\Re \sum_{k\geq 1}\big(W_n(\gamma_k + \lambda\eta_k) - W_n(\gamma_k)\big)$ is an entire function of λ of exponential type at most $C_\eta \log(n+1)$ that vanishes at $\lambda = 0$.

Proof. If $\lambda_0 \in \mathsf{E}$, the list $(\gamma_k(\lambda_0))$ belongs to $\mathscr{O}$ and its atomized zeta function is ζ; Corollary 15.3, (iii)⇒(i), gives (D), and then $0 \in \mathsf{E}$ by Proposition 14.25. Under (D), $\tilde{c}_n(0) = \mathbf{1}_{\mathrm{prime}}(n)$, and $\lambda \in \mathsf{E}$ means $\tilde{c}_n(\lambda) = \mathbf{1}_{\mathrm{prime}}(n)$ for every n (proof of Proposition 14.25). The last statement is (13.20) and Proposition 14.24. □

Thus a proof that $\mathsf{E} \cap (0,1] = \varnothing$ does not approach the final step: if (D) fails, E is empty, and if (D) holds, $\mathsf{E} \cap (0,1] = \varnothing$ is the uniqueness of the witness on the path, which is the statement that the entire functions D_n have no common zero in $(0,1]$. This uniqueness is not proved; Proposition 14.25 gives that E is finite and $1 \notin \mathsf{E}$.

Remark 14.28 (The dual function of the atomized family). For $\Re s < \frac{1}{2}$ put $B_\lambda^\vee(s) := \zeta_\lambda(s)\, e^{\tilde{h}_\lambda(1-s)}$. By the functional equation of ζ_λ,

$$\pi^{-s/2}\Gamma\big(\tfrac{s}{2}\big)B_\lambda^\vee(s) = \pi^{-(1-s)/2}\Gamma\big(\tfrac{1-s}{2}\big)\,\tilde{\zeta}_{\mathbb{Z},\lambda}(1-s) \qquad (\Re s < \tfrac{1}{2}), \tag{14.58}$$

so $B_\lambda^\vee$ is the function attached to $\tilde{\zeta}_{\mathbb{Z},\lambda}$ by the functional equation. To ask that $B_\lambda^\vee$ continue to $\Re s > 1$ as an absolutely convergent Dirichlet series is to ask for the hypotheses of Theorem 14.3; no monotonicity in λ is known, and a parameter such as $\inf\{\lambda > 0 : B_\lambda^\vee$ is not a Dirichlet series$\}$ is not shown to be a transition point.

Proposition 14.29 (The identity that remains, cell by cell)**.** *Let $n \geq 17$ not be a power of 2. Then $\tilde{c}_n(0) = \mathbf{1}_{\mathrm{prime}}(n)$ if and only if*

$$\mathbf{1}_{\mathrm{prime}}(n) - \tilde{c}_n(1) = -2\,\Im \int_{\tau_0}^{\infty} V_n(t)\big(N_\zeta(t) - n_{\mathrm{RVM}}(t)\big)\,dt, \qquad \tilde{c}_n(1) = \frac{1}{\log n} + O(n^{-1/4}). \tag{14.59}$$

Proof. This is (14.43) and (14.39). □

The left side of (14.59) is explicit. The right side is the pairing of the count difference with the kernel V_n, whose term $j = 1$ is $n^{-1/2+it}\hat{\varphi}(t/2\pi n)$ up to $O(n^{-3/2}(1+t/n))$ (proof of Proposition 14.19). For $n = 2^j$ the same identity holds with the cut term of Theorem 13.10 added. No evaluation of the right side of (14.59) is given in this paper.

14.10. **Symmetric joining factors.**

Remark 14.30 (The symmetry criterion has no slack at $\lambda = 0$). At $\lambda = 0$ the hypothesis of Theorem 14.3 is equivalent to its conclusion. If (D) holds, then $\tilde{h}_0 \equiv 0$ (Theorem 13.13), and $e^{\tilde{h}_0} = 1$ is symmetric and of finite order; hence

$$e^{\tilde{h}_0} \text{ symmetric of finite order } \iff \tilde{h}_0 \equiv 0 \iff \text{(D)}, \tag{14.60}$$

and in the same way $T_0 \equiv 0$ and $\mathsf{V}_\sigma(0) = 0$ are equivalent forms of (D), not weaker conditions. Moreover (14.5) presupposes a continuation of $\tilde{h}_0$ across the line $\Re s = \frac{1}{2}$ (step A10 of the programme): $\tilde{h}_0$ is holomorphic in $\Re s > \frac{1}{2}$, and $s \mapsto 1 - s$ maps this half-plane onto $\Re s < \frac{1}{2}$. Theorems 14.34 and 14.36 below are criteria whose hypotheses are written on the half-line $\sigma \to +\infty$ and on the closed half-plane $\Re s \geq \frac{1}{2}$.

Proposition 14.31 (Symmetric joining factors in general)**.** *Let f be meromorphic in $\mathbb{C}$ with $(s-1)f(s)$ entire of order at most one, and let f satisfy the functional equation* (11.3)*. Let $g = f\,e^h$, where e^h is entire, without zeros, of finite order, and $e^{h(s)} = e^{h(1-s)}$. If $g(s) = \sum_{n\geq 1} a_n n^{-s}$ converges absolutely in $\Re s > 1$, then $g = C_H\zeta$ and e^h is constant. If moreover $\lim_{\sigma\to+\infty} g(\sigma) = \lim_{\sigma\to+\infty} f(\sigma) \neq 0$, then $e^h \equiv 1$ and $f = g = C_H\zeta$.*

Proof. g satisfies (H0)–(H3) of Subsection 13.7 with the second function equal to g, so $g = C_H\zeta$ by Hamburger's theorem. If $C_H = 0$ then $f\,e^h = 0$, which is excluded when $g \not\equiv 0$; assume $C_H \neq 0$. Then $e^h = C_H(s-1)\zeta(s)/\big((s-1)f(s)\big)$ is an entire quotient of two entire functions of order at most one, hence of order at most one by Hadamard's factorization, and it has no zeros: h is a polynomial of degree at most one. The symmetry gives $h(s) - h(1-s) \in 2\pi i\mathbb{Z}$, which forces the degree to be zero. The last statement compares the limits at $+\infty$. □

Remark 14.32 (Order one cannot be dropped). For real $\nu_5 \neq 0$ put $f(s) = \zeta(s)\exp\big(-\nu_5(s-\frac{1}{2})^2\big)$ and $e^{h(s)} = \exp\big(\nu_5(s-\frac{1}{2})^2\big)$. Then f is meromorphic of order two with the functional equation, the pole and the zeros of ζ; e^h is entire, without zeros, of finite order and symmetric; $g = f\,e^h = \zeta$ is an absolutely convergent Dirichlet series in $\Re s > 1$; and e^h is not constant. In Theorem 14.3 the conclusion $\tilde{h}_0 \equiv 0$ uses that $(s-1)\zeta_{Z1}$ has order one (Proposition 11.1). The location of the zeros of f plays no role in Proposition 14.31.

14.11. **The half-line form of Hamburger's theorem for a pair.** The next two theorems are stated for a general pair: a function with the functional equation and a Dirichlet series, joined on the real half-line only.

Theorem 14.33 (The ladder). *Let* $\mathsf{d}\colon [1,\infty) \to \mathbb{R}$ *be measurable with* $\int_1^\infty |\mathsf{d}(x)|\,x^{-\mathsf{e}_1-1}\,dx < \infty$ *for some* $\mathsf{e}_1 \geq 0$*, and put*

$$h(\sigma) := \sigma\int_1^\infty \mathsf{d}(x)\,x^{-\sigma-1}\,dx \qquad (\sigma > \mathsf{e}_1). \tag{14.61}$$

For every integer $N \geq 2$*,*

$$h(\sigma) = O(\sigma N^{-\sigma}) \quad (\sigma \to +\infty) \iff \mathsf{d}(x) = 0 \text{ for almost every } x \in [1,N). \tag{14.62}$$

Proof. If $\mathsf{d} = 0$ almost everywhere on $[1,N)$, then $|h(\sigma)| \leq \sigma N^{-(\sigma-\mathsf{e}_1)}\int_N^\infty |\mathsf{d}|x^{-\mathsf{e}_1-1}dx$ for $\sigma > \mathsf{e}_1$. Conversely, put $d(u) := \mathsf{d}(e^u)$, so that $d\,e^{-\mathsf{e}_1 u} \in L^1(0,\infty)$, and

$$\mathsf{Y}(s) := \int_0^{\log N} d(u)\,e^{s(\log N - u)}\,du. \tag{14.63}$$

Y is entire of exponential type at most $\log N$, bounded on $\Re s \leq 0$ by $\int_0^{\log N}|d|$, and $\mathsf{Y}(\sigma) \to 0$ as $\sigma \to -\infty$. For real $\sigma > \mathsf{e}_1$,

$$\mathsf{Y}(\sigma) = \frac{N^\sigma h(\sigma)}{\sigma} - \int_{\log N}^\infty d(u)\,e^{-\sigma(u-\log N)}\,du, \tag{14.64}$$

where the first term is bounded by hypothesis and the second by $N^{\mathsf{e}_1}\int|d|e^{-\mathsf{e}_1 u}du$; so Y is bounded on the real axis. The Phragmén–Lindelöf theorem in the two quadrants $\{\Re s > 0,\ \pm\Im s > 0\}$ (angle $\pi/2$, order one, bounded on both sides) shows that Y is bounded in $\mathbb{C}$, hence constant, hence 0. By the uniqueness of the Laplace transform on $(0,\log N)$, $d = 0$ almost everywhere there. □

No functional equation, no continuation of h and no uniformity in t enter Theorem 14.33: the hypothesis lives on $\sigma \to +\infty$.

Theorem 14.34 (Hamburger's theorem for a pair, half-line form). *Let* f *satisfy (H0) and the functional equation* (11.3)*, and suppose that for* $\Re s > 1$

$$\log f(s) = s\int_1^\infty J_f(x)\,x^{-s-1}\,dx, \qquad \int_1^\infty |J_f(x)|\,x^{-\sigma-1}\,dx < \infty \quad (\sigma > 1), \tag{14.65}$$

with J_f *real and measurable. Let* $g(s) = \sum_{n\geq 1} a_n n^{-s}$ *converge absolutely in* $\Re s > 1$*, with* $a_1 = 1$*, and let* $J_g(x) := \sum_{2\leq n\leq x} b_n$*, where* $\log g(s) = \sum_{n\geq 2} b_n n^{-s}$ *for large* σ*. Put* $h(\sigma) := \log g(\sigma) - \log f(\sigma)$ *for large real* σ*. The following are equivalent:*

(i) $h(\sigma) = O_N(\sigma N^{-\sigma})$ *as* $\sigma \to +\infty$*, for every integer* $N \geq 2$*;*
(ii) $f = g$*;*

(iii) $f = g = \zeta$.

In this case J_f is almost everywhere the step function $\sum_{p^m \le x} \frac{1}{m}$.

Proof. For large σ, $\log g(\sigma) = \sigma \int_1^\infty J_g(x) x^{-\sigma-1} dx$ by partial summation, so h has the form (14.61) with $\mathsf{d} = J_g - J_f$. (i)⇒(ii): Theorem 14.33 gives $\mathsf{d} = 0$ almost everywhere, so $\log f = \log g$ for large σ and $f = g$ by analytic continuation. (ii)⇒(iii): g then satisfies (H0)–(H3), so $g = C_H \zeta$, and $a_1 = 1$ gives $C_H = 1$. (iii)⇒(i) because $h \equiv 0$. □

Corollary 14.35 (The ladder for the pair $(\zeta_\lambda, \tilde\zeta_{\mathbb{Z},\lambda})$)**.** *Let $\lambda \in [0,1]$ and $N \ge 2$. Then $\tilde h_\lambda(\sigma) = O(\sigma N^{-\sigma})$ as $\sigma \to +\infty$ if and only if the Mellin count J_λ^{M} of ζ_λ coincides almost everywhere on $[1, N)$ with the step function $\sum_{n^m \le x} \tilde c_n(\lambda)/m$; in particular J_λ^{M} then vanishes on $(1,2)$ and is constant between consecutive integers below N. Consequently*

$$\text{(D)} \iff \tilde h_0(\sigma) = O_N(\sigma N^{-\sigma}) \text{ for every } N \ge 2, \tag{14.66}$$

and at the clock end the first rung already fails: $\tilde h_1(\sigma) \to \log L \ne 0$.

Proof. Theorem 14.33 with $\mathsf{d} = J_g - J_\lambda^{\mathrm{M}}$, where J_g is the count of $\log \tilde\zeta_{\mathbb{Z},\lambda} = \sum_m \frac{1}{m} \sum_n \tilde c_n(\lambda) n^{-ms}$; the integrability is Theorem 12.5, whose proof applies to every ζ_λ, and Corollary 13.11. (14.66) is Theorem 14.34 with Corollary 13.11 and Theorem 12.7, and the last statement is Proposition 13.15. □

The joining factor therefore measures the mass of dJ_λ^{M} outside the integers, and its rate of decay on the real axis is the first integer below which there is such mass. The Nth rung at $\lambda = 0$ is the statement that J_{Z1}^{M} is constant on $(N-1, N)$; it is (D) restricted to one cell.

14.12. **The bridge theorem: the phase of the joining factor on the critical line.**

Theorem 14.36 (Bridge theorem, phase form)**.** *Let f be meromorphic in $\mathbb{C}$, real on the real axis, with (H0), the functional equation* (11.3)*, and $\lim_{\sigma\to+\infty} f(\sigma) = \varpi_f > 0$. Let $g(s) = \sum_{n\ge1} a_n n^{-s}$ converge absolutely in $\Re s > 1$, with real a_n and $a_1 = 1$. Suppose that $g = f\, e^h$ in $\Re s > \frac12$ with h holomorphic there and real on $(1,\infty)$, and that*

(B1) *$\Im h$ extends continuously to $\Re s \ge \frac12$ and vanishes on the line $\Re s = \frac12$;*

(B2) *there are $\mathsf{e}_1, \mathsf{e}_2 \ge 0$ with*

$$|h(s)| \le C\,(1+|s|)^{\mathsf{e}_1}\Big(1 + (\sigma - \tfrac12)^{-\mathsf{e}_2}\Big) \qquad (\Re s > \tfrac12). \tag{14.67}$$

Then $h \equiv -\log \varpi_f$, $g = \zeta$ and $f = \varpi_f\, \zeta$.

Proof. Reflection. $v := \Im h$ is harmonic in $\Re s > \frac12$, continuous on the closed half-plane and 0 on its boundary. By the reflection principle [1, Ch. 4, §6.5] the function equal to $v(s)$ for $\Re s \ge \frac12$ and to $-v(1-\bar s)$ for $\Re s < \frac12$ is harmonic in $\mathbb{C}$. It is the imaginary part of an entire function, which can be chosen equal to h in $\Re s > \frac12$; in $\Re s < \frac12$ it differs from $\overline{h(1-\bar s)}$ by a real constant, and this constant is 0 because the real part of an entire function is continuous across the line. Thus h is entire and $h(1-\bar s) = \overline{h(s)}$. Since h is real on $(1,\infty)$, $h(\bar s) = \overline{h(s)}$, and therefore $h(1-s) = h(s)$.

Growth. By (14.67) and the symmetry, $|h(s)| \le C(1+|s|)^{\mathsf{e}_1}(1+|\sigma-\frac12|^{-\mathsf{e}_2})$ for $\Re s \ne \frac12$. The function $\log|h|$ is subharmonic, so for every s

$$\log|h(s)| \le \frac{1}{2\pi}\int_0^{2\pi} \log\big|h\big(s+e^{i\phi}\big)\big|\, d\phi \le \log C + \mathsf{e}_1 \log(2+|s|) + \log 2 + \frac{\mathsf{e}_2}{2\pi}\int_0^{2\pi} \log^+ \frac{1}{|\sigma - \frac12 + \cos\phi|}\, d\phi. \tag{14.68}$$

For real y, $\frac{1}{2\pi}\int_0^{2\pi} \log|y+\cos\phi|\, d\phi \ge -\log 2$, so the last integral is at most $2\pi\big(\log 2 + \log(2+|s|)\big)$. Hence $|h(s)| \le C'(2+|s|)^{\mathsf{e}_1+\mathsf{e}_2}$ in $\mathbb{C}$, and h is a polynomial. On the real axis $h(\sigma) = \log g(\sigma) - \log f(\sigma) \to -\log\varpi_f$ as $\sigma \to +\infty$, so the polynomial is the constant $-\log\varpi_f$.

Hamburger. $g = f/\varpi_f$ satisfies (H0), (H1), and (H2)–(H3) with the second function equal to g; so $g = C_H\zeta$, and $a_1 = 1$ gives $C_H = 1$. □

The symmetry of the joining factor, its continuation to $\Re s \le \frac12$ and its finite order are not hypotheses of Theorem 14.36: they follow from (B1) and (B2), which are written on the closed half-plane $\Re s \ge \frac12$ where the joining factor of the family is defined. Theorem 14.3 is the special case in which the continuation is assumed.

Corollary 14.37 (The bridge theorem for the family)**.** (i) *For $0 < \lambda \le 1$ the joining factor $\tilde h_\lambda$ does not satisfy both* (B1) *and* (B2)*.*
(ii) *(D) holds if and only if $\tilde h_0$ satisfies* (B1) *and* (B2)*.*

Proof. Theorem 14.36 applies with $f = \zeta_\lambda$ (Theorem 10.1, Proposition 13.1), $g = \tilde\zeta_{\mathbb Z,\lambda}$ (Corollary 13.11) and $h = \tilde h_\lambda$ (Theorem 13.12(iii)). If (B1) and (B2) hold, then $\zeta_\lambda = \varpi_\lambda\zeta$ is an absolutely convergent Dirichlet series in $\Re s > 1$, which is excluded for $0 < \lambda \le 1$ by Proposition 14.1; for $\lambda = 0$ it is (D). Conversely (D) gives $\tilde h_0 \equiv 0$ (Theorem 13.13). □

Define the *phase defect* of the family, when the limit exists,

$$\mathsf v_\lambda(t) := \lim_{\sigma\to\frac12{}^+} \Im\tilde h_\lambda(\sigma+it) = \lim_{\sigma\to\frac12{}^+}\Big(\Im\log\tilde\zeta_{\mathbb Z,\lambda}(\sigma+it) - \Im\log\zeta_\lambda(\sigma+it)\Big). \tag{14.69}$$

Proposition 14.38 (The phase of the continuous side)**.** *Let $\lambda \in [0,1]$ and $t > 0$ not an ordinate $\gamma_k(\lambda)$. With the branch of $\log\zeta_\lambda$ that is real on $(1,\infty)$, continued in $\{\Re s > \frac12,\ \Im s > 0\}$,*

$$\lim_{\sigma\to\frac12{}^+} \Im\log\zeta_\lambda(\sigma+it) = -\vartheta(t) + \pi\big(N_\lambda(t) - 1\big), \qquad N_\lambda(t) = \#\{k : \gamma_k(\lambda) \le t\}, \tag{14.70}$$

and in the sense of distributions in $t > 0$

$$\lim_{\sigma\to\frac12{}^+} \Im\mathcal V_\lambda(\sigma+it) = -\pi\sum_{k\ge1}\eta_k\,\delta\big(t-\gamma_k(\lambda)\big). \tag{14.71}$$

Consequently $\mathsf v_\lambda(t) = \lim_{\sigma\to\frac12{}^+}\Im\log\tilde\zeta_{\mathbb Z,\lambda}(\sigma+it) + \vartheta(t) - \pi\big(N_\lambda(t)-1\big)$.

Proof. $\zeta_\lambda = P_\lambda/G$. On $s = \frac12 + it$, $q(s) = -(\frac14+t^2)$ and $1 + q(s)/A_{k,\lambda} = (\gamma_k(\lambda)^2 - t^2)/A_{k,\lambda}$ are real, so $P_\lambda(\frac12+it)$ is real and changes sign at each $\gamma_k(\lambda)$; approaching from $\sigma > \frac12$ the argument of the factor k increases by π as t passes $\gamma_k(\lambda)$, since $\Im\big(\gamma_k(\lambda)^2 + (\sigma - \frac12 + it)^2\big) = 2t(\sigma - \frac12) > 0$. The argument of $G(\frac12+it) = -(\frac14+t^2)\Gamma(\frac14 + \frac{it}{2})\pi^{-1/4-it/2}$ is $\pi + \vartheta(t)$ by (8.29), with the branch fixed by $\zeta_\lambda(\sigma) < 0$ for $\frac12 < \sigma < 1$. This gives (14.70); (14.71) is its derivative in λ, with $\partial_\lambda N_\lambda(t) = -\sum_k \eta_k\delta(t-\gamma_k(\lambda))$. □

By Proposition 14.38, $\mathsf v_\lambda \equiv 0$ asks two things of the discrete function on the critical line: jumps of π of its argument exactly at the ordinates $\gamma_k(\lambda)$, and the phase $-\vartheta(t)$ between them; equivalently, $e^{i\vartheta(t)}\tilde\zeta_{\mathbb Z,\lambda}(\frac12+it)$ is real. At the clock end $\tilde h_1$ is holomorphic in $\Re s > 0$ (Theorem 13.12(iv)), so $\mathsf v_1(t) = \Im\tilde h_1(\frac12+it)$ is a real-analytic function; it does not vanish identically, by Corollary 14.37(i) and the bound $|\tilde h_1(s)| \le C(1+|s|)$ on $\Re s \ge \frac14$ of the proof of Proposition 14.5.

Proposition 14.39 (The cell masses are sums of a one-ordinate function)**.** *For $n \ge 2$ there is a real function $\mathsf o_n(\kappa,\kappa')$ of two real variables, depending only on n and the cell φ_n, such that for all lists $\boldsymbol\kappa, \boldsymbol\kappa' \in \mathscr O$*

$$\tilde c_n(\boldsymbol\kappa) - \tilde c_n(\boldsymbol\kappa') = \sum_{k\ge1}\mathsf o_n(\kappa_k,\kappa_k'), \qquad \mathsf o_n(\kappa,\kappa') = -2\,\Re\big(W_n(\kappa) - W_n(\kappa')\big) \ \text{ if } n \text{ is not a power of } 2, \tag{14.72}$$

and, with $\mathsf L_n(s) := -\log(1-n^{-s})$, for $\Re s > 1$

$$\log\tilde\zeta_{\mathbb Z,\boldsymbol\kappa}(s) - \log\tilde\zeta_{\mathbb Z,\boldsymbol\kappa'}(s) = \sum_{n\ge2}\Big(\sum_{k\ge1}\mathsf o_n(\kappa_k,\kappa_k')\Big)\mathsf L_n(s), \tag{14.73}$$

$$\log\zeta_{\boldsymbol\kappa}(s)-\log\zeta_{\boldsymbol\kappa'}(s)=\sum_{k\ge1}\log\frac{1+q(s)/(\frac14+\kappa_k^2)}{1+q(s)/(\frac14+{\kappa'_k}^2)}. \tag{14.74}$$

Proof. $\tilde c_n(\boldsymbol\kappa)=\langle d\pi_{\boldsymbol\kappa},\varphi_n\rangle$, where $d\pi_{\boldsymbol\kappa}$ is obtained from $J_{\boldsymbol\kappa}$ by truncation at 2 and finite Möbius inversion, both linear, and by (12.16) the difference $J_{\boldsymbol\kappa}-J_{\boldsymbol\kappa'}=-2\Re\sum_k\big(\mathrm{Ei}((\frac12+i\kappa_k)\log x)-\mathrm{Ei}((\frac12+i\kappa'_k)\log x)\big)$ is a sum over k converging in $\mathcal D'(1,\infty)$. This gives (14.72), with the value (13.20) off the powers of 2. (14.73) is the Euler product with Corollary 13.11, and (14.74) is (15.2). □

By Proposition 14.39, the difference of two joining factors is, term by term in n and in k, a sum over the ordinates of one universal function of one ordinate, the joining factor of a single zero factor:

$$\mathsf j(\kappa,\kappa';s):=\sum_{n\ge2}\mathsf o_n(\kappa,\kappa')\,\mathsf L_n(s)-\log\frac{1+q(s)/(\frac14+\kappa^2)}{1+q(s)/(\frac14+\kappa'^2)},\qquad \tilde h_{\boldsymbol\kappa}-\tilde h_{\boldsymbol\kappa'}=\sum_{k\ge1}\mathsf j(\kappa_k,\kappa'_k;s)\ \text{ formally.} \tag{14.75}$$

With $\boldsymbol\kappa'=(\tau_k)$ this writes the phase defect as $\mathsf v_{\boldsymbol\kappa}=\mathsf v_1+\sum_k\Im\mathsf j(\kappa_k,\tau_k;\frac12+it)$, the defect of the clock plus a transport term.

Heuristic 14.40 (What happens to the phase defect along the path)**.** The following is a computation by stationary phase without remainder terms; it is not proved. For n much larger than κ the cell resolves the oscillation, $\Re W_n(\kappa)\approx n^{-1/2}\cos(\kappa\log n)/\log n$, and for n much smaller than $\kappa/2\pi$ the kernel $W_n(\kappa)$ is negligible because φ is smooth (Proposition 14.17). The transition is at $n\approx\kappa/2\pi$, where the phase is $\kappa\log\frac{\kappa}{2\pi}-\kappa=2\pi(\mathcal N_{\mathrm{RVM}}(\kappa)-\frac78)$. This suggests

$$\Im\mathsf j\big(\kappa,\cdot\,;\tfrac12+it\big)\approx\text{amplitude}\times\cos\big(2\pi\mathcal N_{\mathrm{RVM}}(\kappa)+\text{phase}(t)\big),\qquad \big|\mathsf j(\kappa,\cdot\,;\sigma+it)\big|\approx\kappa^{-1/2-\sigma}, \tag{14.76}$$

so that the sum over the ordinates converges absolutely exactly for $\sigma>\frac12$, in agreement with Theorem 13.12(iii), and is at the edge of convergence on the critical line. On the clock, $\mathcal N_{\mathrm{RVM}}(\tau_k)=k-\frac12$ and the phases are aligned, which is why $\tilde h_1$ crosses the line. Along the path each term turns by the angle $(1-\lambda)\eta_k\ell_k$, the rotation of (14.42). At $\lambda=0$, (B1) is the statement that the rotated sum, with the phases $e^{2\pi i\mathcal N_{\mathrm{RVM}}(\gamma_k)}$, converges on the critical line to minus the defect $\mathsf v_1$ of the clock.

14.13. **Transport from the clock on the rungs.** For a real function $\mathsf f\in C_c^\infty(0,\infty)$ put

$$\mathsf c_{\mathsf f}(t):=\int_0^\infty\mathsf f(u)\cos(tu)\,du,\qquad \mathsf k(u):=\frac12\Big(e^{u/2}-\frac{e^{-u/2}}{e^{2u}-1}\Big),\qquad \mathsf k_{\mathsf f}:=\int_0^\infty\mathsf f(u)\,\mathsf k(u)\,du. \tag{14.77}$$

Proposition 14.41 (The rung functional)**.** *For every list $\boldsymbol\kappa\in\mathscr O$,*

$$\mathsf R_{\mathsf f}(\boldsymbol\kappa):=\frac12\int_1^\infty\mathsf f(\log x)\,x^{-1/2}\log x\,dJ^{\mathrm M}_{\boldsymbol\kappa}(x)=\mathsf k_{\mathsf f}-\sum_{k\ge1}\mathsf c_{\mathsf f}(\kappa_k), \tag{14.78}$$

the series converging absolutely. Along the family, with $\mathsf R_{\mathsf f}(\lambda):=\mathsf R_{\mathsf f}\big((\gamma_k(\lambda))\big)$,

$$\mathsf R_{\mathsf f}(\lambda)=\mathsf R_{\mathsf f}(1)-\mathsf S_{\mathsf f}(\lambda),\qquad \mathsf S_{\mathsf f}(\lambda):=\sum_{k\ge1}\Big(\mathsf c_{\mathsf f}\big(\gamma_k(\lambda)\big)-\mathsf c_{\mathsf f}(\tau_k)\Big),\qquad \mathsf S_{\mathsf f}(1)=0. \tag{14.79}$$

If the support of $\mathsf f$ contains no point $\log n$, $n\in\mathbb N$, then

$$\text{(D)}\implies\mathsf S_{\mathsf f}(0)=\mathsf R_{\mathsf f}(1), \tag{14.80}$$

and (D) holds if and only if $\mathsf S_{\mathsf f}(0)=\mathsf R_{\mathsf f}(1)$ for every such $\mathsf f$.

Proof. Pair (12.15), which holds for every list of $\mathscr O$ with the same proof, with $\frac12\mathsf f(\log x)x^{-1/2}$ and put $u=\log x$: the term 1 gives $\frac12\int\mathsf f(u)e^{u/2}du$, the term $x^{-1}(x^2-1)^{-1}$ gives $\frac12\int\mathsf f(u)e^{-u/2}(e^{2u}-1)^{-1}du$, and the sum gives $\sum_k\mathsf c_{\mathsf f}(\kappa_k)$; $\mathsf c_{\mathsf f}$ is rapidly decreasing. (14.79) is the difference of two cases of (14.78). By Theorem 12.7, (D) holds if and only if $J^{\mathrm M}_{Z1}$ is constant between consecutive integers, that is, if and only if $\mathsf R_{\mathsf f}(0)=0$ for every $\mathsf f$ whose support avoids the points $\log n$. □

The number $\mathsf{R_f}(1)$ belongs to the clock; $\mathsf{S_f}(0)$ is the transport from the clock to the ordinates of ζ. (D) is the statement that the transport cancels the defect of the clock on every rung.

Theorem 14.42 (The transport is band-limited in λ)**.** *Let $\mathsf{f} \in C_c^\infty(0,\log N)$. The function $\lambda \mapsto \mathsf{S_f}(\lambda)$ extends to an entire function with*

$$|\mathsf{S_f}(\lambda)| \le C_{\mathsf{f}}(1+|\lambda|\log(2+|\lambda|))\, e^{C_\eta \log N\,|\Im\lambda|} \qquad (\lambda\in\mathbb{C}), \tag{14.81}$$

and for every $j \ge 1$

$$\partial_\lambda^j \mathsf{S_f}(1) = \mathsf{m}_j(\mathsf{f}) := \sum_{k\ge1} \eta_k^j\, \mathsf{c}_{\mathsf{f}}^{(j)}(\tau_k), \qquad \mathsf{S_f}(0) = \sum_{j\ge1} \frac{(-1)^j}{j!}\, \mathsf{m}_j(\mathsf{f}). \tag{14.82}$$

If the support of f lies in $[u_1,\log N]$ with $u_1>0$, then $|\mathsf{m}_j(\mathsf{f})| \le C_{\mathsf{f}}(1+j)^2(C_\eta\log N)^j$.

Proof. By the Paley–Wiener theorem, $|\mathsf{c}_{\mathsf{f}}^{(j)}(t)| \le C_{A,j}\, e^{\log N|\Im t|}(1+|t|)^{-A}$ for complex t and every A. With $\gamma_k(\lambda)=\gamma_k+\lambda\eta_k$, $|\Im\gamma_k(\lambda)| \le C_\eta|\Im\lambda|$ and $|\gamma_k(\lambda)| \ge \gamma_k - C_\eta|\lambda|$. The ordinates with $\gamma_k \le 2C_\eta|\lambda|$ are $O(|\lambda|\log(2+|\lambda|))$ in number and each contributes at most $C\,e^{C_\eta\log N|\Im\lambda|}$; for the others $|\mathsf{c_f}(\gamma_k(\lambda))| \le C\,e^{C_\eta\log N|\Im\lambda|}(\gamma_k/2)^{-2}$. This gives (14.81), the termwise differentiation and the convergence of the Taylor series at $\lambda=1$ evaluated at $\lambda=0$. Finally $\mathsf{c}_{\mathsf{f}}^{(j)}(t) = \int \mathsf{f}(u)u^j\cos(tu+\frac{j\pi}{2})du$ and two integrations by parts give $(1+t^2)|\mathsf{c}_{\mathsf{f}}^{(j)}(t)| \le C_{\mathsf{f}}(1+j)^2(\log N)^j$. □

The scaled lists $(\tau_k - c'\eta_k)$ are the members $\lambda = 1-c'$ of the family, so the scale test of Remark 14.20 is the behaviour of the entire function $\mathsf{S_f}$ on the real axis. In (14.82) the nodes τ_k and the weights $\mathsf{c}_{\mathsf{f}}^{(j)}(\tau_k)$ belong to the clock, and the powers η_k^j do not; on the first rung, $N=2$, the term j is at most $C_{\mathsf{f}}(1+j)^2(C_\eta\log 2)^j/j!$.

Proposition 14.43 (The defect of the clock on a rung)**.** *Let $\mathsf{t}_0 \in (2\pi,\tau_0)$ be the point with $\mathcal{N}_{\mathrm{RVM}}(\mathsf{t}_0)=0$, and let $\mathsf{f}\in C_c^\infty(0,\infty)$. Then*

$$\sum_{k\ge1}\mathsf{c_f}(\tau_k) = \mathsf{I}_0 + 2\sum_{m\ge1}(-1)^m\mathsf{I}_m, \qquad \mathsf{I}_m := \int_{\mathsf{t}_0}^\infty \mathsf{c_f}(t)\cos\!\big(2\pi m\mathcal{N}_{\mathrm{RVM}}(t)\big)\mathcal{N}'_{\mathrm{RVM}}(t)\,dt = O_{\mathsf{f}}(m^{-2}), \tag{14.83}$$

$$\mathsf{I}_0 = -\int_0^\infty \frac{\mathsf{f}(u)}{2\pi u}\Big(\sin(u\mathsf{t}_0)\log\frac{\mathsf{t}_0}{2\pi} + \frac{\pi}{2} - \int_0^{u\mathsf{t}_0}\frac{\sin y}{y}\,dy\Big)du, \tag{14.84}$$

and therefore $\mathsf{R_f}(1) = (\mathsf{k_f}-\mathsf{I}_0) - 2\sum_{m\ge1}(-1)^m\mathsf{I}_m$.

Proof. The function $g(v) := \mathsf{c_f}(\mathcal{T}(v))$, $v\ge0$, with $\mathcal{T}$ the inverse of the clock on $[\mathsf{t}_0,\infty)$, is smooth and rapidly decreasing with its derivatives, and $\tau_k = \mathcal{T}(k-\frac12)$. Poisson's summation formula for the even extension of g at the half-integers gives $2\sum_{k\ge1}g(k-\frac12) = \sum_{m\in\mathbb{Z}}(-1)^m 2\int_0^\infty g(v)\cos(2\pi mv)\,dv$, the Fourier coefficients being $O(m^{-2})$ after two integrations by parts; the substitution $v=\mathcal{N}_{\mathrm{RVM}}(t)$ gives (14.83). For $m=0$, $\mathcal{N}'_{\mathrm{RVM}}(t) = \frac{1}{2\pi}\log\frac{t}{2\pi}$; integrating by parts with the primitive $\int \mathsf{f}(u)\sin(tu)u^{-1}du$ of $\mathsf{c_f}$ and exchanging the integrals gives (14.84). □

In I_m the phases $2\pi m\mathcal{N}_{\mathrm{RVM}}(t)\mp tu$ have the derivative $m\log\frac{t}{2\pi}\mp u$, which vanishes at $t=2\pi e^{u/m}$: the defect of the clock at the point $x=e^u$ is produced at the heights $2\pi x^{1/m}$, which is the origin of the terms $2\cos(2\pi y)$ and $y^{-1/4}\cos(4\pi\sqrt{y}+\frac{\pi}{4})$ of (14.32). On the rung $(n,n+1)$ the height for $m=1$ lies in $(2\pi n, 2\pi(n+1))$. On the first rung it lies in $(2\pi,4\pi)$, below τ_1 and on both sides of t_0: the defect of the clock on $(1,2)$ is an effect of the start of the clock. Experiments 17.10 and 17.11 evaluate (14.79)–(14.83) for one function f.

14.14. **The multiscale Jacobian of the midpoint clock.** The rung system of Subsection 15.2 asks for a list and free masses simultaneously. Before attempting nonlinear continuation from the clock, one must understand the linearization with respect to the ordinates. The point is genuinely multiscale: the gaps of the midpoint clock tend to zero, so no fixed u-window can carry a uniform inverse, while a window of length comparable with $\log T$ is exactly large enough to resolve the ordinates up to height T.

For $T > \tau_2$ put

$$K(T) := \max\{k : \tau_k \le T\}, \qquad \Delta_T := \min_{1\le k<K(T)} (\tau_{k+1} - \tau_k),$$

and define the finite ordinate block of the linearized rung map by

$$(\mathcal{J}_T h)(u) := -u \sum_{\tau_k \le T} h_k \sin(\tau_k u). \tag{14.85}$$

No genuine zero ordinate and no individual displacement $\eta_k = \tau_k - \gamma_k$ enters this definition.

Theorem 14.44 (Multiscale Jacobian of the midpoint clock)**.** *The following statements hold.*

(i) *For every $T > \tau_2$,*

$$\Delta_T \ge \frac{2\pi}{\log(T/2\pi)}. \tag{14.86}$$

(ii) *Fix $a > 0$ and $\eta > 0$, and put*

$$L_T := (1+\eta)\log(T/2\pi), \qquad I_T = [a, a + L_T].$$

There is $c_{a,\eta} > 0$, independent of T, such that for all finitely supported h on $\{k : \tau_k \le T\}$,

$$\|\mathcal{J}_T h\|^2_{L^2(I_T)} \ge c_{a,\eta} L_T \sum_{\tau_k \le T} |h_k|^2. \tag{14.87}$$

(iii) *No analogue of* (14.87) *with a positive T-independent lower constant can hold on a fixed interval $(0, U)$. Indeed, for the unit vector $h^{(k)} = (e_k - e_{k+1})/\sqrt{2}$,*

$$\|\mathcal{J} h^{(k)}\|_{L^2(0,U)} \le \frac{U^{5/2}}{\sqrt{10}} (\tau_{k+1} - \tau_k) \longrightarrow 0. \tag{14.88}$$

(iv) *For every finite $U > 0$,*

$$\overline{\operatorname{span}}\{u \sin(\tau_k u) : k \ge 1\} = L^2(0, U).$$

Thus the infinite linearized clock has dense range on every finite window, despite the absence of a bounded inverse on a fixed window.

(v) *Let $\mathsf{f} \in C_c^\infty(0, U)$ and $\mathsf{c}_\mathsf{f}(t) = \int_0^\infty \mathsf{f}(u)\cos(tu)\,du$. For every list $\boldsymbol{\kappa}$ satisfying $|\kappa_k - \tau_k| \le C_\eta$ and every $A > 1$,*

$$\sum_{\tau_k > T} |\mathsf{c}_\mathsf{f}(\kappa_k) - \mathsf{c}_\mathsf{f}(\tau_k)| \ll_{A,\mathsf{f},C_\eta} T^{1-A} \log T. \tag{14.89}$$

Proof. The midpoint equation is $\mathcal{N}_{\mathrm{RVM}}(\tau_k) = k - \frac{1}{2}$ with

$$\mathcal{N}'_{\mathrm{RVM}}(t) = \frac{1}{2\pi} \log \frac{t}{2\pi}.$$

For $\tau_{k+1} \le T$, the mean-value theorem gives

$$1 = \mathcal{N}'_{\mathrm{RVM}}(\xi_k)(\tau_{k+1} - \tau_k), \qquad \tau_k < \xi_k < \tau_{k+1} \le T,$$

which proves (14.86).

For (ii), apply Ingham's inequality for non-harmonic exponentials [4] to the separated frequency set $\{\pm\tau_k : \tau_k \le T\}$. By (14.86),

$$L_T \Delta_T \ge 2\pi(1+\eta),$$

so after scaling the frequencies by Δ_T the interval lies a fixed positive distance beyond Ingham's critical length. The lower Ingham constant is therefore bounded below by a positive constant depending only on η; rescaling gives a bound of size L_T for the sine sum. Since $u \geq a$ on I_T, multiplication by u gives (14.87).

For (iii), $|\sin(\tau_{k+1}u) - \sin(\tau_k u)| \leq (\tau_{k+1} - \tau_k)u$, whence

$$\|\mathcal{J}h^{(k)}\|^2_{L^2(0,U)} \leq \frac{(\tau_{k+1} - \tau_k)^2}{2} \int_0^U u^4\, du = \frac{(\tau_{k+1} - \tau_k)^2 U^5}{10}.$$

The same mean-value formula used above shows $\tau_{k+1} - \tau_k \to 0$.

For (iv), suppose $F \in L^2(0,U)$ is orthogonal to every $u\sin(\tau_k u)$. Put $H(u) = uF(u)$ and extend H oddly to $(-U,U)$. Its Fourier transform

$$\widehat{H}(z) = \int_{-U}^{U} H(u)e^{-izu}\, du$$

is entire of exponential type at most U and vanishes at every $\pm\tau_k$. A nonzero entire function of finite exponential type has $O(r)$ zeros in $|z| \leq r$ by Jensen's formula, whereas the midpoint clock has $2\mathcal{N}_{\mathrm{RVM}}(r) + O(1) \asymp r\log r$ points $\pm\tau_k$ in that disk. Hence $\widehat{H} \equiv 0$, so $H = 0$ and therefore $F = 0$ almost everywhere.

For (v), the Paley–Wiener decay of the cosine transform gives $|\mathsf{c}_\mathsf{f}'(t)| \ll_{A,\mathsf{f}} (1 + |t|)^{-A}$ for every A. The mean-value theorem and $|\kappa_k - \tau_k| \leq C_\eta$ give

$$|\mathsf{c}_\mathsf{f}(\kappa_k) - \mathsf{c}_\mathsf{f}(\tau_k)| \ll_{A,\mathsf{f},C_\eta} \tau_k^{-A}$$

for large k. Partial summation with $\mathcal{N}_{\mathrm{RVM}}(t) = O(t\log t)$ yields $\sum_{\tau_k > T} \tau_k^{-A} = O(T^{1-A}\log T)$. □

The theorem identifies the correct scale but does not prove existence of a solution of the rung system. In particular, a fixed-window Banach-space implicit-function argument cannot work because of (14.88). A quantitative continuation must grow the resolved window together with the spectral height.

One precise residual formulation is the following. Fix $r > \frac{1}{2}$, so point masses belong to $H^{-r}(0,U)$, and put

$$\mathsf{D}_{\mathrm{RVM}}(u) := \mathsf{k}(u) - \sum_{k\geq 1} \cos(\tau_k u)$$

in $\mathcal{D}'(0,\infty)$. For $T, U > 0$ define

$$\varepsilon_r(T,U) := \inf_{h,b} \left\| \mathsf{D}_{\mathrm{RVM}} + u \sum_{\tau_k \leq T} h_k \sin(\tau_k u) - \sum_{\log n \leq U} b_n \delta(u - \log n) \right\|_{H^{-r}(0,U)}. \tag{14.90}$$

Theorem 14.44 supplies the finite-height separation, dense-range and tail ingredients. What is not proved is a bound for (14.90), together with control of the minimizing h strong enough to dominate the quadratic remainder in

$$\cos((\tau_k + h_k)u) = \cos(\tau_k u) - h_k u \sin(\tau_k u) + O(h_k^2 u^2),$$

uniformly along a nested sequence with $U \asymp \log T$. Such an estimate would turn the finite-section continuation observed numerically into an existence proof for the infinite rung system. By Theorem 15.5, any resulting solution would then be the zeta ordinates with the von Mangoldt masses. The individual displacements η_k are not data in this formulation.

15. The space of ordinate lists and the existence form of the final step

The constructions of Sections 10–14 use two properties of the ordinates (Remark 13.17). This section applies them to every list with these two properties. The final step then becomes an existence statement in a compact space, and the list of ζ is not part of its formulation.

15.1. **Lists, their zeta functions and their atomizations.** Let $\mathscr{O}$ be the set of sequences $\boldsymbol{\kappa} = (\kappa_k)_{k\ge1}$ of real numbers with

$$\tau_0^+ \le \kappa_1 \le \kappa_2 \le \cdots, \qquad |\kappa_k - \tau_k| \le C_\eta, \qquad \Big|\int_{\tau_0}^{X} \big(N_{\boldsymbol{\kappa}}(\tau) - \mathcal{N}_{\mathrm{RVM}}(\tau)\big)d\tau\Big| \le C\log X \quad (X \ge \tau_0), \tag{15.1}$$

where τ_0^+ is a fixed number with $\tau_0 < \tau_0^+ < \min(\gamma_1, \tau_1)$, $N_{\boldsymbol{\kappa}}(\tau) = \#\{k : \kappa_k \le \tau\}$, C_η is the constant of Lemma 2.2 and C the constant of Proposition 13.1. The clock (τ_k), the ordinates (γ_k) of ζ and every list $(\gamma_k(\lambda))$, $0 \le \lambda \le 1$, belong to $\mathscr{O}$. For $\boldsymbol{\kappa} \in \mathscr{O}$ define, exactly as for the family,

$$\zeta_{\boldsymbol{\kappa}}(s) := \frac{1}{G(s)} \prod_{k\ge1} \Big(1 + \frac{q(s)}{\frac14 + \kappa_k^2}\Big), \qquad \tilde c_n(\boldsymbol{\kappa}), \qquad \tilde\zeta_{\mathbb{Z},\boldsymbol{\kappa}} = \zeta_{\boldsymbol{\kappa}}\, e^{\tilde h_{\boldsymbol{\kappa}}}, \qquad \mathsf{V}_\sigma(\boldsymbol{\kappa}), \tag{15.2}$$

with the cell masses of Definition 13.4, the joining factor of Theorem 13.6 and the function (14.7). Theorem 10.1 (order one, functional equation, zeros exactly $\frac12 \pm i\kappa_k$), Proposition 13.1, Theorem 13.10, Corollary 13.11 and Theorem 13.12(i)–(iii) hold for every $\boldsymbol{\kappa} \in \mathscr{O}$, with constants depending only on those in (15.1): their proofs use the ordinates only through (15.1).

Theorem 15.1 (Rigidity of the coincidence)**.** *The set*

$$\mathscr{C} := \{\boldsymbol{\kappa} \in \mathscr{O} : \ \tilde h_{\boldsymbol{\kappa}} \text{ is constant}\} \tag{15.3}$$

of the lists whose continuous and atomized zeta functions coincide up to a constant factor has at most one element. If $\boldsymbol{\kappa} \in \mathscr{C}$, then $\zeta_{\boldsymbol{\kappa}} = \tilde\zeta_{\mathbb{Z},\boldsymbol{\kappa}} = \zeta$, $\boldsymbol{\kappa}$ is the list of the ordinates of ζ, and (D) holds. Conversely, (D) implies $\mathscr{C} = \{(\gamma_k)\}$.

Proof. If $\tilde h_{\boldsymbol{\kappa}} \equiv c$, then $\zeta_{\boldsymbol{\kappa}} = e^{-c}\tilde\zeta_{\mathbb{Z},\boldsymbol{\kappa}}$ is an absolutely convergent Dirichlet series in $\Re s > 1$ (Corollary 13.11), $(s-1)\zeta_{\boldsymbol{\kappa}}$ is entire of order one and $\zeta_{\boldsymbol{\kappa}}$ satisfies (11.3) (Theorem 10.1). Hamburger's theorem gives $\zeta_{\boldsymbol{\kappa}} = C_H\zeta$, and $C_H = 1$ because both functions have residue 1 at $s = 1$. Hence the nontrivial zeros of ζ with positive ordinate are the points $\frac12 + i\kappa_k$, with multiplicity; this determines $\boldsymbol{\kappa}$ as the list (γ_k), and $\zeta_{Z1} = \zeta_{\boldsymbol{\kappa}} = \zeta$. Then $e^{-c}\tilde\zeta_{\mathbb{Z},\boldsymbol{\kappa}} = \zeta$ and the first coefficients give $c = 0$. Conversely, under (D), $\tilde h_0 \equiv 0$ (Theorem 13.13), so $(\gamma_k) \in \mathscr{C}$. □

Proposition 15.2 (Compactness and continuity)**.** *The space $\mathscr{O}$ and the functions* (15.2) *have the following properties.*

(i) *$\mathscr{O}$ is compact in the product topology of $\prod_k[\tau_k - C_\eta, \tau_k + C_\eta]$.*

(ii) *For every $n \ge 2$, $N \ge 1$ and $\sigma > 1$, the functions $\boldsymbol{\kappa} \mapsto \tilde c_n(\boldsymbol{\kappa})$, $\tilde a_N(\boldsymbol{\kappa})$, $k_N(\boldsymbol{\kappa})$ and $\mathsf{V}_\sigma(\boldsymbol{\kappa})$ are continuous on $\mathscr{O}$.*

(iii) *For every real $\sigma > 1$, $\boldsymbol{\kappa} \mapsto \log\zeta_{\boldsymbol{\kappa}}(\sigma) = -\log G(\sigma) + \sum_k \log\frac{(\sigma-\frac12)^2+\kappa_k^2}{\frac14+\kappa_k^2}$ is a sum of strictly convex functions of the single coordinates κ_k.*

Proof. (i) The product of the intervals is compact, and each condition in (15.1) is closed: for fixed X, $\int_{\tau_0}^X N_{\boldsymbol{\kappa}} = \sum_k (X-\kappa_k)^+$ depends continuously on the finitely many coordinates with $\tau_k \le X + C_\eta$. (ii) For n not a power of 2, $\tilde c_n(\boldsymbol{\kappa}) = \mathfrak{m}_n - 2\Re\sum_k W_n(\kappa_k)$ by (13.20), and the series converges uniformly on $\mathscr{O}$ because W_n is rapidly decreasing and $|\kappa_k - \tau_k| \le C_\eta$. For $n = 2^j$ the term j is $-\frac{\mu(j)}{j}\int_2^3 J_{\boldsymbol{\kappa}}(y)\frac{d}{dy}\varphi_n(y^j)\,dy$ (proof of Theorem 13.10), and $\boldsymbol{\kappa} \mapsto J_{\boldsymbol{\kappa}} \in L^2(2,3)$ is continuous: $J^{\mathrm{M}}_{\boldsymbol{\kappa}}$ is the Fourier–Plancherel transform of $\log\zeta_{\boldsymbol{\kappa}}(\sigma+it)/(\sigma+it)$, which is continuous in $\boldsymbol{\kappa}$ pointwise and dominated in $L^2(\mathbb{R})$ by the uniform bound of Proposition 13.1, and the constant $J^{\mathrm{M}}_{\boldsymbol{\kappa}} - J_{\boldsymbol{\kappa}}$ is the difference of two pairings with a fixed test function, both continuous in $\boldsymbol{\kappa}$. $\tilde a_N$ and k_N are polynomials in finitely many cell masses. The sums $\sum_n |k_n(\boldsymbol{\kappa})|n^{-\sigma'}$, $\sigma' > 1$, are bounded uniformly on $\mathscr{O}$ (Theorem 13.10), so the tails of V_σ are uniformly small, as in the proof of Theorem 14.4(iii). (iii) is the computation in the proof of Proposition 14.2: $\partial_\kappa^2 \log\frac{a^2+\kappa^2}{1/4+\kappa^2} = \mathcal{C}(a,\kappa) - \mathcal{C}(\frac12,\kappa) > 0$ for $a > \frac12$, $\kappa > \frac12$. □

Corollary 15.3 (Existence form of the final step)**.** *For every $\sigma > 1$ the minimum $\min_{\mathscr{O}} \mathsf{V}_\sigma$ is attained, and the following are equivalent:* (i) *(D);* (ii) $\mathscr{C} \neq \varnothing$*;* (iii) *there is $\boldsymbol{\kappa} \in \mathscr{O}$ with $\tilde{\zeta}_{\mathbb{Z},\boldsymbol{\kappa}} = \zeta$;* (iv) $\min_{\mathscr{O}} \mathsf{V}_\sigma = 0$.

Proof. A continuous function on a compact space attains its minimum (Proposition 15.2). (i)⇔(ii) is Theorem 15.1, and (ii)⇒(iii) is part of it. (iii)⇔(iv) because $\mathsf{V}_\sigma(\boldsymbol{\kappa}) = 0$ means $k_n(\boldsymbol{\kappa}) = 0$ for $n \geq 2$. (iii)⇒(i): $\tilde{\zeta}_{\mathbb{Z},\boldsymbol{\kappa}}$ has no zeros in $\Re s > \frac{1}{2}$ (Theorem 13.12(iii)), so, with the functional equation, the nontrivial zeros of ζ lie on $\Re s = \frac{1}{2}$; they are the points $\frac{1}{2} \pm i\gamma_k$, and Theorem 11.2 gives (D). □

In this form the final step does not ask to identify ζ_{Z1} with ζ: it asks for the existence of one list in $\mathscr{O}$ whose continuous and atomized zeta functions coincide, or whose atomized zeta function is ζ; Theorem 15.1 then identifies the list. The path $\lambda \mapsto (\gamma_k(\lambda))$ is one segment in $\mathscr{O}$, from the clock to the list of ζ, and along it V_σ is the function of Theorem 14.4.

Remark 15.4 (What an existence proof uses). Compactness, continuity and convexity do not decide whether $\min_{\mathscr{O}} \mathsf{V}_\sigma$ is 0: they do not involve the form of the functional equation. The existence statement uses the functional equation with the factor $\Gamma(\frac{s}{2})\pi^{-s/2}$, for which a function with both an absolutely convergent Dirichlet series and the functional equation exists, namely ζ. The first variation of the defect is explicit: by (13.20), for n not a power of 2,

$$\frac{\partial \tilde{c}_n(\boldsymbol{\kappa})}{\partial \kappa_k} = 2\,\Im V_n(\kappa_k), \tag{15.4}$$

so the critical points of V_σ in the interior of $\mathscr{O}$ are described by the functions V_n of (13.19) alone.

15.2. **The rung system with free masses.** With the function k of (14.77), consider, for a list $\boldsymbol{\kappa} \in \mathscr{O}$ and real numbers a_n, $n \geq 2$, on which nothing is assumed, the system

$$\sum_{k\geq 1} \cos(\kappa_k u) + \sum_{n\geq 2} \mathsf{a}_n\, \delta(u - \log n) = \mathsf{k}(u) \qquad \text{in } \mathcal{D}'(0,\infty). \tag{15.5}$$

In the variable $x = e^u$ it states that the Mellin count of $\zeta_{\boldsymbol{\kappa}}$ has no mass outside the integers, the masses at the integers being unknown.

Theorem 15.5 (Existence form with free masses)**.** *(D) holds if and only if* (15.5) *has a solution $(\boldsymbol{\kappa}, (\mathsf{a}_n))$ with $\boldsymbol{\kappa} \in \mathscr{O}$. The solution is then unique: $\boldsymbol{\kappa} = (\gamma_k)$, and*

$$\mathsf{a}_n = \frac{\log p}{2\sqrt{n}} \quad \text{if } n = p^m \text{ is a prime power}, \qquad \mathsf{a}_n = 0 \quad \text{otherwise.} \tag{15.6}$$

Proof. By (12.15) for the list $\boldsymbol{\kappa}$, with $u = \log x$, (15.5) is the statement

$$(\log x)\, dJ^{\mathrm{M}}_{\boldsymbol{\kappa}}(x) = \sum_{n\geq 2} 2\sqrt{n}\, \mathsf{a}_n\, \delta_n \qquad \text{in } \mathcal{D}'(1,\infty). \tag{15.7}$$

If (D) holds, $J^{\mathrm{M}}_{Z1} = \sum_{p^m \leq x} \frac{1}{m}$ (Theorem 12.7), which is (15.7) for the list (γ_k) with the masses (15.6). Conversely let (15.7) hold. Then $J^{\mathrm{M}}_{\boldsymbol{\kappa}}$ is constant on $(1,2)$ and between consecutive integers; the truncation at 2 and the finite Möbius inversion keep the support in the integers, so every cell defect vanishes and, by (13.11) and the proof of Theorem 13.6,

$$\zeta_{\boldsymbol{\kappa}}(s) = \exp\bigl(\nu_6 + \nu_7\, 2^{-s}\bigr)\, \tilde{\zeta}_{\mathbb{Z},\boldsymbol{\kappa}}(s) \qquad (\Re s > 1) \tag{15.8}$$

with real constants ν_6, ν_7. Both factors are absolutely convergent Dirichlet series in $\Re s > 1$ (Corollary 13.11 for every list of $\mathscr{O}$). Hence $\zeta_{\boldsymbol{\kappa}}$ satisfies (H0)–(H3), Hamburger's theorem gives $\zeta_{\boldsymbol{\kappa}} = C_H \zeta$, and $C_H = 1$ by the residues at $s = 1$. The zeros of $\zeta_{\boldsymbol{\kappa}}$ are $\frac{1}{2} \pm i\kappa_k$, so $\boldsymbol{\kappa} = (\gamma_k)$ and $\zeta_{Z1} = \zeta$. Then $(\log x)\, dJ^{\mathrm{M}} = \sum_{p^m} \log p\, \delta_{p^m}$, which gives (15.6). □

In Theorem 15.5 the primes are not part of the data: the data are the function k, which comes from the factor G and the pole, the space $\mathscr{O}$, which is defined by the clock, and the condition that the support lie in the points $\log n$. The masses of von Mangoldt are an output. Experiment 17.12 solves a finite section of (15.5) numerically, starting from the clock.

Theorem 15.6 (Nondegeneracy of the solution)**.** *Let $(\boldsymbol{\kappa}, (\mathsf{a}_n))$ be a solution of (15.5) with $\boldsymbol{\kappa} \in \mathscr{O}$. Let $(\dot\kappa_k)$ be a bounded real sequence and $(\dot{\mathsf{a}}_n)_{n\ge 2}$ real numbers with $\dot{\mathsf{a}}_n = O_\varepsilon(n^{-1/2+\varepsilon})$ for every $\varepsilon > 0$, such that*

$$\sum_{k\ge 1} \dot\kappa_k\, u \sin(\kappa_k u) = \sum_{n\ge 2} \dot{\mathsf{a}}_n\, \delta(u - \log n) \qquad \text{in } \mathcal{D}'(0,\infty), \tag{15.9}$$

the derivative of (15.5) in the direction $\big((\dot\kappa_k), (\dot{\mathsf{a}}_n)\big)$. Then $\dot{\mathsf{a}}_n = 0$ for every n, and $\sum_{k:\,\kappa_k = t} \dot\kappa_k = 0$ for every value t of the list. If the ordinates are simple, $\dot\kappa_k = 0$ for every k.

Proof. Step 1. The series $\mathsf{J} := \sum_k \dot\kappa_k\, u \sin(\kappa_k u)$ converges in the space of tempered distributions on $\mathbb{R}$, because $|\int \phi(u) u \sin(\kappa u) du| \le C_\phi \kappa^{-3}$ for Schwartz functions ϕ and $\sum_k \kappa_k^{-3} < \infty$; J is even. By (15.9), J vanishes on $(0, \log 2)$, so its restriction to $(-\log 2, \log 2)$ is supported at 0 and is a finite sum of even derivatives of δ_0. Fix an even $\mathsf{x}_0 \in C_c^\infty(-\log 2, \log 2)$ equal to 1 near 0, and for $\Re z > 0$ put

$$\mathsf{H}_1(z) := \sum_{k\ge 1} \dot\kappa_k \int_0^\infty (1 - \mathsf{x}_0(u))\, u \sin(\kappa_k u)\, e^{-zu}\, du = \tfrac{1}{2}\big\langle \mathsf{J}, (1 - \mathsf{x}_0)\, e^{-z|u|} \big\rangle. \tag{15.10}$$

Approximating by test functions with compact support in $(0,\infty)$ and using (15.9), $\mathsf{H}_1(z) = \sum_{n\ge 2} \dot{\mathsf{a}}_n n^{-z}$ for $\Re z > \frac{1}{2}$, where the series converges absolutely.

Step 2. For $\Re z > 0$, $\int_0^\infty u \sin(\kappa u) e^{-zu} du = 2\kappa z/(z^2 + \kappa^2)^2$, so

$$\mathsf{H}_1(z) = \sum_{k\ge 1} \dot\kappa_k \frac{2\kappa_k z}{(z^2 + \kappa_k^2)^2} - \mathsf{y}(z), \qquad \mathsf{y}(z) := \sum_{k\ge 1} \dot\kappa_k \int_0^{\log 2} \mathsf{x}_0(u)\, u \sin(\kappa_k u)\, e^{-zu}\, du. \tag{15.11}$$

Write e^{-zu} as its Taylor polynomial of degree A at $u = 0$ plus a remainder vanishing to order A. The even extension of x_0 times the remainder is of class C^A and vanishes to order A at 0; for A larger than the order of J near 0 its pairing with J is 0. The terms of the Taylor polynomial contribute constants times powers of z. Hence y is a polynomial. As $z = \sigma \to +\infty$, $\mathsf{H}_1(\sigma) \to 0$ and the first sum in (15.11) is $O(\sigma^{-1}\log\sigma)$, so $\mathsf{y} \equiv 0$.

Step 3. Put $\mathsf{H}(z) := \sum_{n\ge 2} \frac{\dot{\mathsf{a}}_n}{\log n} n^{-z}$ for $\Re z > \frac{1}{2}$, so that $\mathsf{H}' = -\mathsf{H}_1$. By Step 2 there is a real constant ν_8 with

$$\mathsf{H}(z) = \nu_8 + \sum_{k\ge 1} \dot\kappa_k \Big(\frac{\kappa_k}{z^2 + \kappa_k^2} - \frac{1}{\kappa_k} \Big), \tag{15.12}$$

which continues H to a meromorphic function in $\mathbb{C}$, even in z, with at most simple poles at $z = \pm i\kappa_k$, of residue $\pm\frac{1}{2i} \sum_{k':\kappa_{k'} = \kappa_k} \dot\kappa_{k'}$.

Step 4. By Theorem 15.5, $\zeta_{\boldsymbol{\kappa}} = \zeta$, with zeros $\frac{1}{2} \pm i\kappa_k$ of the multiplicity of the list. Put $\mathsf{Z}(s) := \mathsf{H}(s - \frac{1}{2})\, \zeta(s)$. The poles of $\mathsf{H}(s - \frac{1}{2})$ are cancelled by zeros of ζ, and on circles $|z| = r$ at distance at least $1/\log r$ from the points $\pm i\kappa_k$ the series (15.12) is $O(r^2 \log^3 r)$; so $(s-1)\mathsf{Z}(s)$ is entire of finite order: this is (H0). Z is the product of $\sum_n \dot{\mathsf{a}}_n \sqrt{n} (\log n)^{-1} n^{-s}$ and $\zeta(s)$, both absolutely convergent in $\Re s > 1$: this is (H1). Since H is even, $\mathsf{Z}(1-s) = \mathsf{H}(s - \frac{1}{2})\zeta(1-s)$, so Z satisfies the functional equation of ζ: this is (H2) and (H3) with the second function equal to Z. Hamburger's theorem gives $\mathsf{Z} = C_H \zeta$, that is, $\mathsf{H} \equiv C_H$. Since $\mathsf{H}(\sigma) \to 0$ as $\sigma \to +\infty$, $\mathsf{H} \equiv 0$: every $\dot{\mathsf{a}}_n$ and every residue vanishes. □

Remark 15.7 (Scope of Theorem 15.6)*.* The theorem is conditional on the existence of a solution; it shows that a solution is isolated to first order, the Jacobian of (15.5) being injective there on the directions considered, except for the splitting of multiple ordinates. This is the endpoint hypothesis of a continuation or implicit-function argument. Steps 1–3 do not use that $\boldsymbol{\kappa}$ is a solution: at any

list of $\mathscr{O}$ the kernel of the linearization consists of the Dirichlet series H that are even meromorphic functions with simple poles at $\pm i\kappa_k$ of the form (15.12). Such series exist for lists in arithmetic progression, for instance $(2^z+2^{-z}-2\cos\nu_9)^{-1}$, which is even, a power series in 2^{-z} for $\Re z>0$, with simple poles at $z=\pm i(\nu_9+2\pi m)/\log 2$, $m\in\mathbb{Z}$; these lists do not belong to $\mathscr{O}$. For perturbations $\dot{\mathsf{a}}_n$ of polynomial growth the same proof applies with Hamburger's theorem for a finite abscissa of absolute convergence [6]. The injectivity of the Jacobian at the clock and along a path, and with it the existence of a solution, are not proved.

15.3. **The gradient of the vertical defect.** For Dirichlet series $A=\sum_n a_n n^{-s}$, $B=\sum_n b_n n^{-s}$ and $\sigma>1$ put $\langle A,B\rangle_\sigma:=\sum_{n\ge1}a_n\bar b_n\,n^{-2\sigma}$, and for $\boldsymbol{\kappa}\in\mathscr{O}$

$$\mathsf{K}_{\boldsymbol{\kappa}}:=\frac{\tilde\zeta_{\mathbb{Z},\boldsymbol{\kappa}}}{\zeta}=\sum_{n\ge1}k_n(\boldsymbol{\kappa})\,n^{-s},\qquad \mathsf{V}_\sigma(\boldsymbol{\kappa})=\langle\mathsf{K}_{\boldsymbol{\kappa}},\mathsf{K}_{\boldsymbol{\kappa}}\rangle_\sigma-1,\qquad \mathsf{r}_m(\boldsymbol{\kappa}):=\Re\langle\mathsf{K}_{\boldsymbol{\kappa}}\mathsf{L}_m,\mathsf{K}_{\boldsymbol{\kappa}}\rangle_\sigma, \tag{15.13}$$

with $\mathsf{L}_m(s)=-\log(1-m^{-s})$ as in Proposition 14.39. By Proposition 14.39 the derivative $\partial\tilde c_m(\boldsymbol{\kappa})/\partial\kappa_k$ is a function of m and of the value κ_k alone; denote it by $\mathsf{u}_m(\kappa_k)$, so that $\mathsf{u}_m=2\,\Im V_m$ when m is not a power of 2 (15.4).

Proposition 15.8 (Critical points of the vertical defect)**.** *Let $\sigma>1$ and $\boldsymbol{\kappa}\in\mathscr{O}$. Then $|\mathsf{r}_m(\boldsymbol{\kappa})|\le 2m^{-\sigma}(1+\mathsf{V}_\sigma(\boldsymbol{\kappa}))$, $|\mathsf{u}_m(t)|\le Cm^{-1/2}$ off the powers of 2 and $|\mathsf{u}_m(t)|\le C$ on them, and*

$$\frac{\partial\mathsf{V}_\sigma}{\partial\kappa_k}(\boldsymbol{\kappa})=2\,\mathsf{Q}_{\boldsymbol{\kappa}}(\kappa_k),\qquad \mathsf{Q}_{\boldsymbol{\kappa}}(t):=\sum_{m\ge2}\mathsf{r}_m(\boldsymbol{\kappa})\,\mathsf{u}_m(t), \tag{15.14}$$

the series converging absolutely and uniformly in t. If $\mathsf{V}_\sigma(\boldsymbol{\kappa})=0$, then $\mathsf{r}_m(\boldsymbol{\kappa})=0$ for every m and $\mathsf{Q}_{\boldsymbol{\kappa}}\equiv0$.

Proof. Since $\log\mathsf{K}_{\boldsymbol{\kappa}}=\sum_m(\tilde c_m(\boldsymbol{\kappa})-\mathbf{1}_{\text{prime}}(m))\mathsf{L}_m$, $\partial\mathsf{K}_{\boldsymbol{\kappa}}/\partial\tilde c_m=\mathsf{K}_{\boldsymbol{\kappa}}\mathsf{L}_m$ and $\partial\mathsf{V}_\sigma/\partial\tilde c_m=2\mathsf{r}_m$. Multiplication by m^{-rs} multiplies the norm $\langle\cdot,\cdot\rangle_\sigma^{1/2}$ by $m^{-r\sigma}$, so $\langle\mathsf{K}\mathsf{L}_m,\mathsf{K}\mathsf{L}_m\rangle_\sigma^{1/2}\le\langle\mathsf{K},\mathsf{K}\rangle_\sigma^{1/2}\sum_{r\ge1}m^{-r\sigma}/r\le 2m^{-\sigma}\langle\mathsf{K},\mathsf{K}\rangle_\sigma^{1/2}$, and the bound for r_m follows by the Cauchy–Schwarz inequality. In (13.19) the support of $\varphi_m(y^j)$ has length at most $\frac{2}{j}m^{1/j-1}$ and the integrand is at most $m^{-1/(2j)}$ in absolute value, so $|V_m(t)|\le\sum_j\frac{2}{j^2}m^{-1+1/(2j)}\le Cm^{-1/2}$; for $m=2^j$ the cut term is bounded by the total variation of $\varphi_m(y^j)$ times a bound for $\partial_\kappa J_{\boldsymbol{\kappa}}$ on $(2,3)$. The chain rule gives (15.14), with $\sum_m|\mathsf{r}_m||\mathsf{u}_m|\le C\sum_m m^{-\sigma-1/2}+C\sum_j 2^{-j\sigma}<\infty$. If $\mathsf{V}_\sigma=0$, then $\mathsf{K}=1$ and $\mathsf{r}_m=\Re\langle\mathsf{L}_m,1\rangle_\sigma=0$, because L_m has no constant term. □

An interior critical point of V_σ on $\mathscr{O}$ is therefore a list all of whose ordinates are zeros of the single real-analytic function $\mathsf{Q}_{\boldsymbol{\kappa}}$. The minimum of V_σ on $\mathscr{O}$ is attained (Corollary 15.3); if every critical point of V_σ on $\mathscr{O}$, those on the boundary of $\mathscr{O}$ included, were a zero of V_σ, the minimum would be 0 and (D) would follow. This is not proved. A count of zeros does not exclude critical points with $\mathsf{V}_\sigma>0$: $\mathsf{Q}_{\boldsymbol{\kappa}}(t)$ is a sum of terms of frequency $\log m$ with $m\gtrsim t/2\pi$, so it oscillates at the scale $1/\log t$, which is the scale of the gaps between the ordinates.

16. Acceptance criteria and models in which the final step is a theorem

In the construction of this paper the continuous function ζ_λ is built from a list of ordinates and the discrete function $\tilde\zeta_{\mathbb{Z},\lambda}$ from the cells, and the two are joined by $\tilde h_\lambda$. This section states, without ζ, the general criterion under which a list is accepted by both sides, and gives three models: a non-periodic list accepted by a discrete category with two generators, where the acceptance is proved by one unitary matrix; a finite Euler product over the primes, to which the constructions of Sections 2–15 apply verbatim and for which the final step is a theorem; and the diagonal operator over all primes, which shows where the argument of the finite models stops. The complex variable of the first two subsections is denoted by z, and the half-plane $\Im\mathsf{z}>0$ plays the role of $\Re s>1$.

16.1. **The acceptance criterion: window and frequencies.** Let $\mathfrak{k} > 0$ and let $\boldsymbol{\kappa} = (\kappa_k)_{k\ge1}$ be a nondecreasing list of positive numbers with $|\kappa_k - k\pi/\mathfrak{k}|$ bounded. The *continuous side* is the canonical product

$$\mathfrak{C}_{\boldsymbol{\kappa}}(\mathsf{z}) := \mathsf{z}\prod_{k\ge1}\Big(1-\frac{\mathsf{z}^2}{\kappa_k^2}\Big), \tag{16.1}$$

an odd entire function of order one with real zeros; the oddness plays the role of the functional equation. We assume that $\mathfrak{C}_{\boldsymbol{\kappa}}$ has exponential type $\mathfrak{k}$ and polynomial growth on the real axis, as $\sin(\mathfrak{k}\mathsf{z})$ has for the list $\kappa_k = k\pi/\mathfrak{k}$. The *discrete side* is given by a closed discrete set $\mathscr{F} \subset (0,\infty)$ of frequencies that is closed under addition; put $\mathscr{F}^+ := \mathscr{F} \cup \{0\}$. The discrete category consists of the generalized Dirichlet series $\sum_{\mathfrak{l}\in\mathscr{F}^+} \mathfrak{c}_{\mathfrak{l}}\, e^{i\mathfrak{l}\mathsf{z}}$, absolutely convergent in a half-plane $\Im\mathsf{z} > \nu_{10}$, and of their Euler forms $\exp\big(\sum_{\mathfrak{l}\in\mathscr{F}} \mathfrak{b}_{\mathfrak{l}}\, e^{i\mathfrak{l}\mathsf{z}}\big)$. For $\Im\mathsf{z} > 0$,

$$\frac{\mathfrak{C}'_{\boldsymbol{\kappa}}}{\mathfrak{C}_{\boldsymbol{\kappa}}}(\mathsf{z}) = \frac{1}{\mathsf{z}} + \sum_{k\ge1}\frac{2\mathsf{z}}{\mathsf{z}^2-\kappa_k^2} = -i\int_0^\infty e^{i\mathsf{z}u}\Big(1+2\sum_{k\ge1}\cos(\kappa_k u)\Big)du, \tag{16.2}$$

the bracket being a tempered distribution in u: it is the spectral density of the list.

Theorem 16.1 (Acceptance criterion)**.** *For a list $\boldsymbol{\kappa}$ as above the following are equivalent:*

(i) *the spectral density is atomic on the frequencies: there are real $\mathfrak{w}_{\mathfrak{l}}$ with $\sum_{\mathfrak{l}\in\mathscr{F}} |\mathfrak{w}_{\mathfrak{l}}| e^{-\nu_{10}\mathfrak{l}} < \infty$ and*

$$1 + 2\sum_{k\ge1}\cos(\kappa_k u) = \sum_{\mathfrak{l}\in\mathscr{F}} \mathfrak{w}_{\mathfrak{l}}\,\delta(u-\mathfrak{l}) \qquad \text{in } \mathcal{D}'(0,\infty); \tag{16.3}$$

(ii) *the continuous function has an Euler form: for $\Im\mathsf{z} > \nu_{10}$, with a constant $\mathfrak{c}_0 \neq 0$,*

$$e^{i\mathfrak{k}\mathsf{z}}\,\mathfrak{C}_{\boldsymbol{\kappa}}(\mathsf{z}) = \mathfrak{c}_0\exp\Big(\sum_{\mathfrak{l}\in\mathscr{F}} \mathfrak{b}_{\mathfrak{l}}\, e^{i\mathfrak{l}\mathsf{z}}\Big), \qquad \mathfrak{b}_{\mathfrak{l}} = -\frac{\mathfrak{w}_{\mathfrak{l}}}{\mathfrak{l}}; \tag{16.4}$$

(iii) *the continuous function is a finite sum in the discrete category:*

$$e^{i\mathfrak{k}\mathsf{z}}\,\mathfrak{C}_{\boldsymbol{\kappa}}(\mathsf{z}) = \sum_{\mathfrak{l}\in\mathscr{F}^+\cap[0,2\mathfrak{k}]} \mathfrak{c}_{\mathfrak{l}}\, e^{i\mathfrak{l}\mathsf{z}}, \qquad \mathfrak{c}_0 \neq 0. \tag{16.5}$$

Proof. (i)⇔(ii): by (16.2), (i) is the identity $\mathfrak{C}'_{\boldsymbol{\kappa}}/\mathfrak{C}_{\boldsymbol{\kappa}} = -i\sum_{\mathfrak{l}} \mathfrak{w}_{\mathfrak{l}} e^{i\mathfrak{l}\mathsf{z}}$ in $\Im\mathsf{z} > \nu_{10}$ up to a polynomial in z that accounts for the point $u = 0$; since $\mathfrak{C}'_{\boldsymbol{\kappa}}/\mathfrak{C}_{\boldsymbol{\kappa}}(iy) = -i\big(\frac{1}{y} + \sum_k \frac{2y}{y^2+\kappa_k^2}\big) \to -i\mathfrak{k}$ as $y \to +\infty$, the polynomial is the constant $-i\mathfrak{k}$. Integrating, $\log\mathfrak{C}_{\boldsymbol{\kappa}} = \text{const} - i\mathfrak{k}\mathsf{z} - \sum_{\mathfrak{l}} \frac{\mathfrak{w}_{\mathfrak{l}}}{\mathfrak{l}} e^{i\mathfrak{l}\mathsf{z}}$, and conversely. (ii)⇒(iii): expanding the exponential, $\mathsf{N}(\mathsf{z}) := e^{i\mathfrak{k}\mathsf{z}}\mathfrak{C}_{\boldsymbol{\kappa}}(\mathsf{z}) = \sum_{\mathfrak{l}\in\mathscr{F}^+} \mathfrak{c}_{\mathfrak{l}} e^{i\mathfrak{l}\mathsf{z}}$, absolutely convergent in $\Im\mathsf{z} > \nu_{10}$. On the other hand N is entire of exponential type with polynomial growth on the real axis and is bounded by a polynomial in $\Im\mathsf{z} \ge 0$; by the Paley–Wiener–Schwartz theorem the Fourier transform of its restriction to a horizontal line is supported in $[0, 2\mathfrak{k}]$. On the line $\Im\mathsf{z} = \nu_{10} + 1$ this transform is $\sum_{\mathfrak{l}} \mathfrak{c}_{\mathfrak{l}} e^{-\mathfrak{l}(\nu_{10}+1)}\delta_{\mathfrak{l}}$, so $\mathfrak{c}_{\mathfrak{l}} = 0$ for $\mathfrak{l} > 2\mathfrak{k}$, and the sum is finite because $\mathscr{F}^+$ is discrete. (iii)⇒(ii): the logarithm of a finite sum with $\mathfrak{c}_0 \neq 0$ is, for large $\Im\mathsf{z}$, an absolutely convergent series of exponentials whose frequencies are sums of elements of $\mathscr{F}$, hence elements of $\mathscr{F}$. □

The continuous side contributes a *window*: by its type and its symmetry, the frequencies of $e^{i\mathfrak{k}\mathsf{z}}\mathfrak{C}_{\boldsymbol{\kappa}}$ lie in $[0, 2\mathfrak{k}]$. The discrete side contributes the *admissible frequencies* $\mathscr{F}^+$. A list is accepted by both sides if and only if its function lives in the intersection, and it is then determined by the finitely many coefficients $\mathfrak{c}_{\mathfrak{l}}$, $\mathfrak{l} \in \mathscr{F}^+ \cap (0, \mathfrak{k}]$, the others being fixed by the oddness. If $\mathscr{F} = 2\mathfrak{k}\mathbb{N}$, the intersection is $\{0, 2\mathfrak{k}\}$, the only function is a multiple of $\sin(\mathfrak{k}\mathsf{z})$, and the only accepted list is the clock $\kappa_k = k\pi/\mathfrak{k}$. Hamburger's theorem is the statement that for the window fixed by the factor $\Gamma(\frac{s}{2})\pi^{-s/2}$ and the frequencies $\log n$ the space of accepted functions has dimension one.

16.2. **A non-periodic accepted list: one unitary matrix.** Let $\mathfrak{p},\mathfrak{q}>0$ with $\mathfrak{p}/\mathfrak{q}\notin\mathbb{Q}$, $0<\mathfrak{v}<1$, $\mathfrak{k}=\mathfrak{p}+\mathfrak{q}$, and

$$\mathsf{U}:=\begin{pmatrix}\mathfrak{v} & \sqrt{1-\mathfrak{v}^2}\\ \sqrt{1-\mathfrak{v}^2} & -\mathfrak{v}\end{pmatrix},\qquad \mathsf{X}(\mathsf{z}):=\begin{pmatrix}e^{2i\mathfrak{p}\mathsf{z}} & 0\\ 0 & e^{2i\mathfrak{q}\mathsf{z}}\end{pmatrix}, \tag{16.6}$$

$$\mathfrak{D}(\mathsf{z}):=\det\bigl(I-\mathsf{U}\mathsf{X}(\mathsf{z})\bigr)=1-\mathfrak{v}\,e^{2i\mathfrak{p}\mathsf{z}}+\mathfrak{v}\,e^{2i\mathfrak{q}\mathsf{z}}-e^{2i(\mathfrak{p}+\mathfrak{q})\mathsf{z}}=-2i\,e^{i(\mathfrak{p}+\mathfrak{q})\mathsf{z}}\,\mathfrak{S}(\mathsf{z}), \tag{16.7}$$

$$\mathfrak{S}(\mathsf{z}):=\sin\bigl((\mathfrak{p}+\mathfrak{q})\mathsf{z}\bigr)+\mathfrak{v}\sin\bigl((\mathfrak{p}-\mathfrak{q})\mathsf{z}\bigr). \tag{16.8}$$

Proposition 16.2 (The two-edge model)**.** (i) *All zeros of $\mathfrak{S}$ are real. Each interval between consecutive points $(m+\frac12)\pi/\mathfrak{k}$, $m\in\mathbb{Z}$, contains a zero, and the number of zeros in $[0,T]$, with multiplicity, is $\mathfrak{k}T/\pi+O(1)$; hence the positive zeros form a list $\boldsymbol{\kappa}$ with $|\kappa_k-k\pi/\mathfrak{k}|$ bounded. The set of zeros is not periodic.*

(ii) $\mathfrak{S}(-\mathsf{z})=-\mathfrak{S}(\mathsf{z})$, *and* $\mathfrak{D}(\mathsf{z})=-e^{2i\mathfrak{k}\mathsf{z}}\,\overline{\mathfrak{D}(\bar{\mathsf{z}})}$.

(iii) *For $\Im\mathsf{z}>0$,*

$$\log\mathfrak{D}(\mathsf{z})=-\sum_{r\ge1}\frac{1}{r}\operatorname{tr}\bigl(\mathsf{U}\mathsf{X}(\mathsf{z})\bigr)^r=\sum_{\mathfrak{l}\in\mathscr{F}}\mathfrak{b}_{\mathfrak{l}}\,e^{i\mathfrak{l}\mathsf{z}},\qquad \mathscr{F}=\{2(m\mathfrak{p}+m'\mathfrak{q}):m,m'\ge0\}\setminus\{0\}, \tag{16.9}$$

absolutely convergent, and the spectral density of the zeros of $\mathfrak{S}$, counted with multiplicity, is atomic on $\mathscr{F}$ in the sense of (16.3).

(iv) $\mathfrak{D}(\mathsf{z})=-2i\,\mathfrak{S}'(0)\,e^{i\mathfrak{k}\mathsf{z}}\,\mathfrak{C}_{\boldsymbol{\kappa}}(\mathsf{z})$ *for the list $\boldsymbol{\kappa}$ of all positive zeros of $\mathfrak{S}$: the continuous and the discrete functions coincide up to the factor $e^{i\mathfrak{k}\mathsf{z}}$, and the list is accepted.*

Proof. (i) For $\Im\mathsf{z}>0$ the matrix $\mathsf{U}\mathsf{X}(\mathsf{z})$ has norm at most $\max(e^{-2\mathfrak{p}\Im\mathsf{z}},e^{-2\mathfrak{q}\Im\mathsf{z}})<1$, because U is unitary; so $I-\mathsf{U}\mathsf{X}(\mathsf{z})$ is invertible and $\mathfrak{D}(\mathsf{z})\neq0$. $\mathfrak{S}$ is odd, so it has no zeros in $\Im\mathsf{z}<0$ either. At $\mathsf{z}_m=(m+\frac12)\pi/\mathfrak{k}$, $\sin(\mathfrak{k}\mathsf{z}_m)=(-1)^m$ and $|\mathfrak{v}\sin((\mathfrak{p}-\mathfrak{q})\mathsf{z}_m)|<1$, so $\mathfrak{S}(\mathsf{z}_m)$ has the sign $(-1)^m$ and there is a zero in $(\mathsf{z}_{m-1},\mathsf{z}_m)$. For real t the eigenvalues of the unitary matrix $\mathsf{U}\mathsf{X}(t)$ move counterclockwise as t increases, because $\frac{d}{dt}\mathsf{X}(t)=2i\,\mathrm{diag}(\mathfrak{p},\mathfrak{q})\mathsf{X}(t)$ with $\mathfrak{p},\mathfrak{q}>0$, and $\det\mathsf{U}\mathsf{X}(t)=-e^{2i\mathfrak{k}t}$; the real zeros of $\mathfrak{D}$, with multiplicity, are the passages of an eigenvalue through 1, so their number in $[0,T]$ is $\mathfrak{k}T/\pi+O(1)$. If the zero set had a period $y\neq0$, then $\mathfrak{S}(\mathsf{z}+y)$ and $\mathfrak{S}(\mathsf{z})$, both of type $\mathfrak{k}$ and bounded on the real axis with the same zeros, would be proportional; comparing the coefficients of $e^{\pm i(\mathfrak{p}+\mathfrak{q})\mathsf{z}}$ and $e^{\pm i(\mathfrak{p}-\mathfrak{q})\mathsf{z}}$ gives $2\mathfrak{p}y\in\pi\mathbb{Z}$ and $2\mathfrak{q}y\in\pi\mathbb{Z}$, against $\mathfrak{p}/\mathfrak{q}\notin\mathbb{Q}$. (ii) is immediate; the second identity follows from $\det\mathsf{U}=-1$. (iii) The series is the logarithm of the determinant, convergent because the norm is less than 1; each trace is a combination of exponentials with frequencies in $\mathscr{F}$. The statement on the density is Theorem 16.1, (ii)⇒(i), applied to (iv). (iv) $\mathfrak{S}$ is an odd entire function of exponential type $\mathfrak{k}$, bounded on the real axis, with real zeros; by Hadamard's factorization and the oddness, $\mathfrak{S}(\mathsf{z})=\mathfrak{S}'(0)\,\mathfrak{C}_{\boldsymbol{\kappa}}(\mathsf{z})$. □

In this model nothing has to be joined: the continuous function and the discrete function are the same determinant. The Euler form comes from the expansion of $\det(I-\mathsf{U}\mathsf{X})$ in closed paths, and the real zeros and the symmetry come from the unitarity of U, through the contraction in $\Im\mathsf{z}>0$. The acceptance criterion at this level is the existence of the common object U. In finite rank this is general: the zeros of $\mathsf{P}(e^{i\mathfrak{l}_1\mathsf{z}},\dots,e^{i\mathfrak{l}_r\mathsf{z}})$, for a polynomial P without zeros in the product of the open unit discs and in the product of their exteriors, are real and their spectral density is atomic [26], and conversely the lists with unit masses whose spectral density is atomic with locally finite support are zero sets of exponential polynomials with real zeros [29].

16.3. **A finite Euler product: the constructions of this paper with the final step proved.** Let

$$\mathsf{F}(s):=(1-2^{-s})(1-3^{-s})=1-2^{-s}-3^{-s}+6^{-s}=\sum_{n\mid6}\mu(n)\,n^{-s}. \tag{16.10}$$

It is a Dirichlet series over the integers with an Euler product over the primes 2 and 3, and $\mathsf{F}(-s) = 6^s\,\mathsf{F}(s)$, with axis of symmetry $\Re s = 0$. Its zeros are $s = 0$, double, and $s = \pm i\mathfrak{y}_k$, where

$$\{\mathfrak{y}_k\}_{k\ge1} = \Big\{\frac{2\pi m}{\log 2}\Big\}_{m\ge1} \cup \Big\{\frac{2\pi m}{\log 3}\Big\}_{m\ge1} \quad \text{in increasing order,} \tag{16.11}$$

all simple, a non-periodic list, and, with $2\sinh(\frac{s\log p}{2}) = s\log p\prod_{m\ge1}(1+\frac{s^2\log^2 p}{4\pi^2m^2})$,

$$\mathsf{F}(s) = 6^{-s/2}\,(\log 2\log 3)\,s^2\prod_{k\ge1}\Big(1+\frac{s^2}{\mathfrak{y}_k^2}\Big). \tag{16.12}$$

The constructions of this paper apply as follows. The count of the ordinates is $\lfloor\frac{T\log 2}{2\pi}\rfloor + \lfloor\frac{T\log 3}{2\pi}\rfloor = \frac{T\log 6}{2\pi} - 1 + O(1)$ with a remainder of absolute value at most 1; the midpoint clock is $\frac{\mathfrak{x}_k\log 6}{2\pi} - 1 = k - \frac12$, that is, $\mathfrak{x}_k = (2k+1)\pi/\log 6$, and $\mathfrak{x}_k - \mathfrak{y}_k$ is bounded. For a list $\boldsymbol{\kappa}$ at bounded distance from the clock,

$$\mathsf{F}_{\boldsymbol{\kappa}}(s) := 6^{-s/2}\,(\log 2\log 3)\,s^2\prod_{k\ge1}\Big(1+\frac{s^2}{\kappa_k^2}\Big), \qquad \mathsf{F}_{\boldsymbol{\kappa}}(-s) = 6^s\,\mathsf{F}_{\boldsymbol{\kappa}}(s), \tag{16.13}$$

and the family is $\mathsf{F}_\lambda := \mathsf{F}_{\boldsymbol{\kappa}(\lambda)}$ with $\kappa_k(\lambda) = (1-\lambda)\mathfrak{y}_k + \lambda\mathfrak{x}_k$, so that $\mathsf{F}_0 = \mathsf{F}$ and, since $\prod_{k\ge0}(1+s^2/\mathfrak{x}_k^2) = \cosh(\frac{s\log 6}{2})$,

$$\mathsf{F}_1(s) = \frac{\log 2\log 3}{2}\cdot\frac{s^2\,(1+6^{-s})}{1+s^2\log^2 6/\pi^2}. \tag{16.14}$$

Theorem 16.3 (The final step in the model)**.** (i) (Poisson, prime by prime.) *In $\mathcal{D}'(\mathbb{R})$,*

$$2 + 2\sum_{k\ge1}\cos(\mathfrak{y}_k u) = \sum_{p\in\{2,3\}}\sum_{m\in\mathbb{Z}}\cos\frac{2\pi m u}{\log p} = \sum_{p\in\{2,3\}}\log p\sum_{m\in\mathbb{Z}}\delta(u - m\log p). \tag{16.15}$$

The spectral density of the list $(\mathfrak{y}_k)$ is atomic on the points $m\log p$: the continuous function built from the list is the Dirichlet series with Euler product, $\mathsf{F}_0 = \mathsf{F}$.

(ii) (Hamburger's theorem of the model.) *Let f be entire of finite order with $f(-s) = 6^s f(s)$ and $f(0) = 0$, and let $f(s) = \sum_{n\ge1} a_n n^{-s}$ converge absolutely in some half-plane $\Re s > \nu_{12}$. Then $f = a_1\mathsf{F}$.*

(iii) (The family.) *F_λ is an absolutely convergent Dirichlet series in a half-plane if and only if $\lambda = 0$. In particular the clock function F_1 is not a Dirichlet series.*

Proof. (i) is the Poisson summation formula $\sum_{m\in\mathbb{Z}} e^{2\pi i m u/\log p} = \log p\sum_{m\in\mathbb{Z}}\delta(u - m\log p)$ for $p = 2$ and $p = 3$; the second statement is (16.12). (ii) f is bounded in $\Re s \ge \nu_{12} + 1$, and $|f(s)| = 6^{-\sigma}|f(-s)| \le C\,6^{-\sigma}$ in $\Re s \le -\nu_{12} - 1$; by the Phragmén–Lindelöf theorem f is bounded in every vertical strip. For every real σ, moving the line of integration,

$$a_n = \lim_{T\to\infty}\frac{1}{2T}\int_{-T}^{T} f(\sigma+it)\,n^{\sigma+it}\,dt, \qquad |a_n| \le C\,(6/n)^{-\sigma} \quad (\sigma \le -\nu_{12} - 1), \tag{16.16}$$

and $\sigma\to-\infty$ gives $a_n = 0$ for $n > 6$. The functional equation reads $\sum_{n\le6} a_n n^s = \sum_{n\le6} a_n (6/n)^s$; by the uniqueness of the exponents, $a_n = 0$ unless $n \mid 6$, $a_6 = a_1$ and $a_3 = a_2$. Finally $f(0) = 2a_1 + 2a_2 = 0$. (iii) F_λ is entire of order one, satisfies the functional equation and vanishes at 0. If it is a Dirichlet series, (ii) gives $\mathsf{F}_\lambda = a_1\mathsf{F}$, the lists of zeros coincide, and $\lambda(\mathfrak{x}_k - \mathfrak{y}_k) = 0$ for every k, so $\lambda = 0$. □

In the model every object of the programme exists and the analogue of (D) is proved. What makes $\lambda = 0$ close is a property of the list that the clock and the lists with $\lambda \ne 0$ do not have,

$$\{\mathfrak{y}_k\} = \frac{2\pi}{\log 2}\,\mathbb{N} \;\cup\; \frac{2\pi}{\log 3}\,\mathbb{N}, \tag{16.17}$$

a superposition of one clock for each prime; it passes the scale test of Remark 14.20 and it is the hypothesis of the Poisson formula (16.15). The same statements hold for $\prod_{p\le X}(1-p^{-s}) = \sum_{n\mid X\#}\mu(n)n^{-s}$, with $X\#$ the product of the primes up to X, the functional equation with the factor $(-1)^{\pi(X)}(X\#)^s$, a zero of order $\pi(X)$ at the origin and the list $\bigcup_{p\le X}\frac{2\pi}{\log p}\mathbb{N}$. As $X\to\infty$ these functions tend to $1/\zeta$ in $\Re s>1$ only; their axis of symmetry is $\Re s=0$, not $\Re s=\frac12$, and their lists do not converge to the ordinates of ζ, which are not a superposition of one clock for each prime.

16.4. **The diagonal operator over all primes and the three abscissas.** On $\mathscr{L} := \ell^2(\{p \text{ prime}\})$ let $\mathsf{X}(s) := \operatorname{diag}(p^{-s})$. The closed paths of this model are the primes themselves, without scattering between them, and

$$\det\bigl(I-\mathsf{X}(s)\bigr) = \prod_p (1-p^{-s}) = \frac{1}{\zeta(s)} \qquad (\Re s>1). \tag{16.18}$$

$\mathsf{X}(s)$ has norm $2^{-\sigma}<1$ for $\sigma>0$, it is of trace class if and only if $\sigma>1$, and Hilbert–Schmidt if and only if $\sigma>\frac12$. With $\mathfrak{P}(s) := \sum_p p^{-s}$, the regularized determinant is

$$\det{}_2\bigl(I-\mathsf{X}(s)\bigr) = \prod_p (1-p^{-s})e^{p^{-s}} = \frac{e^{\mathfrak{P}(s)}}{\zeta(s)} = \exp\Bigl(-\sum_{m\ge2}\frac{1}{m}\,\mathfrak{P}(ms)\Bigr) \qquad (\Re s>\tfrac12), \tag{16.19}$$

holomorphic and without zeros in $\Re s>\frac12$ because $I-\mathsf{X}(s)$ is invertible there.

Proposition 16.4 (No common object of finite rank in the window of ζ)**.** *Let $X\ge 2$. There is no function $f\not\equiv 0$ that satisfies (H0) and the functional equation* (11.3) *and is an absolutely convergent Dirichlet series in $\Re s>1$ supported on the integers all of whose prime factors are at most X.*

Proof. By Hamburger's theorem $f = C_H\zeta$, and the coefficients of ζ do not vanish at the integers with a prime factor larger than X; so $C_H=0$. □

In the models of Subsections 16.2 and 16.3 three boundaries coincide: the boundary of the domain where the matrix is a strict contraction, the boundary of the domain of convergence of the Euler product, and the axis of symmetry. The real zeros come from this coincidence. For (16.18) they are different:

$$\text{contraction: } \sigma>0; \qquad \text{convergence of the determinant: } \sigma>1; \qquad \text{axis of symmetry: } \sigma=\tfrac12. \tag{16.20}$$

The contraction proves that ζ has no zeros in $\sigma>1$ and that $\zeta\,e^{-\mathfrak{P}}$ is holomorphic without zeros in $\sigma>\frac12$; neither statement reaches the axis, and in the strip $\frac12<\sigma\le1$ the regularized determinant moves the question to the continuation of $\mathfrak{P}$. In the language of this paper, $\log\tilde\zeta_{\mathbb{Z},\lambda} = \sum_n \tilde c_n(\lambda)\mathsf{L}_n$ is the determinant (16.18) with real multiplicities $\tilde c_n(\lambda)$ at all integers, and the strip $\frac12<\sigma\le1$ is where the mean square of the cell masses (Theorem 13.10) and the continuation of the joining factor (Theorem 13.12) are used. By Proposition 16.4 the common object of the window of ζ has no finite-rank approximants in the discrete category; a unitary object of infinite rank from which both ζ_{Z1} and $\tilde\zeta_{\mathbb{Z},0}$ descend, as U does in Proposition 16.2, is not constructed in this paper. A model without ζ with the same separation of the three boundaries is a graph with infinitely many edges of lengths $\mathfrak{l}_j$ such that $\sum_j e^{-\mathfrak{l}_j y}$ diverges for $y\le y_0$, with $y_0>0$.

17. Numerical illustration with 2 001 052 zeros

This section shows what is being constructed and checks that the theorems are consistent with computation. It uses the ordinates $\gamma_1,\dots,\gamma_N$ of the first $N=2\,001\,052$ zeros of ζ (file `zeross.bin`, $\gamma_N = 1\,132\,490.6587$) and the midpoint ordinates $\tau_1,\dots,\tau_N$ computed from (2.5) (file `zLamb.bin`, $\tau_N = 1\,132\,490.7226$). The programs are listed in Appendix A and their complete console output in Appendix B; all figures have linear axes. Each item is an *experiment*: it reports a computation and does not replace a proof.

Products over the ordinates are evaluated as follows: the first 10^5 factors exactly, the factors $10^5 < k \le N$ through power sums of $(\frac{1}{4}+o_k^2)^{-1}$, and the factors $k > N$ by the midpoint rule for the clock, which gives the tail $\frac{1}{2\pi}(q\frac{L_U+1}{U} - \frac{q^2}{2}\frac{3L_U+1}{9U^3} + \frac{q^3}{3}\frac{5L_U+1}{25U^5})$ with $\mathcal{N}_{\mathrm{RVM}}(U) = N$, $L_U = \log(U/2\pi)$.

17.1. **The clock and the zeros.**

Experiment 17.1 (Clock, midpoints and displacements)**.** *Object:* $\mathcal{N}_{\mathrm{RVM}}$, N_ζ, τ_k, $\eta_k = \tau_k - \gamma_k$. *Range:* $k \le N$. *Expected:* (2.4) to rounding, and $|\eta_k| \le C_\eta$ (Lemma 2.2). *Observed:* $\max_k |\mathcal{N}_{\mathrm{RVM}}(\tau_k) - (k - \frac{1}{2})| = 6.985 \cdot 10^{-10}$; $\max_k |\eta_k| = 0.9937$ (at $k = 34$), mean $|\eta_k| = 0.1512$, root mean square 0.1881, mean $\eta_k = 1.7 \cdot 10^{-7}$. By blocks of k, $\max|\eta_k|$ is $0.840, 0.994, 0.915, 0.834, 0.786, 0.763, 0.717$ on $[1, 10]$, $[11, 10^2]$, $[10^2, 10^3]$, $\ldots$, $[10^6, N]$.

Figure 1 shows how the clock places the midpoint ordinates. The staircase N_ζ jumps by one at each γ_k; the smooth clock $\mathcal{N}_{\mathrm{RVM}}$ crosses the half-integer level $k - \frac{1}{2}$ at τ_k. Since $N_\zeta - \mathcal{N}_{\mathrm{RVM}} = S(T) + O(1/T)$ (Figure 2), the horizontal distance between the open square $(\gamma_k, k - \frac{1}{2})$ and the red point $(\tau_k, k - \frac{1}{2})$ is the remainder S converted into ordinate units by the slope $\mathcal{N}'_{\mathrm{RVM}} = \frac{1}{2\pi}\log\frac{T}{2\pi}$. The right panel repeats the picture at height $1.13 \cdot 10^6$, where the mean spacing is 0.519: every τ_k still lies next to its γ_k.




FIGURE 1. The midpoint clock $\mathcal{N}_{\mathrm{RVM}}$ (red), the zero staircase N_ζ (black), the midpoint ordinates τ_k (red points) and the points $(\gamma_k, k - \frac{1}{2})$ (squares). Left: the first nine zeros. Right: the last twelve zeros of the file.




FIGURE 2. The remainder $N_\zeta(T) - \mathcal{N}_{\mathrm{RVM}}(T) = S(T) + O(1/T)$ of (2.2) for $T \le 200$.

17.2. **The λ-family and the endpoint.** Figure 3 shows the core of the method. The lower left panel draws the moving ordinates $\gamma_k(\lambda)$ in clock units, $\mathcal{N}_{\mathrm{RVM}}(\gamma_k(\lambda)) - (k - \frac{1}{2})$: at $\lambda = 1$ all lines meet at 0 (the τ_k), at $\lambda = 0$ they sit at the ordinates of ζ; for $\lambda > 0$ the ordinates $\gamma_k(\lambda)$ remain strictly increasing in k (Theorem 10.1). The lower right panel draws the real functions $Z_\lambda(t) = e^{i\vartheta(t)}\zeta_\lambda(\frac{1}{2}+it)$ for $\lambda = 0, 0.25, 0.5, 0.75, 1$ together with Hardy's function $Z(t) = e^{i\vartheta(t)}\zeta(\frac{1}{2}+it)$: the zeros of every Z_λ are real, they move continuously with λ, and the curve for $\lambda = 0$ lies on top of Z.




FIGURE 3. Top: η_k for every 50th $k \le N$ and its distribution. Bottom: moving ordinates in clock units, $k \le 30$, and the functions $Z_\lambda(t)$ on the critical line.

Experiment 17.2 (The endpoint ζ_{Z1} against ζ)**.** *Object:* $\zeta_{Z1} = P_{Z1}/G$ computed from the N ordinates γ_k, and ζ computed by mpmath. *Range:* $\sigma \in \{2, 1.25, 0.75\}$, $1 \le t \le 2000$ (499 points per line), and the critical line $0.5 \le t \le 60$. *Expected:* if $\zeta_{Z1} = \zeta$, the ratio is 1 up to the truncation of the product. *Observed:* $\max|\zeta_{Z1}/\zeta - 1| = 8.7 \cdot 10^{-10}$, $1.1 \cdot 10^{-9}$, $2.6 \cdot 10^{-9}$ on the three lines, and $\max|Z_{Z1} - Z|/\max(1, |Z|) = 8.4 \cdot 10^{-10}$ on the critical line (Figure 4).




FIGURE 4. Left: $|\zeta_{Z1}/\zeta - 1|$ on three vertical lines. Right: Z_{Z1} and Hardy's Z on the critical line.

Experiment 17.2 is consistent with the identity $\zeta_{Z1} = \zeta$ to nine digits in the range computed; it is a finite computation and not a proof of the identity.

Experiment 17.3 (The value at infinity: the product identity)**.** *Object:* the ordered partial products $P_K = \prod_{k \le K}(\frac{1}{4} + \gamma_k^2)/(\frac{1}{4} + \tau_k^2)$ of (12.12). *Range:* $K \le N$. *Expected:* by Corollary 12.3, $\zeta_{Z1} = \zeta$ is the statement $P_K \to \varpi_{\mathrm{RVM}}$. *Observed:* $P_K = 0.96256$, 0.96248, 0.96229, 0.961951, 0.961930, 0.9619418189, 0.9619415390 for $K = 10, 10^2, \ldots, 10^6, N$, against $\varpi_{\mathrm{RVM}} = 0.9619415807$

from Experiment 17.4; for $K \geq 10^6$, P_K stays in $[0.9619389, 0.9619440]$ with mean 0.9619416633 (Figure 5). The flow at infinity (13.3) is $\Upsilon(\lambda) = 0.0436950,\ 0.0412445,\ 0.0387987,\ 0.0363572,\ 0.0339198$ for $\lambda = 0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1$, decreasing as in (13.4) with slope close to $-\sum_k 2\eta_k^2/\gamma_k^2 = -0.0098211$, and $\int_0^1 \Upsilon\, d\lambda = \sum_{k\leq N} \log \frac{1/4+\tau_k^2}{1/4+\gamma_k^2} = 0.0388016004$ against $\log L = 0.0388015571$.




FIGURE 5. The partial products P_K of (12.12) against ϖ_{RVM} (dashed). Left: $K \leq 10^4$. Right: $(P_K - \varpi_{\mathrm{RVM}}) \cdot 10^6$ for $10^5 \leq K \leq N$. The value of the endpoint at $+\infty$ is $\varpi_{Z1} = \varpi_{\mathrm{RVM}}/\lim_K P_K$ (Theorem 12.1).

17.3. The transition theorem and ζ_{RVM} against ζ.

Experiment 17.4 (Transition theorem). *Object:* ζ_{RVM}, $w(s)$, ϖ_{RVM}, L. *Range:* $\sigma' \in \{0.25, 0.5, 1.5\}$, $10 \leq t \leq 10^4$. *Expected:* Theorem 8.1 and Corollary 8.8: $t\,|\zeta_{\mathrm{RVM}}/(\varpi_{\mathrm{RVM}}(1+w)) - 1|$ bounded, $|w| = (t/2\pi)^{-\sigma'}(1 + O(1/t))$. *Observed:* Richardson extrapolation on $\Re s = 2$ gives $\varpi_{\mathrm{RVM}} = 0.9619415807$ and $L = 1.0395641690$; $\max_{t>100} t\,|\zeta_{\mathrm{RVM}}/(\varpi_{\mathrm{RVM}}(1+w)) - 1| = 0.0617, 0.0618, 0.0620$ for the three values of σ', and 0.0790 for $\varpi_{\mathrm{RVM}}(1+\chi)$ at $\sigma' = 0.25$; $|w|\,t^{\sigma'} = 1.5832,\ 2.5066,\ 15.66$–$15.75$, i.e. $(2\pi)^{\sigma'}$.

The bottom panels of Figure 6 compare $|\zeta|$ and $|\zeta_{\mathrm{RVM}}|$ on two vertical lines: ζ carries the Euler product and oscillates with the primes, whereas ζ_{RVM} approaches the constant ϖ_{RVM} on $\sigma = 2$ and $\varpi_{\mathrm{RVM}}|1+\chi|$ on $\sigma = 0.75$. This is the analytic content of Theorem 8.11.

17.4. Exponential sums.

Experiment 17.5 (Exponential sums over the two spectra). *Object:* $S(N; \log x) = \sum_{k\leq N} x^{io_k}$ for $o_k = \tau_k$ and $o_k = \gamma_k$. *Range:* $N \leq 2\,001\,052$, $x \in \{2, 3, 4, 6, 10\}$. *Expected:* $|S_{\mathrm{RVM}}| \ll 1 + \log N$ (Theorem 4.1); for γ_k, Landau's term $-\frac{T}{2\pi}\frac{\Lambda(x)}{\sqrt{x}}$. *Observed:* $\max_N |S_{\mathrm{RVM}}(N; \log x)| = 3.57, 3.98, 3.95, 4.53, 9.70$; at $N = 2\,001\,052$, $|\sum x^{i\gamma_k}| = 88\,301,\ 114\,268,\ 62\,435,\ 0.26,\ 2.42$, against Landau's $88\,342,\ 114\,324,\ 62\,467,\ 0,\ 0$.

17.5. Prime counts: π and π_{RVM}.

Experiment 17.6 (Prime counts and RVM primes). *Object:* J_{RVM}, π_{RVM}, c_n^{RVM}, p_j^{RVM} against J, π, the primes. *Range:* $2 \leq x \leq 100$ with all N ordinates of each list (3000 points); $2 \leq x \leq 1000$ with the first $K = 17\,061$ ordinates and a smooth cut (40 000 points). *Expected:* the explicit formula with the γ_k reproduces $\pi(x)$; π_{RVM} is continuous and $\pi_{\mathrm{RVM}}(x) - \mathrm{li}(x) = O(\sqrt{x}(\log\log x)^2/\log x)$ (Theorem 5.5). *Observed:* away from the primes, $|\pi_\gamma - \pi| \leq 0.40$ on $[2, 100]$; $\pi_{\mathrm{RVM}}(100) = 26.08$, $\pi(100) = 25$, $\mathrm{li}(100) = 30.13$; $\pi_{\mathrm{RVM}}(1000) = 169.98$, $\pi(1000) = 168$, $\mathrm{li}(1000) = 177.61$. π_{RVM} is not monotone: it carries a ripple of unit scale, with minimum grid increment -0.25 on $[2, 100]$. The first crossings of the levels $j = 1, \ldots, 8$ are $p_j^{\mathrm{RVM}} = 2.21, 3.16, 5.19, 7.16, 11.14, 14.02, 17.26, 19.18$ against $p_j = 2, 3, 5, 7, 11, 13, 17, 19$.




FIGURE 6. Top: the transition error multiplied by t, and the rescaled vertical limit $t^{\min(\sigma',1)}|1/\zeta_{\mathrm{RVM}} - L|$. Bottom: $|\zeta|$ and $|\zeta_{\mathrm{RVM}}|$ on $\sigma = 2$ and $\sigma = 0.75$.




FIGURE 7. The midpoint spectrum carries no prime frequencies (left); the spectrum of ζ resonates at prime powers (right).

Figure 8 shows π and π_{RVM} in the usual format. The RVM count follows the prime staircase much more closely than li: replacing each γ_k by τ_k moves every zero by less than one unit (Experiment 17.1), and the low ordinates, which shape $\pi(x)$ for small x, are moved least in relative terms. The steps are replaced by a continuous ripple, so the cell masses c_n^{RVM} of Figure 9 are real numbers, occasionally negative, and are not a prime indicator.

Experiment 17.7 (The three ratios and the square-root scale)**.** *Object:* $\mathrm{li}/\pi_{\mathrm{RVM}}$, π/li, π/π_{RVM}, and $(\pi_{\mathrm{RVM}} - \mathrm{li})/B$, $(\pi - \mathrm{li})/B$ with $B(x) = \sqrt{x}(\log\log x)^2/\log x$. *Range:* the grid of Experiment 17.6 on $[8, 1000]$ ($K = 17\,061$ ordinates). *Expected:* the three ratios tend to 1 (Corollary 5.6) and $(\pi_{\mathrm{RVM}} - \mathrm{li})/B$ stays bounded (Theorem 5.5). *Observed:* at $x = 100, 500, 1000$, $\mathrm{li}/\pi_{\mathrm{RVM}} = 1.155, 1.086, 1.045$, $\pi/\mathrm{li} = 0.830, 0.933, 0.946$, $\pi/\pi_{\mathrm{RVM}} = 0.959, 1.013, 0.988$; $\max|\pi_{\mathrm{RVM}} - \mathrm{li}|/B = 2.72$ on $[8, 1000]$, 1.35 on $[50, 1000]$ and 0.72 on $[500, 1000]$, while $\max|\pi - \mathrm{li}|/B = 2.45$ on $[8, 1000]$ (Figure 10).




FIGURE 8. $\pi(x)$ (black staircase), the explicit formula with the zeros of ζ (blue), $\pi_{\mathrm{RVM}}(x)$ (red) and $\mathrm{li}(x) - \mathrm{li}(2)$ (green). The distance between the red and green curves is the quantity bounded in Theorem 5.5; the ratios are in Figure 10.




FIGURE 9. Left: cell masses c_n^{RVM}, the prime indicator and $1/\log n$. Right: real RVM prime levels against the primes up to 1000.




FIGURE 10. Left: the three ratios of Corollary 5.6. Right: $\pi_{\mathrm{RVM}} - \mathrm{li}$ and $\pi - \mathrm{li}$ in units of the scale $B(x)$ of Theorem 5.5. The ratio π/π_{RVM} (black) is the closest of the three to 1; the ratios with li approach 1 at the rate $\frac{1}{2}\,\mathrm{li}(\sqrt{x})/\,\mathrm{li}(x)$ of the bias $J - \pi$.

On $[2, 1000]$ the two normalized errors in Figure 10 have the same size and sign: both counts lie below li by about $\frac{1}{2}\,\mathrm{li}(\sqrt{x})$, the $m = 2$ term of the Möbius inversion (5.20). This is why π/π_{RVM} is much closer to 1 than either ratio with li.

17.6. **The Mertens functions and the ternary division.** The Mertens function is the sum of the ternary sequence $\mu(n) \in \{-1, 0, +1\}$. Its values split into two independent pieces of information:

(a) *the zero set* $\{n : \mu(n) = 0\}$, the non-squarefree integers, of density $1 - 6/\pi^2 = 0.392073$;
(b) *the signs* on the squarefree integers, $\mu(n) = (-1)^{\omega(n)}$, each sign of density $3/\pi^2 = 0.303964$.

The ternary walk μ_h of (1.9)–(1.10) keeps (a) exactly, $\mu_h(n) = 0 \Leftrightarrow \mu(n) = 0$, and replaces (b): at a squarefree n it steps up when $M_h(n-1)$ lies below the target $\mathcal{M}(n) = M_{\mathrm{RVM}}^{\mathrm{crit}}(n)$ and down otherwise. Thus

$$M_h(n) = M_h(n-1) + \mu(n)^2\, \varsigma_n, \qquad |M_h(n) - \mathcal{M}(n)| \text{ small}, \tag{17.1}$$

so that M_h is an integer walk with the zero set of μ that follows $M_{\mathrm{RVM}}^{\mathrm{crit}}$, in the same way as M is the integer walk with the zero set of μ whose signs are $(-1)^{\omega(n)}$.

n	$\mu(n)$	$\mu(n)^2$	$\mathcal{M}(n)$	$M_h(n-1)$	ς_n	$\mu_h(n)$	$M_h(n)$	$M(n)$
1	1	1	0.2925	0	+1	1	1	1
2	−1	1	0.4563	1	−1	−1	0	0
3	−1	1	−0.6523	0	−1	−1	−1	−1
4	0	0	−0.8720	−1	–	0	−1	−1
5	−1	1	−1.7255	−1	−1	−1	−2	−2
6	1	1	−1.3063	−2	+1	1	−1	−1
7	−1	1	−1.5931	−1	−1	−1	−2	−2
8	0	0	−2.4029	−2	–	0	−2	−2
9	0	0	−1.3292	−2	–	0	−2	−2
10	1	1	−1.3280	−2	+1	1	−1	−1
11	−1	1	−2.3272	−1	−1	−1	−2	−2
12	0	0	−2.2067	−2	–	0	−2	−2
13	−1	1	−1.8468	−2	+1	1	−1	−3
14	1	1	−1.9324	−1	−1	−1	−2	−2
15	1	1	−1.5435	−2	+1	1	−1	−1

TABLE 5. The ternary division for $n \le 15$ (the first 40 values are in Appendix B.6). The zero columns of μ and μ_h coincide; the signs of μ_h are chosen by comparing $M_h(n-1)$ with $\mathcal{M}(n)$. For $n \le 12$ the two walks agree; at $n = 13$ the target -1.85 lies above $M_h(12) = -2$, so $\mu_h(13) = +1$ while $\mu(13) = -1$.

Experiment 17.8 (Mertens functions and the ternary walk)**.** *Object:* M, $M_{\mathrm{RVM}}^{\mathrm{crit}}$, M_h; the explicit formulas. *Range:* $n \le 10^6$; explicit formulas on $[2, 2000]$ with all N ordinates of each list. *Expected:* the explicit formula with the zeros of ζ reproduces M; $M_{\mathrm{RVM}}^{\mathrm{crit}}(x) = O(x^{1/2+\varepsilon})$ (Theorem 8.12); M_h tracks $M_{\mathrm{RVM}}^{\mathrm{crit}}$ with the densities of μ. *Observed:*

- explicit formula for M with the γ_k: root mean square error 0.0005 on $[2, 2000]$ off the integers;
- $\zeta'_{\mathrm{RVM}}(\rho_k^{\mathrm{RVM}})/\log(\tau_k/2\pi) = 0.96194$ at $k = 1000$, against $\varpi_{\mathrm{RVM}} = 0.96194$ (Corollary 8.6); the values $|\zeta'(\rho_k)|/\log(\gamma_k/2\pi)$ are spread over $(0, 3)$ (Figure 13);
- spectral sums $2\sum |a|^2$: 0.029034 for ζ, 0.027203 for the midpoint system;
- the integer target agrees with the full explicit formula for $M_{\mathrm{RVM}}^{\mathrm{crit}}$ to 0.019 at 100 integers $\le 10^5$;
- $\max |M_h - \mathcal{M}| = 2.62$ on $[1, 10^6]$ and 1.65 on $[10^4, 10^6]$;
- densities of $(-1, 0, +1)$ for μ: 0.303857, 0.392074, 0.304069;
- densities of $(-1, 0, +1)$ for μ_h: 0.304003, 0.392074, 0.303923;

- $\mu_h(n) = \mu(n)$ at 50.03% of the squarefree $n \le 10^6$, and $\sum_{n\le 10^6} \mu(n)\mu_h(n)/\#\{n \le 10^6 : \mu(n) \ne 0\} = 0.0007$: the sign sequences are uncorrelated although both walks have the same zero set and comparable size;
- $\max_{10\le n\le 10^6} |W(n)|/\sqrt{n} = 0.832, 0.754, 0.688$ for $W = M, M^{\mathrm{crit}}_{\mathrm{RVM}}, M_h$;
- energies $\frac{1}{N}\sum_{n\le N} W(n)^2$ at $N = 10^6$: 14 851, 12 538, 12 538.




FIGURE 11. Left: M (black), $M^{\mathrm{crit}}_{\mathrm{RVM}}$ (red) and M_h (green) at every integer $n \le 2000$. Right: range of M and mean of $M^{\mathrm{crit}}_{\mathrm{RVM}}$ in blocks of 500 integers up to 10^6, with $\pm\sqrt{n}$.




FIGURE 12. Left: $\mu(n)$ and $\mu_h(n)$ for $n \le 60$: same zeros, different signs. Right: M, the target $M^{\mathrm{crit}}_{\mathrm{RVM}}$ and M_h for $n \le 300$.

Figures 11–13 summarize the Mertens side. $M^{\mathrm{crit}}_{\mathrm{RVM}}$ has the size and the visual texture of M, it stays inside $\pm\sqrt{n}$ as Theorem 8.12 requires, and its residues are deterministic; M_h is its ternary integer version with the zero set of μ. The passage from these synthetic objects to M itself is not numerical: it is the identity $\zeta_{Z1} = \zeta$ of Step 3, which Experiment 17.2 is consistent with.

Remark 17.9 (The opposite phase of M and $M^{\mathrm{crit}}_{\mathrm{RVM}}$ between 10^3 and $2\cdot 10^3$). In the left panel of Figure 11 the curves of M and $M^{\mathrm{crit}}_{\mathrm{RVM}}$ are in opposite phase between $n \approx 10^3$ and $n \approx 2\cdot 10^3$. Both functions are dominated there by their first mode, $2\sqrt{x}\,\Re(x^{i\gamma_1}/(\rho_1\zeta'(\rho_1)))$ and $2\sqrt{x}\,\Re(x^{i\tau_1}/(\rho_1^{\mathrm{RVM}}\zeta'_{\mathrm{RVM}}(\rho_1^{\mathrm{RVM}})))$, two oscillations in $\log x$ with the frequencies γ_1 and τ_1. Their phase difference is $\eta_1 \log x$ with $\eta_1 = \tau_1 - \gamma_1 = 14.5213 - 14.1347 = 0.3866$, up to the constant difference of the arguments of the two residues, which is not computed here; $\eta_1 \log x$ equals π at $x = e^{\pi/\eta_1} \approx 3.4\cdot 10^3$ and lies between 2.67 and 2.94 on $[10^3, 2\cdot 10^3]$. The beat has period $2\pi/\eta_1 \approx 16.3$ in $\log x$. This is the rotation (14.42) seen on the Mertens side: the passage from the clock to ζ turns the mode k by the angle $\eta_k \log x$. The higher modes turn at different rates, and at large scale neither




FIGURE 13. Left: $M(x)$, its explicit formula from the zeros of ζ and $M^{\mathrm{crit}}_{\mathrm{RVM}}(x)$ from the τ_k on $[2, 300]$. Right: normalized derivatives at the zeros: erratic for ζ, constant ϖ_{RVM} for ζ_{RVM}.

a correlation nor an anticorrelation remains, in agreement with the value 0.0007 of Experiment 17.8. The effect depends on η_1 alone and changes sign with x; it does not distinguish $\lambda = 0$.

17.7. **The rung equations: transport, the defect of the clock, and the solution from the clock.** The three experiments of this subsection evaluate the objects of Subsections 14.13 and 15.2. In the first two, f is the function $\exp(-1/(1-y^2))$, $y = (u - 0.36)/0.26$, on $[0.10, 0.62] \subset (0, \log 2)$, normalized to $\int \mathsf{f} = 1$; it tests the first rung. In the programs f, $\mathsf{k_f}$ and $\mathsf{c_f}$ are called `psi`, `W_psi` and `c_psi`.

Experiment 17.10 (Transport on the first rung)**.** *Object:* $\mathsf{R_f}(1)$, $\mathsf{S_f}(\lambda)$ and the moments $\mathsf{m}_j(\mathsf{f})$ of (14.78)–(14.82). *Range:* the first 20 000 ordinates of each list; $\max_{k\ge1000} |\mathsf{c_f}(\tau_k)| = 2.5 \cdot 10^{-10}$ and $\max_{k\ge3000} |\mathsf{c_f}(\tau_k)| = 2.1 \cdot 10^{-13}$. *Expected:* if $\zeta_{Z1} = \zeta$, then $\mathsf{S_f}(0) = \mathsf{R_f}(1)$ (Proposition 14.41); the series (14.82) converges to $\mathsf{S_f}(0)$ with terms bounded by $C(1+j)^2(C_\eta \log 2)^j/j!$ (Theorem 14.42). *Observed:*

- $\mathsf{k_f} = 0.133643530916$, $\sum_k \mathsf{c_f}(\tau_k) = 0.176106813076$, $\sum_k \mathsf{c_f}(\gamma_k) = 0.133643530929$;
- defect of the clock $\mathsf{R_f}(1) = -0.042463282161$; transport $\mathsf{S_f}(0) = -0.042463282147$; $\mathsf{R_f}(0) = \mathsf{R_f}(1) - \mathsf{S_f}(0) = -1.4 \cdot 10^{-11}$;
- moment series (Table 6): the term $j = 1$ is 95.1% of $\mathsf{R_f}(1)$, the partial sum with $j \le 2$ is 100.8%, and the partial sums with $j \le 3$, $j \le 5$, $j \le 8$ differ from $\mathsf{R_f}(1)$ by $3.7 \cdot 10^{-6}$, $6.5 \cdot 10^{-9}$, $1.0 \cdot 10^{-11}$;
- along the family (Table 6) $\mathsf{R_f}(\lambda)$ is monotone and almost linear on $[-1, 2]$, with slope about -0.044 and a single zero, at $\lambda = 0$;
- the partial sums $\sum_{k\le K_0}(\mathsf{c_f}(\gamma_k) - \mathsf{c_f}(\tau_k))$ are -0.02086, -0.04381, -0.04649, -0.04221, -0.04249, -0.0424634 for $K_0 = 1, 5, 10, 50, 100, 400$: the first ordinate gives half of the transport, and the first rung is decided by about fifty ordinates.

Experiment 17.11 (The defect of the clock by Poisson summation)**.** *Object:* the two parts of $\mathsf{R_f}(1)$ in Proposition 14.43. *Expected:* $\mathsf{R_f}(1) = (\mathsf{k_f} - \mathsf{l}_0) - 2\sum_{m\ge1}(-1)^m \mathsf{l}_m$. *Observed:* $\mathsf{t}_0 = 9.676906787$; $\mathsf{l}_0 = 0.063533498702$ from (14.84); the mismatch of the smooth densities is $\mathsf{k_f} - \mathsf{l}_0 = +0.070110032214$, and the oscillating part, obtained as the difference $\mathsf{R_f}(1) - (\mathsf{k_f} - \mathsf{l}_0)$ and not by summing the l_m, is -0.112573314375. The stationary heights $2\pi e^u$, u in the support of f, fill $[6.944, 11.680]$; only those with $u > 0.4319$ lie above t_0.

Experiment 17.12 (From the clock to the ordinates and to the masses of the primes)**.** *Object:* a finite section of the system (15.5): the unknowns are the ordinates κ_k below a height and the masses a_n, $2 \le n \le 11$; the equations are the pairings of (15.5) with $w(u)\cos(\omega u)$ and $w(u)\sin(\omega u)$,

j	$\mathsf{m}_j(\mathsf{f})$	$(-1)^j\mathsf{m}_j(\mathsf{f})/j!$	partial sum $-\ \mathsf{R}_\mathsf{f}(1)$	λ	$\mathsf{R}_\mathsf{f}(\lambda)$
1	$+4.0395\cdot10^{-2}$	$-4.0395\cdot10^{-2}$	$+2.07\cdot10^{-3}$	-1.00	$+0.045303554710$
2	$-4.8098\cdot10^{-3}$	$-2.4049\cdot10^{-3}$	$-3.37\cdot10^{-4}$	-0.50	$+0.022413444752$
3	$-2.0432\cdot10^{-3}$	$+3.4053\cdot10^{-4}$	$+3.69\cdot10^{-6}$	-0.25	$+0.011133149816$
4	$-5.5184\cdot10^{-5}$	$-2.2993\cdot10^{-6}$	$+1.39\cdot10^{-6}$	0.00	-0.000000000014
5	$+1.6734\cdot10^{-4}$	$-1.3945\cdot10^{-6}$	$-6.47\cdot10^{-9}$	$+0.25$	-0.010956709511
6	$+5.5721\cdot10^{-6}$	$+7.7390\cdot10^{-9}$	$+1.27\cdot10^{-9}$	$+0.50$	-0.021706821780
7	$+7.2608\cdot10^{-6}$	$-1.4406\cdot10^{-9}$	$-1.69\cdot10^{-10}$	$+1.00$	-0.042463282161
8	$+6.4039\cdot10^{-6}$	$+1.5883\cdot10^{-10}$	$-9.77\cdot10^{-12}$	$+2.00$	-0.080112190380

TABLE 6. Experiment 17.10. Left: the moments $\mathsf{m}_j(\mathsf{f})=\sum_k \eta_k^j \mathsf{c}_\mathsf{f}^{(j)}(\tau_k)$ and the partial sums of the series (14.82) for $\mathsf{S}_\mathsf{f}(0)$, compared with the defect $\mathsf{R}_\mathsf{f}(1)=-0.042463282161$ of the clock. Right: the rung functional along the family.

where w is a fixed smooth function supported in $(0,\log 13.5)$ and ω runs over a grid of step 0.1 in $[0,\Omega]$. The ordinates above $\Omega+12$ are kept on the clock. *Method:* Levenberg–Marquardt iteration started at $\kappa_k=\tau_k$, the masses being eliminated linearly at each step, with continuation in height $\Omega=18,22,\dots,50$. The file `zeross.bin` is read only after the iteration, for the comparison. *Expected:* by Theorem 15.5, if $\zeta_{Z1}=\zeta$ the system has the unique solution $\kappa_k=\gamma_k$, $\mathsf{a}_n=\log p/(2\sqrt{n})$ at prime powers and 0 elsewhere. *Observed:* the iteration converges at every stage (residual norms between $4\cdot10^{-4}$ and $5\cdot10^{-3}$, of the size of the contribution of the ordinates kept on the clock); the solution is in Table 7: the first ten ordinates agree with $\gamma_1,\dots,\gamma_{10}$ within $4.4\cdot10^{-4}$, the eleventh within $2.2\cdot10^{-3}$, and the masses agree with (15.6) within $8\cdot10^{-4}$, including the values 0 at $n=6,10$ and the prime-power values at $n=4,8,9$. The ordinates $k=12,13,14$, which lie above the band $\omega\le50$, are not determined (errors 0.10, 0.77, 1.73). With masses up to $n\le34$ and the band $\omega\le60$ the section is not resolved and the iteration does not converge: resolving the ordinates up to a height Ω requires $\log N\ge\log\frac{\Omega}{2\pi}$ and resolving the masses up to N requires Ω of the order of πN, in agreement with the heights $2\pi x^{1/m}$ of Proposition 14.43. The experiment illustrates the existence form of the final step; it is a computation and not a proof of existence.

k	τ_k (start)	solution	γ_k	n	a_n (solution)	(15.6)
1	14.521346953	14.134660056	14.134725142	2	$+0.245076213$	0.245064536
2	20.655740356	21.022091631	21.022039639	3	$+0.317145240$	0.317142050
3	25.492675432	25.010827129	25.010857580	4	$+0.173268937$	0.173286795
4	29.739411632	30.424917668	30.424876126	5	$+0.359884977$	0.359881258
5	33.624531889	32.935072953	32.935061588	6	-0.000004422	0
6	37.257370087	37.586146548	37.586178159	7	$+0.367785301$	0.367742452
7	40.700551604	40.918919600	40.918719012	8	$+0.122615622$	0.122532268
8	43.994013953	43.327322117	43.327073281	9	$+0.183143344$	0.183102048
9	47.165128427	48.004716753	48.005150881	10	-0.000286111	0
10	50.233653391	49.773657726	49.773832478	11	$+0.362276961$	0.361496314
11	53.214422874	52.972535790	52.970321478			

TABLE 7. Experiment 17.12. Solution of a finite section of the rung system (15.5), started at the clock. Left: the ordinates. Right: the masses at the points $\log n$, which are not assumed, against $\log p/(2\sqrt{n})$ at prime powers $n=p^m$.

Experiment 17.13 (Multiscale Jacobian of the clock)**.** *Object:* the minimum midpoint-clock gap, its bound (14.86), the Ingham critical window $2\pi/\Delta_T$, and the adjacent-pair bound (14.88). *Input:*

none of the genuine zeta ordinates; the program `s12_multiscale_jacobian.py` generates the τ_k directly from the Lambert formula up to $K = 2\,001\,052$. *Observed:*

K	$T = \tau_K$	Δ_T	$2\pi/\log(T/2\pi)$
10^2	235.987267	1.73462445	1.73286190
10^3	1419.517765	1.15930485	1.15921748
10^4	9877.629616	0.85368118	0.85367617
10^5	74920.891033	0.66939912	0.66939880
$5 \cdot 10^5$	319387.322632	0.57982852	0.57982848
$2\,001\,052$	1132490.722588	0.51918344	0.51918343

At the last height, $2\pi/\Delta_T = 12.102053$ while $\log(T/2\pi) = 12.10205288$. Thus the rigorous scale $U \asymp \log T$ is numerically almost exact for the midpoint clock. On every fixed window the adjacent-pair upper bound decreases with the clock gap, illustrating the obstruction in Theorem 14.44(iii). These calculations use no γ_k and no η_k.




FIGURE 14. Minimum gap of the midpoint clock up to height T and the explicit lower bound $2\pi/\log(T/2\pi)$. Their ratio is already $1 + 2 \cdot 10^{-8}$ at $K = 2\,001\,052$.




FIGURE 15. Upper bound in (14.88) for adjacent clock frequencies on four fixed u-windows. The decay is slow, of order $1/\log \tau_k$, but it is enough to exclude a height-uniform lower frame bound on any fixed window.

18. Conclusions and open problems

The following statements are proved:

(1) the midpoint ordinates τ_k are given by the Lambert formula (2.5) and stay within a bounded distance of the ordinates γ_k of ζ (Lemma 2.2);
(2) $P^{\mathrm{crit}}_{\mathrm{RVM}}$ is entire and symmetric, and ζ_{RVM} has exactly the simple zeros $\frac{1}{2} \pm i\tau_k$, the trivial zeros and a residue-one pole at 1 (Theorem 3.1);
(3) $S_{\mathrm{RVM}}(N;u) \ll 1 + \log N$ uniformly on compact positive frequency intervals, so the point centroid and $M_H^{\mathrm{RVM}}(x)$ tend to 0 (Section 4);
(4) J_{RVM} exists compact-uniformly and is related to π_{RVM} by exact finite Möbius inversion (Section 5);
(5) $J_{\mathrm{RVM}}(x) - \mathrm{li}(x)$ and $\pi_{\mathrm{RVM}}(x) - \mathrm{li}(x)$ are $O(\sqrt{x}(\log\log x)^2/\log x)$, so $\mathrm{li}/\pi_{\mathrm{RVM}} \to 1$, and with the prime number theorem $\pi/\mathrm{li} \to 1$ and $\pi/\pi_{\mathrm{RVM}} \to 1$ (Theorem 5.5, Corollary 5.6);
(6) J_{RVM} is Lipschitz on $[2,\infty)$, $c_n^{\mathrm{RVM}} = O(1)$, and $\zeta_{\mathbb{Z},\mathrm{RVM}}$ is an absolutely convergent Dirichlet series in $\Re s > 1$ (Section 6);
(7) $\zeta_{\mathrm{RVM}} = \varpi_{\mathrm{RVM}}(1+w)(1+O(1/|t|))$ on every band to the right of $\frac{1}{2}$ and $\zeta_{\mathrm{RVM}} = \varpi_{\mathrm{RVM}}(1+\chi)(1+O(1/|t|))$ on closed substrips of $\frac{1}{2} < \Re s < 1$ (Theorem 8.1);
(8) $1/\zeta_{\mathrm{RVM}} \to L$ on every vertical line to the right of $\frac{1}{2}$, $\zeta_{\mathrm{RVM}}^{\pm 1}$ have no absolutely convergent Dirichlet expansion, and $M^{\mathrm{crit}}_{\mathrm{RVM}}(x) = L + O_\varepsilon(x^{1/2+\varepsilon})$ (Section 8);
(9) 2ξ equals its Hadamard product over $\rho(1-\rho)$ exactly (Theorem 9.1), and $|\zeta| \asymp |\mathcal{Q}_{\mathrm{RVM}}|$ on $\frac{1}{2} + \varepsilon \le \sigma \le c'$, $|t| \ge t_0$ (Corollary 9.3);
(10) the λ-family from ζ_{RVM} to ζ_{Z1} converges jointly, has simple zeros for $\lambda > 0$, an explicit generator and total multiplier, and its members differ only through $\gamma_k - \tau_k$ (Section 10);
(11) ζ_{Z1} has the functional equation, the pole, the residue and the order of ζ, and $\zeta_{Z1} = \zeta \Leftrightarrow$(D)$\Leftrightarrow$RH (Theorem 11.2);
(12) ζ_{RVM} is bounded above and below on every half-plane $\Re s \ge \sigma_0 > 1$ and tends to ϖ_{RVM} as $\Re s \to +\infty$ (Theorem 8.10); $1/\zeta_{\mathbb{Z},\mathrm{RVM}}$ is an absolutely convergent Dirichlet series in $\Re s > 1$, so (D) is equivalent to an absolutely convergent Dirichlet expansion of $\mathcal{K}$ in $\Re s > 1$ (Theorem 11.3);
(13) no member of the λ-family, ζ_{Z1} included, can have a zero off the critical line other than the trivial zeros (Theorem 10.1);
(14) the endpoint has the value $\varpi_{Z1} = \varpi_{\mathrm{RVM}} \prod_k (\frac{1}{4} + \tau_k^2)/(\frac{1}{4} + \gamma_k^2) \ge 1$ at $+\infty$, and $\zeta_{Z1} = \zeta$ if and only if $\varpi_{Z1} = 1$ (Theorems 12.1, 12.2, Corollary 12.3);
(15) the prime-power counts of ζ_{RVM} and ζ_{Z1} are the Mellin transforms of their logarithms and satisfy a distributional explicit formula; this proves the Mellin–Stieltjes identification and the weighted factorization $\zeta_{\mathbb{Z},\mathrm{RVM}} = \zeta_{\mathrm{RVM}} e^{h_{\mathrm{RVM}}}$, and (D) holds if and only if the prime-power count of ζ_{Z1} is constant between consecutive integers (Theorems 12.5, 12.6, 6.4, 12.7);
(16) along the family $\frac{d}{d\lambda}\log\varpi_\lambda = -\Upsilon(\lambda)$, and (D) is the identity $\int_{0^+}^1 \Upsilon\, d\lambda = \log L$; with the verified zeros, $0 \le \log\varpi_{Z1} < 4 \cdot 10^{-13}$ (Propositions 13.2, 13.3);
(17) the atomized family $\tilde{\zeta}_{\mathbb{Z},\lambda} = \zeta_\lambda e^{\tilde{h}_\lambda}$ consists of absolutely convergent Dirichlet series in $\Re s > 1$ for every λ, the endpoint included, because the cell masses have mean square $X^{1+\varepsilon}$ on dyadic blocks uniformly in λ; the joining factor is explicit cell by cell and holomorphic in $\Re s > \frac{1}{2}$ (in $\Re s > 0$ for $\lambda = 1$), and (D) is equivalent to the vanishing of every cell defect at $\lambda = 0$ and of J_{Z1} on the first cell $(1,2)$ (Proposition 13.5, Theorems 13.6, 13.10, 13.12, 13.13);
(18) at the clock endpoint the joining factor is not constant, $\tilde{h}_1(\sigma) \to \log L$; the atomized reciprocals have Möbius coefficients $\tilde{\mu}_\lambda$, and the endpoint Mertens function has a Mellin form B_{Z1} with $B_{Z1} = M$ if and only if (D) (Propositions 13.15, 13.16);
(19) Step 3: $\zeta_{Z1} = \zeta$ implies $M(x) = O_\varepsilon(x^{1/2+\varepsilon})$ (Theorem 11.8); equivalently, the synthetic bound $M^{\mathrm{crit}}_{\mathrm{RVM}}(x) = L + O_\varepsilon(x^{1/2+\varepsilon})$ transfers to M through $1/\zeta = F_{\mathrm{RVM}}\Pi_{\mathrm{tr}}^{-1}$ (Theorem 11.9);

(20) for $0 < \lambda \le 1$ no ζ_λ is an absolutely convergent Dirichlet series in $\Re s > 1$ and no joining factor $\tilde h_\lambda$ is constant, and $\lambda \mapsto \log \zeta_\lambda(\sigma)$ is strictly convex for every real $\sigma > 1$ (Propositions 14.1, 14.2);

(21) if the joining factor $e^{\tilde h_\lambda}$ is meromorphic of finite order with finitely many poles and symmetric under $s \mapsto 1 - s$, Hamburger's theorem gives $\tilde\zeta_{\mathbb{Z},\lambda} = \zeta$, and for $\lambda = 0$ also $\zeta_{Z1} = \zeta$ (Theorem 14.3);

(22) if ζ_λ is the Mellin–Stieltjes transform of a positive measure, then $\varpi_\lambda \le 1$; (D) holds if and only if ζ_{Z1} has such a representation, if and only if $J^{\mathrm M}_{Z1}$ is non-decreasing (Theorem 14.7, Corollary 14.8);

(23) J_{RVM} is continuously differentiable on $(2,\infty)$ with $J'_{\mathrm{RVM}}(y) = \frac{1}{\log y} + 2\cos(2\pi y) - 2^{-1/2} y^{-1/4}\cos(4\pi\sqrt{y} + \frac{\pi}{4}) + O(y^{-1/3})$; consequently $\tilde c_n(1) = \frac{1}{\log n} + O(n^{-1/4})$ off the powers of 2, and Hypothesis 6.7 holds off the integers 2^j with $\mu(j) \neq 0$ (Theorem 14.15, Corollary 14.16);

(24) the cell masses are sums over the ordinates with the height kernel $\hat\varphi(\gamma/2\pi n)$, which vanishes at the heights $2\pi q n$; their flow is the pairing of V_n with the count difference $N_\zeta - n_{\mathrm{RVM}}$; and the Gaussian defect of the atomization is $\frac12(e^{\tilde h_\lambda(1)} - 1) y^{-1/2} + O(y^{-1/4-\delta})$ (Propositions 14.17, 14.18, 14.22);

(25) $\zeta_{Z1}(\sigma) \ge \zeta(\sigma)$ for every real $\sigma > 1$, with equality at one point if and only if $\zeta_{Z1} = \zeta$; in particular $\prod_k (1 + 2/(\frac14 + \gamma_k^2)) \ge \pi/3$ (Theorem 14.11);

(26) every ζ_λ is the Mellin transform of a function A_λ that satisfies the summation formula of ζ, and $\varpi_\lambda - 1 = -\mathsf{W}(\lambda)/\hat{\mathsf{g}}(1)$ for one explicit number $\mathsf{W}(\lambda)$; (D) is equivalent to $\mathsf{W}(0) \ge 0$ and to $\varpi_\lambda < 1$ for all $\lambda \in (0,1]$ (Lemma 14.9, Propositions 14.10, 14.12);

(27) Hamburger's theorem for the pair holds without continuation of the joining factor: $\tilde\zeta_{\mathbb{Z},\lambda} = \zeta$ if and only if the atomized coefficients satisfy the theta relation, if and only if the vertical mean square of $\tilde\zeta_{\mathbb{Z},\lambda}/\zeta$ on a line $\Re s = \sigma > 1$ is 1; both defects are continuous in λ (Theorem 14.4);

(28) at the clock end the joining factor has no symmetric meromorphic continuation, and $\tilde\zeta_{\mathbb{Z},\lambda} = \zeta$ for at most finitely many $\lambda \in [0,1]$, not for $\lambda = 1$ (Propositions 14.5, 14.25);

(29) the cell masses are entire functions of λ of exponential type $O(\log n)$, and the cell defects have the height kernel $\hat\varphi'(\gamma/2\pi n)$ (Propositions 14.24, 14.19);

(30) in the compact space $\mathscr{O}$ of ordinate lists at bounded distance from the clock, at most one list has coinciding continuous and atomized zeta functions; such a list exists if and only if $\zeta_{Z1} = \zeta$, and it is then the list of the ordinates of ζ (Theorem 15.1, Proposition 15.2, Corollary 15.3);

(31) (D) holds if and only if $\tilde c_n(0) = \mathbf{1}_{\mathrm{prime}}(n)$ for every $n \ge 2$, one real number for each cell; every λ with $\tilde\zeta_{\mathbb{Z},\lambda} = \zeta$ is a witness of (D), so the exceptional set is empty or contains 0; cell by cell the remaining identity is (14.59) (Corollary 14.26, Propositions 14.27, 14.29);

(32) for a pair joined by a symmetric factor Hamburger's theorem gives a constant factor when the continuous function has order one, and not in general; at $\lambda = 0$ the symmetry of the joining factor is equivalent to (D) (Proposition 14.31, Remarks 14.30, 14.32);

(33) the ladder: for every $N \ge 2$, $\tilde h_\lambda(\sigma) = O(\sigma N^{-\sigma})$ as $\sigma \to +\infty$ if and only if the Mellin count of ζ_λ has no mass outside the integers below N; (D) holds if and only if this holds at $\lambda = 0$ for every N, and at the clock end the first rung fails (Theorems 14.33, 14.34, Corollary 14.35);

(34) the bridge theorem: if the imaginary part of the joining factor of a pair extends continuously to $\Re s \ge \frac12$, vanishes on the critical line, and the factor satisfies the temperate bound (14.67), then the factor is constant, the discrete function is ζ and the continuous one is a multiple of ζ; for the family the two conditions hold for no $\lambda \in (0,1]$, and at $\lambda = 0$ they are equivalent to (D) (Theorem 14.36, Corollary 14.37);

(35) on the critical line $\Im \log \zeta_\lambda = -\vartheta(t) + \pi(N_\lambda(t) - 1)$, and the cell masses of two lists differ by a sum over the ordinates of one function of one ordinate (Propositions 14.38, 14.39);

(36) on each rung the passage from the clock to the ordinates of ζ is the transport $\mathsf{S}_{\mathsf{f}}(\lambda)$, an entire function of λ of exponential type at most $C_\eta \log N$ whose value at $\lambda = 0$ is the sum of the series of the moments $\sum_k \eta_k^j \mathsf{c}_{\mathsf{f}}^{(j)}(\tau_k)$; (D) holds if and only if the transport equals the defect of the clock on every rung, and this defect is given exactly by Poisson summation, with stationary heights $2\pi x^{1/m}$ (Proposition 14.41, Theorem 14.42, Proposition 14.43);

(37) (D) holds if and only if the rung system (15.5), with free masses at the points $\log n$, has a solution in $\mathscr{O}$; the solution is then the list of ζ with the masses of von Mangoldt, and its linearization is injective up to the splitting of multiple ordinates (Theorems 15.5, 15.6);

(38) at the midpoint clock, the finite ordinate Jacobian up to height T has a uniform lower frame bound on windows of length $(1+\eta)\log(T/2\pi)$, no uniform lower bound is possible on a fixed window, the infinite ordinate block has dense range on every finite window, and smooth bounded-displacement transport tails are $O_{A,\mathsf{f}}(T^{1-A}\log T)$ (Theorem 14.44);

(39) the gradient of the vertical defect on $\mathscr{O}$ is $\partial \mathsf{V}_\sigma / \partial \kappa_k = 2\mathsf{Q}_{\boldsymbol{\kappa}}(\kappa_k)$ with one real-analytic function $\mathsf{Q}_{\boldsymbol{\kappa}}$ (Proposition 15.8);

(40) a list is accepted by a continuous side and a discrete side if and only if its function lies in the intersection of the window of the first with the frequencies of the second; a non-periodic list is accepted through one unitary matrix; for the finite Euler products $\prod_{p \le X}(1-p^{-s})$ the constructions of the paper apply and the endpoint $\lambda = 0$ is the only member of the family that is a Dirichlet series (Theorem 16.1, Proposition 16.2, Theorem 16.3);

(41) there is no function with the functional equation of ζ and an absolutely convergent Dirichlet series supported on the integers with bounded prime factors, and for the diagonal operator over the primes the contraction, the convergence of the determinant and the symmetry hold in $\sigma > 0$, in $\sigma > 1$ and on $\sigma = \frac{1}{2}$ respectively (Proposition 16.4, Subsection 16.4).

Not proved. The identity $\zeta_{Z1} = \zeta$ is not proved. The statements proved to be equivalent to it are listed in Subsection 1.4; in the form of Corollary 15.3 it is the existence of one list in $\mathscr{O}$ whose continuous and atomized zeta functions coincide, and in the form of Corollary 14.37 it is the vanishing on the critical line of the imaginary part of the joining factor $\tilde{h}_0$, with the bound (14.67). The theorems of Subsections 14.9–14.13 and of Sections 15 and 16 are formulations of this step and tools for it; none of them evaluates the pairing (14.59) of the count difference $N_\zeta - n_{\mathrm{RVM}}$, on which every one of them depends at $\lambda = 0$.

The computations of Section 17 with 2 001 052 zeros are consistent with each of these statements.

Remark 18.1 (Open problems).

(i) Prove the identity $\zeta_{Z1} = \zeta$, equivalently condition (D), equivalently an absolutely convergent Dirichlet expansion of $\mathcal{K} = \zeta_{Z1}/\zeta_{\mathbb{Z},\mathrm{RVM}}$ in $\Re s > 1$, equivalently the product identity (12.12), equivalently the cell identity (12.18) with the jump bound of Theorem 12.7, equivalently step A7 of the programme in Subsection 13.8, equivalently the positivity of the measure representing ζ_{Z1} (Corollary 14.8), equivalently the existence of a coincidence list in $\mathscr{O}$ (Corollary 15.3, step A19). This is the final step of the method.

(ii) Hypothesis 6.7 is proved for the integers n that are not of the form 2^j with $\mu(j) \neq 0$ (Corollary 14.16); at $n = 2^j$ the cell mass is $(1 + \mu(j) J_{\mathrm{RVM}}(2)\log 2)/\log n + O(n^{-1/4})$. Determine the sign of $1 \pm J_{\mathrm{RVM}}(2)\log 2$.

(iii) Decide, for $0 < \lambda < 1$ outside the finite set E, which hypothesis of Theorem 14.3 fails, and which of (B1), (B2) fails in Theorem 14.36; prove that every critical point of V_σ on $\mathscr{O}$, the boundary included, is a zero of V_σ (Proposition 15.8); and decide whether the atomized coefficients $\tilde{a}_N(0)$ control the sign of $\mathsf{W}(0)$ in Proposition 14.10.

(iv) Determine the monotone branches of π_{RVM}, on which the real prime levels p_j^{RVM} are defined, and explain the observed proximity $p_j^{\mathrm{RVM}} \approx p_j$ for small j (Experiment 17.6).

(v) Derive the bound (B2) for $\tilde{h}_\lambda$ from the bounds of the cell defects, and turn Heuristic 14.40 into a lemma with remainder; decide (B1) at $\lambda = 0$.

(vi) Evaluate from the construction the right side of (14.59), or the moments $\mathsf{m}_j(\mathsf{f})$ of (14.82); on the first rung the moments $j = 1, 2$ carry the transport up to $4 \cdot 10^{-6}$ (Experiment 17.10).

(vii) Construct a quantitative multiscale continuation from the clock to a solution of the rung system. Theorem 14.44 proves the finite-height lower frame bound on the natural scale $U \asymp \log T$, the failure of fixed-window uniform invertibility, dense range on every fixed window, and rapidly summable smooth tails. What remains is to bound the residual (14.90) and a compatible finite-section right inverse strongly enough to dominate the quadratic remainder along a nested sequence $T \to \infty$ (Experiment 17.12 follows such a path numerically).

(viii) Under (D), decide whether $\mathsf{E} = \{0\}$, that is, whether the functions D_n of (14.57) have a common zero in $(0, 1]$.

(ix) Construct an object of infinite rank from which ζ_{Z1} and $\tilde{\zeta}_{\mathbb{Z},0}$ both descend, as the matrix U of Proposition 16.2 does for the two-edge model (Subsection 16.4).

Remark 18.2 (Statements derived in part or conjectural, not proved here). The following statements arose in the work on the final step; none of them is used in this paper.

(a) *Rates in λ.* Off the powers of 2 the formulas of Proposition 14.24 suggest

$$\tilde{h}_\lambda(s) - \tilde{h}_0(s) = O_s(\lambda), \qquad \bigl|\mathfrak{h}_n(\lambda; s) - \mathfrak{h}_n(0; s)\bigr| \ll_{s,\varepsilon} \lambda\, n^{-\sigma-\frac{1}{2}+\varepsilon}; \tag{18.1}$$

the cells $n = 2^j$ and the uniformity in n of the first-cell term are not written.

(b) *Flow form of the density condition.* $\tilde{h}_0(1) = 0$ if and only if $\int_{0^+}^1 \partial_\lambda \tilde{h}_\lambda(1)\, d\lambda = \tilde{h}_1(1)$; the differentiability at $s = 1$ uniformly in λ is not written.

(c) *Pointwise bound.* $|\tilde{c}_n(\lambda)| \le C_\varepsilon n^\varepsilon$ uniformly for λ in a neighbourhood of 0 is open; by Corollary 14.26 it is not needed for the final step.

(d) *Leading term of the clock defect.* The form $\mathfrak{h}_n(1; s) \approx \sqrt{2}\,\pi\,\mathsf{M}_2\, s\, n^{-s-7/4} \sin(4\pi\sqrt{n} + \frac{\pi}{4})$ is not proved; the bound (14.40) is.

(e) *A heuristic model of the atomized clock.* By (14.39), $\tilde{\zeta}_{\mathbb{Z},1}(s)$ is close to $\prod_{n\ge2}(1 - n^{-s})^{-1/\log n}$, whose logarithm has the leading part $\sum_n n^{-s}/\log n = \int_s^\infty (\zeta(y) - 1)\, dy$; no error term is claimed.

(f) The comparison between the two columns of Subsection 13.7 and the use of the classification of [28] remain as in Remark 18.1(iii) and are not proved.

Appendix A. Python programs

The programs below produce every number, table and figure of Section 17. They read the two data files `data/zeross.bin` (ordinates γ_k of the zeros of ζ) and `data/zLamb.bin` (midpoint ordinates τ_k), both little-endian float64 with 2 001 052 entries, and write their console output, CSV, MAT and binary files to `output/` and the figures to `figures/`. They require Python 3 with NumPy, SciPy, mpmath and Matplotlib, and are run from the directory `code/` in the order `s01`–`s10`; the programs `s08`–`s10` use NumPy only and read the same two data files.

A.1. `rvm_common.py`.

```
"""Common routines for the midpoint Riemann-von Mangoldt (RVM) zeta system.

Data files (little-endian float64, one ordinate per entry):
    data/zeross.bin  ordinates gamma_k of the nontrivial zeros of zeta (2,001,052 values)
    data/zLamb.bin   midpoint RVM ordinates tau_k, N_RVM(tau_k) = k - 1/2

Notation follows the paper: N_RVM is the clock, tau_k the midpoint ordinates,
gamma_k the ordinates of zeta, eta_k = tau_k - gamma_k, q(s) = s(s-1),
G(s) = s(s-1) Gamma(s/2) pi^(-s/2), zeta_lambda = P_lambda / G with ordinates
gamma_k(lambda) = (1-lambda) gamma_k + lambda tau_k.
"""
import os
import sys
```

```
import time
import numpy as np
from scipy import special

ROOT = os.path.dirname(os.path.dirname(os.path.abspath(__file__)))
DATA = os.path.join(ROOT, "data")
OUT = os.path.join(ROOT, "output")
FIG = os.path.join(ROOT, "figures")
os.makedirs(OUT, exist_ok=True)
os.makedirs(FIG, exist_ok=True)

TWO_PI = 2.0 * np.pi


class Tee:
    """Write console output both to the screen and to output/<name>_console.txt."""

    def __init__(self, name):
        self.f = open(os.path.join(OUT, name + "_console.txt"), "w")
        self.t0 = time.time()

    def __call__(self, fmt, *args):
        s = fmt % args if args else fmt
        sys.stdout.write(s + "\n")
        sys.stdout.flush()
        self.f.write(s + "\n")
        self.f.flush()

    def close(self):
        self("total run time %.1f s" % (time.time() - self.t0))
        self.f.close()


def read_bin(name):
    return np.fromfile(os.path.join(DATA, name), dtype="<f8")


def load_ordinates():
    gam = read_bin("zeross.bin")
    tau = read_bin("zLamb.bin")
    n = min(gam.size, tau.size)
    return gam[:n], tau[:n]


def clock(T):
    """N_RVM(T) = T/(2 pi) log(T/(2 pi e)) + 7/8 (real or complex T, principal log)."""
    return T / TWO_PI * (np.log(T / TWO_PI) - 1.0) + 0.875


def tail_coefficients(U):
    """Midpoint-rule tail of sum_{k>N} log(1 + q/tau_k^2), N = N_RVM(U):
    returns (c1, c2, c3) with tail(q) = c1 q - c2 q^2 + c3 q^3."""
    LU = np.log(U / TWO_PI)
    c1 = (LU + 1.0) / U / TWO_PI
    c2 = (3.0 * LU + 1.0) / (9.0 * U ** 3) / 2.0 / TWO_PI
    c3 = (5.0 * LU + 1.0) / (25.0 * U ** 5) / 3.0 / TWO_PI
    return c1, c2, c3


def clock_inverse(y):
    """Solve N_RVM(U) = y by Newton iteration."""
    U = TWO_PI * max(y, 2.0) / np.log(max(y, 3.0))
    for _ in range(80):
        U -= (clock(U) - y) / (np.log(U / TWO_PI) / TWO_PI)
    return U
```

```
class Product:
    """log P(s) = sum_k log(1 + q(s)/(1/4 + o_k^2)) for an ordinate list o_k.

    Exact summation for k <= K1, power sums for K1 < k <= N, and the clock tail
    for k > N.  Accurate for |q(s)| << o_{K1}^2."""

    def __init__(self, ords, K1=100000, nterms=6):
        self.o = np.asarray(ords, dtype=float)
        self.N = self.o.size
        self.K1 = min(K1, self.N)
        A = 0.25 + self.o[self.K1:] ** 2
        self.pw = [np.sum(A ** (-j)) for j in range(1, nterms + 1)]
        self.A1 = 0.25 + self.o[: self.K1] ** 2
        self.tail = tail_coefficients(clock_inverse(float(self.N)))

    def logP(self, s):
        s = np.atleast_1d(np.asarray(s, dtype=complex))
        q = s * (s - 1.0)
        out = np.empty(s.shape, dtype=complex)
        for i, qi in enumerate(q):
            v = np.sum(np.log1p(qi / self.A1))
            for j, p in enumerate(self.pw, start=1):
                v += (-1) ** (j + 1) * qi ** j / j * p
            c1, c2, c3 = self.tail
            v += c1 * qi - c2 * qi ** 2 + c3 * qi ** 3
            out[i] = v
        return out


def logG(s):
    s = np.asarray(s, dtype=complex)
    return np.log(s) + np.log(s - 1.0) + special.loggamma(s / 2.0) - s / 2.0 * np.log(np.pi)


def zeta_from_product(prod, s):
    s = np.atleast_1d(np.asarray(s, dtype=complex))
    return np.exp(prod.logP(s) - logG(s))


def chi(s):
    s = np.asarray(s, dtype=complex)
    return np.exp((s - 0.5) * np.log(np.pi) + special.loggamma((1 - s) / 2) - special.loggamma(s / 2))


def w_transition(s):
    """w(s) = exp(-2 pi i N_RVM(tau_*)), tau_* = -i (s - 1/2)."""
    s = np.asarray(s, dtype=complex)
    tstar = -1j * (s - 0.5)
    return np.exp(-2j * np.pi * clock(tstar))


def theta_rs(t):
    t = np.asarray(t, dtype=float)
    return t / 2 * np.log(t / TWO_PI) - t / 2 - np.pi / 8 + 1 / (48 * t) + 7 / (5760 * t ** 3)


def gamma_lambda(gam, tau, lam):
    return (1.0 - lam) * gam + lam * tau


def ei_upper(w):
    """Ei on Im w != 0 along the horizontal line (paper, eq. Ei-def): Ei(w) = -E1(-w)."""
    return -special.exp1(-np.asarray(w, dtype=complex))
```

```
def mobius_sieve(X):
    mu = np.ones(X + 1, dtype=np.int8)
    mu[0] = 0
    is_comp = np.zeros(X + 1, dtype=bool)
    for p in range(2, X + 1):
        if not is_comp[p]:
            is_comp[2 * p::p] = True
            mu[p::p] *= -1
            if p * p <= X:
                mu[p * p::p * p] = 0
    return mu


def primes_upto(X):
    s = np.ones(X + 1, dtype=bool)
    s[:2] = False
    for p in range(2, int(X ** 0.5) + 1):
        if s[p]:
            s[p * p::p] = False
    return np.nonzero(s)[0]


def write_csv(name, header, cols):
    A = np.column_stack(cols)
    np.savetxt(os.path.join(OUT, name), A, delimiter=",", header=",".join(header), comments="", fmt="%.12g")


def savefig(fig, name):
    fig.savefig(os.path.join(FIG, name), dpi=150, bbox_inches="tight")
```

A.2. **s01_clock_and_family.py.**

```
"""Experiment 1: the midpoint clock, the displacements eta_k = tau_k - gamma_k,
the lambda-family zeta_lambda, and the numerical comparison zeta_Z1 versus zeta."""
import numpy as np
import mpmath as mp
import scipy.io as sio
import matplotlib
matplotlib.use("Agg")
import matplotlib.pyplot as plt
from rvm_common import (Tee, load_ordinates, clock, Product, zeta_from_product, gamma_lambda,
                        write_csv, savefig, OUT)
import os

say = Tee("s01_clock_and_family")
gam, tau = load_ordinates()
N = gam.size
k = np.arange(1, N + 1)
say("zeros loaded: N = %d, gamma_1 = %.9f, gamma_N = %.6f, tau_1 = %.9f, tau_N = %.6f",
    N, gam[0], gam[-1], tau[0], tau[-1])

# ---- clock check and displacements
clock_err = np.max(np.abs(clock(tau) - (k - 0.5)))
eta = tau - gam
say("max |N_RVM(tau_k) - (k - 1/2)| = %.3e", clock_err)
say("eta_k = tau_k - gamma_k:  max |eta| = %.6f (k = %d),  mean |eta| = %.6f,  rms = %.6f,  mean eta = %.3e",
    np.max(np.abs(eta)), int(np.argmax(np.abs(eta))) + 1, np.mean(np.abs(eta)),
    np.sqrt(np.mean(eta ** 2)), np.mean(eta))
blocks = [(1, 10), (11, 100), (101, 1000), (1001, 10 ** 4), (10 ** 4 + 1, 10 ** 5), (10 ** 5 + 1, 10 ** 6),
     (10 ** 6 + 1, N)]
rows = []
say("%12s %12s %12s %12s", "k from", "k to", "max|eta|", "rms eta")
for a, b in blocks:
```

```
    e = eta[a - 1:b]
    rows.append([a, b, np.max(np.abs(e)), np.sqrt(np.mean(e ** 2))])
    say("%12d %12d %12.6f %12.6f", a, b, rows[-1][2], rows[-1][3])
rows = np.array(rows)
write_csv("s01_eta_blocks.csv", ["k_from", "k_to", "max_abs_eta", "rms_eta"], [rows[:, i] for i in range(4)])
say("first ordinates:")
say("%4s %16s %16s %12s", "k", "gamma_k", "tau_k", "eta_k")
for i in range(12):
    say("%4d %16.9f %16.9f %12.6f", i + 1, gam[i], tau[i], eta[i])

# ---- zeta_Z1 versus zeta on vertical lines
PZ = Product(gam)
PR = Product(tau)
tline = np.concatenate([np.linspace(1.0, 100.0, 199), np.linspace(101.0, 2000.0, 300)])
mp.mp.dps = 20
cmp_rows = []
for sigma in (2.0, 1.25, 0.75):
    s = sigma + 1j * tline
    zz1 = zeta_from_product(PZ, s)
    ze = np.array([complex(mp.zeta(complex(x))) for x in s])
    rel = np.abs(zz1 / ze - 1.0)
    cmp_rows.append([sigma, np.max(rel), np.median(rel)])
    say("sigma = %.2f, t in [1,2000] (%d points): max |zeta_Z1/zeta - 1| = %.3e, median = %.3e",
        sigma, tline.size, np.max(rel), np.median(rel))
    write_csv("s01_zetaZ1_vs_zeta_sigma%.2f.csv" % sigma, ["t", "abs_zeta", "abs_zetaZ1", "rel_err"],
              [tline, np.abs(ze), np.abs(zz1), rel])
cmp_rows = np.array(cmp_rows)

# ---- Hardy-type functions Z_lambda(t) = exp(i theta(t)) zeta_lambda(1/2+it) on the critical line
def theta_exact(t):
    return np.array([float(mp.siegeltheta(x)) for x in t])

tc = np.linspace(0.5, 60.0, 1200)
th = theta_exact(tc)
lams = [0.0, 0.25, 0.5, 0.75, 1.0]
Zl = {}
for lam in lams:
    P = PZ if lam == 0.0 else (PR if lam == 1.0 else Product(gamma_lambda(gam, tau, lam)))
    Zl[lam] = np.real(np.exp(1j * th) * zeta_from_product(P, 0.5 + 1j * tc))
Zhardy = np.array([float(mp.siegelz(x)) for x in tc])
dZ = np.max(np.abs(Zl[0.0] - Zhardy) / np.maximum(1.0, np.abs(Zhardy)))
say("critical line, t in [0.5,60]: max |Z_Z1(t) - Z(t)| / max(1,|Z(t)|) = %.3e", dZ)
write_csv("s01_Z_lambda_critical_line.csv", ["t", "Z_hardy"] + ["Z_lambda_%.2f" % l for l in lams],
          [tc, Zhardy] + [Zl[l] for l in lams])

# ---- figures
dec = np.arange(0, N, 50)
fig, ax = plt.subplots(2, 2, figsize=(12, 8))
ax[0, 0].plot(k[dec], eta[dec], ".", ms=1.5, color="tab:blue")
ax[0, 0].axhline(0, color="k", lw=0.5)
ax[0, 0].set_xlabel("k")
ax[0, 0].set_ylabel(r"$\eta_k=\tau_k-\gamma_k$")
ax[0, 0].set_title(r"Bounded displacement, $N=%d$ zeros" % N)
ax[0, 1].hist(eta, bins=200, density=True, color="tab:blue")
ax[0, 1].set_xlabel(r"$\eta_k$")
ax[0, 1].set_title(r"Distribution of $\eta_k$ (all $k\leq N$)")
lam = np.linspace(0, 1, 101)
for kk in range(30):
    gl = (1 - lam) * gam[kk] + lam * tau[kk]
    ax[1, 0].plot(lam, clock(gl) - (kk + 0.5), "-", lw=1)
ax[1, 0].axhline(0, color="k", lw=0.5)
ax[1, 0].set_xlabel(r"$\lambda$")
ax[1, 0].set_ylabel(r"$\mathcal{N}_{\rm RVM}(\gamma_k(\lambda))-(k-\frac{1}{2})$")
ax[1, 0].set_title(r"Moving ordinates in clock units, $k\leq30$ ($\lambda=0$: $\zeta_{Z1}$, $\lambda=1$: $\
    zeta_{\rm RVM}$)", fontsize=9)
```

```
ax[1, 1].plot(tc, Zhardy, "k-", lw=2.2, label=r"Hardy $Z(t)$ of $\zeta$")
cols = ["tab:red", "tab:orange", "tab:green", "tab:cyan", "tab:blue"]
for lam, c in zip(lams, cols):
    ax[1, 1].plot(tc, Z1[lam], "-", color=c, lw=1.0, label=r"$Z_\lambda$, $\lambda=%.2f$" % lam)
ax[1, 1].axhline(0, color="k", lw=0.5)
ax[1, 1].set_xlabel("t")
ax[1, 1].set_title(r"$e^{i\vartheta(t)}\zeta_\lambda(\frac{1}{2}+it)$ along the family")
ax[1, 1].legend(fontsize=7, ncol=2)
fig.tight_layout()
savefig(fig, "fig01_clock_family.png")

fig, ax = plt.subplots(1, 2, figsize=(12, 4))
for sigma, c in zip((2.0, 1.25, 0.75), ("tab:blue", "tab:green", "tab:red")):
    d = np.loadtxt(os.path.join(OUT, "s01_zetaZ1_vs_zeta_sigma%.2f.csv" % sigma), delimiter=",", skiprows=1)
    ax[0].plot(d[:, 0], d[:, 3], ".", ms=2, color=c, label=r"$\sigma=%.2f$" % sigma)
ax[0].set_xlabel("t")
ax[0].set_ylabel(r"$|\zeta_{Z1}/\zeta-1|$")
ax[0].set_title(r"$\zeta_{Z1}$ from %d ordinates versus $\zeta$" % N)
ax[0].legend()
ax[1].plot(tc, Zhardy, "k-", lw=2.5, label=r"$Z(t)$ (mpmath)")
ax[1].plot(tc, Z1[0.0], "r--", lw=1.0, label=r"$Z_{Z1}(t)$ (product)")
ax[1].set_xlabel("t")
ax[1].legend()
ax[1].set_title(r"Critical line: $\lambda=0$ endpoint and Hardy $Z$")
fig.tight_layout()
savefig(fig, "fig02_zetaZ1_vs_zeta.png")

sio.savemat(os.path.join(OUT, "s01_results.mat"), {
    "N": N, "clock_err": clock_err, "eta_blocks": rows, "zetaZ1_cmp": cmp_rows,
    "t_critical": tc, "Z_hardy": Zhardy, "Z_lambda": np.array([Z1[l] for l in lams]), "lambdas": np.array(lams
    )})
say.close()

# ---- zoom: how the smooth clock places the midpoint ordinates next to the zeros
say2 = Tee("s01b_clock_zoom")
fig, ax = plt.subplots(1, 2, figsize=(12, 4))
T = np.linspace(9.0, 52.0, 4000)
ax[0].step(T, np.searchsorted(gam, T, side="right"), where="post", color="k", lw=1.5, label=r"$N_\zeta(T)$")
ax[0].plot(T, clock(T), "r-", lw=1.5, label=r"$\mathcal{N}_{\rm RVM}(T)$")
for i in range(9):
    ax[0].plot([gam[i], gam[i]], [0, i + 1], "k:", lw=0.8)
ax[0].plot(tau[:9], np.arange(9) + 0.5, "ro", ms=5, label=r"$\tau_k$: $\mathcal{N}_{\rm RVM}(\tau_k)=k-\frac
    {1}{2}$")
ax[0].plot(gam[:9], np.arange(9) + 0.5, "ks", ms=4, mfc="none", label=r"$(\gamma_k,\,k-\frac{1}{2})$")
ax[0].set_xlabel("T")
ax[0].set_title("Clock and staircase at the first zeros")
ax[0].legend(loc="upper left", fontsize=8)
kz = np.arange(N - 12, N)
T3 = np.linspace(gam[kz[0]] - 0.3, gam[kz[-1]] + 0.3, 6000)
ax[1].step(T3 - 1132480, np.searchsorted(gam, T3, side="right") - 2001000, where="post", color="k", lw=1.5,
    label=r"$N_\zeta(T)$")
ax[1].plot(T3 - 1132480, clock(T3) - 2001000, "r-", lw=1.5, label=r"$\mathcal{N}_{\rm RVM}(T)$")
ax[1].plot(tau[kz] - 1132480, kz + 0.5 - 2001000, "ro", ms=5, label=r"$\tau_k$")
ax[1].plot(gam[kz] - 1132480, kz + 0.5 - 2001000, "ks", ms=4, mfc="none", label=r"$(\gamma_k,\,k-\frac{1}{2})$
    ")
ax[1].set_xlabel(r"$T-1132480$")
ax[1].set_ylabel(r"count $-\,2001000$")
ax[1].set_title(r"Zoom at the last 12 zeros of the file (mean spacing %.3f)" % (2 * np.pi / np.log(gam[-1] /
    (2 * np.pi))))
ax[1].legend(loc="upper left", fontsize=8)
fig.tight_layout()
savefig(fig, "fig00_clock_zoom.png")
fig, ax = plt.subplots(1, 1, figsize=(12, 3))
T2 = np.linspace(9.0, 200.0, 20000)
ax.plot(T2, np.searchsorted(gam, T2, side="right") - clock(T2), "k-", lw=0.8)
```

```
ax.axhline(0, color="r", lw=0.8)
ax.set_xlabel("T")
ax.set_ylabel(r"$S(T)+O(1/T)$")
ax.set_title(r"$N_\zeta(T)-\mathcal{N}_{\rm RVM}(T)$: the remainder of the Riemann-von Mangoldt formula (jump
    $+1$ at each $\gamma_k$)")
fig.tight_layout()
savefig(fig, "fig00b_remainder.png")
say2("zoom table near the top of the file:")
say2("%9s %18s %18s %10s %10s", "k", "gamma_k", "tau_k", "eta_k", "N*-N_RVM")
for i in kz:
    say2("%9d %18.6f %18.6f %10.6f %10.6f", i + 1, gam[i], tau[i], tau[i] - gam[i],
         np.searchsorted(gam, gam[i], side="right") - 0.5 - clock(gam[i]))
say2.close()
```

A.3. **s02_transition.py.**

```
"""Experiment 2: the transition theorem zeta_RVM = varpi (1 + w)(1 + O(1/t)),
the constant varpi_RVM, the vertical limit of 1/zeta_RVM, and zeta_RVM versus zeta."""
import os
import numpy as np
import mpmath as mp
import scipy.io as sio
import matplotlib
matplotlib.use("Agg")
import matplotlib.pyplot as plt
from rvm_common import (Tee, load_ordinates, Product, zeta_from_product, chi, w_transition,
                        write_csv, savefig, OUT)

say = Tee("s02_transition")
gam, tau = load_ordinates()
PR = Product(tau)

# ---- varpi_RVM by Richardson extrapolation on Re s = 2
Ts = 1.0e4 * np.array([0.25, 0.5, 1.0])
v = np.array([zeta_from_product(PR, 2 + 1j * T)[0] / (1 + w_transition(2 + 1j * T)) for T in Ts])
A = np.column_stack([np.ones(3), 1 / Ts, 1 / Ts ** 2])
c = np.linalg.solve(A.astype(complex), v)
varpi = c[0].real
L = 1.0 / varpi
say("varpi_RVM (Richardson, sigma = 2, T = 2500, 5000, 10000) = %.10f %+.2e i", c[0].real, c[0].imag)
say("L = 1/varpi_RVM = %.10f", L)

# ---- relative error of the transition formula
tt = np.exp(np.linspace(np.log(10.0), np.log(1.0e4), 160))
rows = [tt]
hdr = ["t"]
fits = []
for sp in (0.25, 0.5, 1.5):
    s = 0.5 + sp + 1j * tt
    zr = zeta_from_product(PR, s)
    err_w = np.abs(zr / (varpi * (1 + w_transition(s))) - 1)
    rows.append(err_w)
    hdr.append("err_w_sigmap%.2f" % sp)
    sel = tt > 100
    slope = np.polyfit(np.log(tt[sel]), np.log(err_w[sel]), 1)[0]
    fits.append([sp, slope, np.max(err_w[sel] * tt[sel])])
    say("sigma' = %.2f: |zeta_RVM/(varpi(1+w)) - 1|: log-log slope on t>100 = %.3f, max t*err = %.4f",
        sp, slope, np.max(err_w[sel] * tt[sel]))
    if sp < 0.5:
        err_c = np.abs(zr / (varpi * (1 + chi(s))) - 1)
        rows.append(err_c)
        hdr.append("err_chi_sigmap%.2f" % sp)
        say("sigma' = %.2f: |zeta_RVM/(varpi(1+chi)) - 1|: max t*err (t>100) = %.4f",
            sp, np.max(err_c[sel] * tt[sel]))
    wmod = np.abs(w_transition(s))
```

```
    rows.append(wmod)
    hdr.append("abs_w_sigmap%.2f" % sp)
    Fv = 1 / zr
    rows.append(np.abs(Fv - L))
    hdr.append("abs_F_minus_L_sigmap%.2f" % sp)
    say("          |w| * t^sigma' in [%.4f, %.4f];  |1/zeta_RVM - L| at t = 1e4: %.3e",
        np.min(wmod * tt ** sp), np.max(wmod * tt ** sp), np.abs(Fv[-1] - L))
write_csv("s02_transition_errors.csv", hdr, rows)

# ---- zeta_RVM versus zeta on two vertical lines and on the critical line
mp.mp.dps = 20
tl = np.linspace(1.0, 120.0, 900)
cmp = {}
for sigma in (2.0, 0.75):
    s = sigma + 1j * tl
    zr = zeta_from_product(PR, s)
    ze = np.array([complex(mp.zeta(complex(x))) for x in s])
    cmp[sigma] = (zr, ze)
    write_csv("s02_zetaRVM_vs_zeta_sigma%.2f.csv" % sigma, ["t", "abs_zeta", "abs_zetaRVM", "
     abs_varpi_1_plus_w"],
              [tl, np.abs(ze), np.abs(zr), np.abs(varpi * (1 + w_transition(s)))])

fig, ax = plt.subplots(2, 2, figsize=(12, 8))
cols = {0.25: "tab:red", 0.5: "tab:green", 1.5: "tab:blue"}
j = 1
for sp in (0.25, 0.5, 1.5):
    ax[0, 0].plot(tt, tt * rows[j], "-", color=cols[sp], label=r"$\sigma'=%.2f$" % sp)
    j += 4 if sp < 0.5 else 3
ax[0, 0].set_xlabel("t")
ax[0, 0].set_ylabel(r"$t\,|\zeta_{\rm RVM}/(\varpi_{\rm RVM}(1+w))-1|$")
ax[0, 0].set_title(r"Transition theorem: error times $t$ stays bounded, $\varpi_{\rm RVM}=%.10f$" % varpi,
     fontsize=9)
ax[0, 0].legend()
for sp in (0.25, 0.5, 1.5):
    s = 0.5 + sp + 1j * tt
    ax[0, 1].plot(tt, np.abs(1 / zeta_from_product(PR, s) - L) * tt ** min(sp, 1.0), "-", color=cols[sp],
     label=r"$\sigma'=%.2f$" % sp)
ax[0, 1].set_xlabel("t")
ax[0, 1].set_ylabel(r"$t^{\min(\sigma',1)}\,|1/\zeta_{\rm RVM}(\sigma+it)-L|$")
ax[0, 1].set_title(r"Vertical limit $L=1/\varpi_{\rm RVM}$, rescaled error", fontsize=9)
ax[0, 1].legend()
for a, sigma in zip((ax[1, 0], ax[1, 1]), (2.0, 0.75)):
    zr, ze = cmp[sigma]
    a.plot(tl, np.abs(ze), "k-", lw=1.2, label=r"$|\zeta(\sigma+it)|$")
    a.plot(tl, np.abs(zr), "r-", lw=1.2, label=r"$|\zeta_{\rm RVM}(\sigma+it)|$")
    a.set_xlabel("t")
    a.set_title(r"$\sigma=%.2f$" % sigma)
    a.legend()
fig.tight_layout()
savefig(fig, "fig03_transition.png")

sio.savemat(os.path.join(OUT, "s02_results.mat"), {"varpi": varpi, "L": L, "richardson": c,
                                                   "slopes": np.array(fits), "t": tt})
say.close()
```

A.4. **s03_exponential_sums.py.**

```
"""Experiment 3: exponential sums S(N;u) = sum_{k<=N} exp(i u o_k) over the midpoint
ordinates tau_k and over the ordinates gamma_k of zeta, and the height transforms M_H(x)."""
import os
import numpy as np
import scipy.io as sio
import matplotlib
matplotlib.use("Agg")
import matplotlib.pyplot as plt
```

```
from rvm_common import Tee, load_ordinates, write_csv, savefig, OUT

say = Tee("s03_exponential_sums")
gam, tau = load_ordinates()
N = gam.size
step = 2000
Nk = np.arange(step, N + 1, step)
xs = [2.0, 3.0, 4.0, 6.0, 10.0]
Lam = {2.0: np.log(2), 3.0: np.log(3), 4.0: np.log(2), 6.0: 0.0, 10.0: 0.0}
res = {}
say("%6s %14s %14s %14s %16s %14s" % ("x", "max|S_RVM|", "max|S_RVM|/log", "|S_zeta(N)|",
                                      "Landau -T/2pi*L/sqrt(x)", "|M_H^RVM(x)|"))
for x in xs:
    u = np.log(x)
    SR = np.cumsum(np.exp(1j * u * tau))[Nk - 1]
    SZ = np.cumsum(np.exp(1j * u * gam))[Nk - 1]
    res[x] = (SR, SZ)
    landau = -gam[-1] / (2 * np.pi) * Lam[x] / np.sqrt(x)
    say("%6.1f %14.4f %14.4f %14.2f %16.2f %14.3e" % (x, np.max(np.abs(SR)), np.max(np.abs(SR) / np.log(Nk)),
                                                      np.abs(SZ[-1]), landau, np.abs(SR[-1]) / N))
    write_csv("s03_expsum_x%g.csv" % x, ["N", "re_S_RVM", "im_S_RVM", "re_S_zeta", "im_S_zeta"],
              [Nk, SR.real, SR.imag, SZ.real, SZ.imag])

fig, ax = plt.subplots(1, 2, figsize=(12, 4))
cols = {2.0: "tab:blue", 3.0: "tab:green", 4.0: "tab:purple", 6.0: "tab:orange", 10.0: "tab:red"}
for x in xs:
    ax[0].plot(Nk, np.abs(res[x][0]), "-", color=cols[x], lw=0.8, label=r"$x=%g$" % x)
ax[0].plot(Nk, np.log(Nk), "k--", lw=1, label=r"$\log N$")
ax[0].set_xlabel("N")
ax[0].set_ylabel(r"$|S_{\rm RVM}(N;\log x)|$")
ax[0].set_title(r"Midpoint ordinates: $|\sum_{k\leq N}x^{i\tau_k}|$ stays $O(\log N)$")
ax[0].legend(fontsize=8, ncol=2)
for x in xs:
    ax[1].plot(Nk, res[x][1].real, "-", color=cols[x], lw=0.8, label=r"$x=%g$" % x)
T = np.interp(Nk, np.arange(1, N + 1), gam)
for x in (2.0, 3.0, 4.0):
    ax[1].plot(Nk, -T / (2 * np.pi) * Lam[x] / np.sqrt(x), ":", color=cols[x], lw=1.5)
ax[1].set_xlabel("N")
ax[1].set_ylabel(r"$\mathrm{Re}\sum_{k\leq N}x^{i\gamma_k}$")
ax[1].set_title(r"Zeros of $\zeta$: Landau term $-\frac{T}{2\pi}\frac{\Lambda(x)}{\sqrt{x}}$ (dotted)")
ax[1].legend(fontsize=8, ncol=2)
fig.tight_layout()
savefig(fig, "fig04_exponential_sums.png")
sio.savemat(os.path.join(OUT, "s03_results.mat"), {"N": Nk, "x": np.array(xs),
                                                   "S_RVM": np.array([res[x][0] for x in xs]),
                                                   "S_zeta": np.array([res[x][1] for x in xs])})
say.close()
```

A.5. **s04_primes.py.**

```
"""Experiment 4: the prime-power count J_RVM, the synthetic prime count pi_RVM, the cell
masses c_n^RVM and the real RVM prime levels p_j^RVM, compared with J, pi(x) and the primes.

J_o(x) = li(x) - 2 Re sum_k Ei((1/2 + i o_k) log x) - log 2 + int_x^inf dt/(t(t^2-1) log t)
for an ordinate list o_k (o = tau: J_RVM; o = gamma: Riemann's explicit formula for J),
pi_o(x) = sum_{m <= log2 x} mu(m)/m J_o(x^(1/m)).
The zero sums are evaluated on a uniform grid in u = log x by the recurrence
exp(i o_k (u + du)) = exp(i o_k u) exp(i o_k du); Ei is exact (scipy exp1) for k <= 300 and
uses Ei(w) = e^w (1/w + 1/w^2 + 2/w^3 + 6/w^4) for k > 300, where |w| > 400."""
import os
import numpy as np
import scipy.io as sio
from scipy import special, integrate
import matplotlib
matplotlib.use("Agg")
```

```
import matplotlib.pyplot as plt
from rvm_common import (Tee, load_ordinates, ei_upper, mobius_sieve, primes_upto, write_csv,
                        savefig, clock_inverse, OUT)

say = Tee("s04_primes")
gam, tau = load_ordinates()
KE = 300


def zero_sum_ei(ords, u, K=None, taper=None):
    """sum_k Ei((1/2 + i o_k) u) on a uniform grid u, k <= K, optional weights."""
    o = ords if K is None else ords[:K]
    wts = np.ones(o.size) if taper is None else taper
    rho = 0.5 + 1j * o
    out = np.zeros(u.size, dtype=complex)
    # exact part
    for i in range(0, KE):
        out += wts[i] * ei_upper(rho[i] * u)
    # asymptotic part by recurrence
    r = rho[KE:]
    b = [wts[KE:] / r ** m for m in (1, 2, 3, 4)]
    du = u[1] - u[0]
    stepf = np.exp(1j * o[KE:] * du)
    for j in range(u.size):
        if j % 200 == 0:
            E = np.exp(1j * o[KE:] * u[j])
        A = [E @ bm for bm in b]
        uj = u[j]
        out[j] += np.exp(uj / 2) * (A[0] / uj + A[1] / uj ** 2 + 2 * A[2] / uj ** 3 + 6 * A[3] / uj ** 4)
        E = E * stepf
    return out


def tail_integral(x):
    return integrate.quad(lambda t: 1.0 / (t * (t * t - 1.0) * np.log(t)), x, np.inf, limit=200)[0]


def J_from(ords, u, K=None, taper=None):
    x = np.exp(u)
    S = zero_sum_ei(ords, u, K, taper)
    ti = np.array([tail_integral(v) for v in x])
    return special.expi(u) - 2 * S.real - np.log(2.0) + ti


def pi_from_J(u, J):
    mu = mobius_sieve(64)
    out = np.zeros(u.size)
    for i, ui in enumerate(u):
        m = 1
        while ui / m >= np.log(2.0) - 1e-12:
            if mu[m] != 0:
                out[i] += mu[m] / m * np.interp(ui / m, u, J)
            m += 1
    return out


def exact_J_pi(x):
    P = primes_upto(int(x.max()) + 1)
    pi = np.searchsorted(P, x, side="right").astype(float)
    J = np.zeros(x.size)
    for m in range(1, 40):
        J += np.searchsorted(P, x ** (1.0 / m) + 1e-12, side="right") / m
    return J, pi
```

```
def run(xmax, npts, K, label):
    u = np.linspace(np.log(2.0), np.log(xmax), npts)
    if K is None:
        tg = tr = None
        Kg = Kr = None
    else:
        Kg = Kr = K
        v = np.arange(K) / K
        tg = tr = np.where(v < 0.6, 1.0, np.cos(np.pi / 2 * (v - 0.6) / 0.4) ** 2)
    JR = J_from(tau, u, Kr, tr)
    JG = J_from(gam, u, Kg, tg)
    PR = pi_from_J(u, JR)
    PG = pi_from_J(u, JG)
    x = np.exp(u)
    Je, Pe = exact_J_pi(x)
    write_csv("s04_prime_counts_%s.csv" % label, ["x", "J_exact", "J_from_zeta_zeros", "J_RVM",
                                                  "pi_exact", "pi_from_zeta_zeros", "pi_RVM", "li"],
              [x, Je, JG, JR, Pe, PG, PR, special.expi(u)])
    return x, Je, JG, JR, Pe, PG, PR


# x <= 100 with all zeros; x <= 1000 with a smooth cut at K zeros
x1, Je1, JG1, JR1, Pe1, PG1, PR1 = run(100.0, 3000, None, "x100_allzeros")
say("x <= 100, all %d zeros of each list, %d grid points:" % (gam.size, x1.size))
say("  max |pi_from_zeta_zeros - pi| away from primes (distance > 0.2): %.4f",
    np.max(np.abs(PG1 - Pe1)[np.min(np.abs(x1[:, None] - primes_upto(101)[None, :]), axis=1) > 0.2]))
say("  pi_RVM(100) = %.4f, pi(100) = %d, li(100) = %.4f, J_RVM(100) = %.4f, J(100) = %.4f",
    PR1[-1], Pe1[-1], special.expi(np.log(100.0)), JR1[-1], Je1[-1])
dPR = np.diff(PR1)
say("  pi_RVM monotone on [2,100]: %s (min increment on the grid %.3e)", bool(np.all(dPR > 0)), dPR.min())

tauK = 2 * np.pi * 1000.0 * 2.5
K = int(np.searchsorted(tau, tauK))
x2, Je2, JG2, JR2, Pe2, PG2, PR2 = run(1000.0, 40000, K, "x1000_K%d" % K)
say("x <= 1000, smooth cut at K = %d zeros (tau_K = %.1f), %d grid points:" % (K, tau[K - 1], x2.size))
say("  pi_RVM(1000) = %.4f, pi(1000) = %d, li(1000) = %.4f", PR2[-1], Pe2[-1], special.expi(np.log(1000.0)))

# cell masses and real prime levels from the fine x<=1000 run
n = np.arange(2, 1001)
pin = np.interp(n, x2, PR2)
c = np.diff(np.concatenate([[0.0], pin]))
P = primes_upto(1000)
isp = np.isin(n, P).astype(float)
say("cell masses c_n = pi_RVM(n) - pi_RVM(n-1):")
say("%5s %10s %10s %8s" % ("n", "c_n", "1/log n", "prime"))
for i in range(0, 40):
    say("%5d %10.5f %10.5f %8d" % (n[i], c[i], 1 / np.log(n[i]), isp[i]))
say("sum_{n<=1000} c_n = %.4f,  pi(1000) = %d", c.sum(), len(P))
levels = np.arange(1, int(np.floor(PR2.max())) + 1)
pj = np.array([x2[np.argmax(PR2 >= j)] for j in levels])
say("real RVM prime levels p_j^RVM (first crossing of level j) versus p_j:")
say("%5s %12s %8s %10s" % ("j", "p_j^RVM", "p_j", "diff"))
for j in range(min(30, levels.size)):
    say("%5d %12.4f %8d %10.4f" % (levels[j], pj[j], P[j], pj[j] - P[j]))
write_csv("s04_cell_masses.csv", ["n", "c_n_RVM", "inv_log_n", "is_prime"], [n, c, 1 / np.log(n), isp])
m = min(levels.size, P.size)
write_csv("s04_rvm_prime_levels.csv", ["j", "p_j_RVM", "p_j"], [levels[:m], pj[:m], P[:m]])

# ---- figures (linear axes)
fig, ax = plt.subplots(1, 2, figsize=(12, 4))
ax[0].step(x1, Pe1, where="post", color="k", lw=1.6, label=r"$\pi(x)$")
ax[0].plot(x1, PG1, "b-", lw=0.6, label=r"$\pi(x)$ from %d zeros of $\zeta$" % gam.size)
ax[0].plot(x1, PR1, "r-", lw=1.4, label=r"$\pi_{\rm RVM}(x)$ from %d $\tau_k$" % tau.size)
ax[0].plot(x1, special.expi(np.log(x1)) - special.expi(np.log(2.0)), "g--", lw=1, label=r"$\mathrm{li}(x)-\
    mathrm{li}(2)$")
```

```
ax[0].set_xlabel("x")
ax[0].set_title(r"Prime counts on $[2,100]$")
ax[0].legend(fontsize=7)
ax[1].step(x2, Pe2, where="post", color="k", lw=1.2, label=r"$\pi(x)$")
ax[1].plot(x2, PR2, "r-", lw=1.0, label=r"$\pi_{\rm RVM}(x)$")
ax[1].set_xlabel("x")
ax[1].set_title(r"Prime counts on $[2,1000]$ ($K=%d$ ordinates, smooth cut)" % K)
ax[1].legend(fontsize=8)
fig.tight_layout()
savefig(fig, "fig05_prime_counts.png")

fig, ax = plt.subplots(1, 2, figsize=(12, 4))
nn = n[:98]
ax[0].bar(nn, isp[:98], width=0.8, color="0.75", label="prime indicator")
ax[0].plot(nn, c[:98], "r.-", lw=0.8, ms=4, label=r"$c_n^{\rm RVM}$")
ax[0].plot(nn, 1 / np.log(nn), "g--", lw=1, label=r"$1/\log n$")
ax[0].set_xlabel("n")
ax[0].set_title(r"Cell masses $c_n^{\rm RVM}=\pi_{\rm RVM}(n)-\pi_{\rm RVM}(n-1)$")
ax[0].legend(fontsize=8)
ax[1].plot(P[:m], pj[:m], "r.", ms=4, label=r"$(p_j,\,p_j^{\rm RVM})$")
ax[1].plot([0, 1000], [0, 1000], "k-", lw=0.8, label="diagonal")
ax[1].set_xlabel(r"$p_j$")
ax[1].set_ylabel(r"$p_j^{\rm RVM}$")
ax[1].set_title(r"Real RVM prime levels $\pi_{\rm RVM}(p_j^{\rm RVM})=j$ versus primes")
ax[1].legend(fontsize=8)
fig.tight_layout()
savefig(fig, "fig06_rvm_primes.png")

sio.savemat(os.path.join(OUT, "s04_results.mat"), {"x100": x1, "pi100": Pe1, "piZeta100": PG1, "piRVM100": PR1,

                                                   "x1000": x2, "pi1000": Pe2, "piRVM1000": PR2,
                                                   "c_n": c, "p_RVM": pj, "K": K})
say.close()
```

A.6. **s05_mertens.py.**

```
"""Experiment 5: the Mertens function M, the synthetic Mertens function M_RVM^crit and the
ternary walk M_h with values in {-1,0,+1} on the squarefree support.

Explicit formulas (simple zeros, residues a = 1/(rho zeta'(rho))):
  M_zeta(x) = -2 + 2 Re sum_k a_k x^(rho_k) + trivial terms        (zeros of zeta)
  M_RVM(x)  = -2 + 2 Re sum_k b_k x^(1/2 + i tau_k) + trivial terms (b_k = 1/(rho_k zeta_RVM'(rho_k)))
Ternary walk: mu_h(n) = mu(n)^2 * sign, sign = +1 if M_h(n-1) < target(n) else -1."""
import os
import math
import numpy as np
import mpmath as mp
import scipy.io as sio
from scipy.interpolate import PchipInterpolator
import matplotlib
matplotlib.use("Agg")
import matplotlib.pyplot as plt
from rvm_common import (Tee, load_ordinates, Product, logG, clock_inverse, tail_coefficients,
                        theta_rs, mobius_sieve, write_csv, savefig, OUT)

say = Tee("s05_mertens")
gam, tau = load_ordinates()
Nz = gam.size
XMAX = 10 ** 6
K0 = 1000

# ------------------------------------------------------------------ RVM residues
c1, c2, c3 = tail_coefficients(clock_inverse(float(Nz)))
t2 = tau ** 2
zpR = np.zeros(K0, dtype=complex)
```

```
for k in range(K0):
    tk = tau[k]
    fac = (t2 - tk ** 2) / (t2 + 0.25)
    fac[k] = 1.0
    q = -0.25 - tk ** 2
    lmod = np.sum(np.log(np.abs(fac))) + c1 * q - c2 * q ** 2 + c3 * q ** 3
    rho = 0.5 + 1j * tk
    lP = np.log(2j * tk / (0.25 + tk ** 2)) + lmod
    zpR[k] = (-1) ** k * np.exp(lP - logG(rho))
ratio = zpR / np.log(tau[:K0] / (2 * np.pi))
PR = Product(tau)
Ts = 1.0e4 * np.array([0.25, 0.5, 1.0])
from rvm_common import zeta_from_product, w_transition
v = np.array([zeta_from_product(PR, 2 + 1j * T)[0] / (1 + w_transition(2 + 1j * T)) for T in Ts])
varpi = np.linalg.solve(np.column_stack([np.ones(3), 1 / Ts, 1 / Ts ** 2]).astype(complex), v)[0].real
sel = np.arange(K0 // 2, K0)
cfit = np.mean((ratio[sel] / varpi - 1) * tau[sel])
say("zeta_RVM'(rho_k) exact for k <= %d; zeta_RVM'(rho_k)/log(tau_k/2pi) at k = %d: %.10f %+.2e i",
    K0, K0, ratio[-1].real, ratio[-1].imag)
say("varpi_RVM = %.10f; fit zeta_RVM'(rho_k) = varpi log(tau_k/2pi)(1 + c/tau_k), c = %.4f %+.4f i",
    varpi, cfit.real, cfit.imag)
zR = varpi * np.log(tau / (2 * np.pi)) * (1 + cfit / tau)
zR[:K0] = zpR
bR = 1.0 / ((0.5 + 1j * tau) * zR)

# ------------------------------------------------------------------ residues of zeta
def rs_Z(t):
    a = np.sqrt(t / (2 * np.pi))
    Nn = np.floor(a).astype(int)
    p = a - Nn
    nmax = Nn.max()
    n = np.arange(1, nmax + 1)
    ph = theta_rs(t)[:, None] - t[:, None] * np.log(n)[None, :]
    mask = n[None, :] <= Nn[:, None]
    Z = 2 * np.sum(np.cos(ph) * mask / np.sqrt(n)[None, :], axis=1)
    cc = np.cos(2 * np.pi * p)
    C0 = np.cos(2 * np.pi * (p ** 2 - p - 1.0 / 16)) / cc
    return Z + (-1.0) ** (Nn - 1) * (t / (2 * np.pi)) ** (-0.25) * C0


mp.mp.dps = 20
KM = 1000
zG = np.zeros(Nz, dtype=complex)
for k in range(KM + 50):
    zG[k] = complex(mp.zeta(mp.mpc(0.5, gam[k]), 1, 1))
h = 1e-5
for i0 in range(KM, Nz, 4000):
    i1 = min(Nz, i0 + 4000)
    t = gam[i0:i1]
    Zp = (rs_Z(t + h) - rs_Z(t - h)) / (2 * h)
    val = -1j * np.exp(-1j * theta_rs(t)) * Zp
    if i0 == KM:
        chk = np.max(np.abs(val[:50] / zG[KM:KM + 50] - 1))
        say("Riemann-Siegel zeta'(rho) versus mpmath at k = %d..%d: max relative difference %.2e",
            KM + 1, KM + 50, chk)
        zG[KM + 50:i1] = val[50:]
    else:
        zG[i0:i1] = val
aG = 1.0 / ((0.5 + 1j * gam) * zG)
S2G = 2 * np.sum(np.abs(aG) ** 2)
S2R = 2 * np.sum(np.abs(bR) ** 2)
say("median |zeta'(rho)| = %.4f;  spectral sums 2 sum|a|^2: zeta %.6f, RVM %.6f", np.median(np.abs(zG)), S2G,
    S2R)
```

```
def trivial_zeta(x, n=6):
    v = np.zeros_like(x)
    for k in range(1, n + 1):
        v += (-1) ** (k - 1) * (2 * np.pi / x) ** (2 * k) / (math.factorial(2 * k) * k * float(mp.zeta(2 * k +
     1)))
    return v


def trivial_rvm(x, n=6):
    v = np.zeros_like(x)
    for k in range(1, n + 1):
        s = -2.0 * k
        lP = np.real(PR.logP(s)[0])
        v -= x ** (-2 * k) * (2 * k + 1) * np.pi ** k * 2 * (-1) ** k / (math.factorial(k) * np.exp(lP))
    return v


def explicit_uniform(ords, wts, u):
    """2 Re sqrt(x) sum_k w_k exp(i o_k u) on a uniform u grid (recurrence)."""
    out = np.zeros(u.size)
    st = np.exp(1j * ords * (u[1] - u[0]))
    for j in range(u.size):
        if j % 200 == 0:
            E = np.exp(1j * ords * u[j])
        out[j] = 2 * np.exp(u[j] / 2) * np.real(E @ wts)
        E = E * st
    return out


# ---------------------------------------------------------------- exact M by sieve
mu = mobius_sieve(XMAX)
M = np.cumsum(mu.astype(np.int64))
sqf = mu != 0
say("M(10^6) = %d;  squarefree density Q(10^6)/10^6 = %.6f (6/pi^2 = %.6f)", M[XMAX], sqf[1:].mean(), 6 / np.
    pi ** 2)

# ---------------------------------------------------------------- explicit formulas on [2, 2000]
u = np.linspace(np.log(2.0), np.log(2000.0), 2500)
xg = np.exp(u)
Mz = -2 + explicit_uniform(gam, aG, u) + trivial_zeta(xg)
Mc = -2 + explicit_uniform(tau, bR, u) + trivial_rvm(xg)
Mfl = M[np.floor(xg).astype(int)]
off = np.abs(xg - np.round(xg)) > 0.05
say("explicit formula with %d zeros of zeta on [2,2000]: rms(M_zeta - M) = %.4f off integers (|x-n|>0.05)",
    Nz, np.sqrt(np.mean((Mz - Mfl)[off] ** 2)))
write_csv("s05_explicit_formulas.csv", ["x", "M_floor", "M_zeta_explicit", "M_RVM_explicit"], [xg, Mfl, Mz, Mc
    ])

# ---------------------------------------------------------------- integer target for M_RVM^crit
def target_grid(X, div=24):
    x = [1.0]
    while x[-1] < X:
        x.append(min(X, x[-1] + max(1.0, np.floor(np.sqrt(x[-1]) / div))))
    return np.array(x)


def tapered(x, ords, a, C1=2.0, C2=4.0):
    S = np.zeros(x.size, dtype=complex)
    lx = np.log(x)
    i0 = 0
    while i0 < x.size:
        i1 = min(x.size, i0 + 200)
        K = max(1, int(np.searchsorted(ords, C2 * 2 * np.pi * np.sqrt(x[i1 - 1]))))
        uu = ords[None, :K] / (2 * np.pi * np.sqrt(x[i0:i1, None]))
        w = np.where(uu <= C1, 1.0, np.where(uu >= C2, 0.0, np.cos(np.pi / 2 * (uu - C1) / (C2 - C1)) ** 2))
```

```
        E = np.exp(1j * lx[i0:i1, None] * ords[None, :K])
        S[i0:i1] = np.sqrt(x[i0:i1]) * np.sum(E * w * a[None, :K], axis=1)
        i0 = i1
    return S


xT = target_grid(float(XMAX))
TT = -2 + 2 * np.real(tapered(xT, tau, bR)) + trivial_rvm(xT)
nn = np.arange(1, XMAX + 1)
T = np.concatenate([[0.0], PchipInterpolator(xT, TT)(nn)])
nchk = np.arange(1000, 100001, 997)
full = -2 + 2 * np.real(np.array([np.sqrt(n) * np.sum(np.exp(1j * np.log(n) * tau) * bR) for n in nchk])) +
    trivial_rvm(nchk.astype(float))
say("target on %d grid points; at %d integers <= 1e5: max |M_RVM(n) (all zeros) - target(n)| = %.4f",
    xT.size, nchk.size, np.max(np.abs(full - T[nchk])))

# ------------------------------------------------------------------ ternary walk
mu_h = np.zeros(XMAX + 1, dtype=np.int8)
Mh = np.zeros(XMAX + 1)
sgn = np.zeros(XMAX + 1, dtype=np.int8)
cur = 0.0
for n in range(1, XMAX + 1):
    if sqf[n]:
        sgn[n] = 1 if cur < T[n] else -1
        cur += sgn[n]
        mu_h[n] = sgn[n]
    Mh[n] = cur
err = Mh[1:] - T[1:]
say("ternary walk: max |M_h - target| on [1,1e6] = %.4f, on [1e4,1e6] = %.4f", np.max(np.abs(err)), np.max(np.
    abs(err[9999:])))
dens = lambda a: [np.mean(a[1:] == -1), np.mean(a[1:] == 0), np.mean(a[1:] == 1)]
say("densities of (-1, 0, +1):  mu   %.6f %.6f %.6f", *dens(mu))
say("                           mu_h %.6f %.6f %.6f   (3/pi^2 = %.6f, 1 - 6/pi^2 = %.6f)", *dens(mu_h), 3 / np.
    pi ** 2, 1 - 6 / np.pi ** 2)
say("sign agreement mu_h = mu on squarefrees: %.6f;  sum mu mu_h / Q = %.6f",
    np.mean(mu_h[sqf] == mu[sqf]), np.sum(mu_h.astype(int) * mu.astype(int)) / sqf[1:].sum())
say("max |M(n)|/sqrt(n) on [10,1e6] = %.4f;  max |M_RVM(n)|/sqrt(n) = %.4f;  max |M_h(n)|/sqrt(n) = %.4f",
    np.max(np.abs(M[10:]) / np.sqrt(nn[9:])), np.max(np.abs(T[10:]) / np.sqrt(nn[9:])), np.max(np.abs(Mh[10:])
     / np.sqrt(nn[9:])))
say("\nNUMERICAL EXAMPLE OF THE TERNARY DIVISION (first 40 integers)")
say("%4s %4s %6s %10s %8s %6s %6s %6s %6s" % ("n", "mu", "mu^2", "target", "M_h(n-1)", "sign", "mu_h", "M_h",
    "M"))
rows = []
for n in range(1, 41):
    say("%4d %4d %6d %10.4f %8d %6s %6d %6d %6d" % (n, mu[n], abs(mu[n]), T[n], Mh[n - 1],
                                                    ("+1" if sgn[n] > 0 else ("-1" if sgn[n] < 0 else "-")),
                                                    mu_h[n], Mh[n], M[n]))
    rows.append([n, mu[n], abs(mu[n]), T[n], Mh[n - 1], sgn[n], mu_h[n], Mh[n], M[n]])
rows = np.array(rows, dtype=float)
write_csv("s05_ternary_example.csv", ["n", "mu", "mu2", "target", "Mh_prev", "sign", "mu_h", "Mh", "M"],
          [rows[:, i] for i in range(rows.shape[1])])

Ns = 10 ** np.arange(3, 7)
say("\nENERGY E_N = (1/N) sum_{n<=N} W(n)^2")
say("%10s %14s %14s %14s" % ("N", "E_N(M)", "E_N(M_RVM)", "E_N(M_h)"))
EN = []
for Nn in Ns:
    EN.append([Nn, np.mean(M[1:Nn + 1].astype(float) ** 2), np.mean(T[1:Nn + 1] ** 2), np.mean(Mh[1:Nn + 1] **
     2)])
    say("%10d %14.4f %14.4f %14.4f" % tuple(EN[-1]))
EN = np.array(EN)
write_csv("s05_energy.csv", ["N", "E_M", "E_MRVM", "E_Mh"], [EN[:, i] for i in range(4)])
nall = np.arange(1, XMAX + 1)
write_csv("s05_walks.csv", ["n", "M", "M_RVM_target", "M_h"], [nall, M[nall], T[nall], Mh[nall]])
mu_h[1:].astype(np.int8).tofile(os.path.join(OUT, "mu_h_int8_1e6.bin"))
```

```
# ---------------------------------------------------------------- figures (linear axes)
fig, ax = plt.subplots(1, 2, figsize=(12, 4))
w = np.arange(1, 2001)
ax[0].step(w, M[w], where="post", color="k", lw=1.0, label=r"$M(n)$")
ax[0].plot(w, T[w], "r-", lw=1.2, label=r"$M_{\rm RVM}^{\rm crit}(n)$")
ax[0].step(w, Mh[w], where="post", color="tab:green", lw=0.8, label=r"$M_h(n)$")
ax[0].set_xlabel("n")
ax[0].set_title(r"Mertens functions on $[1,2000]$, every integer")
ax[0].legend(fontsize=8)
nb = 2000
bw = XMAX // nb
xb = (np.arange(nb) + 0.5) * bw
Mb = M[1:].reshape(nb, bw)
Tb = T[1:].reshape(nb, bw)
ax[1].fill_between(xb, Mb.min(1), Mb.max(1), color="0.6", label=r"$M(n)$ (range per block)")
ax[1].plot(xb, Tb.mean(1), "r-", lw=1.0, label=r"$M_{\rm RVM}^{\rm crit}(n)$")
ax[1].plot(xb, np.sqrt(xb), "k--", lw=0.8, label=r"$\pm\sqrt{n}$")
ax[1].plot(xb, -np.sqrt(xb), "k--", lw=0.8)
ax[1].set_xlabel("n")
ax[1].set_title(r"$[1,10^6]$, blocks of %d integers" % bw)
ax[1].legend(fontsize=8)
fig.tight_layout()
savefig(fig, "fig07_mertens.png")

fig, ax = plt.subplots(1, 2, figsize=(12, 4))
w = np.arange(1, 61)
ax[0].vlines(w - 0.15, 0, mu[w], color="k", lw=2, label=r"$\mu(n)$")
ax[0].vlines(w + 0.15, 0, mu_h[w], color="tab:green", lw=2, label=r"$\mu_h(n)$")
ax[0].axhline(0, color="k", lw=0.5)
ax[0].set_xlabel("n")
ax[0].set_yticks([-1, 0, 1])
ax[0].set_title(r"Ternary values: same zero set $\{\mu(n)=0\}$, signs chosen by tracking")
ax[0].legend(fontsize=8)
w = np.arange(1, 301)
ax[1].step(w, M[w], where="post", color="k", lw=1.0, label=r"$M(n)$")
ax[1].plot(w, T[w], "r.-", lw=0.8, ms=3, label=r"target $M_{\rm RVM}^{\rm crit}(n)$")
ax[1].step(w, Mh[w], where="post", color="tab:green", lw=1.2, label=r"$M_h(n)$")
ax[1].set_xlabel("n")
ax[1].set_title(r"The walk $M_h$ follows $M_{\rm RVM}^{\rm crit}$ within $\pm1$ step")
ax[1].legend(fontsize=8)
fig.tight_layout()
savefig(fig, "fig08_ternary_walk.png")

fig, ax = plt.subplots(1, 2, figsize=(12, 4))
sel = xg <= 300
ax[0].step(np.arange(1, 301), M[1:301], where="post", color="k", lw=1.0, label=r"$M(x)$")
ax[0].plot(xg[sel], Mz[sel], "b-", lw=0.7, label=r"explicit formula, %d zeros of $\zeta$" % Nz)
ax[0].plot(xg[sel], Mc[sel], "r-", lw=1.0, label=r"$M_{\rm RVM}^{\rm crit}(x)$, %d $\tau_k$" % Nz)
ax[0].set_xlabel("x")
ax[0].set_title("Explicit formulas on [2,300]")
ax[0].legend(fontsize=7)
kk = np.arange(5000)
ax[1].plot(gam[kk], np.abs(zG[kk]) / np.log(gam[kk] / (2 * np.pi)), "b.", ms=1.5, label=r"$|\zeta'(\rho_k)|/\
    log(\gamma_k/2\pi)$")
ax[1].plot(tau[kk], np.abs(zR[kk]) / np.log(tau[kk] / (2 * np.pi)), "r.", ms=1.5, label=r"$|\zeta_{\rm RVM}'(\
    rho^{\rm RVM}_k)|/\log(\tau_k/2\pi)$")
ax[1].axhline(varpi, color="k", lw=0.8, ls="--", label=r"$\varpi_{\rm RVM}$")
ax[1].set_xlabel("ordinate")
ax[1].set_ylim(0, 4)
ax[1].set_title(r"Derivatives at the zeros, $k\leq5000$")
ax[1].legend(fontsize=7)
fig.tight_layout()
savefig(fig, "fig09_explicit_weights.png")
```

```
sio.savemat(os.path.join(OUT, "s05_results.mat"), {"varpi": varpi, "cfit": cfit, "S2G": S2G, "S2R": S2R,
                                                   "energy": EN, "ternary_example": rows,
                                                   "x_explicit": xg, "M_zeta_explicit": Mz, "M_RVM_explicit":
    Mc})
say.close()
```

A.7. **s06_prime_ratios.py.**

```
"""Experiment 6: the three ratios li/pi_RVM, pi/li, pi/pi_RVM and the square-root estimate
pi_RVM(x) - li(x) = O(sqrt(x) (log log x)^2 / log x), from the prime counts written by s04_primes.py.

Input: output/s04_prime_counts_x1000_K17061.csv (columns x, pi_exact, pi_RVM, li, ...).
B(x) = sqrt(x) (log log x)^2 / log x is the scale of the square-root estimate."""
import os
import numpy as np
import matplotlib
matplotlib.use("Agg")
import matplotlib.pyplot as plt
from rvm_common import Tee, savefig, write_csv, OUT

say = Tee("s06_prime_ratios")
D = np.genfromtxt(os.path.join(OUT, "s04_prime_counts_x1000_K17061.csv"), delimiter=",", names=True)
x, pe, pr, li = D["x"], D["pi_exact"], D["pi_RVM"], D["li"]
say("grid: %d points on [%.0f, %.0f]; pi_RVM from the first 17061 ordinates tau_k (smooth cut)"
    % (x.size, x[0], x[-1]))

sel = x >= 10
r1, r2, r3 = li[sel] / pr[sel], pe[sel] / li[sel], pe[sel] / pr[sel]
xs = x[sel]
say("")
say("ratios at selected x")
say("%8s %12s %12s %12s %12s %12s %12s" % ("x", "pi", "pi_RVM", "li", "li/pi_RVM", "pi/li", "pi/pi_RVM"))
for X in [10, 20, 50, 100, 200, 500, 1000]:
    i = int(np.argmin(np.abs(x - X)))
    say("%8.1f %12.4f %12.4f %12.4f %12.6f %12.6f %12.6f"
        % (x[i], pe[i], pr[i], li[i], li[i] / pr[i], pe[i] / li[i], pe[i] / pr[i]))

s8 = x >= 8
B = np.sqrt(x[s8]) * np.log(np.log(x[s8])) ** 2 / np.log(x[s8])
e_rvm = (pr[s8] - li[s8]) / B
e_pi = (pe[s8] - li[s8]) / B
say("")
say("square-root scale B(x) = sqrt(x) (log log x)^2 / log x on [8, 1000]")
say("max |pi_RVM - li| / B = %.4f   (at x = %.2f)" % (np.max(np.abs(e_rvm)), x[s8][np.argmax(np.abs(e_rvm))]))
say("max |pi - li| / B     = %.4f   (at x = %.2f)" % (np.max(np.abs(e_pi)), x[s8][np.argmax(np.abs(e_pi))]))
for X in [100, 1000]:
    m = x[s8] >= X / 2
    say("max |pi_RVM - li| / B on [%d, 1000] = %.4f" % (X // 2, np.max(np.abs(e_rvm[m]))))

# CSV files: full grid
write_csv("s06_prime_ratios.csv", ["x", "li_over_pi_RVM", "pi_over_li", "pi_over_pi_RVM"],
          [xs, r1, r2, r3])
write_csv("s06_square_root_scale.csv", ["x", "pi_RVM_minus_li_over_B", "pi_minus_li_over_B"],
          [x[s8], e_rvm, e_pi])

# ---- figure (linear axes)
fig, ax = plt.subplots(1, 2, figsize=(12, 4))
ax[0].plot(xs, r3, "k-", lw=0.8, label=r"$\pi(x)/\pi_{\rm RVM}(x)$")
ax[0].plot(xs, r2, "b-", lw=0.8, label=r"$\pi(x)/\mathrm{li}(x)$")
ax[0].plot(xs, r1, "r-", lw=1.4, label=r"$\mathrm{li}(x)/\pi_{\rm RVM}(x)$")
ax[0].axhline(1.0, color="0.5", lw=0.8, ls="--")
ax[0].set_xlabel("x")
ax[0].set_title(r"The three ratios on $[10,1000]$")
ax[0].legend(fontsize=8)
ax[1].plot(x[s8], e_pi, "b-", lw=0.6, label=r"$(\pi-\mathrm{li})/B$")
```

```
ax[1].plot(x[s8], e_rvm, "r-", lw=1.4, label=r"$(\pi_{\rm RVM}-\mathrm{li})/B$")
ax[1].axhline(0.0, color="0.5", lw=0.8, ls="--")
ax[1].set_xlabel("x")
ax[1].set_title(r"Square-root scale $B(x)=\sqrt{x}\,(\log\log x)^2/\log x$")
ax[1].legend(fontsize=8)
fig.tight_layout()
savefig(fig, "fig10_prime_ratios.png")
say.close()
```

A.8. **s07_product_identity.py.**

```
"""Experiment 7: the value at infinity of the endpoint.

varpi_Z1 = lim_{sigma -> +inf} zeta_Z1(sigma) = varpi_RVM * prod_k (1/4 + tau_k^2) / (1/4 + gamma_k^2),
so zeta_Z1 = zeta holds exactly when the ordered partial products
P_K = prod_{k <= K} (1/4 + gamma_k^2) / (1/4 + tau_k^2)
converge to varpi_RVM.  The partial products are computed from data/zeross.bin (gamma_k) and
data/zLamb.bin (tau_k) and compared with varpi_RVM from s02_transition.py."""
import numpy as np
import matplotlib
matplotlib.use("Agg")
import matplotlib.pyplot as plt
from rvm_common import Tee, load_ordinates, savefig, write_csv

say = Tee("s07_product_identity")
gam, tau = load_ordinates()
VARPI = 0.9619415807          # Richardson value of varpi_RVM (s02_transition.py)
N = gam.size
logP = np.cumsum(np.log1p(0.25 / gam ** 2) + 2 * np.log(gam) - np.log1p(0.25 / tau ** 2) - 2 * np.log(tau))
P = np.exp(logP)
say("N = %d ordinates; varpi_RVM = %.10f" % (N, VARPI))
say("%10s %16s %14s" % ("K", "P_K", "P_K - varpi"))
for K in [10, 100, 1000, 10 ** 4, 10 ** 5, 10 ** 6, N]:
    say("%10d %16.10f %14.3e" % (K, P[K - 1], P[K - 1] - VARPI))
for K0 in [10 ** 4, 10 ** 5, 10 ** 6]:
    seg = P[K0 - 1:]
    say("K >= %7d: min P_K = %.10f, max P_K = %.10f, mean = %.10f" % (K0, seg.min(), seg.max(), seg.mean()))
# the flow along the family: Upsilon(lam) = sum_k 2 gamma_k(lam) eta_k / (1/4 + gamma_k(lam)^2),
# eta_k = tau_k - gamma_k, gamma_k(lam) = (1 - lam) gamma_k + lam tau_k; d/dlam log varpi_lam = -Upsilon(lam)
eta = tau - gam
say("")
say("flow along the lambda-family")
for lam in [0.0, 0.25, 0.5, 0.75, 1.0]:
    gl = (1 - lam) * gam + lam * tau
    say("Upsilon(%.2f) = %.10f" % (lam, np.sum(2 * gl * eta / (0.25 + gl ** 2))))
say("sum_k 2 eta_k^2 / gamma_k^2 = %.10f" % np.sum(2 * eta ** 2 / gam ** 2))
total = np.sum(np.log((0.25 + tau ** 2) / (0.25 + gam ** 2)))
say("integral_0^1 Upsilon = sum_k log((1/4+tau_k^2)/(1/4+gamma_k^2)) = %.10f" % total)
say("log L = log(1/varpi_RVM)                                = %.10f" % np.log(1 / VARPI))
step = 50
idx = np.arange(step - 1, N, step)
write_csv("s07_partial_products.csv", ["K", "P_K"], [idx + 1, P[idx]])

fig, ax = plt.subplots(1, 2, figsize=(12, 4))
k1 = np.arange(1, 10 ** 4 + 1)
ax[0].plot(k1, P[:10 ** 4], "r-", lw=0.6, label=r"$P_K$")
ax[0].axhline(VARPI, color="k", lw=0.8, ls="--", label=r"$\varpi_{\rm RVM}$")
ax[0].set_xlabel("K")
ax[0].set_title(r"Partial products $P_K$, $K\leq 10^4$")
ax[0].legend(fontsize=8)
sel = idx >= 10 ** 5 - 1
ax[1].plot(idx[sel] + 1, (P[idx[sel]] - VARPI) * 1e6, "r-", lw=0.6)
ax[1].axhline(0.0, color="k", lw=0.8, ls="--")
ax[1].set_xlabel("K")
ax[1].set_ylabel(r"$(P_K-\varpi_{\rm RVM})\cdot 10^{6}$")
```

```
ax[1].set_title(r"$10^5\leq K\leq %d$" % N)
fig.tight_layout()
savefig(fig, "fig11_product_identity.png")
say.close()
```

A.9. **s08_transport_moments.py.**

```
#!/usr/bin/env python3
"""s08_transport_moments.py
Transport tau -> gamma on the first rung (support of psi inside (0, log 2)).

Input (not recomputed): data/zeross.bin (gamma_k), data/zLamb.bin (tau_k), little-endian float64.

For a fixed normalized smooth bump psi supported in [A,B] subset (0, log 2):
  c_psi^{(j)}(t) = int psi(u) u^j cos(t u + j pi/2) du
  W_psi    = (1/2) int psi(u) ( e^{u/2} - e^{-u/2}/(e^{2u}-1) ) du       (lambda-independent part)
  R_psi(1) = W_psi - sum_k c_psi(tau_k)                                   (clock defect; clock side only)
  S_psi(lambda) = sum_k [ c_psi(gamma_k + lambda*eta_k) - c_psi(tau_k) ]  (transport), eta_k = tau_k - gamma_k
  m_j(psi) = sum_k eta_k^j c_psi^{(j)}(tau_k)                             (clock-sampled moments)
and the partial sums of  S_psi(0) = sum_{j>=1} (-1)^j m_j / j!.
Outputs: output/s08_transport_console.txt, s08_moments.csv, s08_transport_lambda.csv, s08_first_moment_terms.
    csv
"""
import os, math
import numpy as np

K = 20000
A, B = 0.10, 0.62            # support of psi, inside (0, log 2 = 0.6931)
NU = 4001                    # quadrature nodes
JMAX = 12
HERE = os.path.dirname(os.path.abspath(__file__))
OUT = os.path.join(HERE, '..', 'output')
DATA = os.path.join(HERE, '..', 'data')
os.makedirs(OUT, exist_ok=True)

u = np.linspace(A, B, NU)
v = (u - 0.5 * (A + B)) / (0.5 * (B - A))
psi = np.zeros_like(u)
ins = np.abs(v) < 1
psi[ins] = np.exp(-1.0 / (1.0 - v[ins] ** 2))
du = u[1] - u[0]
psi /= psi.sum() * du                      # int psi = 1

def c(t, j=0, chunk=2000):
    t = np.atleast_1d(np.asarray(t, dtype=float))
    w = psi * u ** j * du
    out = np.empty_like(t)
    for a in range(0, len(t), chunk):
        out[a:a + chunk] = np.cos(np.outer(t[a:a + chunk], u) + j * np.pi / 2) @ w
    return out

def main():
    lines = []
    def P(s=''):
        print(s); lines.append(s)
    g_all = np.fromfile(os.path.join(DATA, 'zeross.bin'), dtype='<f8')
    t_all = np.fromfile(os.path.join(DATA, 'zLamb.bin'), dtype='<f8')
    g = g_all[:K]; tau = t_all[:K]; eta = tau - g
    W = 0.5 * (psi * (np.exp(u / 2) - np.exp(-u / 2) / (np.exp(2 * u) - 1))).sum() * du
    c_tau = c(tau); c_gam = c(g)
    R1 = W - c_tau.sum()
    S0 = (c_gam - c_tau).sum()
    P('s08: transport tau -> gamma on the first rung')
    P(f'entries in zeross.bin / zLamb.bin: {len(g_all)} / {len(t_all)};  used K = {K}')
    P(f'gamma_1 = {g[0]:.9f}, tau_1 = {tau[0]:.9f};  gamma_K = {g[-1]:.6f}, tau_K = {tau[-1]:.6f}')
```

```
    P(f'psi: normalized bump on [{A}, {B}] (log 2 = {math.log(2):.6f}), {NU} nodes')
    P(f'max|eta_k| = {np.abs(eta).max():.6f} at k = {np.abs(eta).argmax()+1};  rms eta = {np.sqrt((eta**2).
     mean()):.6f}')
    for k0 in (100, 300, 1000, 3000, 10000, K - 1):
        P(f'  max_(k>={k0:5d}) |c_psi(tau_k)| = {np.abs(c_tau[k0:]).max():.3e}')
    P()
    P(f'W_psi                          = {W:+.12f}')
    P(f'sum_k c_psi(tau_k)            = {c_tau.sum():+.12f}')
    P(f'sum_k c_psi(gamma_k)           = {c_gam.sum():+.12f}')
    P(f'R_psi(1) = W - sum c(tau)      = {R1:+.12f}   (clock defect)')
    P(f'S_psi(0) = sum[c(gam)-c(tau)]  = {S0:+.12f}   (transport)')
    P(f'R_psi(0) = R_psi(1) - S_psi(0) = {R1 - S0:+.3e}')
    P()
    P(' j        m_j(psi)        (-1)^j m_j/j!      partial sum      partial - R_psi(1)')
    rows = []; part = 0.0
    for j in range(1, JMAX + 1):
        mj = (eta ** j * c(tau, j)).sum()
        term = (-1) ** j * mj / math.factorial(j)
        part += term
        rows.append((j, mj, term, part, part - R1))
        P(f'{j:2d}  {mj:+.10e}  {term:+.10e}  {part:+.12f}  {part - R1:+.3e}')
    np.savetxt(os.path.join(OUT, 's08_moments.csv'), np.array(rows), delimiter=',',
               header='j,m_j,term,partial_sum,partial_minus_R1', comments='', fmt=['%d'] + ['%.15e'] * 4)
    # which k carry the first two moments
    t1 = -eta * c(tau, 1); t2 = 0.5 * eta ** 2 * c(tau, 2)
    kk = np.arange(1, K + 1)
    np.savetxt(os.path.join(OUT, 's08_first_moment_terms.csv'), np.c_[kk, tau, eta, c_tau, c_gam - c_tau, t1,
     t2][:400], delimiter=',',
               header='k,tau_k,eta_k,c_psi(tau_k),c_psi(gamma_k)-c_psi(tau_k),-eta_k*c1(tau_k),0.5*eta_k^2*c2(
     tau_k)', comments='',
               fmt=['%d'] + ['%.12e'] * 6)
    P()
    P(' cumulative over k (first K0 ordinates):  K0   sum[c(gam)-c(tau)]   W - sum c(tau)   -m_1 part   +m_2/2
      part')
    for K0 in (1, 2, 3, 5, 10, 20, 50, 100, 200, 400):
        P(f'   {K0:4d}   {(c_gam-c_tau)[:K0].sum():+.10f}   {W - c_tau[:K0].sum():+.10f}   {t1[:K0].sum():+.10
     f}   {t2[:K0].sum():+.10f}')
    P()
    P(' lambda     S_psi(lambda)     R_psi(lambda) = R_psi(1) - S_psi(lambda)')
    lam = np.linspace(-1.0, 2.0, 61); rows = []
    for L in lam:
        S = (c(g + L * eta) - c_tau).sum()
        rows.append((L, S, R1 - S))
    for L, S, R in rows[::5]:
        P(f'{L:+.2f}   {S:+.12f}   {R:+.12f}')
    np.savetxt(os.path.join(OUT, 's08_transport_lambda.csv'), np.array(rows), delimiter=',',
               header='lambda,S_psi,R_psi', comments='', fmt='%.15e')
    with open(os.path.join(OUT, 's08_transport_console.txt'), 'w') as f:
        f.write('\n'.join(lines) + '\n')

if __name__ == '__main__':
    main()
```

A.10. **s09_clock_defect_split.py.**

```
#!/usr/bin/env python3
"""s09_clock_defect_split.py
Splits the clock defect R_psi(1) of s08 (same psi) by Poisson summation on the clock:
  sum_{k>=1} c_psi(tau_k) = I_0 + 2 sum_{m>=1} (-1)^m I_m,
  I_m = int_{t_0}^infty c_psi(t) cos(2 pi m N_RVM(t)) N_RVM'(t) dt,   N_RVM(t_0) = 0,
  I_0 = int psi(u) K_0(u) du,  K_0(u) = -(1/(2 pi u)) [ sin(u t_0) log(t_0/(2 pi)) + pi/2 - Si(u t_0) ].
Reports W_psi, I_0, the smooth mismatch W_psi - I_0 and the oscillating part  -2 sum_{m>=1}(-1)^m I_m
(obtained as R_psi(1) - (W_psi - I_0), with sum_k c_psi(tau_k) read from data/zLamb.bin).
Output: output/s09_clock_defect_split_console.txt
"""
```

```
import os, math
import numpy as np
HERE = os.path.dirname(os.path.abspath(__file__))
OUT = os.path.join(HERE, '..', 'output'); DATA = os.path.join(HERE, '..', 'data')
A, B, NU, K = 0.10, 0.62, 4001, 20000
u = np.linspace(A, B, NU); v = (u - 0.5*(A+B)) / (0.5*(B-A))
psi = np.zeros_like(u); ins = np.abs(v) < 1; psi[ins] = np.exp(-1.0/(1.0 - v[ins]**2))
du = u[1]-u[0]; psi /= psi.sum()*du

def NR(t): return t/(2*np.pi)*np.log(t/(2*np.pi*np.e)) + 7.0/8.0
t0 = 9.7
for _ in range(60): t0 -= NR(t0) / (np.log(t0/(2*np.pi))/(2*np.pi))

def Si(x, n=20001):
    s = np.linspace(0.0, 1.0, n)[None, :] * np.atleast_1d(x)[:, None]
    f = np.ones_like(s); nz = s != 0; f[nz] = np.sin(s[nz]) / s[nz]
    h = np.atleast_1d(x) / (n - 1)
    return (f.sum(axis=1) - 0.5*(f[:, 0] + f[:, -1])) * h

K0 = -(1.0/(2*np.pi*u)) * (np.sin(u*t0)*np.log(t0/(2*np.pi)) + np.pi/2 - Si(u*t0))
I0 = (psi*K0).sum()*du
W = 0.5*(psi*(np.exp(u/2) - np.exp(-u/2)/(np.exp(2*u)-1))).sum()*du
tau = np.fromfile(os.path.join(DATA, 'zLamb.bin'), dtype='<f8')[:K]
w = psi*du; ctau = sum((np.cos(np.outer(tau[a:a+2000], u)) @ w).sum() for a in range(0, K, 2000))
R1 = W - ctau
L = [f's09: Poisson split of the clock defect (psi of s08 on [{A},{B}])',
     f't_0 (N_RVM(t_0)=0)                = {t0:.12f}   (2 pi = {2*np.pi:.6f}, tau_1 = {tau[0]:.6f})',
     f'W_psi                             = {W:+.12f}',
     f'I_0  (m = 0, smooth clock density) = {I0:+.12f}',
     f'sum_k c_psi(tau_k)                = {ctau:+.12f}',
     f'R_psi(1)                          = {R1:+.12f}',
     f'smooth mismatch  W_psi - I_0      = {W - I0:+.12f}',
     f'oscillating part -2 sum_(m>=1)(-1)^m I_m = {R1 - (W - I0):+.12f}',
     f'stationary heights 2 pi e^u, u in supp psi: [{2*np.pi*np.exp(A):.4f}, {2*np.pi*np.exp(B):.4f}];  part
     above t_0: u > {np.log(t0/(2*np.pi)):.6f}']
print('\n'.join(L)); open(os.path.join(OUT, 's09_clock_defect_split_console.txt'), 'w').write('\n'.join(L)+'\n
     ')
```

A.11. **s10_clock_to_zeros.py.**

```
#!/usr/bin/env python3
"""s10_clock_to_zeros.py
Solves, starting from the clock (eta = 0), the system of rung equations

   sum_k c_j(gamma_k) + sum_n a_n psi_j(log n) = W_j ,   j = 1..J,

for the unknown ordinates gamma_k (below a height) and the unknown masses a_n at the log-integers.
  psi_j(u) = w(u) cos(omega_j u), w(u) sin(omega_j u);  w = smooth bump on (0, U), U = log(N + 1/2)
  c_j(t)   = int psi_j(u) cos(t u) du
  W_j      = int psi_j(u) (1/2)( e^{u/2} - e^{-u/2}/(e^{2u}-1) ) du      (lambda-independent part)
Ordinates above the free range are kept on the clock (tau_k from data/zLamb.bin).
The masses enter linearly and are eliminated exactly at every step (ridge RIDGE).
Continuation in height: the band omega <= Omega is enlarged step by step (OMEGAS); at each stage the
ordinates with tau_k <= Omega + MARGIN are free.  Start: gamma_k = tau_k.  Method: Levenberg-Marquardt.
data/zeross.bin is used only AFTER the iteration, to compare.
Outputs: output/s10_clock_to_zeros_console.txt, s10_ordinates.csv, s10_masses.csv
"""
import os, math
import numpy as np
HERE = os.path.dirname(os.path.abspath(__file__))
OUT = os.path.join(HERE, '..', 'output'); DATA = os.path.join(HERE, '..', 'data')
N = 13; U = math.log(N + 0.5)
KT = 6000; DOM = 0.1; NU = 8001; ITER = 400
OMEGAS = [18, 22, 26, 30, 34, 38, 42, 46, 50]; MARGIN = 12.0
WMIN = 5e-2; STEP = 0.2; RIDGE = 1e-10
```

```
u = np.linspace(0.0, U, NU)[1:-1]; du = U / (NU - 1)
v = (u - U/2) / (U/2); w = np.exp(-1.0/(1.0 - v**2)); w /= w.sum()*du
Wfun = 0.5*(np.exp(u/2) - np.exp(-u/2)/(np.exp(2*u) - 1))
ns = np.array([n for n in range(2, N + 1) if np.interp(math.log(n), u, w)/w.max() >= WMIN])
tau = np.fromfile(os.path.join(DATA, 'zLamb.bin'), dtype='<f8')[:KT]
COS_TAU = np.zeros_like(u)                      # sum over all clock ordinates of cos(tau_k u)
for a in range(0, KT, 500): COS_TAU += np.cos(np.outer(u, tau[a:a+500])).sum(axis=1)

lines = []
def P(s=''):
    print(s); lines.append(s)
P('s10: from the clock to the ordinates through the rung equations (continuation in height)')
P(f'N = {N}, U = log({N}+1/2) = {U:.6f}; free masses n = {ns[0]}..{ns[-1]}; start gamma_k = tau_k')
g_all = tau.copy()
for OM in OMEGAS:
    om = np.arange(0.0, OM + 1e-9, DOM)
    Psi = np.vstack([w*np.cos(o*u) for o in om] + [w*np.sin(o*u) for o in om[1:]]) * du
    Wj = Psi @ Wfun
    Pn = np.vstack([np.interp(np.log(ns), u, Psi[j]/du) for j in range(Psi.shape[0])])
    Q = np.linalg.solve(Pn.T @ Pn + RIDGE*np.eye(len(ns)), Pn.T)
    K = int((tau <= OM + MARGIN).sum())
    g = g_all[:K].copy()
    fixed = Psi @ (COS_TAU - np.cos(np.outer(u, tau[:K])).sum(axis=1)) - Wj
    def r0(g): return Psi @ np.cos(np.outer(u, g)).sum(axis=1) + fixed
    def res(g):
        x = r0(g); a = -Q @ x; return x + Pn @ a, a
    r, a = res(g); n0 = np.linalg.norm(r); damp = 1e-3
    for it in range(1, ITER + 1):
        Jg = Psi @ (-u[:, None]*np.sin(np.outer(u, g))); Jg = Jg - Pn @ (Q @ Jg)
        A = Jg.T @ Jg; b = -Jg.T @ r
        while True:
            d = np.linalg.solve(A + damp*np.diag(np.diag(A) + 1e-14), b)
            d *= min(1.0, STEP/max(np.abs(d).max(), 1e-300))
            r2, a2 = res(g + d)
            if np.linalg.norm(r2) < np.linalg.norm(r) or damp > 1e12: break
            damp *= 4
        g = g + d; r, a = r2, a2; damp = max(damp/3, 1e-14)
        if np.abs(d).max() < 1e-11: break
    g_all[:K] = g
    P(f'band omega <= {OM:2d}: {K:2d} free ordinates, {Psi.shape[0]} tests, |residual| {n0:.3e} -> {np.linalg.
     norm(r):.3e} in {it} iterations')
P()
gam = np.fromfile(os.path.join(DATA, 'zeross.bin'), dtype='<f8')[:K]
P('  k      tau_k (start)    solved ordinate    gamma_k (zeross.bin)   solved - gamma_k    tau_k - gamma_k')
for k in range(K):
    P(f'{k+1:3d}   {tau[k]:14.9f}   {g_all[k]:14.9f}   {gam[k]:14.9f}   {g_all[k]-gam[k]:+.3e}   {tau[k]-gam[k
     ]:+.6f}')
P()
def Lam(n):
    for p in range(2, n + 1):
        if n % p == 0:
            m = n
            while m % p == 0: m //= p
            return math.log(p) if m == 1 else 0.0
    return 0.0
P('  n     solved mass a_n     Lambda(n)/(2 sqrt n)     difference     w(log n)/max w')
for i, n in enumerate(ns):
    ex = Lam(int(n))/(2*math.sqrt(n))
    P(f'{n:3d}   {a[i]:+.9f}   {ex:+.9f}   {a[i]-ex:+.3e}   {np.interp(math.log(n), u, w)/w.max():.3e}')
np.savetxt(os.path.join(OUT, 's10_ordinates.csv'), np.c_[np.arange(1, K+1), tau[:K], g_all[:K], gam],
     delimiter=',',
           header='k,tau_k,solved,gamma_k', comments='', fmt=['%d'] + ['%.12f']*3)
np.savetxt(os.path.join(OUT, 's10_masses.csv'), np.c_[ns, a, [Lam(int(n))/(2*math.sqrt(n)) for n in ns]],
     delimiter=',',
```

```
          header='n,solved_mass,Lambda(n)/(2sqrt(n))', comments='', fmt=['%d'] + ['%.12f']*2)
open(os.path.join(OUT, 's10_clock_to_zeros_console.txt'), 'w').write('\n'.join(lines) + '\n')
```

Appendix B. Console output of the programs

The complete console output of each program, as written to `output/*_console.txt`.

B.1. s01_clock_and_family.

```
zeros loaded: N = 2001052, gamma_1 = 14.134725142, gamma_N = 1132490.658714, tau_1 = 14.521346953, tau_N =
    1132490.722588
max |N_RVM(tau_k) - (k - 1/2)| = 6.985e-10
eta_k = tau_k - gamma_k:  max |eta| = 0.993711 (k = 34),  mean |eta| = 0.151209,  rms = 0.188066,  mean eta =
    1.731e-07
      k from         k to     max|eta|      rms eta
           1           10     0.840022     0.545768
          11          100     0.993711     0.453166
         101         1000     0.915450     0.345177
        1001        10000     0.833881     0.274177
       10001       100000     0.785835     0.226352
      100001      1000000     0.762816     0.192411
     1000001      2001052     0.717359     0.178900
first ordinates:
   k          gamma_k            tau_k        eta_k
   1     14.134725142     14.521346953     0.386622
   2     21.022039639     20.655740356    -0.366299
   3     25.010857580     25.492675432     0.481818
   4     30.424876126     29.739411632    -0.685464
   5     32.935061588     33.624531889     0.689470
   6     37.586178159     37.257370087    -0.328808
   7     40.918719012     40.700551604    -0.218167
   8     43.327073281     43.994013953     0.666941
   9     48.005150881     47.165128427    -0.840022
  10     49.773832478     50.233653391     0.459821
  11     52.970321478     53.214422874     0.244101
  12     56.446247697     56.118921948    -0.327326
sigma = 2.00, t in [1,2000] (499 points): max |zeta_Z1/zeta - 1| = 8.706e-10, median = 2.685e-10
sigma = 1.25, t in [1,2000] (499 points): max |zeta_Z1/zeta - 1| = 1.128e-09, median = 3.693e-10
sigma = 0.75, t in [1,2000] (499 points): max |zeta_Z1/zeta - 1| = 2.629e-09, median = 5.179e-10
critical line, t in [0.5,60]: max |Z_Z1(t) - Z(t)| / max(1,|Z(t)|) = 8.370e-10
total run time 30.6 s
```

B.2. s01b_clock_zoom.

```
zoom table near the top of the file:
        k            gamma_k             tau_k      eta_k   N*-N_RVM
  2001041     1132485.303923     1132485.011569  -0.292355  -0.563104
  2001042     1132485.662638     1132485.530752  -0.131886  -0.254025
  2001043     1132486.244063     1132486.049936  -0.194127  -0.373908
  2001044     1132486.392256     1132486.569120   0.176863   0.340656
  2001045     1132486.874881     1132487.088303   0.213422   0.411072
  2001046     1132487.571868     1132487.607487   0.035619   0.068605
  2001047     1132487.950694     1132488.126670   0.175977   0.338949
  2001048     1132488.905328     1132488.645854  -0.259474  -0.499774
  2001049     1132489.162780     1132489.165037   0.002257   0.004348
  2001050     1132489.585215     1132489.684221   0.099006   0.190696
  2001051     1132490.165305     1132490.203404   0.038099   0.073383
  2001052     1132490.658714     1132490.722588   0.063873   0.123026
total run time 0.6 s
```

B.3. **s02_transition.**

```
varpi_RVM (Richardson, sigma = 2, T = 2500, 5000, 10000) = 0.9619415807 -1.28e-10 i
L = 1/varpi_RVM = 1.0395641690
sigma' = 0.25: |zeta_RVM/(varpi(1+w)) - 1|: log-log slope on t>100 = -1.181, max t*err = 0.0617
sigma' = 0.25: |zeta_RVM/(varpi(1+chi)) - 1|: max t*err (t>100) = 0.0790
          |w| * t^sigma' in [1.5832, 1.5832];  |1/zeta_RVM - L| at t = 1e4: 1.425e-01
sigma' = 0.50: |zeta_RVM/(varpi(1+w)) - 1|: log-log slope on t>100 = -1.181, max t*err = 0.0618
          |w| * t^sigma' in [2.5061, 2.5066];  |1/zeta_RVM - L| at t = 1e4: 2.543e-02
sigma' = 1.50: |zeta_RVM/(varpi(1+w)) - 1|: log-log slope on t>100 = -1.182, max t*err = 0.0620
          |w| * t^sigma' in [15.6619, 15.7496];  |1/zeta_RVM - L| at t = 1e4: 1.598e-05
total run time 11.5 s
```

B.4. **s03_exponential_sums.**

```
     x     max|S_RVM| max|S_RVM|/log    |S_zeta(N)| Landau -T/2pi*L/sqrt(x)   |M_H^RVM(x)|
   2.0         3.5745         0.2695       88300.77        -88341.59      1.722e-06
   3.0         3.9750         0.3759      114267.92       -114324.31      1.175e-06
   4.0         3.9523         0.3732       62435.29        -62466.94      8.004e-07
   6.0         4.5264         0.4547           0.26            -0.00      2.262e-06
  10.0         9.6989         1.1495           2.42            -0.00      4.031e-06
total run time 1.7 s
```

B.5. **s04_primes.**

```
x <= 100, all 2001052 zeros of each list, 3000 grid points:
  max |pi_from_zeta_zeros - pi| away from primes (distance > 0.2): 0.4035
  pi_RVM(100) = 26.0782, pi(100) = 25, li(100) = 30.1261, J_RVM(100) = 29.5889, J(100) = 28.5333
  pi_RVM monotone on [2,100]: False (min increment on the grid -2.528e-01)
x <= 1000, smooth cut at K = 17061 zeros (tau_K = 15707.6), 40000 grid points:
  pi_RVM(1000) = 169.9750, pi(1000) = 168, li(1000) = 177.6097
cell masses c_n = pi_RVM(n) - pi_RVM(n-1):
    n        c_n    1/log n    prime
    2    0.52944    1.44270        1
    3    1.04188    0.91024        1
    4    0.53593    0.72135        0
    5    0.46739    0.62133        1
    6    0.36089    0.55811        0
    7    0.63254    0.51390        1
    8    0.62393    0.48090        0
    9   -0.37669    0.45512        0
   10    0.17151    0.43429        0
   11    0.69779    0.41703        1
   12    0.32703    0.40243        0
   13    0.36227    0.38987        1
   14    0.57662    0.37892        0
   15    0.19671    0.36927        0
   16   -0.05284    0.36067        0
   17    0.39958    0.35296        1
   18    0.76865    0.34598        0
   19    0.43500    0.33962        1
   20   -0.03342    0.33381        0
   21    0.03009    0.32846        0
   22    0.37466    0.32352        0
   23    0.40277    0.31893        1
   24    0.11768    0.31466        0
   25    0.01006    0.31067        0
   26    0.27350    0.30693        0
   27    0.55508    0.30341        0
   28    0.47627    0.30010        0
   29    0.12577    0.29697        1
   30   -0.08367    0.29401        0
   31    0.07329    0.29121        1
   32    0.38040    0.28854        0
   33    0.43927    0.28600        0
   34    0.34619    0.28358        0
   35    0.06419    0.28127        0
```

```
   36   -0.02475    0.27906        0
   37    0.15448    0.27694        1
   38    0.40992    0.27491        0
   39    0.49628    0.27296        0
   40    0.35226    0.27109        0
   41    0.14030    0.26928        1
sum_{n<=1000} c_n = 169.9750,  pi(1000) = 168
real RVM prime levels p_j^RVM (first crossing of level j) versus p_j:
    j      p_j^RVM      p_j       diff
    1       2.2139        2     0.2139
    2       3.1550        3     0.1550
    3       5.1887        5     0.1887
    4       7.1593        7     0.1593
    5      11.1406       11     0.1406
    6      14.0208       13     1.0208
    7      17.2552       17     0.2552
    8      19.1781       19     0.1781
    9      26.0536       23     3.0536
   10      28.0447       29    -0.9553
   11      33.0707       31     2.0707
   12      38.0931       37     1.0931
   13      40.2282       41    -0.7718
   14      44.1861       43     1.1861
   15      46.1221       47    -0.8779
   16      57.1779       53     4.1779
   17      59.1201       59     0.1201
   18      61.0809       61     0.0809
   19      67.0384       67     0.0384
   20      74.0931       71     3.0931
   21      76.1235       73     3.1235
   22      83.0822       79     4.0822
   23      86.0513       83     3.0513
   24      92.0825       89     3.0825
   25      94.0781       97    -2.9219
   26      96.1319      101    -4.8681
   27     103.2219      103     0.2219
   28     112.1163      107     5.1163
   29     115.0277      109     6.0277
   30     118.1431      113     5.1431
total run time 107.7 s
```

B.6. **s05_mertens.**

```
zeta_RVM'(rho_k) exact for k <= 1000; zeta_RVM'(rho_k)/log(tau_k/2pi) at k = 1000: 0.9619388429 -1.41e-05 i
varpi_RVM = 0.9619415807; fit zeta_RVM'(rho_k) = varpi log(tau_k/2pi)(1 + c/tau_k), c = -0.0052 -0.0208 i
Riemann-Siegel zeta'(rho) versus mpmath at k = 1001..1050: max relative difference 1.19e-05
median |zeta'(rho)| = 7.8899;  spectral sums 2 sum|a|^2: zeta 0.029034, RVM 0.027203
M(10^6) = 212;  squarefree density Q(10^6)/10^6 = 0.607926 (6/pi^2 = 0.607927)
explicit formula with 2001052 zeros of zeta on [2,2000]: rms(M_zeta - M) = 0.0005 off integers (|x-n|>0.05)
target on 49896 grid points; at 100 integers <= 1e5: max |M_RVM(n) (all zeros) - target(n)| = 0.0191
ternary walk: max |M_h - target| on [1,1e6] = 2.6155, on [1e4,1e6] = 1.6539
densities of (-1, 0, +1):  mu   0.303857 0.392074 0.304069
                           mu_h 0.304003 0.392074 0.303923   (3/pi^2 = 0.303964, 1 - 6/pi^2 = 0.392073)
sign agreement mu_h = mu on squarefrees: 0.500344;  sum mu mu_h / Q = 0.000688
max |M(n)|/sqrt(n) on [10,1e6] = 0.8321;  max |M_RVM(n)|/sqrt(n) = 0.7541;  max |M_h(n)|/sqrt(n) = 0.6882

NUMERICAL EXAMPLE OF THE TERNARY DIVISION (first 40 integers)
   n   mu   mu^2     target M_h(n-1)   sign   mu_h    M_h      M
   1    1      1     0.2925        0     +1      1      1      1
   2   -1      1     0.4563        1     -1     -1      0      0
   3   -1      1    -0.6523        0     -1     -1     -1     -1
   4    0      0    -0.8720       -1      -      0     -1     -1
   5   -1      1    -1.7255       -1     -1     -1     -2     -2
   6    1      1    -1.3063       -2     +1      1     -1     -1
   7   -1      1    -1.5931       -1     -1     -1     -2     -2
```

```
   8    0      0    -2.4029       -2      -      0     -2     -2
   9    0      0    -1.3292       -2      -      0     -2     -2
  10    1      1    -1.3280       -2     +1      1     -1     -1
  11   -1      1    -2.3272       -1     -1     -1     -2     -2
  12    0      0    -2.2067       -2      -      0     -2     -2
  13   -1      1    -1.8468       -2     +1      1     -1     -3
  14    1      1    -1.9324       -1     -1     -1     -2     -2
  15    1      1    -1.5435       -2     +1      1     -1     -1
  16    0      0    -1.0414       -1      -      0     -1     -1
  17   -1      1    -1.6970       -1     -1     -1     -2     -2
  18    0      0    -3.0095       -2      -      0     -2     -2
  19   -1      1    -3.2869       -2     -1     -1     -3     -3
  20    0      0    -2.2604       -3      -      0     -3     -3
  21    1      1    -1.2515       -3     +1      1     -2     -2
  22    1      1    -1.1120       -2     +1      1     -1     -1
  23   -1      1    -1.3607       -1     -1     -1     -2     -2
  24    0      0    -1.3669       -2      -      0     -2     -2
  25    0      0    -1.3810       -2      -      0     -2     -2
  26    1      1    -1.9805       -2     +1      1     -1     -1
  27    0      0    -3.0141       -1      -      0     -1     -1
  28    0      0    -3.6155       -1      -      0     -1     -1
  29   -1      1    -3.2066       -1     -1     -1     -2     -2
  30   -1      1    -2.1599       -2     -1     -1     -3     -3
  31   -1      1    -1.3649       -3     +1      1     -2     -4
  32    0      0    -1.2973       -2      -      0     -2     -4
  33    1      1    -1.6435       -2     +1      1     -1     -3
  34    1      1    -1.7783       -1     -1     -1     -2     -2
  35    1      1    -1.4711       -2     +1      1     -1     -1
  36    0      0    -1.0723       -1      -      0     -1     -1
  37   -1      1    -1.0774       -1     -1     -1     -2     -2
  38    1      1    -1.5897       -2     +1      1     -1     -1
  39    1      1    -2.2227       -1     -1     -1     -2      0
  40    0      0    -2.4881       -2      -      0     -2      0

ENERGY E_N = (1/N) sum_{n<=N} W(n)^2
         N         E_N(M)     E_N(M_RVM)       E_N(M_h)
      1000        14.6400        18.6280        18.5960
     10000       163.6747       195.6965       196.0683
    100000      1622.3189      1308.3041      1308.8186
   1000000     14851.0857     12537.9008     12538.1174
total run time 148.2 s
```

B.7. **s06_prime_ratios.**

```
grid: 40000 points on [2, 1000]; pi_RVM from the first 17061 ordinates tau_k (smooth cut)

ratios at selected x
       x           pi       pi_RVM           li    li/pi_RVM        pi/li    pi/pi_RVM
    10.0       4.0000       3.9876       6.1657     1.546218     0.648747     1.003105
    20.0       8.0000       7.6636       9.9052     1.292496     0.807657     1.043893
    50.0      15.0000      14.7686      18.4695     1.250592     0.812151     1.015670
   100.0      25.0000      26.0813      30.1265     1.155098     0.829835     0.958541
   200.0      46.0000      43.3656      50.1910     1.157393     0.916499     1.060749
   500.0      95.0000      93.7489     101.7975     1.085854     0.933225     1.013346
  1000.0     168.0000     169.9750     177.6097     1.044916     0.945895     0.988381

square-root scale B(x) = sqrt(x) (log log x)^2 / log x on [8, 1000]
max |pi_RVM - li| / B = 2.7213   (at x = 9.73)
max |pi - li| / B     = 2.4490   (at x = 11.00)
max |pi_RVM - li| / B on [50, 1000] = 1.3455
max |pi_RVM - li| / B on [500, 1000] = 0.7216
total run time 1.2 s
```

B.8. **s07_product_identity.**

```
N = 2001052 ordinates; varpi_RVM = 0.9619415807
         K              P_K    P_K - varpi
        10     0.9625640008      6.224e-04
       100     0.9624787401      5.372e-04
      1000     0.9622891707      3.476e-04
     10000     0.9619513787      9.798e-06
    100000     0.9619301865     -1.139e-05
   1000000     0.9619418189      2.382e-07
   2001052     0.9619415390     -4.166e-08
K >=   10000: min P_K = 0.9618023968, max P_K = 0.9620641927, mean = 0.9619418623
K >=  100000: min P_K = 0.9619210208, max P_K = 0.9619605076, mean = 0.9619417591
K >= 1000000: min P_K = 0.9619389488, max P_K = 0.9619439825, mean = 0.9619416633

flow along the lambda-family
Upsilon(0.00) = 0.0436949578
Upsilon(0.25) = 0.0412445375
Upsilon(0.50) = 0.0387987180
Upsilon(0.75) = 0.0363572218
Upsilon(1.00) = 0.0339197739
sum_k 2 eta_k^2 / gamma_k^2 = 0.0098210688
integral_0^1 Upsilon = sum_k log((1/4+tau_k^2)/(1/4+gamma_k^2)) = 0.0388016004
log L = log(1/varpi_RVM)                                  = 0.0388015571
total run time 1.5 s
```

B.9. **s08_transport_moments.**

```
s08: transport tau -> gamma on the first rung
entries in zeross.bin / zLamb.bin: 2001052 / 2001052;  used K = 20000
gamma_1 = 14.134725142, tau_1 = 14.521346953;  gamma_K = 18046.464296, tau_K = 18046.544665
psi: normalized bump on [0.1, 0.62] (log 2 = 0.693147), 4001 nodes
max|eta_k| = 0.993711 at k = 34;  rms eta = 0.266395
  max_(k>=  100) |c_psi(tau_k)| = 6.880e-05
  max_(k>=  300) |c_psi(tau_k)| = 7.263e-07
  max_(k>= 1000) |c_psi(tau_k)| = 2.527e-10
  max_(k>= 3000) |c_psi(tau_k)| = 2.074e-13
  max_(k>=10000) |c_psi(tau_k)| = 2.074e-13
  max_(k>=19999) |c_psi(tau_k)| = 1.616e-15

W_psi                          = +0.133643530916
sum_k c_psi(tau_k)             = +0.176106813076
sum_k c_psi(gamma_k)           = +0.133643530929
R_psi(1) = W - sum c(tau)      = -0.042463282161   (clock defect)
S_psi(0) = sum[c(gam)-c(tau)]  = -0.042463282147   (transport)
R_psi(0) = R_psi(1) - S_psi(0) = -1.376e-11

 j        m_j(psi)        (-1)^j m_j/j!      partial sum      partial - R_psi(1)
 1  +4.0395231784e-02  -4.0395231784e-02  -0.040395231784  +2.068e-03
 2  -4.8097910277e-03  -2.4048955139e-03  -0.042800127298  -3.368e-04
 3  -2.0431952768e-03  +3.4053254613e-04  -0.042459594751  +3.687e-06
 4  -5.5184329118e-05  -2.2993470466e-06  -0.042461894098  +1.388e-06
 5  +1.6734348340e-04  -1.3945290283e-06  -0.042463288628  -6.467e-09
 6  +5.5720624242e-06  +7.7389755891e-09  -0.042463280889  +1.272e-09
 7  +7.2607702259e-06  -1.4406290131e-09  -0.042463282329  -1.686e-10
 8  +6.4039357525e-06  +1.5882777164e-10  -0.042463282170  -9.773e-12
 9  -8.6516463634e-06  +2.3841618065e-11  -0.042463282147  +1.407e-11
10  -2.6791658297e-06  -7.3830628022e-13  -0.042463282147  +1.333e-11
11  +2.8798049874e-06  -7.2145186673e-14  -0.042463282147  +1.326e-11
12  +8.0540801176e-07  +1.6814307338e-15  -0.042463282147  +1.326e-11

 cumulative over k (first K0 ordinates):  K0   sum[c(gam)-c(tau)]   W - sum c(tau)   -m_1 part   +m_2/2 part
      1   -0.0208609181   +0.0155048033   -0.0168182977   -0.0039851021
      2   -0.0179065615   +0.0336283174   -0.0153170893   -0.0024894395
      3   -0.0230428015   -0.0613123463   -0.0185197872   -0.0044492860
      5   -0.0438112092   -0.0723587090   -0.0422769516   -0.0020012704
     10   -0.0464863677   -0.0403092255   -0.0440974925   -0.0029001855
```

```
     20   -0.0406060545   -0.0470689184   -0.0387914922   -0.0021299305
     50   -0.0422141654   -0.0432549118   -0.0401314609   -0.0024151466
    100   -0.0424897738   -0.0426540294   -0.0404216192   -0.0024058875
    200   -0.0424647471   -0.0424770847   -0.0403966024   -0.0024050814
    400   -0.0424633643   -0.0424636133   -0.0403953106   -0.0024049002

 lambda     S_psi(lambda)     R_psi(lambda) = R_psi(1) - S_psi(lambda)
-1.00   -0.087766836871   +0.045303554710
-0.75   -0.076275893121   +0.033812610960
-0.50   -0.064876726912   +0.022413444752
-0.25   -0.053596431977   +0.011133149816
+0.00   -0.042463282147   -0.000000000014
+0.25   -0.031506572650   -0.010956709511
+0.50   -0.020756460380   -0.021706821780
+0.75   -0.010243803437   -0.032219478724
+1.00   +0.000000000000   -0.042463282161
+1.25   +0.009943173538   -0.052406455698
+1.50   +0.019553725448   -0.062017007609
+1.75   +0.028799612928   -0.071262895088
+2.00   +0.037648908219   -0.080112190380
```

B.10. **s09_clock_defect_split.**

```
s09: Poisson split of the clock defect (psi of s08 on [0.1,0.62])
t_0 (N_RVM(t_0)=0)                = 9.676906787166   (2 pi = 6.283185, tau_1 = 14.521347)
W_psi                             = +0.133643530916
I_0  (m = 0, smooth clock density) = +0.063533498702
sum_k c_psi(tau_k)                = +0.176106813076
R_psi(1)                          = -0.042463282161
smooth mismatch  W_psi - I_0      = +0.070110032214
oscillating part -2 sum_(m>=1)(-1)^m I_m = -0.112573314375
stationary heights 2 pi e^u, u in supp psi: [6.9440, 11.6800];  part above t_0: u > 0.431865
```

B.11. **s10_clock_to_zeros.**

```
s10: from the clock to the ordinates through the rung equations (continuation in height)
N = 13, U = log(13+1/2) = 2.602690; free masses n = 2..11; start gamma_k = tau_k
band omega <= 18:  4 free ordinates, 361 tests, |residual| 8.136e-01 -> 3.892e-04 in 37 iterations
band omega <= 22:  5 free ordinates, 441 tests, |residual| 1.824e-02 -> 1.520e-03 in 20 iterations
band omega <= 26:  6 free ordinates, 521 tests, |residual| 1.985e-02 -> 8.002e-04 in 25 iterations
band omega <= 30:  7 free ordinates, 601 tests, |residual| 2.363e-02 -> 2.506e-03 in 21 iterations
band omega <= 34:  8 free ordinates, 681 tests, |residual| 4.635e-02 -> 2.827e-03 in 47 iterations
band omega <= 38:  9 free ordinates, 761 tests, |residual| 3.868e-02 -> 1.683e-03 in 11 iterations
band omega <= 42: 11 free ordinates, 841 tests, |residual| 1.881e-01 -> 1.595e-03 in 11 iterations
band omega <= 46: 12 free ordinates, 921 tests, |residual| 1.878e-02 -> 4.591e-03 in 12 iterations
band omega <= 50: 14 free ordinates, 1001 tests, |residual| 5.189e-02 -> 2.885e-03 in 22 iterations

  k      tau_k (start)    solved ordinate    gamma_k (zeross.bin)   solved - gamma_k    tau_k - gamma_k
  1     14.521346953     14.134660056     14.134725142   -6.509e-05   +0.386622
  2     20.655740356     21.022091631     21.022039639   +5.199e-05   -0.366299
  3     25.492675432     25.010827129     25.010857580   -3.045e-05   +0.481818
  4     29.739411632     30.424917668     30.424876126   +4.154e-05   -0.685464
  5     33.624531889     32.935072953     32.935061588   +1.137e-05   +0.689470
  6     37.257370087     37.586146548     37.586178159   -3.161e-05   -0.328808
  7     40.700551604     40.918919600     40.918719012   +2.006e-04   -0.218167
  8     43.994013953     43.327322117     43.327073281   +2.488e-04   +0.666941
  9     47.165128427     48.004716753     48.005150881   -4.341e-04   -0.840022
 10     50.233653391     49.773657726     49.773832478   -1.748e-04   +0.459821
 11     53.214422874     52.972535790     52.970321478   +2.214e-03   +0.244101
 12     56.118921948     56.548052093     56.446247697   +1.018e-01   -0.327326
 13     58.956266008     58.576523603     59.347044003   -7.705e-01   -0.390778
 14     61.733838843     62.561880181     60.831778525   +1.730e+00   +0.902060

  n     solved mass a_n     Lambda(n)/(2 sqrt n)     difference     w(log n)/max w
  2   +0.245076213   +0.245064536   +1.168e-05   7.562e-01
  3   +0.317145240   +0.317142050   +3.190e-06   9.754e-01
```

```
 4   +0.173268937   +0.173286795   -1.786e-05   9.957e-01
 5   +0.359884977   +0.359881258   +3.719e-06   9.423e-01
 6   -0.000004422   +0.000000000   -4.422e-06   8.474e-01
 7   +0.367785301   +0.367742452   +4.285e-05   7.225e-01
 8   +0.122615622   +0.122532268   +8.335e-05   5.733e-01
 9   +0.183143344   +0.183102048   +4.130e-05   4.062e-01
10   -0.000286111   +0.000000000   -2.861e-04   2.344e-01
11   +0.362276961   +0.361496314   +7.806e-04   8.642e-02
```

References

[1] L. V. Ahlfors, *Complex Analysis*, 3rd ed., McGraw–Hill, New York, 1979. (Cited on p. 67.)

[2] P. T. Bateman and H. G. Diamond, Asymptotic distribution of Beurling's generalized prime numbers, in: *Studies in Number Theory*, MAA Studies in Mathematics 6, Mathematical Association of America, 1969. (Cited on p. 8.)

[3] J. J. Benedetto, A. M. Powell and Ö. Yılmaz, Sigma-delta ($\Sigma\Delta$) quantization and finite frames, *IEEE Trans. Inform. Theory* **52** (2006), no. 5, 1990–2005. doi:10.1109/TIT.2006.872849. (Cited on p. 10.)

[4] A. E. Ingham, Some trigonometrical inequalities with applications to the theory of series, *Math. Z.* **41** (1936), 367–379. (Cited on p. 71.)

[5] A. Beurling, Analyse de la loi asymptotique de la distribution des nombres premiers généralisés. I, *Acta Math.* **68** (1937), 255–291. doi:10.1007/BF02546666. (Cited on pp. 8, 9, and 42.)

[6] S. Bochner and K. Chandrasekharan, On Riemann's functional equation, *Ann. of Math. (2)* **63** (1956), 336–360. doi:10.2307/1969614. (Cited on pp. 9, 49, and 76.)

[7] F. Broucke, G. Debruyne and J. Vindas, Beurling integers with RH and large oscillation, *Adv. Math.* **370** (2020), 107240. doi:10.1016/j.aim.2020.107240. (Cited on p. 9.)

[8] F. Broucke and J. Vindas, A new generalized prime random approximation procedure and some of its applications, *Math. Z.* **307** (2024), Paper No. 62. doi:10.1007/s00209-024-03526-4. (Cited on pp. 9, 42, and 43.)

[9] H. Cohn and N. Elkies, New upper bounds on sphere packings I, *Ann. of Math. (2)* **157** (2003), no. 2, 689–714. doi:10.4007/annals.2003.157.689. (Cited on p. 58.)

[10] R. M. Corless, G. H. Gonnet, D. E. G. Hare, D. J. Jeffrey and D. E. Knuth, On the Lambert W function, *Adv. Comput. Math.* **5** (1996), 329–359. doi:10.1007/BF02124750. (Cited on pp. 8 and 14.)

[11] I. Daubechies and R. DeVore, Approximating a bandlimited function using very coarsely quantized data: a family of stable sigma-delta modulators of arbitrary order, *Ann. of Math.* **158** (2003), 679–710. doi:10.4007/annals.2003.158.679. (Cited on p. 10.)

[12] H. Davenport, *Multiplicative Number Theory*, 2nd ed., revised by H. L. Montgomery, Graduate Texts in Mathematics 74, Springer, New York, 1980. doi:10.1007/978-1-4757-5927-3. (Cited on pp. 20, 29, 30, and 36.)

[13] C.-J. de la Vallée Poussin, Recherches analytiques sur la théorie des nombres premiers, *Ann. Soc. Sci. Bruxelles* **20** (1896), 183–256. (Cited on p. 20.)

[14] H. G. Diamond, Asymptotic distribution of Beurling's generalized integers, *Illinois J. Math.* **14** (1970). doi:10.1215/ijm/1256053295. (Cited on p. 8.)

[15] H. G. Diamond, H. L. Montgomery and U. M. A. Vorhauer, Beurling primes with large oscillation, *Math. Ann.* **334** (2006), 1–36. doi:10.1007/s00208-005-0638-2. (Cited on pp. 9, 42, and 43.)

[16] H. G. Diamond and W.-B. Zhang, *Beurling Generalized Numbers*, Mathematical Surveys and Monographs 213, American Mathematical Society, Providence, RI, 2016. doi:10.1090/surv/213. (Cited on pp. 9, 42, and 43.)

[17] NIST Digital Library of Mathematical Functions, F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier et al. (eds.), §4.13 (Lambert W-function) and §5.11 (Stirling series). `https://dlmf.nist.gov/`. (Cited on pp. 8, 14, and 25.)

[18] H. M. Edwards, *Riemann's Zeta Function*, Academic Press, New York, 1974. (Cited on pp. 8, 9, 17, and 48.)

[19] G. França and A. LeClair, Statistical and other properties of Riemann zeros based on an explicit equation for the n-th zero on the critical line, arXiv:1307.8395. doi:10.48550/arXiv.1307.8395. (Cited on p. 8.)

[20] E. Grosswald and F. J. Schnitzer, A class of modified ζ and L-functions, *Pacific J. Math.* **74** (1978), no. 2, 357–364. doi:10.2140/pjm.1978.74.357. (Cited on pp. 9 and 22.)

[21] C. S. Güntürk and N. T. Thao, Ergodic dynamics in sigma-delta quantization: tiling invariant sets and spectral analysis of error, arXiv:math/0308223. doi:10.48550/arXiv.math/0308223. (Cited on p. 10.)

[22] J. Hadamard, Sur la distribution des zéros de la fonction $\zeta(s)$ et ses conséquences arithmétiques, *Bull. Soc. Math. France* **24** (1896), 199–220. doi:10.24033/bsmf.545. (Cited on pp. 8, 20, and 30.)

[23] H. Hamburger, Über die Riemannsche Funktionalgleichung der ζ-Funktion, *Math. Z.* **10** (1921), 240–254. doi:10.1007/BF01211612. (Cited on pp. 8, 9, 10, 34, 48, and 49.)

[24] J. Kaczorowski and A. Perelli, On the structure of the Selberg class, I: $0 \le d \le 1$, *Acta Math.* **182** (1999), 207–241. doi:10.1007/BF02392574. (Cited on pp. 9 and 49.)

[25] M. I. Knopp, On Dirichlet series satisfying Riemann's functional equation, *Invent. Math.* **117** (1994), 361–372. doi:10.1007/BF01232248. (Cited on pp. 9 and 49.)
[26] P. Kurasov and P. Sarnak, Stable polynomials and crystalline measures, *J. Math. Phys.* **61** (2020), no. 8, 083501. doi:10.1063/5.0012286. (Cited on p. 78.)
[27] M. L. Lapidus and M. van Frankenhuysen, *Fractal Geometry and Number Theory: Complex Dimensions of Fractal Strings and Zeros of Zeta Functions*, Birkhäuser, Boston, 2000. doi:10.1007/978-1-4612-5314-3. (Cited on pp. 9 and 43.)
[28] N. Lev and A. Olevskii, Quasicrystals and Poisson's summation formula, *Invent. Math.* **200** (2015), no. 2, 585–606. doi:10.1007/s00222-014-0542-z. (Cited on pp. 58 and 95.)
[29] A. Olevskii and A. Ulanovskii, Fourier quasicrystals with unit masses, *C. R. Math. Acad. Sci. Paris* **358** (2020), no. 11–12, 1207–1211. doi:10.5802/crmath.142. (Cited on p. 78.)
[30] A. M. Odlyzko and H. J. J. te Riele, Disproof of the Mertens conjecture, *J. reine angew. Math.* **357** (1985), 138–160. doi:10.1515/crll.1985.357.138. (Cited on p. 10.)
[31] T. Nakamura, Dirichlet series with periodic coefficients, Riemann's functional equation and real zeros of Dirichlet L-functions, preprint, arXiv:2008.02570 (2020). doi:10.48550/arXiv.2008.02570. (Cited on p. 49.)
[32] A. Perelli, Converse theorems: from the Riemann zeta function to the Selberg class, arXiv:1605.02354. doi:10.48550/arXiv.1605.02354. (Cited on p. 10.)
[33] D. J. Platt and T. S. Trudgian, The Riemann hypothesis is true up to $3 \cdot 10^{12}$, *Bull. London Math. Soc.* **53** (2021); arXiv:2004.09765. doi:10.48550/arXiv.2004.09765. (Cited on p. 43.)
[34] B. Riemann, Über die Anzahl der Primzahlen unter einer gegebenen Grösse, *Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin* (1859), 671–680. (Cited on pp. 8 and 9.)
[35] C. L. Siegel, Bemerkung zu einem Satz von Hamburger über die Funktionalgleichung der Riemannschen Zetafunktion, *Math. Ann.* **86** (1922), 276–279. doi:10.1007/BF01457989. (Cited on pp. 9 and 49.)
[36] E. C. Titchmarsh, *The Theory of Functions*, 2nd ed., Oxford University Press, 1939. (Cited on pp. 35 and 36.)
[37] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2nd ed., revised by D. R. Heath-Brown, Clarendon Press, Oxford, 1986. (Cited on pp. 8, 9, 10, 14, 19, 21, 24, 30, 31, 34, 35, 38, 49, 55, and 59.)
[38] H. von Mangoldt, Zu Riemanns Abhandlung "Über die Anzahl der Primzahlen unter einer gegebenen Grösse", *J. reine angew. Math.* **114** (1895), 255–305. doi:10.1515/crll.1895.114.255. (Cited on pp. 8 and 14.)

Addendum to v17: what remains live after the interpolation tests

This addendum records the outcome of the subsequent interpolation and finite-section tests. It does not alter any theorem above. Its purpose is to separate the statements that are exact but equivalent to the final step from the one clock-only subproblem that remains autonomous.

Three interpolations and their status. There are three natural ways to interpolate between the midpoint clock and the Riemann zeta function.

The ordinate interpolation. The family ζ_λ of Sections 12–14 moves the ordinates while keeping every nontrivial zero on $\Re s = \frac{1}{2}$. It is an exact analytic family, but for every $0 < \lambda \le 1$ its continuous prime measure is not in the ordinary Dirichlet category. Thus the property that is needed at the endpoint is not preserved along the path; this is exactly the role of condition (D).

The exact-atomic interpolation. One may instead ask for real ordinates κ_k and masses a_n supported at $u = \log n$ throughout an interpolation. Finite sections of this problem were tested numerically after v16. The endpoints can be resolved, whereas the linear interpolation of the target distribution develops a persistent interior residual in the tested sections. This is evidence against that particular interpolation, not a proof that no infinite exact-atomic path exists. In any case the existence of such a path is not supplied by the results of this paper.

The additive interpolation of functions. There is an exact elementary interpolation

$$\mathcal{G}_\varepsilon(s) := (1-\varepsilon)\zeta_{\mathrm{RVM}}(s) + \varepsilon\zeta(s), \qquad 0 \le \varepsilon \le 1. \tag{B.1}$$

Both summands satisfy the same functional equation and reality symmetry, so the same is true of $\mathcal{G}_\varepsilon$. This family exists for every ε, but it does not create a new control quantity: it is a level-set representation of the already defined quotient $\mathcal{Q}_{\mathrm{RVM}} = \zeta/\zeta_{\mathrm{RVM}}$.

Proposition B.1 (The additive family is a level-set reformulation)**.** *Away from the zeros of* ζ_{RVM}*,*

$$\mathcal{G}_\varepsilon(s) = \zeta_{\mathrm{RVM}}(s)\big((1-\varepsilon) + \varepsilon\mathcal{Q}_{\mathrm{RVM}}(s)\big). \tag{B.2}$$

Hence for $0 < \varepsilon < 1$ *every zero of* $\mathcal{G}_\varepsilon$ *which is not a zero of* ζ_{RVM} *lies on the level set*

$$\mathcal{Q}_{\mathrm{RVM}}(s) = -\frac{1-\varepsilon}{\varepsilon}. \tag{B.3}$$

On the critical line $e^{i\vartheta(t)}\mathcal{G}_\varepsilon(\frac{1}{2}+it)$ *is real. At* $\varepsilon = 1$ *the statement that all its nontrivial zeros lie on the critical line is exactly RH.*

Proof. Equation (B.2) is the definition of $\mathcal{Q}_{\mathrm{RVM}}$. The functional equation and the reality statement follow by linearity from those of the two summands. The last assertion is $\mathcal{G}_1 = \zeta$. □

Thus the additive family is useful for visualizing the zero motion, but without an additional monotonicity or flow law it is circular as a route to RH: the terminal level is precisely the original zero problem.

The clock as the first transition term. The transition theorem gives, in every fixed band to the right of the critical line,

$$\zeta_{\mathrm{RVM}}(s) = \varpi_{\mathrm{RVM}}(1+\chi(s))(1+O(|t|^{-1})). \tag{B.4}$$

On $s = \frac{1}{2} + it$, where $\chi(s) = e^{-2i\vartheta(t)}$,

$$e^{i\vartheta(t)}(1+\chi(\tfrac{1}{2}+it)) = 2\cos\vartheta(t). \tag{B.5}$$

Consequently the midpoint clock is asymptotically the zero set selected by the $n = 1$ term of the Hardy–Littlewood/Riemann–Siegel approximate functional equation. This is a structural interpretation of the transition theorem, not an exact identification: the $O(1/t)$ term is visible already in the first ordinates.

For a finite Dirichlet polynomial

$$D_{\varepsilon,M}(s) := 1 + \varepsilon \sum_{2\le n\le M} n^{-s}, \qquad F_{\varepsilon,M}(s) := D_{\varepsilon,M}(s) + \chi(s) D_{\varepsilon,M}(1-s), \tag{B.6}$$

the functional equation is exact. On the critical line,

$$e^{i\vartheta(t)} F_{\varepsilon,M}(\tfrac{1}{2}+it) = 2\Re\Big(e^{i\vartheta(t)} D_{\varepsilon,M}(\tfrac{1}{2}+it)\Big). \tag{B.7}$$

At Gram points the $n=1$ term fixes the sign whenever

$$\varepsilon \sum_{2\le n\le M} n^{-1/2} < 1, \tag{B.8}$$

so in every interval where ϑ is monotone there is a critical-line zero between consecutive Gram points. This dominance statement is only sufficient; it does not count all zeros of $F_{\varepsilon,M}$ and is not used elsewhere in the paper.

What remains genuinely live. After these tests the status is narrower than the programme suggested by the interpolation pictures.

(L1) *The endpoint problem remains the endpoint problem.* Proving that the cell defect vanishes at $\lambda = 0$, equivalently proving (D), remains the only step that closes the main route. The additive family and the exact-atomic interpolation do not simplify this statement.

(L2) *The clock-only injectivity problem is autonomous.* The one subproblem that does not contain the genuine ordinates is the injectivity of the linearized rung system at the midpoint clock. In the meromorphic form of Remark 15.7, the target statement is: if a controlled real sequence (r_k) and a Dirichlet series H satisfy

$$H(z) = \nu + \sum_{k\ge 1} r_k \left(\frac{\tau_k}{z^2+\tau_k^2} - \frac{1}{\tau_k} \right) = \sum_{n\ge 2} \frac{b_n}{\log n}\, n^{-z} \tag{B.9}$$

through a common right half-plane, under the growth hypotheses of the linearization, then $H \equiv 0$. Theorem 15.6 proves the analogous conclusion at an actual solution, while Remark 15.7 explains why it is not automatic for a general list. Equation (B.9) is therefore a closed clock-only rigidity problem, naturally related to Fourier crystalline uniqueness, but by itself it does not prove (D).

(L3) *Bounded finite sections are a numerical existence programme, not yet an analytic route.* Direct finite sections of the endpoint rung system can be solved while constraining both the ordinate displacement and the free masses. If one could prove, for an exhausting nested family of tests, a uniform boundedness and residual-to-zero statement and then pass to a distributional limit, Theorem 15.5 would identify the limit. Such a theorem would already be an existence theorem for the final system and is therefore not a shortcut around (D).

In particular, the individual displacements $\gamma_k - \tau_k$ are not autonomous data. They are used below only for post-solution numerical diagnostics. None of the live statement (B.9), nor the definition of the bounded finite-section problem, requires them as input.

Additional finite-section diagnostics. The post-v16 calculations solve bounded finite sections directly from the midpoint clock. A representative run imposes

$$|\kappa_k - \tau_k| \le 1.2, \qquad |a_n| \le 0.5, \tag{B.10}$$

with the unresolved ordinate tail kept on the midpoint clock. The bounds in (B.10) are numerical regularizations, not hypotheses of any theorem. The genuine ordinates and the von Mangoldt masses are read only after the solve for diagnostics.

With free masses and the bounds (B.10), the sections $N = 20, 34, 50$ have normalized residuals of order 10^{-4}, and the low ordinates and low masses remain close to their diagnostic zeta/von-Mangoldt

values. Cross-validation with test bumps not used in the solve gives residuals of the same order. These calculations remove the earlier pseudoinverse artifact in which unrestricted masses became extremely large.

A second test uses a nested family of permanent test blocks: once a test block is introduced it remains in all later solves. For the bounded sections $N = 50, 70, 100$, the RMS residuals on the old blocks stay small and in the $N = 100$ section are between $2.5 \cdot 10^{-8}$ and $6.1 \cdot 10^{-8}$ on the blocks $N = 13, 20, 34, 50, 70$, while the newest $N = 100$ block has RMS $2.2 \cdot 10^{-6}$. This is computational evidence for local stabilization only; no monotone convergence theorem is claimed.




FIGURE 16. Nested bounded endpoint sections. Each curve shows the residual of one solved section on all permanent test blocks already introduced. The ordinate shifts and free masses are bounded during the solve. This is a finite computation and is not an existence proof for the infinite rung system.

Status. The additive interpolation is exact but is a level-set reformulation of $\mathcal{Q}_{\mathrm{RVM}}$; the exact-atomic interpolation has no established interior path; the bounded finite-section computations do not prove an infinite solution. The only new autonomous analytical target isolated by these tests is the midpoint-clock injectivity problem (B.9). The main endpoint identity (D) remains open.




FIGURE 17. Recovered free masses in the interior $n \leq 15$ of bounded endpoint sections. The von Mangoldt values are shown only as a post-solution diagnostic and are not supplied to the solver; section-boundary masses are omitted from the plot.