THE AREA FUNCTION OF THE MERTENS EXCURSION SKELETON AND HORIZON-LOCALIZED CRITERIA FOR THE RIEMANN HYPOTHESIS

V. CASAÑA-GINER

ABSTRACT. We decompose the Mertens function $M(x)=\sum_{n\le x}\mu(n)$ into its excursions between zero runs and study a regularized triangular skeleton of these excursions, its area curves, and exact quadratic and correlation identities inside each completed excursion.

For any ordered family of disjoint integer intervals $(x_0,x_1]$, elementary counting shows that the members starting at most X with $x_1>x_0^A$ $(A>1)$ are $O_A(\log\log X)$, those with $x_1>cx_0^2$ are $O_c(\log\log X)$, those with $x_1>x_0\psi(x_0)$ are $o(\log X)$ whenever $\psi\to\infty$, and those with $x_1>\lambda x_0$ are at most $\log X/\log\lambda+O(1)$. Combined with the Kaczorowski–Pintz oscillation theorem, this gives a horizon-localization principle: if the excursions starting at most X that belong to an exceptional family or cross a horizon $\mathfrak{h}$ number at most $(\kappa+o(1))\log X$ with $\kappa<\gamma_1/\pi$, then the Riemann hypothesis is equivalent to Littlewood's bound $|M(t)|\ll_\varepsilon t^{1/2+\varepsilon}$ restricted to $x_0<t\le\min\{x_1,\mathfrak{h}(x_0)\}$ on the remaining excursions. The horizon can be lowered from x_0^2 through cx_0^2 and $x_0\psi(x_0)$ to λx_0 for every $\lambda>\lambda_*:=e^{\pi/\gamma_1}$, the exact threshold of this counting mechanism. The same input shows that late excursions end at $x_1=x_0^{1+o(1)}$, so that their length satisfies $\ell=x_1-x_0\le x_0^{1+o(1)}$. On the resulting window every $\theta>\theta_*:=\frac12 e^{\pi/\gamma_1}$ is an admissible prefix-dependency cutoff: with $M(u)=0$ and $u<t\le\lambda u<2u$,

$$M(t)=-\sum_{d\le\lambda u/2}\mu(d)\Big(\Big\lfloor\frac{t}{d}\Big\rfloor-\Big\lfloor\frac{u}{d}\Big\rfloor\Big).$$

After deleting zero increments, a completed excursion is a primitive Dyck path on its fixed support mask. Under the uniform primitive Catalan reference law on that mask, a value $|M(t)|\ge t^{1/2+\varepsilon}$ in an excursion with $x_1\le\lambda x_0$ has probability at most $x_0^{O(1)}e^{-x_0^{2\varepsilon}/2}$, and polynomial lower bounds for the complete-excursion or open-prefix tail weights imply $|M(t)|\ll\sqrt{t\log t}$ on the corresponding ranges.

The one-step recursion concentrates, for every fixed $K\ge2$, all dependence above $\lambda u/(K+1)$ in the $K-1$ channels $M(t/k)-M(u/k)$, $2\le k\le K$, and a persistence lemma quantifies how a large value must be carried by one of them or by the deeper remainder. A multiplicative-layer filtration shows that for every fixed depth k the first k Möbius layers determine the $(k+1)$-st layer of the future window; the determined positions have density zero, while their signed contribution is of order $u(\log\log u)^k/\log u$. Thus substantial cancellation with the remaining layers is necessary to reach the RH scale; no assertion on the number of layers required is made.

Inside a completed excursion, signed lag averages of two-point correlations are forced by the return to zero, while the weighted identity $Q=-\sum_h hC_h$ holds globally at every zero of M with exact cancellation of the cross-excursion terms; this yields an RH criterion in terms of h-weighted correlations at zeros of M, showing that RH is a statement of cancellation between lags. Combining the Fourier-uniform estimate of Matomäki–Teräväinen with Parseval, we obtain uniform two-point Cesàro cancellation in every interval $(u,u+L]$ with $L\ge u^{3/5+\eta}$, averaged over lags $h\le R$ with $R\ge L(\log u)^{-2A}$, $A<1/3$; it applies to all Kaczorowski–Pintz excursions if $\Theta>3/5$, and the geometry of an excursion alone cannot replace this arithmetic input.

Under a uniform sub-Weibull tail hypothesis for a normalized excursion statistic, the counting bounds give extreme-value scales by a union bound and a Borel–Cantelli argument. We also give an exact bilinear representation of M on the square range of an excursion, an exact Mellin decomposition of $1/\zeta$ into completed excursion blocks, and asymptotic decompositions of M into smooth and large-prime parts.

2020 *Mathematics Subject Classification.* Primary 11N37; Secondary 11M26, 60G70.

Key words and phrases. Mertens function, Möbius function, excursions, Riemann hypothesis, Kaczorowski–Pintz oscillation, lacunary sequences, Dyck paths, Catalan tails, extreme values.

Contents

1. Introduction 2
Main results 2
Conventions 4
Notation 4
2. Excursions and the regularized skeleton 5
3. Exact formulas 6
4. Internal correlations of an excursion 7
5. Where ζ enters 9
6. Bounds for the area curves 10
7. Global criteria equivalent to RH 11
8. Lacunary excursion families and horizon-localized criteria 13
8.1. Geometric counts 13
8.2. Arithmetic input: Kaczorowski–Pintz 14
8.3. Horizon localization 15
8.4. Fixed-mask Catalan compression inside fractional square horizons 17
Wide peaks from cancellation in all short intervals 22
8.5. The linear window and the one-step recursion 23
8.6. Conditional extreme-value consequence of excursion sparsity 30
8.7. The square range as an exact bilinear form of the prefix 32
8.8. Mellin mass of a complete excursion 32
9. Two-point correlations and the energy of excursions 33
9.1. Signed lag averages are forced; weighted ones are not 33
9.2. Lag localization of the energy of an excursion 34
9.3. Uniform two-point cancellation in sublinear intervals 35
10. Trial-division horizon of an excursion 36
11. Large-prime decomposition at level U^2 37
12. Local large-prime decomposition of an excursion 39
13. A one-frequency model for the black curve 40
14. Numerical data 41
15. Concluding remarks 41
16. Open problems 41
References 42

1. Introduction

The object of this paper is the Mertens function M, decomposed into excursions between zero runs (Section 2). Two kinds of information about M are separated throughout: what is forced by the ordering and local geometry of the excursions alone, and what requires arithmetic input.

Main results. The central results are in Section 8.

Geometric counts. For any ordered family of disjoint integer intervals $(x_0, x_1]$, the members with $x_1 > x_0^A$ $(A > 1)$ starting below X are $O_A(\log\log X)$, those with $x_1 > cx_0^2$ are $O_c(\log\log X)$, those with $x_1 > x_0\psi(x_0)$ are $o(\log X)$ whenever $\psi \to \infty$, and those with $x_1 > \lambda x_0$ are at most $\log X/\log\lambda + O(1)$ (Lemmas 8.1, 8.2, 8.3, 8.4). No arithmetic enters.

Horizon localization. By Kaczorowski–Pintz [9], if $\Theta > 1/2$ then at least $(\gamma_1/\pi - o(1))\log X$ distinct excursions starting below X contain values of size $t^{\Theta-\eta}$. Theorem 8.11 turns every geometric count below this threshold into an equivalent form of RH: for any horizon h whose crossing family has at

most $(\kappa+o(1))\log X$ members with $\kappa<\gamma_1/\pi$,

$$\mathrm{RH}\iff |M(t)|\ll_\varepsilon t^{1/2+\varepsilon}\quad\text{for } x_0<t\le\min\{x_1,\mathfrak{h}(x_0)\}.$$

In particular (Corollary 8.13)

$$\mathrm{RH}\iff \text{Littlewood's bound holds on } x_0<t\le\min\{x_1,\lambda x_0\},\qquad \lambda>\lambda_*:=e^{\pi/\gamma_1}=1.2488\ldots,$$

and the square horizon x_0^2 is eroded from above through cx_0^2 and $x_0\psi(x_0)$. If RH fails, a positive proportion (on the scale $\log X$) of the Kaczorowski–Pintz excursions carry their large value in $[x_0+\frac12 x_0^{\Theta-\eta},\lambda x_0]$ (Corollary 8.14). The threshold λ_* is the multiplicative distance between consecutive sign changes of $\cos(\gamma_1\log x)$ (Remark 8.29). It is the exact threshold of this counting mechanism, not a proved limit for other localization methods. The local form of the Kaczorowski–Pintz theorem also shows that for every fixed $A>1$ all late excursions satisfy $x_1\le x_0^A$ (Corollary 8.8): the endpoints of late Mertens excursions satisfy $x_1=x_0^{1+o(1)}$, and hence $\ell=x_1-x_0\le x_0^{1+o(1)}$; no lower bound on ℓ follows. The fractional-square lemma makes the intermediate erosion explicit:

$$x_0^2\longrightarrow\frac{x_0^2}{2}\longrightarrow\frac{x_0^2}{4}\longrightarrow\frac{x_0^2}{8}\longrightarrow\cdots\longrightarrow x_0\psi(x_0)\longrightarrow\lambda x_0.$$

For each fixed dyadic level the crossing family is $O(\log\log X)$; the passage to a level depending on x_0 is supplied by the superlinear-horizon lemma, not by a uniform use of the fixed-c estimate.

Kaczorowski–Pintz [9] provide the oscillation-density input. The excursion-counting lemmas, the horizon-localization theorem, the descending-horizon principle, the threshold $\lambda_*=e^{\pi/\gamma_1}$, and the resulting prefix-dependency cutoff $\theta_*=\frac12 e^{\pi/\gamma_1}$ are developed here from that input.

Linear window. For $u<t\le 2u$ one has $\mu(n)=-\sum_{d\mid n,\,d\le u}\mu(d)$. After horizon localization, if $u<t\le\lambda u<2u$, the terms with $d>\lambda u/2$ have no multiple in $(u,t]$, and therefore

$$M(t)=-\sum_{d\le\lambda u/2}\mu(d)\Big(\Big\lfloor\frac{t}{d}\Big\rfloor-\Big\lfloor\frac{u}{d}\Big\rfloor\Big).$$

Thus every $\theta>\theta_*:=\frac12 e^{\pi/\gamma_1}$ is an admissible prefix cutoff for the RH-equivalent excursion window; the convenient choice $\lambda=5/4$ uses only $d\le 5u/8$. This is a descent of dependency, not of size. More sharply, for every fixed $K\ge 2$ the dependence above $\lambda u/(K+1)$ is concentrated in the $K-1$ increments $M(t/k)-M(u/k)$, $2\le k\le K$; for $K=2$ and $\lambda=5/4$, the only channel above $5u/12$ is the increment across $(u/2,t/2]$. The one-step recursion does not by itself give an iterable maximum principle, and a termwise absolute-value contraction gives no nontrivial gain. The exact intra-excursion formulation, the scale cascade, a persistence extraction lemma, and a ballistic counterexample to the relaxed divisor-sum version are given below. A bounded number of multiplicative layers determines only a density-zero set of increments in the localized window, but for every fixed depth k the signed contribution of the determined layers is of order $u(\log\log u)^k/\log u$, far above the RH scale; reaching the RH scale requires substantial cancellation with the remaining layers, and the present arguments do not determine how many layers are needed.

Section 9 connects the excursions with averaged forms of Chowla's conjecture. Signed lag averages of correlations inside a bridge are forced by the return to zero, while the weighted identity $Q_\nu=-\sum_h hC_h^{(\nu)}$ holds globally at every zero of M with exact cancellation of the cross-excursion terms. This gives an RH criterion in terms of h-weighted correlations at zeros of M and shows that RH is a statement of cancellation between lags. Combined with the averaged Chowla theorem of Matomäki–Radziwiłł–Tao, it forces the energy of linear-scale high excursions onto large lags, and it reduces the remaining case to a uniform, translated, sublinear averaged Chowla estimate on blocks selected by the zeros of M.

Extreme values. If a normalized excursion statistic has a uniform tail $\Pr(Z>z)\le Ce^{-\kappa_0 z^{\varpi}}$, the counting bounds force the maximum over a family with at most $\Lambda(X)$ members starting below

X to be of order $(\log\Lambda(X))^{1/\varpi}$ in probability, with an almost sure lim sup constant $\kappa_0^{-1/\varpi}$ (Theorems 8.68 and 8.70). Each level of lacunarity removes one logarithm from the extreme scale: $\sqrt{\log X}$, $\sqrt{\log\log X}$, $\sqrt{\log\log\log X}$.

The remaining sections contain exact identities for the skeleton and its area curves (Sections 2–3), exact correlation identities and sharp bounds inside an excursion (Section 4), the relation with ζ (Section 5), area bounds and global criteria equivalent to RH (Sections 6–7), the exact bilinear form of M on the square range of an excursion and the Mellin mass of an excursion (Section 8), and the decomposition of M by large prime factors, globally and inside an excursion (Sections 10–12). Section 14 records numerical data on the excursion census.

Conventions. ρ runs over nontrivial zeros of ζ, with ordinate $\gamma = \Im\rho$,

$$\Theta = \sup\{\Re\rho\}, \qquad \gamma_1 = 14.134725\ldots$$

is the ordinate of the lowest zero, and RH means $\Theta = 1/2$. Littlewood's criterion [20, §14.25] states that RH holds if and only if $M(x) \ll_\varepsilon x^{1/2+\varepsilon}$ for every $\varepsilon > 0$. Statements labelled *Heuristic* are not used in any proof.

Notation. The following symbols keep one meaning throughout; symbols defined and used only inside a single proof are not listed. The letters c, C (with or without subscripts other than those listed) denote positive constants whose value may change from line to line.

Zeros and constants. ρ, $\gamma = \Im\rho$: nontrivial zeros of ζ; $\Theta = \sup\Re\rho$; $\gamma_1 = 14.1347\ldots$; $\lambda_* = e^{\pi/\gamma_1} = 1.24889\ldots$, the threshold of the counting mechanism of Theorem 8.11; $\theta_* = \lambda_*/2 = 0.62444\ldots$, the prefix cutoff; $M_0 = M + 2$.

Excursions. $I_\nu = (x_{0,\nu}, x_{1,\nu}]$ an excursion, also written $(u_\nu, v_\nu]$ or $(u, v]$; $\sigma_\nu \in \{\pm 1\}$ its orientation (always σ, never a real part); $\ell_\nu = x_{1,\nu} - x_{0,\nu}$; H_ν its height; τ_ν its apex; $a_\nu = \tau_\nu - x_{0,\nu}$, $b_\nu = x_{1,\nu} - \tau_\nu$; $u(x)$, L_x: last zero $\le x$ and length of the terminal block.

Skeleton and curves. M_{s}, M_{sk}: regularized and unregularized skeletons; $\mathcal{A}_+, \mathcal{A}_-, \mathcal{A}$: area curves; $\widetilde{A}(x) = \int_0^x M$.

Quadratic data. S_t, S_r: oriented partial sums inside a block or excursion; C_h, $C_h^{(\nu)}$: lag correlations; q_ν: number of nonzero steps; Q_ν, Q_{term}: quadratic masses; β_ν, β_{term}: absolute masses $\sum|M|$; $P_h^{(\nu)}, D_h^{(\nu)}$: support and opposite-sign pair counts; $\mathcal{E}$: energy.

Families and counts. $\mathcal{F}$ a family of excursions; $E_{\mathcal{F}}(X)$ the number of its members starting at most X; $\mathfrak{h}$ a horizon function and $\mathcal{F}_{\mathfrak{h}}$ the family crossing it; E_A, $E_\square = E_2$, E_λ the counts for $x_1 > x_0^A$, $x_1 > x_0^2$, $x_1 > \lambda x_0$; c in cx_0^2 is a fixed horizon fraction; κ the density constant in (8.7); $N_\eta(Y)$ the Kaczorowski–Pintz excursion count.

Extreme values. Z_k normalized excursion statistics; $Z_{\max,\mathcal{F}}(X)$ their family maximum; C, κ_0, ϖ the constants of (SG_ϖ); $\Lambda(X)$ a deterministic bound for $E_{\mathcal{F}}(X)$.

Linear window and recursion. $\mu_u = \mu\mathbf{1}_{[1,u]}$; $\mathcal{B}_u(t)$ the bilinear form; $c_q(t,u)$ the kernel coefficients, with e_q and $\mathrm{Err}(t,u)$ their remainder; $\Delta_k(u,t) = M(t/k) - M(u/k)$ the channels and $\mathcal{R}_K(u,t)$ the deep remainder; X_k, Y_K their oriented values; $\alpha \in (0,1)$ the exponent in the normalized ratios $|M(q)|/q^\alpha$; w_r, K_q weights and combined kernel; δ_d a relaxed prefix and $D_\delta(n)$ its divisor sums; Ω, ω the number of prime factors with and without multiplicity.

Catalan comparison. $\mathcal{S}(x) = \#\{n \le x : \mu(n) \ne 0\}$ the squarefree-compressed clock; $L^{\#}(I)$ the compressed length; $r_I(t)$ the support time; Cat_n the Catalan numbers; $p_{\mathrm{Cat}}(I,t)$, $p_{\mathrm{Cat}}^{\max}(I)$, $p_{\mathrm{open}}(t)$ reference tail weights; $\mathcal{I}_{\mathrm{Cat}}$ selection energy; $\mathcal{M}_r$ positive meanders.

Large-prime decomposition. $U \geq 2$ a level (not an excursion start); M_U^{sm}, M_U^{lp} the U^2-smooth and large-prime parts; $R = t/U^2$; $N_y(t,U)$ copy multiplicities; $\mathcal{C}(R)$, $\Upsilon(R)$ the coherent and multiplicity-weighted coefficients; $\widetilde{m}(y) = \sum_{a \leq y} \mu(a)/a$; $P^+(n)$ the largest prime factor; $\Pi_a(t)$ local prime counts.

Mellin. $s = \varsigma + it$ in Section 7 and in Proposition 8.75; $D_I(s)$ the excursion block of $1/\zeta$; J a leg in Proposition 5.1.

2. Excursions and the regularized skeleton

Let μ be the Möbius function and

$$M(x) = \sum_{n \leq x} \mu(n), \qquad x \geq 0, \tag{2.1}$$

piecewise constant and right-continuous, with jumps $\mu(n) \in \{-1, 0, 1\}$ at the integers. The segmentation of M by zero runs and the record of its extrema are standard computational objects; Hurst [6] computed $M(x)$ for all $x \leq 10^{16}$. The area and quadratic functionals of lattice excursions are classical [19, 16, 18, 8, 2, 1]; here they are applied to the deterministic Möbius walk.

Definition 2.1 (Excursions and zero runs)**.** An *excursion* is an interval

$$I_\nu = (x_{0,\nu}, x_{1,\nu}]$$

with integer endpoints, $M(x_{0,\nu}) = M(x_{1,\nu}) = 0$, $M(x_{0,\nu}+1) = \sigma_\nu \in \{\pm 1\}$, and $\sigma_\nu M(n) \geq 1$ for every integer $x_{0,\nu} < n < x_{1,\nu}$. Thus $x_{1,\nu}$ is the first return to zero after the departure from $x_{0,\nu}$. Consecutive excursions may be separated by a *zero run* $[x_{1,\nu}, x_{0,\nu+1}]$, a maximal set of consecutive integers on which $M = 0$; it may consist of a single point. The first excursion is $(0, 2]$. Since M takes values of both signs of size $\gg \sqrt{x}$ infinitely often (Remark 6.6) and has unit steps, it has infinitely many zeros and every excursion is finite. We also write $u_\nu = x_{0,\nu}$ and $v_\nu = x_{1,\nu}$.

Put

$$\ell_\nu = x_{1,\nu} - x_{0,\nu}, \qquad H_\nu = \max_{x_{0,\nu} < n < x_{1,\nu}} |M(n)|,$$

$$\tau_\nu = \min\{n \in I_\nu : |M(n)| = H_\nu\}, \qquad a_\nu = \tau_\nu - x_{0,\nu}, \qquad b_\nu = x_{1,\nu} - \tau_\nu.$$

Every excursion has $H_\nu \geq 1$ and $\ell_\nu \geq 2$.

Definition 2.2 (Completed excursions and open prefixes)**.** An excursion in the sense of Definition 2.1 is always *completed*: both $M(x_{0,\nu}) = 0$ and $M(x_{1,\nu}) = 0$ are part of the definition. A segment $(u, x]$ with $M(u) = 0$ and $M(n) \neq 0$ for $u < n \leq x$, with no return condition at x, is an *open prefix*; the terminal block of Definition 2.6 is one. Any estimate whose proof uses the backward bound $|M(u+r)| \leq x_{1,\nu} - u - r$ holds for completed excursions only and is never applied to an open prefix.

Lemma 2.3 (Height and step count)**.** *For every excursion,*

$$H_\nu \leq \min(a_\nu, b_\nu) \leq \frac{\ell_\nu}{2}. \tag{2.2}$$

Proof. The steps of M are 0 or ± 1, so at least H_ν steps are needed to reach height H_ν from zero and at least H_ν more to return to zero. □

Definition 2.4 (Crossing and touching zero runs)**.** The zero run $[x_{1,\nu}, x_{0,\nu+1}]$ between consecutive excursions is a *crossing run* if $\sigma_{\nu+1} = -\sigma_\nu$ and a *touching run* if $\sigma_{\nu+1} = \sigma_\nu$.

Definition 2.5 (Regularized skeleton)**.** The *skeleton* M_{s} is the continuous, piecewise linear function defined on every excursion by

$$M_{\mathrm{s}}(t) = \begin{cases} \sigma_\nu \sqrt{H_\nu}\, \dfrac{t - x_{0,\nu}}{a_\nu}, & x_{0,\nu} \le t \le \tau_\nu, \\ \sigma_\nu \sqrt{H_\nu}\, \dfrac{x_{1,\nu} - t}{b_\nu}, & \tau_\nu \le t \le x_{1,\nu}, \end{cases} \tag{2.3}$$

and by $M_{\mathrm{s}} \equiv 0$ on zero runs. The *unregularized skeleton* M_{sk} is defined in the same way with peak $\sigma_\nu H_\nu$.

Definition 2.6 (Terminal block of a truncated computation)**.** Let $N \ge 2$ and $u = \max\{n \le N : M(n) = 0\}$. If $u = N$ there is no terminal block. Otherwise $M(n) \ne 0$ for $u < n \le N$, and since M has steps $0, \pm 1$ all these values have the sign of $M(N)$. Put

$$\sigma_{\mathrm{term}} = \operatorname{sgn} M(N), \qquad H_{\mathrm{term}} = \max_{u<n\le N} |M(n)|, \qquad \tau_{\mathrm{term}} = \min\{n \in (u, N] : |M(n)| = H_{\mathrm{term}}\}.$$

On $[u, N]$ the regularized skeleton is the segment from $(u, 0)$ to $(\tau_{\mathrm{term}}, \sigma_{\mathrm{term}}\sqrt{H_{\mathrm{term}}})$ followed, if $\tau_{\mathrm{term}} < N$, by the segment ending at $(N, \sigma_{\mathrm{term}}\sqrt{|M(N)|})$. Its signed area is

$$\Delta\mathcal{A}(u, N] = \frac{\sigma_{\mathrm{term}}}{2}\Big[\sqrt{H_{\mathrm{term}}}(N - u) + \sqrt{|M(N)|}(N - \tau_{\mathrm{term}})\Big].$$

The terminal skeleton is provisional: after the next return to zero the completed excursion acquires a different descending side. Independently of it, for real x with last zero $u(x) \le \lfloor x \rfloor$ define

$$\beta_{\mathrm{term}}(x) := \sum_{u(x)<n\le\lfloor x\rfloor} |M(n)|, \qquad Q_{\mathrm{term}}(x) := \sum_{u(x)<n\le\lfloor x\rfloor} M(n)^2, \qquad L_x := \lfloor x \rfloor - u(x).$$

These depend only on M.

Definition 2.7 (The three curves)**.** With $M_{\mathrm{s}}^{\pm} = \max(\pm M_{\mathrm{s}}, 0)$,

$$\mathcal{A}_+(x) = \int_0^x M_{\mathrm{s}}^{+}(t)\, dt, \qquad \mathcal{A}_-(x) = -\int_0^x M_{\mathrm{s}}^{-}(t)\, dt, \qquad \mathcal{A}(x) = \int_0^x M_{\mathrm{s}}(t)\, dt. \tag{2.4}$$

$\mathcal{A}_+$ is nondecreasing, $\mathcal{A}_-$ is nonincreasing, and both are constant on zero runs and on excursions of the opposite sign.

Remark 2.8 (The family of g-skeletons). For any $g : \mathbb{N}_0 \to [0, \infty)$ with $g(0) = 0$ and $g(h) > 0$ for $h \ge 1$, one may replace the apex height $\sqrt{H}$ by $g(H)$. The signed area increment becomes $\frac{1}{2}\sigma_\nu g(H_\nu)\ell_\nu$. The correlation identities of Section 4 concern the Möbius increments and do not depend on g. The choice $g(H) = \sqrt{H}$ is a normalization.

3. Exact formulas

Lemma 3.1 (Increment per excursion)**.** *For every complete excursion,*

$$\mathcal{A}(x_{1,\nu}) - \mathcal{A}(x_{0,\nu}) = \sigma_\nu \frac{\sqrt{H_\nu}\,\ell_\nu}{2}, \qquad \sum_{n\in I_\nu} M_{\mathrm{s}}(n) = \sigma_\nu \sqrt{H_\nu}\frac{\ell_\nu}{2}. \tag{3.1}$$

Proof. The two pieces are triangles of height $\sqrt{H_\nu}$ and bases a_ν, b_ν. For the discrete sum, the ascending values at $j = 1, \dots, a_\nu$ sum to $\sqrt{H_\nu}\sum_{j\le a_\nu} j/a_\nu = \sqrt{H_\nu}(a_\nu + 1)/2$; the descending values at $\tau_\nu + r$, $r = 1, \dots, b_\nu$, sum to $\sqrt{H_\nu}\sum_{r\le b_\nu}(b_\nu - r)/b_\nu = \sqrt{H_\nu}(b_\nu - 1)/2$. □

Proposition 3.2 (Explicit form of the three curves)**.** *Let $x \in I_\nu$. Then*

$$\mathcal{A}(x) = \begin{cases} \mathcal{A}(x_{0,\nu}) + \sigma_\nu \sqrt{H_\nu} \dfrac{(x - x_{0,\nu})^2}{2a_\nu}, & x \le \tau_\nu, \\ \mathcal{A}(x_{0,\nu}) + \sigma_\nu \sqrt{H_\nu} \left(\dfrac{a_\nu}{2} + (x - \tau_\nu) - \dfrac{(x - \tau_\nu)^2}{2b_\nu} \right), & x \ge \tau_\nu. \end{cases} \tag{3.2}$$

At excursion endpoints,

$$\mathcal{A}(x_{1,k}) = \sum_{\nu \le k} \sigma_\nu \frac{\sqrt{H_\nu}\ell_\nu}{2}, \qquad \mathcal{A}_+(x_{1,k}) = \sum_{\substack{\nu \le k \\ \sigma_\nu = +1}} \frac{\sqrt{H_\nu}\ell_\nu}{2}, \qquad \mathcal{A}_-(x_{1,k}) = -\sum_{\substack{\nu \le k \\ \sigma_\nu = -1}} \frac{\sqrt{H_\nu}\ell_\nu}{2}. \tag{3.3}$$

The curves are piecewise quadratic and C^1.

Proof. Integrate (2.3); the endpoint formulas follow from Lemma 3.1. □

A point y is a *strict inflection point* of a function if there exists $\delta > 0$ such that the function is strictly convex on one side of y and strictly concave on the other, within distance δ. The second derivative need not exist at y.

Lemma 3.3 (Critical geometry of $\mathcal{A}$)**.** *Let $R = [p, q]$ be the zero run between an excursion of sign σ_- ending at p and the next excursion of sign σ_+ starting at q.*

(i) *Every apex τ_ν is a strict inflection point.*
(ii) *If $\sigma_+ = -\sigma_-$, then R is an extremal plateau; it is a strict local extremum exactly when $p = q$.*
(iii) *If $\sigma_+ = \sigma_- =: \sigma$, then $\mathcal{A}$ has no local extremum on R. Its concavity has sign $-\sigma$ immediately to the left and $+\sigma$ immediately to the right. If $p = q$ the point is a strict inflection; if $p < q$, no point of R is a strict inflection, and the change of concavity is carried by the whole plateau.*
(iv) *Strict local extrema occur exactly at isolated crossing zeros, and plateaus of extrema exactly at crossing runs of positive length; every point of such a plateau is a non-strict local extremum.*

Proof. On the descending side of the left excursion, $\operatorname{sgn} \mathcal{A}' = \sigma_-$ and $\operatorname{sgn} \mathcal{A}'' = -\sigma_-$. On the ascending side of the right excursion, $\operatorname{sgn} \mathcal{A}' = \sigma_+$ and $\operatorname{sgn} \mathcal{A}'' = \sigma_+$. On R, $\mathcal{A}' = 0$. The assertions follow, and at every apex the sign of $\mathcal{A}''$ reverses. □

4. Internal correlations of an excursion

For a real block $x_1, \dots, x_L$ put

$$S_t = \sum_{j=1}^{t} x_j, \qquad C_h = \sum_{j=1}^{L-h} x_j x_{j+h}. \tag{4.1}$$

Lemma 4.1 (Triangular identity)**.** *For every real finite sequence,*

$$\sum_{t=1}^{L-1} S_t^2 + \sum_{h=1}^{L-1} h C_h = S_L \sum_{t=1}^{L-1} S_t. \tag{4.2}$$

Proof. Expanding the partial sums,

$$\sum_{t=1}^{L} S_t^2 = \sum_{i,j=1}^{L} x_i x_j \big(L + 1 - \max(i, j)\big).$$

Write $\max(i, j) = \frac{1}{2}(i + j + |i - j|)$. The part containing $|i - j|$ is $\frac{1}{2} \sum_{i,j} x_i x_j |i - j| = \sum_{h=1}^{L-1} h C_h$. The remaining part is $(L + 1) S_L^2 - S_L \sum_i i x_i$, and $\sum_i i x_i = L S_L - \sum_{t=1}^{L-1} S_t$. Hence $\sum_{t=1}^{L} S_t^2 = S_L^2 + S_L \sum_{t<L} S_t - \sum_h h C_h$; subtract S_L^2. □

Corollary 4.2 (Bridge case)**.** *If $S_L = 0$, then*

$$\sum_{t=1}^{L-1} S_t^2 = -\sum_{h=1}^{L-1} hC_h. \tag{4.3}$$

Remark 4.3 (Hermitian form). For complex x_j and $C_h = \sum_{j=1}^{L-h} x_j\overline{x_{j+h}}$, the same computation gives

$$\sum_{t=1}^{L-1} |S_t|^2 + \sum_{h=1}^{L-1} h\Re C_h = \Re\left(\overline{S_L}\sum_{t=1}^{L-1} S_t\right).$$

For a complete excursion orient the increments by

$$x_j = \sigma_\nu \mu(x_{0,\nu} + j), \qquad S_t = |M(x_{0,\nu} + t)|. \tag{4.4}$$

Then $S_t > 0$ for $1 \le t < \ell_\nu$ and $S_{\ell_\nu} = 0$. Put

$$Q_\nu = \sum_{t=1}^{\ell_\nu - 1} S_t^2, \qquad C_h^{(\nu)} = \sum_{j=1}^{\ell_\nu - h} x_j x_{j+h}, \qquad \text{so that} \qquad Q_\nu = -\sum_{h=1}^{\ell_\nu - 1} hC_h^{(\nu)}. \tag{4.5}$$

Lemma 4.4 (Zeroth moment and extreme lag)**.** *With $q_\nu = \sum_j x_j^2$,*

$$\sum_{h=1}^{\ell_\nu - 1} C_h^{(\nu)} = -\frac{q_\nu}{2}, \qquad C_{\ell_\nu - 1}^{(\nu)} = -1, \tag{4.6}$$

and q_ν is even and at least 2*.*

Proof. $0 = (\sum_j x_j)^2 = q_\nu + 2\sum_h C_h^{(\nu)}$. The first oriented step is $+1$; immediately before the first return the path is at height 1, so the last step is -1. The bridge condition forces equally many positive and negative active steps. □

Proposition 4.5 (Quadratic decomposition by excursions)**.** *For real $x \ge 1$ let $u(x)$ and L_x be as in Definition 2.6, and for $1 \le j \le L_x$ put $y_j = \mu(u(x) + j)$ and $C_h^{\mathrm{term}} = \sum_{j=1}^{L_x - h} y_j y_{j+h}$. Then*

$$\sum_{n \le x} M(n)^2 = \sum_{\nu : x_{1,\nu} \le u(x)} Q_\nu + Q_{\mathrm{term}}(x), \tag{4.7}$$

$$Q_{\mathrm{term}}(x) = M(\lfloor x \rfloor)^2 + M(\lfloor x \rfloor)\sum_{t=1}^{L_x - 1} M(u(x) + t) - \sum_{h=1}^{L_x - 1} hC_h^{\mathrm{term}}, \tag{4.8}$$

with $Q_{\mathrm{term}}(x) = 0$ if $L_x = 0$. At a zero x of M,

$$\sum_{n \le x} M(n)^2 = \sum_{x_{1,\nu} \le x} Q_\nu = -\sum_{x_{1,\nu} \le x} \sum_h hC_h^{(\nu)}. \tag{4.9}$$

Proof. Every integer up to $u(x)$ lies either in a zero run or in a unique excursion interior. Apply Lemma 4.1 to the terminal block and add its final term $S_{L_x}^2$. □

Lemma 4.6 (Elementary bound for an open terminal block)**.** *Let u be a zero of M and $L \ge 0$. Then $|M(u + r)| \le r$ for $1 \le r \le L$, and*

$$\sum_{r=1}^{L} M(u + r)^2 \le \frac{L(L+1)(2L+1)}{6}. \tag{4.10}$$

In particular $Q_{\mathrm{term}}(x) \ll L_x^3$. The bridge bound (4.17)*, whose leading coefficient is* $1/12$ *rather than* $1/3$*, applies only to complete excursions.*

Proof. $M(u + r) = \sum_{j \le r} \mu(u + j)$ with $|\mu| \le 1$; square and sum. □

Thus the quadratic sum sees only lag correlations internal to excursions: pairs of Möbius increments in different completed excursions do not contribute.

Let

$$\beta_\nu = \sum_{t=1}^{\ell_\nu - 1} S_t = \sum_{n \in I_\nu} |M(n)|. \tag{4.11}$$

Cauchy–Schwarz gives

$$\beta_\nu^2 \le (\ell_\nu - 1) Q_\nu, \tag{4.12}$$

with equality exactly for a flat excursion of height 1. To separate support from sign geometry, let $P_h^{(\nu)} = \#\{j : x_j x_{j+h} \neq 0\}$ and let $D_h^{(\nu)}$ count the opposite-sign pairs among them. Then

$$C_h^{(\nu)} = P_h^{(\nu)} - 2D_h^{(\nu)}, \qquad Q_\nu = \sum_{h=1}^{\ell_\nu - 1} h\big(2D_h^{(\nu)} - P_h^{(\nu)}\big). \tag{4.13}$$

Proposition 4.7 (Sharp bounds at fixed length and height)**.** *Let $S_0 = S_L = 0$, $S_t \ge 1$ for $0 < t < L$, $S_t - S_{t-1} \in \{-1, 0, 1\}$ and $H = \max S_t$. Then $L \ge 2H$ and*

$$H^2 + L - 2H \le \beta \le H(L - H), \tag{4.14}$$

$$\frac{2H^3 + H}{3} + L - 2H \le Q \le \frac{2H^3 + H}{3} + (L - 2H)H^2. \tag{4.15}$$

All four bounds are attained. In particular

$$Q \ge \frac{2H^3 + H}{3}, \qquad Q \ge L - 1. \tag{4.16}$$

Proof. Before and after the first visit to H the path visits each level $1, \dots, H-1$. Together with one visit to H, these $2H - 1$ designated interior times contribute $2(1 + \cdots + (H-1)) + H = H^2$ to β and $2\sum_{j<H} j^2 + H^2 = (2H^3 + H)/3$ to Q. The remaining $L - 2H$ interior times lie at heights between 1 and H; placing them all at height 1, or all at height H, attains the bounds. □

Corollary 4.8 (Sharp fixed-length bounds)**.** *For $L \ge 2$,*

$$\beta \le \left\lfloor \frac{L^2}{4} \right\rfloor, \qquad Q \le \begin{cases} L(L^2+2)/12, & L \text{ even}, \\ L(L^2-1)/12, & L \text{ odd}, \end{cases}$$

and in particular

$$Q \le \frac{L^3 + 2L}{12}. \tag{4.17}$$

Both bounds are attained by the tent path, with one horizontal step at maximal height when L is odd.

Proof. Use $S_t \le \min(t, L - t)$ and sum this bound and its square. □

5. Where ζ enters

The black curve depends on M nonlinearly, through $\sqrt{H_\nu}$ and through the selection of $x_{0,\nu}, \tau_\nu, x_{1,\nu}$, and no closed expression of its Mellin transform in terms of ζ is known. The nodes $x_{0,\nu}, x_{1,\nu}$ are zeros of M. If RH holds and all zeros are simple, there is a sequence $T_k \to \infty$ such that, for non-integer x,

$$M(x) = \lim_{k \to \infty} \sum_{|\gamma| < T_k} \frac{x^\rho}{\rho\, \zeta'(\rho)} - 2 + \sum_{n \ge 1} \frac{(-1)^{n-1} (2\pi/x)^{2n}}{(2n)!\, n\, \zeta(2n+1)} \tag{5.1}$$

[20, §14.27]. For $\widetilde{A}(x) := \int_0^x M(t)\,dt$ one has, for $\Re s > 1$,

$$\int_1^\infty \widetilde{A}(x) x^{-s-2}\,dx = \frac{1}{s(s+1)\zeta(s)}. \tag{5.2}$$

Shifting the contour formally, the residues are

$$\sum_\rho \frac{x^{\rho+1}}{\rho(\rho+1)\zeta'(\rho)} - 2x + 12 + (\text{residues at the trivial zeros}), \tag{5.3}$$

where $x/\zeta(0) = -2x$ is the residue at $s = 0$ and $1/((-1)\zeta(-1)) = 12$ the residue at $s = -1$. The factor $(\rho+1)^{-1}$ improves convergence, but RH and simplicity alone do not control $1/\zeta'(\rho)$ well enough to turn (5.3) into an identity with an untruncated sum; results of this type require additional input of Gonek–Hejhal type [15]. The area of M_{sk} differs from $\widetilde{A}$ by explicit local terms.

Proposition 5.1 (Dipole identity)**.** *For each excursion let its* legs *be the integer sets $J = (u_\nu, \tau_\nu] \cap \mathbb{Z}$ and $(\tau_\nu, v_\nu] \cap \mathbb{Z}$, with midpoints $c_J = (u_\nu + \tau_\nu)/2$ and $(\tau_\nu + v_\nu)/2$. Then for every zero v_k,*

$$\int_0^{v_k} M_{\mathrm{sk}}(t)\,dt = \widetilde{A}(v_k) + \sum_{J \subset (0, v_k]} \sum_{m \in J} \mu(m)\,(m - c_J). \tag{5.4}$$

Proof. Since M is constant on each $[n, n+1)$, $\int_a^b M(t)\,dt = \sum_{n=a}^{b-1} M(n)$ for integers $a < b$. On an ascent leg, $\int_u^\tau M_{\mathrm{sk}} = \sigma H a/2$ and $\int_u^\tau M = \sum_{m=u+1}^{\tau} \mu(m)(\tau - m)$, while $\sigma H = \sum_{m=u+1}^{\tau} \mu(m)$; the difference is $\sum_m \mu(m)(a/2 - \tau + m) = \sum_m \mu(m)(m - c_J)$. On a descent leg, $\int_\tau^v M_{\mathrm{sk}} = \sigma H b/2$ and $\int_\tau^v M = b\,\sigma H + \sum_{m=\tau+1}^{v} \mu(m)(v - m)$ with $\sum_{m=\tau+1}^{v} \mu(m) = -\sigma H$; the difference is $\sum_m \mu(m)(m - (v - b/2))$, and $v - b/2 = (\tau + v)/2 = c_J$. On zero runs both integrands vanish. Sum over legs. □

6. Bounds for the area curves

Lemma 6.1 (Transfer)**.** *Let $x \geq 1$ and let $x^* \geq x$ be a zero of M (in a computation truncated at $N \geq x$, take $x^* = N$ and the terminal skeleton of Definition 2.6). If $|M(n)| \leq C_0\, n^{1/2}$ for all $1 \leq n \leq x^*$ (so necessarily $C_0 \geq 1$), then*

$$|M_{\mathrm{s}}(t)| \leq C_0^{1/2} t^{1/4} \quad (0 \leq t \leq x), \qquad \mathcal{A}_+(x) + |\mathcal{A}_-(x)| \leq \tfrac{4}{5}\, C_0^{1/2} x^{5/4}. \tag{6.1}$$

Proof. For $t \geq \tau_\nu$, $|M_{\mathrm{s}}(t)| \leq \sqrt{H_\nu} \leq C_0^{1/2} \tau_\nu^{1/4} \leq C_0^{1/2} t^{1/4}$. For $u_\nu \leq t \leq \tau_\nu$ the chord from $(u_\nu, 0)$ to $(\tau_\nu, \sqrt{H_\nu})$ lies below the chord of $C_0^{1/2} t^{1/4}$ over $[u_\nu, \tau_\nu]$, whose endpoints $C_0^{1/2} u_\nu^{1/4} \geq 0$ and $C_0^{1/2} \tau_\nu^{1/4} \geq \sqrt{H_\nu}$ (as $\tau_\nu \leq x^*$) are at least as high; that chord lies below $C_0^{1/2} t^{1/4}$ by concavity. Integrate. □

Lemma 6.2 (A concave majorant)**.** *Let $\phi(y) = (\log y)^{3/5} (\log\log y)^{-1/5}$ and, for $\kappa > 0$, $F(y) = y^{1/2} e^{-\kappa\phi(y)}$. There is $y_0 = y_0(\kappa)$ such that F is increasing and concave on $[y_0, \infty)$.*

Proof. With $y = e^s$ and $g(s) = s^{3/5}(\log s)^{-1/5}$, one has $F = \exp(s/2 - \kappa g(s))$. Put $h(s) = \frac{1}{2} - \kappa g'(s)$. Then

$$y F'(y) = F\, h(s), \qquad y^2 F''(y) = F\big(h(s)^2 - h(s) - \kappa g''(s)\big).$$

Since $g'(s) \to 0$ and $g''(s) \to 0$, $h(s) \to \frac{1}{2}$ and $h^2 - h - \kappa g'' \to -\frac{1}{4}$. Hence $F' > 0$ and $F'' < 0$ for large y. □

Corollary 6.3. *From the Vinogradov–Korobov bound* [21]

$$M(x) \ll x \exp\big(-c\,\phi(x)\big) \tag{6.2}$$

one obtains

$$\mathcal{A}_+(x) + |\mathcal{A}_-(x)| \ll x^{3/2} \exp\big(-\tfrac{c}{2}\,\phi(x)\big). \tag{6.3}$$

Proof. Let K be such that $|M(y)| \le Ky\,e^{-c\phi(y)}$ for $y \ge y_1$, and let F be as in Lemma 6.2 with $\kappa = c/2$, so that $\sqrt{|M(y)|} \le K^{1/2}F(y)$ for $y \ge y_1$. Only finitely many excursions start below $Y_0 := \max(y_0, y_1)$; they are finite, so they contribute $O(1)$ to the area. Consider an excursion with $u_\nu \ge Y_0$. Its apex satisfies $\sqrt{H_\nu} = \sqrt{|M(\tau_\nu)|} \le K^{1/2}F(\tau_\nu)$; the bound is taken at the apex τ_ν, which lies in the excursion, and not at its future endpoint v_ν. For $t \ge \tau_\nu$ monotonicity of F gives $|M_{\mathrm{s}}(t)| \le \sqrt{H_\nu} \le K^{1/2}F(t)$. For $u_\nu \le t \le \tau_\nu$, $|M_{\mathrm{s}}(t)|$ is the chord from $(u_\nu, 0)$ to $(\tau_\nu, \sqrt{H_\nu})$. This chord lies below the chord from $(u_\nu, K^{1/2}F(u_\nu))$ to $(\tau_\nu, K^{1/2}F(\tau_\nu))$, which has both endpoints at least as high, and the latter lies below $K^{1/2}F$ on $[u_\nu, \tau_\nu] \subset [y_0, \infty)$ by concavity. Hence $|M_{\mathrm{s}}(t)| \le K^{1/2}F(t)$ for all $t \ge Y_0$ outside the finitely many early excursions, and since F is increasing,

$$\mathcal{A}_+(x) + |\mathcal{A}_-(x)| \le O(1) + K^{1/2}\int_{Y_0}^{x} F(t)\,dt \le O(1) + K^{1/2}xF(x) = O(1) + K^{1/2}x^{3/2}e^{-\frac{c}{2}\phi(x)}. \quad \square$$

Corollary 6.4. *If RH holds, then*

$$\mathcal{A}_+(x) + |\mathcal{A}_-(x)| \ll_\varepsilon x^{5/4+\varepsilon}. \tag{6.4}$$

Consequently, if $\mathcal{A}_+(x) + |\mathcal{A}_-(x)| \gg x^{5/4+\delta}$ along an unbounded sequence for some fixed $\delta > 0$, then RH is false.

Remark 6.5 (Endpoint-indexed bounds). From $\ell_\nu \ge 2H_\nu$ and $\sqrt{H_\nu}\ell_\nu \ll v_\nu^{5/4+\varepsilon}$ one deduces only $H_\nu \ll v_\nu^{5/6+O(\varepsilon)}$. This bound is indexed by the *future* endpoint v_ν of the excursion; it gives no bound for $M(x)$ in terms of x at interior points, and hence no zero-free half-plane. Whether (6.4) is equivalent to RH is open.

Remark 6.6 (Oscillation of the heights). The Mertens conjecture is false [17, 10]; the current bounds are due to Hurst [6], who proves $\limsup M(x)/\sqrt{x} > 1.826054$ and $\liminf M(x)/\sqrt{x} < -1.837625$. The excursion containing a point n with $M(n) > 1.826054\sqrt{n}$ has $u_\nu < n$, so $H_\nu > 1.826054\,u_\nu^{1/2}$ for infinitely many positive excursions, and similarly with 1.837625 for negative ones. A lower bound indexed by v_ν does not follow, since v_ν may be much larger than n.

7. Global criteria equivalent to RH

The Mellin identity and the global L^1/L^2 equivalences below are consequences of Littlewood's criterion and partial summation; the excursion decompositions and pruning lemmas repackage them in terms of excursions. For $\Re s > 1$,

$$\int_1^\infty M(x)x^{-s-1}\,dx = \frac{1}{s\,\zeta(s)}. \tag{7.1}$$

If $\int_1^\infty |M(x)|x^{-\varsigma-1}dx < \infty$ for every $\varsigma > 1/2$, the left side continues analytically to $\Re s > 1/2$, so ζ has no zeros there.

Proposition 7.1 (L^1 criterion)**.** *RH holds if and only if, for every $\varepsilon > 0$,*

$$\sum_{v_\nu \le x} \beta_\nu + \beta_{\mathrm{term}}(x) = \sum_{n \le x} |M(n)| \ll x^{3/2+\varepsilon}. \tag{7.2}$$

Proof. RH gives $M(x) \ll_\varepsilon x^{1/2+\varepsilon}$. Conversely, for $\varsigma > 1/2$ choose $0 < \varepsilon < \varsigma - 1/2$; partial summation gives $\sum_n |M(n)|n^{-\varsigma-1} < \infty$, and since M is constant on $[n, n+1)$ the remark after (7.1) applies. $\square$

Lemma 7.2 (Pruning)**.** $\beta_\nu \le H_\nu(\ell_\nu - 1) \le \ell_\nu^2/2$, *hence*

$$\sum_{v_\nu \le x,\ \ell_\nu \le L_0} \beta_\nu \le \frac{xL_0}{2}. \tag{7.3}$$

Corollary 7.3 (Reduction to long excursions)**.** *For fixed $\delta \geq 0$ and $L_0 = x^{1/2-\delta}$, RH holds if and only if*

$$\sum_{\substack{v_\nu \leq x \\ \ell_\nu > L_0}} \beta_\nu + \mathbf{1}_{\{L_x > L_0\}} \beta_{\mathrm{term}}(x) \ll x^{3/2+\varepsilon} \quad \text{for every } \varepsilon > 0. \tag{7.4}$$

At most x/L_0 long excursions are involved.

Proof. If RH holds, the left side is at most $\sum_{n\leq x} |M(n)| \ll x^{3/2+\varepsilon}$. Conversely, the complete excursions with $\ell_\nu \leq L_0$ contribute at most $xL_0/2 \leq x^{3/2}$ by Lemma 7.2, and if $L_x \leq L_0$ the terminal block contributes $\beta_{\mathrm{term}}(x) \leq \sum_{r\leq L_x} r = L_x(L_x+1)/2 \leq L_0^2 \leq x$. Hence $\sum_{n\leq x} |M(n)| \ll x^{3/2+\varepsilon}$, and Proposition 7.1 applies. □

Remark 7.4 (The $\sqrt{x}$ frontier). The cut may depend on ε: with $L_0 = x^{1/2+\varepsilon/2}$ the discarded part is $\leq x^{3/2+\varepsilon/2}/2$. The frontier of pruning is therefore $x^{1/2+o(1)}$; what requires new information is a cut x^θ with a fixed $\theta > 1/2$. For $\ell_\nu \asymp x$ the trivial bound gives $\beta_\nu \ll x^2$ against a budget $x^{3/2+\varepsilon}$, and there are only $O(1)$ such excursions.

Heuristic 7.5 (Renewal tail)**.** If excursion lengths behave like those of a recurrent random walk,

$$\#\{v_\nu \leq x : \ \ell_\nu > L\} \asymp \sqrt{x}\, L^{-1/2}, \tag{7.5}$$

and with $\beta_\nu \asymp \ell_\nu^{3/2}$ the dyadic range $[L, 2L]$ contributes $\asymp \sqrt{x}\, L$ to (7.2), so the terminal scales $L \asymp x$ dominate the L^1 sum.

Theorem 7.6 (Correlation criterion)**.** *RH holds if and only if, for every $\varepsilon > 0$,*

$$\sum_{n\leq x} M(n)^2 = \sum_{v_\nu \leq x} Q_\nu + Q_{\mathrm{term}}(x) = -\sum_{v_\nu \leq x} \sum_h h C_h^{(\nu)} + Q_{\mathrm{term}}(x) \ll x^{2+\varepsilon}. \tag{7.6}$$

Proof. Under RH, $M(n)^2 \ll_\varepsilon n^{1+\varepsilon}$. Conversely, by Cauchy–Schwarz,

$$\sum_{n\leq x} |M(n)| \leq x^{1/2} \Big(\sum_{n\leq x} M(n)^2 \Big)^{1/2} \ll x^{3/2+\varepsilon/2};$$

apply Proposition 7.1. □

Corollary 7.7 (Local sufficient conditions)**.** *Suppose that for every $\varepsilon > 0$ there is C_ε such that*

$$Q_\nu \leq C_\varepsilon\, \ell_\nu^{2+\varepsilon} \quad \text{for every complete excursion with } \ell_\nu > v_\nu^{1/2}, \tag{7.7}$$

Then RH holds.

Proof. Let $\varepsilon_0 \in (0,1)$, put $\varepsilon = \varepsilon_0/4$, and let C_ε be as in the hypothesis. Split the completed excursions at $\ell_\nu = x^{1/2}$. If $\ell_\nu \leq x^{1/2}$, then (4.17) gives $Q_\nu \leq \ell_\nu x/12 + \ell_\nu/6$, and disjointness bounds the total by $x^2/12 + x/6$. If $\ell_\nu > x^{1/2}$ then $\ell_\nu > v_\nu^{1/2}$ since $v_\nu \leq x$, and the hypothesis bounds the total by $C_\varepsilon \sum \ell_\nu^{2+\varepsilon} \leq C_\varepsilon (\sum \ell_\nu)^{2+\varepsilon} \leq C_\varepsilon x^{2+\varepsilon}$. The terminal block is not a complete excursion, and (4.17) does not apply to it. By Lemma 4.6, $Q_{\mathrm{term}}(x) \leq L_x(L_x+1)(2L_x+1)/6$, which is $O(x^{3/2})$ if $L_x \leq x^{1/2}$. If $L_x > x^{1/2}$, let ν be the complete excursion containing x, starting at $u(x)$; then $Q_{\mathrm{term}}(x) \leq Q_\nu$. Fix $A = 1+\varepsilon$. By Corollary 8.8, $v_\nu \leq u(x)^A \leq x^A$ once x is large, so $\ell_\nu \leq x^A$. If $\ell_\nu > v_\nu^{1/2}$ the hypothesis gives $Q_\nu \leq C_\varepsilon x^{A(2+\varepsilon)}$; otherwise (4.17) gives $Q_\nu \ll \ell_\nu^3 \leq v_\nu^{3/2} \leq x^{3A/2}$. In all cases $Q_{\mathrm{term}}(x) \ll_\varepsilon x^{(1+\varepsilon)(2+\varepsilon)} \leq x^{2+4\varepsilon} = x^{2+\varepsilon_0}$. This step uses the arithmetic input of Corollary 8.8: for a general family of disjoint intervals, the future return of the terminal block is not controlled by x, and a bound for complete excursions would not control $Q_{\mathrm{term}}(x)$. Altogether $\sum_{n\leq x} M(n)^2 \ll_{\varepsilon_0} x^{2+\varepsilon_0}$ for every $\varepsilon_0 \in (0,1)$, and Theorem 7.6 gives RH. □

Heuristic 7.8 (Ballistic excursions)**.** In a model where, after a zero of M, the next values of μ behave locally at random with squarefree density $6/\pi^2$, the pattern "k non-zero steps of one sign followed by k of the other" follows a given zero with probability $\geq c_0^{-k}$. If excursions with $H_\nu = k$ and $\ell_\nu \asymp k$ occur for unbounded k, then by (4.16) $Q_\nu/\ell_\nu^{2+\varepsilon} \gg k^{1-\varepsilon} \to \infty$, so the unpruned version of (7.7) fails for every $\varepsilon \in (0,1)$. Such excursions have $\ell_\nu \ll v_\nu^{1/2}$ and do not affect the pruned condition (7.7); this is why the pruning in Corollary 7.7 is needed.

Theorem 7.9 (Energy criterion)**.** *With u the last zero $\leq x$ and $H_{\mathrm{run}}(x) = \max_{u<n\leq x}|M(n)|$, define*

$$\mathcal{E}(x) := \frac{1}{4}\Big(\sum_{v_\nu \leq x} H_\nu \ell_\nu + (x-u)H_{\mathrm{run}}(x)\Big). \tag{7.8}$$

For a complete excursion let J be the chord joining $(u_\nu, \mathcal{A}(u_\nu))$ to $(v_\nu, \mathcal{A}(v_\nu))$. By (3.1), $J' = \sigma_\nu\sqrt{H_\nu}/2$, so $\frac{1}{4}H_\nu\ell_\nu = \int_{I_\nu}(J')^2$ is the Dirichlet energy of the chord. Then

$$\sum_{n\leq x}|M(n)| \leq 4\mathcal{E}(x) \leq x\max_{n\leq x}|M(n)|, \tag{7.9}$$

and RH holds if and only if $\mathcal{E}(x) \ll x^{3/2+\varepsilon}$ for every $\varepsilon > 0$.

Proof. $\beta_\nu \leq H_\nu\ell_\nu$, $\beta_{\mathrm{term}}(x) \leq (x-u)H_{\mathrm{run}}(x)$ and $\sum \ell_\nu + (x-u) \leq x$; apply Proposition 7.1. □

8. Lacunary excursion families and horizon-localized criteria

8.1. **Geometric counts.** The lemmas of this subsection use only that excursions are disjoint integer intervals ordered by their starting points, the next one starting at or after the end of the previous one. They therefore hold verbatim for every sequence of disjoint integer intervals, deterministic or random, and in particular pathwise for any stochastic model of excursions (Section 8.6). For a family $\mathcal{F}$ of excursions and $X \geq 2$ put

$$E_{\mathcal{F}}(X) := \#\{\nu \in \mathcal{F} : x_{0,\nu} \leq X\}.$$

For a function $\mathfrak{h} : [0,\infty) \to [0,\infty]$, the family of excursions *crossing the horizon h* is

$$\mathcal{F}_{\mathfrak{h}} := \{\nu : x_{1,\nu} > \mathfrak{h}(x_{0,\nu})\}.$$

For $A > 1$ the excursion I_ν is *A-exceptional* if $x_{1,\nu} > x_{0,\nu}^A$, i.e. $\nu \in \mathcal{F}_{x^A}$, and *$A$-ordinary* otherwise; for $A = 2$ we say *square-exceptional* and *square-ordinary*. For $\lambda > 1$ it is *λ-long* if $x_{1,\nu} > \lambda x_{0,\nu}$. We write $E_A = E_{\mathcal{F}_{x^A}}$, $E_\square = E_2$ and $E_\lambda = E_{\mathcal{F}_{\lambda x}}$; the context separates E_A (A an exponent) from E_λ (λ a factor).

Lemma 8.1 (Power lacunarity)**.** *For every $A > 1$ and $X \geq 2$,*

$$E_A(X) \leq 2 + \frac{1}{\log A}\log\frac{\log X}{\log 2}. \tag{8.1}$$

In particular

$$E_\square(X) \leq 2 + \log_2\frac{\log X}{\log 2}, \qquad E_\square(X) \leq 4\log\log X \quad (X \geq e^e). \tag{8.2}$$

Proof. No excursion starts at 1, since $M(1) = 1$. List the A-exceptional starts ≥ 2 as $2 \leq a_1 < a_2 < \cdots$. The next excursion starts at or after the end of the preceding one, so $a_{j+1} \geq x_{1,j} > a_j^A$ and $\log a_j \geq A^{j-1}\log 2$. If $a_j \leq X$ then $j \leq 1 + \log(\log X/\log 2)/\log A$; add 1 for the excursion $(0,2]$. For the second bound in (8.2), $\log_2(\log X/\log 2) = (\log\log X - \log\log 2)/\log 2 \leq 1.443\log\log X + 0.529$. □

The next lemma is the partition of the square horizon $[x_0, x_0^2] = [x_0, cx_0^2] \cup [cx_0^2, x_0^2]$: the upper part is accessible only to a doubly lacunary family, for every fixed $c > 0$.

Lemma 8.2 (Fractional square horizon). *For $0 < c \le 1$ and $X \ge 2$,*

$$E_{\mathcal{F}_{cx^2}}(X) = \#\{\nu : x_{0,\nu} \le X,\ x_{1,\nu} > cx_{0,\nu}^2\} \le \frac{2}{c} + 2 + \log_2 \frac{\log X}{\log 2}. \tag{8.3}$$

Consequently, the range $cx_0^2 < t \le x_0^2$ of the excursions starting at most X is visited by at most $O_c(\log\log X)$ excursions.

Proof. At most $\lceil 2/c \rceil \le 2/c + 1$ members start below $2/c$. List the others as $a_1 < a_2 < \cdots$ and put $b_j = ca_j \ge 2$. Then $a_{j+1} \ge x_{1,j} > ca_j^2$, so

$$b_{j+1} = ca_{j+1} > c^2 a_j^2 = b_j^2,$$

and $\log b_j \ge 2^{j-1} \log 2$. Since $b_j \le cX \le X$, the number of such j is at most $1 + \log_2(\log X/\log 2)$. An excursion containing a point $t > cx_0^2$ has $x_1 \ge t > cx_0^2$. □

Lemma 8.3 (Superlinear horizons). *Let $\psi : [1,\infty) \to (0,\infty)$ satisfy $\psi(x) \to \infty$. Then*

$$E_{\mathcal{F}_{x\psi(x)}}(X) = \#\{\nu : x_{0,\nu} \le X,\ x_{1,\nu} > x_{0,\nu}\psi(x_{0,\nu})\} = o(\log X).$$

Proof. Given $K > 0$ choose x_K with $\psi(x) \ge e^K$ for $x \ge x_K$. At most $x_K + 1$ members start below x_K. For consecutive members $a_j < a_{j+1}$ starting at or above x_K, $a_{j+1} \ge x_{1,j} > a_j\psi(a_j) \ge e^K a_j$, so $\log a_j$ increases by more than K at each step and there are at most $1 + \log X/K$ of them. Hence $\limsup E(X)/\log X \le 1/K$ for every K. □

Lemma 8.4 (Linear lacunarity). *For every $\lambda > 1$ and $X \ge 2$,*

$$E_\lambda(X) = \#\{\nu : x_{0,\nu} \le X,\ x_{1,\nu} > \lambda x_{0,\nu}\} \le 2 + \frac{\log(X/2)}{\log\lambda}. \tag{8.4}$$

Proof. For starts $a_j \ge 2$, $a_{j+1} \ge x_{1,j} > \lambda a_j$, hence $a_j > 2\lambda^{j-1}$ for $j \ge 2$; add 1 for $(0,2]$. □

All four bounds are sharp up to constants for general families of disjoint intervals, as geometric and doubly geometric chains of intervals show.

8.2. **Arithmetic input: Kaczorowski–Pintz.** We separate the published statement, its translation from $M_0 = M + 2$ to M, and the consequences for excursions.

Theorem 8.5 (Kaczorowski–Pintz [9]). *Let $M_0(x) = M(x) - 1/\zeta(0) = M(x) + 2$, the function M_a of [9] at $a = 0$. Let $V(f,Y)$ be defined as in [9], through the largest number of points of $[1,Y]$ at which f is nonzero with alternating signs; the number of sign changes of f on $[1,Y]$ is at least $V(f,Y) - 1$, which does not affect the asymptotic statements below. Then:*

(i) [9, Corollary 4, p. 178, (2.13)–(2.15)]

$$\liminf_{Y\to\infty} \frac{V(M_0,Y)}{\log Y} \ge \frac{\gamma_1}{\pi}, \tag{8.5}$$

with combined oscillation of size $x^{\Theta-\eta}$ for every $\eta > 0$, in the sense of their definition following (1.1): the alternating points t_j counted by V can be chosen with $|M_0(t_j)| \ge t_j^{\Theta-\eta}$;

(ii) *Corollary 4 is deduced from their Corollary 1, which asserts relations (2.1)–(2.3) of their Theorem 1* [9, pp. 175–176]*; in particular, for every $\delta > 0$ and $\eta > 0$ there is $Y_0(\delta,\eta)$ such that for $Y > Y_0(\delta,\eta)$ the interval $[Y^{1-\delta}, Y]$ contains two points t_+, t_- with $M_0(t_+) \ge t_+^{\Theta-\eta}$ and $M_0(t_-) \le -t_-^{\Theta-\eta}$. (In* [9] *both losses are governed by a single parameter ε; the form above follows by taking $\varepsilon = \min\{\delta,\eta\}$.)*

Lemma 8.6 (From M_0 to M). *If $|M_0(t)| > 2$, then $\operatorname{sgn} M(t) = \operatorname{sgn} M_0(t)$ and $|M(t)| \ge |M_0(t)| - 2$. In particular, for points with $|M_0(t)| \ge t^{\Theta-\eta}$ and t large, $\operatorname{sgn} M(t) = \operatorname{sgn} M_0(t)$ and $|M(t)| \ge \frac{1}{2}t^{\Theta-\eta}$.*

Proof. $M_0 > 2$ gives $M > 0$; $M_0 < -2$ gives $M < -4$. The second assertion uses $t^{\Theta-\eta} \to \infty$ for $\eta < \Theta$. □

Corollary 8.7 (Distinct large excursions)**.** *If $\Theta > 1/2$ and $0 < \eta < \Theta - 1/2$, then*

$$N_\eta(Y) \geq \Big(\frac{\gamma_1}{\pi} - o(1)\Big) \log Y, \tag{8.6}$$

where $N_\eta(Y)$ counts distinct excursions starting at most Y that contain a point $t \leq Y$ with $|M(t)| \geq \frac{1}{2}t^{\Theta-\eta}$.

Proof. Take the alternating points of Theorem 8.5(i) in $[1, Y]$, discarding the boundedly many below the threshold of Lemma 8.6. By that lemma, consecutive points $t < t'$ have $M(t)$ and $M(t')$ of opposite signs. Since M is integer valued with steps $0, \pm 1$, it vanishes at some integer in (t, t'), so t and t' lie in different excursions. Each point with $M(t) \neq 0$ belongs to exactly one excursion, and excursions are ordered, so the excursions containing the ordered points are pairwise distinct. Their starts are below the points, hence at most Y, and $|M(t)| \geq \frac{1}{2}t^{\Theta-\eta}$ at each point. □

Corollary 8.8 (Multiplicative localization of Mertens returns)**.** *For every fixed $A > 1$ there exists U_A such that every excursion starting at $u \geq U_A$ ends at $v \leq u^A$. Equivalently, M has only finitely many A-exceptional excursions; in particular only finitely many square-exceptional ones.*

Proof. Put $\delta = 1 - 1/A$ and $Y = u^A$, so that $[Y^{1-\delta}, Y] = [u, u^A]$, and fix $0 < \eta < \Theta$ (possible since $\Theta \geq 1/2$). For u large, Theorem 8.5(ii) and Lemma 8.6 give $t_+, t_- \in [u, u^A]$ with $M(t_+) > 0 > M(t_-)$. Neither point lies in $[u, u+1)$, because $M(u) = 0$; so $u < \lfloor t_\pm \rfloor$. Since M changes by $0, \pm 1$ at each integer, it vanishes at some integer n strictly between $\lfloor t_+ \rfloor$ and $\lfloor t_- \rfloor$, and $u < n < u^A$. If $u = x_{0,\nu}$, the first return satisfies $x_{1,\nu} \leq n < u^A$. □

Remark 8.9. A bare sign change of $M_0 = M + 2$ would not force a zero of M. The decisive point is that the local sign change is combined with oscillation of size $t^{\Theta-\eta} \to \infty$, so that Lemma 8.6 applies to both witnesses.

Remark 8.10 (Geometric counts and arithmetic localization). For the actual Mertens excursions Corollary 8.8 is stronger than Lemma 8.1: each A-exceptional family is finite. The geometric lemmas remain separate results because they are elementary, independent of ζ, and valid for every ordered family of disjoint intervals. This is exactly what the localization theorem below and the extreme-value theorems of Section 8.6 use: the former needs only a count below $(\gamma_1/\pi)\log X$, the latter needs a count valid pathwise in a model where no analogue of Kaczorowski–Pintz is available.

8.3. **Horizon localization.**

Theorem 8.11 (Horizon-localized RH criterion)**.** *Let $\mathfrak{h} : [0, \infty) \to [0, \infty]$ be a function, let $\mathcal{F}$ be a family of excursions, and suppose that for some $\kappa < \gamma_1/\pi$*

$$E_{\mathcal{F} \cup \mathcal{F}_\mathfrak{h}}(X) \leq (\kappa + o(1)) \log X \qquad (X \to \infty). \tag{8.7}$$

Then RH is equivalent to the following statement: for every $\varepsilon > 0$ there is C_ε such that

$$|M(t)| \leq C_\varepsilon t^{1/2+\varepsilon} \qquad \text{for all } t \in I_\nu,\ \nu \notin \mathcal{F},\ t \leq \mathfrak{h}(x_{0,\nu}). \tag{8.8}$$

No condition is imposed on the excursions in $\mathcal{F}$, nor on the part of any excursion lying beyond its horizon $\mathfrak{h}(x_0)$.

Proof. RH implies (8.8) by Littlewood's criterion. Conversely, suppose $\Theta > 1/2$ and fix $\eta, \varepsilon > 0$ with $1/2 + \varepsilon < \Theta - \eta$. By (8.6) there are at least $(\gamma_1/\pi - o(1)) \log X$ distinct excursions starting at most X, each containing a witness t with $|M(t)| \geq \frac{1}{2}t^{\Theta-\eta}$. Consider such an excursion ν with witness t. If $\nu \notin \mathcal{F}$ and $t \leq \mathfrak{h}(x_{0,\nu})$, the witness lies in the range of (8.8). Otherwise $\nu \in \mathcal{F}$, or $t > \mathfrak{h}(x_{0,\nu})$, in which case $x_{1,\nu} > t > \mathfrak{h}(x_{0,\nu})$ (as $M(t) \neq 0$) and $\nu \in \mathcal{F}_\mathfrak{h}$. By (8.7) at least

$$\Big(\frac{\gamma_1}{\pi} - \kappa - o(1)\Big) \log X \longrightarrow \infty$$

witnesses lie in the range of (8.8). They belong to distinct excursions, so they are unbounded, and at them $t^{\Theta-\eta}/t^{1/2+\varepsilon} \to \infty$, contradicting (8.8). □

The case $\mathfrak{h}(x) = \infty$ excludes a sparse family $\mathcal{F}$ of whole excursions; the case $\mathcal{F} = \emptyset$ truncates every excursion at its horizon.

Corollary 8.12 (Square horizon)**.** *For every fixed $A > 1$, RH is equivalent to* (8.8) *restricted to A-ordinary excursions. In particular*

$$\boxed{\text{RH} \iff |M(t)| \ll_\varepsilon t^{1/2+\varepsilon} \text{ for all } t \text{ in excursions with } x_1 \le x_0^2,} \tag{8.9}$$

and, equivalently, RH holds if and only if $|M(t)| \ll_\varepsilon t^{1/2+\varepsilon}$ for $x_{0,\nu} < t \le \min\{x_{1,\nu}, x_{0,\nu}^2\}$ over all excursions.

Proof. Take $\mathcal{F} = \mathcal{F}_{x^A}$, $\mathfrak{h} = \infty$, respectively $\mathcal{F} = \emptyset$, $\mathfrak{h}(x) = x^2$, in Theorem 8.11; Lemma 8.1 gives $\kappa = 0$. □

The count $E_\square(X) = O(\log\log X)$ is thus what makes the square horizon admissible. The square horizon can be lowered.

Corollary 8.13 (Descending horizons)**.** *RH is equivalent to Littlewood's bound $|M(t)| \ll_\varepsilon t^{1/2+\varepsilon}$ on the ranges*

$$x_{0,\nu} < t \le \min\{x_{1,\nu}, \mathfrak{h}(x_{0,\nu})\}$$

over all excursions, for each of the following horizons:

(i) $\mathfrak{h}(x) = cx^2$*, for any fixed $c > 0$;*
(ii) $\mathfrak{h}(x) = x\psi(x)$*, for any ψ with $\psi(x) \to \infty$;*
(iii) $\mathfrak{h}(x) = \lambda x$*, for any fixed $\lambda > \lambda_* := e^{\pi/\gamma_1} = 1.24889\ldots$*

The same holds with "truncate at $\mathfrak{h}(x_0)$" replaced by "discard the excursions with $x_1 > \mathfrak{h}(x_0)$". In particular it is enough to control the excursions with $x_1 \le \frac{5}{4}x_0$.

Proof. Apply Theorem 8.11 with $\mathcal{F} = \emptyset$ (respectively $\mathcal{F} = \mathcal{F}_\mathfrak{h}$, $\mathfrak{h} = \infty$). For (i), $c \ge 1$ reduces to Lemma 8.1 and $0 < c < 1$ to Lemma 8.2; $\kappa = 0$. For (ii), Lemma 8.3 gives $\kappa = 0$. For (iii), Lemma 8.4 gives $\kappa = 1/\log\lambda$, and $1/\log\lambda < \gamma_1/\pi$ exactly when $\lambda > \lambda_*$. Finally $5/4 > \lambda_*$. □

Case (iii) implies (i) and (ii): in both, $\mathfrak{h}(x_0) \ge \lambda x_0$ for large x_0, so the bound on the larger ranges implies it on the ranges of (iii), the finitely many remaining excursions affecting only C_ε. The horizon has been lowered from x_0^2 to λx_0: the possible failure of Littlewood's bound, if any, is visible within a bounded multiplicative distance of the start of an excursion. Quantitatively:

Corollary 8.14 (Two-sided localization of large values)**.** *Suppose $\Theta > 1/2$, and fix $0 < \eta < \Theta - \frac{1}{2}$ and $\lambda > \lambda_*$. Among the excursions starting at $x_0 \le X$, at least*

$$\Big(\frac{\gamma_1}{\pi} - \frac{1}{\log\lambda} - o(1)\Big)\log X$$

contain a point t_* with $|M(t_*)| \ge \frac{1}{2}t_*^{\Theta-\eta}$ and

$$x_0 + \tfrac{1}{2}x_0^{\Theta-\eta} \le t_* \le \lambda x_0.$$

Proof. Of the excursions of Corollary 8.7, those whose witness exceeds λx_0 are λ-long, hence at most $\log X/\log\lambda + 2$ by Lemma 8.4. For the others, $M(x_0) = 0$ and unit steps give $|M(t_*)| \le t_* - x_0$, so $t_* - x_0 \ge \frac{1}{2}t_*^{\Theta-\eta} \ge \frac{1}{2}x_0^{\Theta-\eta}$. □

Remark 8.15 (Dyadic erosion and the uniform transition). For every fixed integer $j \ge 0$, Lemma 8.2 with $c = 2^{-j}$ gives

$$\#\left\{\nu : x_{0,\nu} \le X,\ x_{1,\nu} > \frac{x_{0,\nu}^2}{2^j}\right\} = O_{2^j}(\log\log X),$$

so Theorem 8.11 successively permits the horizons

$$x_0^2, \quad \frac{x_0^2}{2}, \quad \frac{x_0^2}{4}, \quad \frac{x_0^2}{8}, \dots.$$

The constant in the fixed-c estimate depends on c; hence one may not set $j = j(x_0) \sim \log_2 x_0$ by that lemma alone. If instead $2^{j(x)} = x/\psi(x)$ with $\psi(x) \to \infty$, then the corresponding horizon is $x\psi(x)$ and Lemma 8.3 supplies the uniform $o(\log X)$ crossing count. Thus the rigorous erosion chain is

$$x_0^2 \to \frac{x_0^2}{2} \to \frac{x_0^2}{4} \to \cdots \to x_0\psi(x_0) \to \lambda x_0, \qquad \lambda > \lambda_*.$$

8.4. **Fixed-mask Catalan compression inside fractional square horizons.** The fractional-square localization has a useful interaction with the exact combinatorics of a completed excursion. The relevant Catalan time is not physical time: zero Möbius increments must first be removed. This support-time distinction is essential below.

Definition 8.16 (Support-time path and fixed-mask Catalan law)**.** Let $I = (x_0, x_1]$ be a complete excursion, and list the nonzero increments in I as

$$x_0 < a_1 < a_2 < \cdots < a_L \le x_1, \qquad \mu(a_j) \neq 0.$$

Put

$$r_I(t) := \#\{j : a_j \le t\}, \qquad x_0 < t \le x_1.$$

If $\sigma \in \{\pm 1\}$ is the orientation of the excursion, define

$$S_0 = 0, \qquad S_r := \sigma \sum_{j=1}^{r} \mu(a_j) \qquad (1 \le r \le L).$$

Then $L = 2n$ is even and

$$S_0 = S_L = 0, \qquad S_r > 0 \quad (1 \le r < L), \qquad S_r - S_{r-1} \in \{-1, +1\}.$$

Thus $(S_r)_{0 \le r \le L}$ is a primitive Dyck path. The *fixed-mask primitive Catalan law* keeps the physical support $\{a_1, \dots, a_L\}$ fixed and assigns to those support positions a uniformly chosen primitive Dyck sign sequence of length L. We denote its probability by $\mathbb{P}_{\mathrm{pCat},I}$.

For the observed excursion one has, exactly,

$$|M(t)| = S_{r_I(t)}, \qquad r_I(t) \le t - x_0 < t. \tag{8.10}$$

The first equality uses the orientation of the excursion and the fact that M is constant across the zero increments.

We first record the global Catalan estimate, since it is useful in its own right and clarifies why a different estimate is needed at the square horizon.

Lemma 8.17 (Catalan maximum tail)**.** *Let $L = 2n$ and let $H_{\max}$ be the maximum of a uniformly chosen Dyck path of length $2n$. For $1 \le h \le n$,*

$$\mathbb{P}_{\mathrm{Cat},L}\{H_{\max} \ge h\} \le (n+1)\frac{\binom{2n}{n+h}}{\binom{2n}{n}} \le (n+1)\exp\left(-\frac{h^2}{n+h}\right) \le \left(\frac{L}{2}+1\right)e^{-h^2/L}. \tag{8.11}$$

For a uniformly chosen primitive Dyck path of length $L = 2n$,

$$\mathbb{P}_{\mathrm{pCat},L}\{H_{\max} \ge h\} \le \frac{L}{2}\exp\left(-\frac{(h-1)^2}{L-2}\right), \qquad 2 \le h \le n. \tag{8.12}$$

Proof. There are $\mathrm{Cat}_n = (n+1)^{-1}\binom{2n}{n}$ Dyck paths of length $2n$ and $\binom{2n}{n}$ unrestricted bridges. Reflecting a bridge after its first visit to level h gives a bijection between bridges that reach h and unrestricted walks ending at $2h$, hence their number is $\binom{2n}{n+h}$. Therefore

$$\mathbb{P}_{\mathrm{Cat},2n}\{H_{\max} \geq h\} \leq (n+1)\frac{\binom{2n}{n+h}}{\binom{2n}{n}}.$$

Moreover

$$\frac{\binom{2n}{n+h}}{\binom{2n}{n}} = \prod_{j=1}^{h} \frac{n-j+1}{n+j},$$

and $\log(1-x) \leq -x$ gives

$$\log \frac{\binom{2n}{n+h}}{\binom{2n}{n}} \leq -\sum_{j=1}^{h} \frac{2j-1}{n+j} \leq -\frac{h^2}{n+h}.$$

Since $n+h \leq 2n = L$, (8.11) follows. A primitive path of length $2n$ has the form $+1$, followed by a Dyck path of length $2n-2$, followed by -1; its maximum is one plus the maximum of the interior Dyck path. Applying the preceding bound with $n-1$ and $h-1$ proves (8.12). □

Remark 8.18 (Why the maximum tail is not the square-horizon estimate). The exponent in (8.12) is of order H^2/L. For a general family of disjoint intervals retained by the square horizon one may have $L \asymp x_0^2$ (for M itself such excursions are finitely many by Corollary 8.8), while the relevant witness lies much earlier, at physical time $t \asymp x_0$. Then a violation of size $t^{1/2+\varepsilon}$ need not make H^2/L large. Thus the global maximum tail, although correct, is not the right Catalan quantity for combining with the fractional-square horizon. The correct quantity is the pointwise height at the support time $r_I(t)$.

Lemma 8.19 (Pointwise primitive-Catalan tail)**.** *Let $L = 2n \geq 4$ and let (S_r) be a uniformly chosen primitive Dyck path of length L. For $1 \leq r \leq L-1$ and $h \geq 1$,*

$$\mathbb{P}_{\mathrm{pCat},L}\{S_r \geq h\} \leq 8n^{3/2}\exp\left(-\frac{h^2}{2r}\right) \leq 3L^{3/2}\exp\left(-\frac{h^2}{2r}\right). \tag{8.13}$$

By time reversal the same estimate holds with r replaced by $L-r$; hence

$$\mathbb{P}_{\mathrm{pCat},L}\{S_r \geq h\} \leq 3L^{3/2}\exp\left[-\frac{h^2}{2\min\{r, L-r\}}\right]. \tag{8.14}$$

If $h > \min\{r, L-r\}$ the probability is 0, so the bound is only relevant for $h \leq \min\{r, L-r\}$.

Proof. Ignore both positivity and the endpoint constraint. Among all 2^L sign words, the proportion whose first r signs have sum at least h is at most $\exp(-h^2/(2r))$ by the elementary Chernoff bound for a Rademacher sum. Therefore the number of primitive Dyck paths satisfying $S_r \geq h$ is at most

$$2^L \exp\left(-\frac{h^2}{2r}\right).$$

The total number of primitive Dyck paths of length $2n$ is Cat_{n-1}. The standard central-binomial lower bound $\binom{2m}{m} \geq 4^m/(2\sqrt{m})$ gives

$$\mathrm{Cat}_{n-1} = \frac{1}{n}\binom{2n-2}{n-1} \geq \frac{4^{n-1}}{2n\sqrt{n-1}}.$$

Since $2^L = 4^n$, division gives the first inequality in (8.13); the second follows from $8n^{3/2} < 3(2n)^{3/2}$. Reversing the path and changing the signs of the reversed increments preserves the primitive Dyck class and carries height S_r to height S_{L-r}, which proves (8.14). □

For an observed excursion I define its pointwise Catalan upper-tail value by

$$p_{\mathrm{Cat}}(I,t) := \mathbb{P}_{\mathrm{pCat},I}\left\{S_{r_I(t)} \geq |M(t)|\right\}. \tag{8.15}$$

This is a tail probability under the fixed-mask Catalan reference law; it is not a probability assigned to the arithmetic Möbius path itself.

Corollary 8.20 (Fractional-square Catalan rarity)**.** *Fix $c > 0$ and $\varepsilon > 0$. Let $I = (x_0, x_1]$ be a complete excursion with $x_1 \leq cx_0^2$, and keep its zero mask fixed. For all sufficiently large x_0,*

$$\mathbb{P}_{\mathrm{pCat},I}\left\{\exists\, t \in I :\ S_{r_I(t)} \geq t^{1/2+\varepsilon}\right\} \leq 3c^{5/2}x_0^5 \exp\left(-\frac{1}{2}x_0^{2\varepsilon}\right). \tag{8.16}$$

In particular the right-hand side is smaller than x_0^{-A} for every fixed $A > 0$ once x_0 is large enough.

Proof. At a support position $t = a_r$ one has $r \leq t - x_0 < t$. Hence Lemma 8.19 with $h = t^{1/2+\varepsilon}$ gives

$$\mathbb{P}_{\mathrm{pCat},I}\{S_r \geq t^{1/2+\varepsilon}\} \leq 3L^{3/2}\exp\left(-\frac{1}{2}t^{2\varepsilon}\right) \leq 3L^{3/2}\exp\left(-\frac{1}{2}x_0^{2\varepsilon}\right).$$

There are at most L support positions. Since $L \leq x_1 - x_0 \leq x_1 \leq cx_0^2$, the union bound gives (8.16). Between support positions the path is constant, so no additional physical times have to be counted. □

The preceding statement is purely combinatorial: it measures how far a large value lies in the tail of a reference law, not how the Möbius constraints select a path. Combined with the localization theorem it gives the following quantitative form.

Corollary 8.21 (Catalan surprise of a violation)**.** *Suppose $\Theta > 1/2$, fix $0 < \eta < \Theta - \frac{1}{2}$, put $\delta = \Theta - \eta - \frac{1}{2}$, and fix $\lambda > \lambda_*$. Then there are infinitely many excursions $I = (x_0, x_1]$ with $x_1 \leq \lambda x_0$ containing a point t with*

$$-\log p_{\mathrm{Cat}}(I,t) \ \geq\ \frac{1}{8}x_0^{2\delta} - O(\log x_0). \tag{8.17}$$

The same holds with $x_1 \leq \lambda x_0$ replaced by $x_1 \leq cx_0^2$ for any fixed $c > 0$.

Proof. By Corollary 8.7 and Lemma 8.4, at least $(\gamma_1/\pi - 1/\log\lambda - o(1))\log X$ excursions starting at most X have $x_1 \leq \lambda x_0$ and contain t with $h = |M(t)| \geq \frac{1}{2}t^{1/2+\delta}$; for cx_0^2 use Lemma 8.2 instead. With $r = r_I(t) < t$ and $L \leq x_1 - x_0$, Lemma 8.19 gives $p_{\mathrm{Cat}}(I,t) \leq 3L^{3/2}\exp(-h^2/(2r)) \leq 3L^{3/2}\exp(-t^{2\delta}/8)$, and $t > x_0$, $L \leq \lambda x_0$ (respectively $L \leq cx_0^2$). □

Remark 8.22 (Selector bounds are pointwise bounds). Up to polynomial factors, $p_{\mathrm{Cat}}(I,t)$ is a decreasing function of $|M(t)|^2/r_I(t)$: by Lemma 8.19,

$$-\log p_{\mathrm{Cat}}(I,t) \ \geq\ \frac{|M(t)|^2}{2r_I(t)} - \log\bigl(3L^{3/2}\bigr).$$

Hence a selector bound $-\log p_{\mathrm{Cat}}(I,t) \leq \Phi(x_0)$ with $\Phi(x) = x^{o(1)}$ on the excursions with $x_1 \leq \lambda x_0$ implies $|M(t)| \leq t^{1/2}x_0^{o(1)}$ there, and RH then follows from Corollary 8.13; it is not an independent criterion. The selector bound is strictly stronger than Littlewood's bound on these ranges: it controls $|M(t)|$ at the diffusive scale $r_I(t)^{1/2}$ of the support time, which may be much smaller than $t^{1/2}$, and a polynomial selector $p_{\mathrm{Cat}} \geq x_0^{-A}$ would give $|M(t)| \ll \sqrt{t\log t}$ on the retained ranges, beyond any bound known to follow from RH. Turning p_{Cat} into an exact measure of deviation would require the matching lower bound $-\log p_{\mathrm{Cat}}(I,t) \ll |M(t)|^2/r_I(t) + \log L$ for the primitive Catalan tail, which is not proved here.

Remark 8.23 (The correct quantifier). A single value $|M(t)| \geq t^{1/2+\varepsilon}$, or many values with bounded ratio $|M(t)|/t^{1/2+\varepsilon}$, is compatible with RH, since Littlewood's criterion allows a constant C_ε. The negation of the criterion is

$$\exists\, \varepsilon_0 > 0 : \qquad \sup_{t\geq 1} \frac{|M(t)|}{t^{1/2+\varepsilon_0}} = \infty,$$

equivalently there are $t_j \to \infty$ and $A_j \to \infty$ with $|M(t_j)| \geq A_j t_j^{1/2+\varepsilon_0}$. Accordingly, the rarity bound (8.16) is a statement about the reference law, and the arithmetic content lies in the sequence form below. If $\Theta > 1/2$ and $0 < \delta < \Theta - \frac{1}{2}$, the oscillation theorem gives $|M(t_j)| \geq \frac{1}{2} t_j^{1/2+\delta}$ along a sequence, and then $|M(t_j)|/t_j^{1/2+\varepsilon} \geq \frac{1}{2} t_j^{\delta-\varepsilon} \to \infty$ for every $0 < \varepsilon < \delta$.

Corollary 8.24 (Catalan cost along a Littlewood-violating sequence). *Let $t_j \to \infty$, $A_j \to \infty$ and $\varepsilon_0 > 0$ with $|M(t_j)| \geq A_j t_j^{1/2+\varepsilon_0}$, and let $I_j = (x_{0,j}, x_{1,j}]$ be the excursion containing t_j. If $x_{1,j} \leq x_{0,j}^B$ for a fixed B, then*

$$-\log p_{\mathrm{Cat}}(I_j, t_j) \;\geq\; \frac{A_j^2}{2}\, t_j^{2\varepsilon_0} - O_B(\log x_{0,j}) \;\geq\; \frac{A_j^2}{2}\, x_{0,j}^{2\varepsilon_0} - O_B(\log x_{0,j}).$$

Proof. With $r = r_{I_j}(t_j) < t_j$ and $h = |M(t_j)|$, $h^2/(2r) \geq A_j^2 t_j^{1+2\varepsilon_0}/(2t_j)$; apply Lemma 8.19 with $L \leq x_{1,j} \leq x_{0,j}^B$. □

Selection energies: complete excursions and open prefixes. The support-time clock of Definition 8.16 is the squarefree-compressed clock

$$\mathcal{S}(x) = \#\{n \leq x : \mu(n) \neq 0\}, \qquad L^{\#}(I) = \mathcal{S}(x_1) - \mathcal{S}(x_0) \leq x_1 - x_0,$$

and $r_I(t) = \mathcal{S}(t) - \mathcal{S}(x_0)$. Two tail quantities are natural: one for the height of a complete excursion, and one for an open prefix, which does not require knowledge of the future return.

Definition 8.25 (Selection energies). For a complete excursion I with height $H(I)$ and compressed length $L^{\#} = L^{\#}(I)$, put

$$p_{\mathrm{Cat}}^{\max}(I) := \mathbb{P}_{\mathrm{pCat},L^{\#}}\{H_{\max} \geq H(I)\}, \qquad \mathcal{I}_{\mathrm{Cat}}(I) := -\log p_{\mathrm{Cat}}^{\max}(I).$$

For $t \geq 1$ with $M(t) \neq 0$ let $a(t)$ be the last zero of M before t and $r = \mathcal{S}(t) - \mathcal{S}(a(t))$. After orientation, the compressed segment from $a(t)$ to t is a positive meander of length r: a ± 1 path $(S_k)_{k\leq r}$ with $S_0 = 0$ and $S_k > 0$ for $1 \leq k \leq r$. Let $\mathcal{M}_r$ be the set of such paths and

$$p_{\mathrm{open}}(t) := \mathbb{P}_{\mathcal{M}_r}\{S_r \geq |M(t)|\},$$

the uniform measure on $\mathcal{M}_r$.

Lemma 8.26 (Height–entropy inequalities). (i) *If $L^{\#} = 2$, the compressed path is $0, 1, 0$ and $H(I) = 1$. For every complete excursion with $L^{\#} \geq 4$,*

$$\mathcal{I}_{\mathrm{Cat}}(I) \;\geq\; \frac{(H(I)-1)^2}{L^{\#}-2} - \log\frac{L^{\#}}{2}.$$

(ii) $\#\mathcal{M}_r = \binom{r-1}{\lfloor (r-1)/2 \rfloor} \geq 2^r/(4\sqrt{r})$, *and for $h \geq 1$*

$$\mathbb{P}_{\mathcal{M}_r}\{S_r \geq h\} \leq 4\sqrt{r}\, e^{-h^2/(2r)}, \qquad -\log p_{\mathrm{open}}(t) \;\geq\; \frac{M(t)^2}{2r} - \log\bigl(4\sqrt{r}\bigr).$$

Proof. (i) is (8.12). (ii) The count of positive meanders is classical (ballot theorem; the number ending at s is $\frac{s}{r}\binom{r}{(r+s)/2}$). From $\binom{2m}{m} \geq 4^m/(2\sqrt{m})$ and $\binom{2m+1}{m} = \frac{1}{2}\binom{2m+2}{m+1}$ one gets $\#\mathcal{M}_r \geq 2^r/(4\sqrt{r})$. Among all 2^r words the proportion with $S_r \geq h$ is at most $e^{-h^2/(2r)}$ by the Chernoff bound; divide. □

Proposition 8.27 (Selector criteria). (i) Complete excursions. *Fix $\lambda > \lambda_*$. If there is A such that $p^{\max}_{\mathrm{Cat}}(I) \geq x_0^{-A}$ for every sufficiently large complete excursion with $x_1 \leq \lambda x_0$, then $H(I) \ll_{A,\lambda} \sqrt{x_0 \log x_0}$ on that family, and RH holds.*

(ii) Open prefixes. *If there is A such that $p_{\mathrm{open}}(t) \geq t^{-A}$ for all large t with $M(t) \neq 0$, then $|M(t)| \ll_A \sqrt{t \log t}$, and RH holds.*

Proof. (i) If $L^{\#} = 2$ then $H(I) = 1$. Otherwise $L^{\#} \geq 4$ and, by Lemma 8.26(i), $(H-1)^2 \leq (L^{\#}-2)(A \log x_0 + \log L^{\#})$, and $L^{\#} \leq (\lambda - 1)x_0$. Littlewood's bound therefore holds on every excursion with $x_1 \leq \lambda x_0$, and Corollary 8.13(iii), in the form that discards the λ-long excursions, gives RH. (ii) By Lemma 8.26(ii), $M(t)^2 \leq 2r(A \log t + \log(4\sqrt{r}))$ with $r \leq t$. □

Remark 8.28 (Status of the two selectors). Both hypotheses are stronger than RH: each yields $|M| \ll \sqrt{t \log t}$ on its range, beyond any bound known to follow from RH. Neither is known to follow from RH alone: a ballistic stretch of compressed length r has $-\log p \asymp r$, and RH does not exclude such stretches with r a small power of t. The open selector controls every point without reference to the future return; the complete selector uses only the horizon-localized family, where $L^{\#} \leq (\lambda - 1)x_0$ is controlled. By contraposition and Corollary 8.24, failure of RH forces the compressed Möbius path to realize, along an unbounded sequence, sectors of Catalan or meander weight $\exp(-c\,t^{2\delta})$ with $\delta > 0$, not merely polynomially rare ones.

Remark 8.29 (The threshold λ_*). The function $\cos(\gamma_1 \log x)$ changes sign γ_1/π times per unit of $\log x$, i.e. once in every multiplicative interval of ratio $e^{\pi/\gamma_1} = \lambda_*$; the constant in (8.5) is this density. On the geometric side, a chain of λ-long intervals realizes the bound of Lemma 8.4. The method of Theorem 8.11 compares these two counts, and λ_* is the exact point where they balance: for $\lambda \leq \lambda_*$ the λ-long family can, as far as counting alone is concerned, absorb every Kaczorowski–Pintz excursion. Going below λ_* requires information on where inside an excursion the large values occur, not only how many excursions carry them.

Proposition 8.30 (Logarithmic-length budget). *Let $\mathcal{F}$ be a family of excursions and let $u_1 < \cdots < u_N$ be the starts of its members with $2 \leq u_j \leq X$, with ends v_j. Then*

$$\sum_{j=1}^{N-1} \log \frac{v_j}{u_j} \leq \log \frac{X}{2}.$$

If moreover $v_N \leq u_N^A$ for some $A \geq 1$, then $\sum_{j=1}^{N} \log(v_j/u_j) \leq A \log X$.

Proof. Disjointness gives $u_{j+1} \geq v_j$, so $\log(v_j/u_j) \leq \log u_{j+1} - \log u_j$; the sum telescopes to at most $\log u_N - \log u_1 \leq \log(X/2)$. The last term is at most $(A-1)\log u_N \leq (A-1)\log X$. □

Lemmas 8.1–8.4 are the special cases in which every term is bounded below. The budget form shows what the horizon theorem actually measures.

Corollary 8.31 (Mean logarithmic length of Kaczorowski–Pintz excursions). *Suppose $\Theta > 1/2$, fix $0 < \eta < \Theta - \frac{1}{2}$, and let $\mathcal{K}(X)$ be any set of distinct excursions starting at most X, each containing a point t with $|M(t)| \geq \frac{1}{2}t^{\Theta-\eta}$, such that $\#\mathcal{K}(X) \geq (\gamma_1/\pi - o(1)) \log X$ (such sets exist by Corollary 8.7). Then*

$$\frac{1}{\#\mathcal{K}(X)} \sum_{\nu \in \mathcal{K}(X)} \log \frac{v_\nu}{u_\nu} \;\leq\; \frac{\pi}{\gamma_1} + o(1) = \log \lambda_* + o(1).$$

Proof. Fix $A > 1$. By Corollary 8.8, all but finitely many excursions satisfy $v \leq u^A$; the finitely many others contribute $O_A(1)$. Proposition 8.30 gives a total of at most $A \log X + O_A(1)$. Divide by $\#\mathcal{K}(X)$ and let $A \downarrow 1$. □

Remark 8.32 (What lowering λ_* requires). Corollary 8.13(iii) is the statement that the excursions carrying Kaczorowski–Pintz witnesses cannot all have $\log(v/u) > \log\lambda$ with $\lambda > \lambda_*$; Corollary 8.31 is its averaged form. Consequently, any proof that such excursions have mean logarithmic length at least $\pi/\gamma_1 + \delta$ for some fixed $\delta > 0$ would contradict Corollary 8.31 and prove RH, and a surplus $\delta > \log(\lambda_*/\lambda)$ among the excursions whose witness lies beyond λx_0 would extend Corollary 8.13(iii) to that $\lambda < \lambda_*$. Kaczorowski–Pintz, as stated, gives neither: it supplies at least one return in each band $[Y^{1-\delta}, Y]$ but no upper separation between consecutive witnesses. Nor do short-interval estimates for μ supply it: $|M(x+h) - M(x)| = o(h)$ is compatible with M staying positive throughout $(x, x+h]$, whereas a return requires a decrement of size at least $M(x)$ in a prescribed direction. Energy bounds do not help either, since a flat excursion of height 1 has $\beta_I, Q_I \asymp \ell_I$.

Wide peaks from cancellation in all short intervals. The only arithmetic input of the following lemma is the theorem of Matomäki–Teräväinen [13]: for every fixed $\theta > 0.55$, $\sum_{x<n\le x+x^\theta} \mu(n) = o(x^\theta)$ as $x \to \infty$.

Lemma 8.33 (Mesoscopic Lipschitz bound). *Fix $\Lambda > 1$ and $\vartheta > 0.55$. For large u put $h_u = \lceil (\Lambda u)^\vartheta \rceil$ and*

$$\eta_u := \max\Big\{ h_u^{-1},\ \max_{u\le a\le \Lambda u} \frac{|M(a+h_u) - M(a)|}{h_u} \Big\}.$$

Then $\eta_u \to 0$, and for all integers $u \le a < b \le \Lambda u$,

$$|M(b) - M(a)| \le \eta_u (b-a) + h_u. \tag{8.18}$$

Proof. Fix $\theta' \in (0.55, \vartheta)$. For $u \le a \le \Lambda u$ cover $(a, a+h_u]$ by consecutive intervals $(b, b + \lceil b^{\theta'} \rceil]$, $b \ge a$, and a final piece of length at most $\lceil (2\Lambda u)^{\theta'} \rceil = o(h_u)$. By [13] each full piece has Möbius sum $o(b^{\theta'})$, uniformly as $u \to \infty$, so $|M(a+h_u) - M(a)| = o(h_u)$ uniformly in a, and $\eta_u \to 0$. For (8.18), write $b - a = q h_u + r$ with $0 \le r < h_u$: the q full blocks start in $[u, \Lambda u]$ and contribute at most $q\eta_u h_u$, and the remainder at most $r < h_u$. □

Proposition 8.34 (Wide peaks). *Let $(u, v]$ be an excursion with $v \le \Lambda u$, let $u < t < v$ with $|M(t)| = H$, and put $A = (H - h_u)_+$ and $m = \lfloor A/\eta_u \rfloor$. Then*

$$t - u \ge \frac{A}{\eta_u}, \qquad v - t \ge \frac{A}{\eta_u},$$

$$\beta \ge H + 2\sum_{j=1}^{m} (A - \eta_u j), \qquad Q \ge H^2 + 2\sum_{j=1}^{m} (A - \eta_u j)^2.$$

Consequently, uniformly for such excursions with $H/h_u \to \infty$,

$$v - u \ge (2 + o(1))\frac{H}{\eta_u}, \qquad \beta \ge (1 + o(1))\frac{H^2}{\eta_u}, \qquad Q \ge \Big(\frac{2}{3} + o(1)\Big)\frac{H^3}{\eta_u},$$

so that $(v-u)/H$, β/H^2 and Q/H^3 tend to infinity, whereas the unit-step bounds of Lemma 2.3 and Proposition 4.7 give only 2, 1 and $\frac{2}{3}$.

Proof. By (8.18) applied to $(u, t]$ and $(t, v]$, $H \le \eta_u(t-u) + h_u$ and $H \le \eta_u(v-t) + h_u$. For $1 \le j \le m$ the points $t \pm j$ lie in $[u, v]$, and (8.18) gives $\sigma M(t \pm j) \ge A - \eta_u j \ge 0$. Summing t and the pairs $t \pm j$ gives the bounds for β and Q. Finally $2\sum_{j\le m}(A - \eta_u j) = A^2/\eta_u + O(A)$, $2\sum_{j\le m}(A - \eta_u j)^2 = \frac{2}{3}A^3/\eta_u + O(A^2)$, and $A = (1 + o(1))H$. □

Corollary 8.35 (Packing of high excursions). *Let $Y = Y(U)$ with $Y/U^\vartheta \to \infty$. The number of excursions with $U \le u \le 2U$, $v \le \Lambda u$ and height at least Y is $o(U/Y)$, whereas the unit-step bound gives $O(U/Y)$.*

Proof. They are disjoint, contained in $[U, 2\Lambda U]$, and each has length at least $(2+o(1))Y/\max_{U\le u\le 2U}\eta_u$ by Proposition 8.34. □

Remark 8.36 (Scope). Proposition 8.34 is informative only for heights above $u^{0.55+o(1)}$. If $\Theta > 0.55$ and $0.55 < \vartheta < \Theta - \eta$, it applies to the Kaczorowski–Pintz witness excursions of Corollary 8.14 with $v \le \lambda u$: their peaks are wide in the sense above. It does not lower λ_*: the resulting lower bound for $\log(v/u)$ is of order $H/(\eta_u u)$, which may still tend to 0, whereas Remark 8.32 requires a fixed surplus. A quantitative rate for η_u would turn the $o(\cdot)$ gains into explicit logarithmic factors.

8.5. **The linear window and the one-step recursion.** Once the horizon is below $2x_0$, the square-range bilinear structure of Proposition 8.72 is no longer needed: on $(u, 2u]$ the Möbius function is a *linear* expression in the prefix $\mu \mathbf{1}_{[1,u]}$.

Proposition 8.37 (Linear window). *Let $u \ge 1$. For every integer $u < n \le 2u$,*

$$\mu(n) = -\sum_{\substack{d \mid n \\ d \le u}} \mu(d). \tag{8.19}$$

Consequently, if $M(u) = 0$ and $u < t \le 2u$,

$$M(t) = -\sum_{d \le u} \mu(d) \left(\left\lfloor \frac{t}{d} \right\rfloor - \left\lfloor \frac{u}{d} \right\rfloor \right). \tag{8.20}$$

Proof. Since $n \ge 2$, $\sum_{d \mid n} \mu(d) = 0$. Every proper divisor of n is at most $n/2 \le u$, and every divisor $d \le u$ is proper because $n > u$. Summing (8.19) over $u < n \le t$, the number of multiples of d in $(u, t]$ is $\lfloor t/d \rfloor - \lfloor u/d \rfloor$. □

Corollary 8.38 (Localized prefix formula). *Let $M(u) = 0$, let $1 < \lambda < 2$, and let $u < t \le \lambda u$. Then*

$$M(t) = -\sum_{d \le \lambda u/2} \mu(d) \left(\left\lfloor \frac{t}{d} \right\rfloor - \left\lfloor \frac{u}{d} \right\rfloor \right). \tag{8.21}$$

Indeed, if $\lambda u/2 < d \le u$, then $d > u/2$ and the first multiple of d strictly larger than u is $2d > \lambda u \ge t$.

Corollary 8.39 (Prefix-dependency cutoff). *Put*

$$\theta_* := \frac{1}{2} e^{\pi/\gamma_1} = 0.6244\ldots.$$

For every fixed θ with $\theta_* < \theta < 1$, RH is equivalent to Littlewood's bound on every excursion starting at u throughout

$$u < t \le \min\{x_1, 2\theta u\},$$

where, throughout that window, $M(t)$ is given by the sum (8.21) using only the prefix $\mu(d)$ with $d \le \theta u$. In particular, the choice $\theta = 5/8$ corresponds to $\lambda = 5/4$ and gives the RH-equivalent window

$$1 \le r = t - u \le \min\{\ell - 1, u/4\}$$

using only $d \le 5u/8$.

The number θ_* is an infimum, not an attained endpoint, because the horizon theorem requires $\lambda > \lambda_*$.

Proof. Take $\lambda = 2\theta > \lambda_*$ in Corollary 8.13(iii), then use Corollary 8.38. □

Corollary 8.40 (Algebraic dependency gap). *Let $\theta_* < \theta < 1$. If u is a zero of M, then every increment $\mu(n)$ with*

$$u < n \le 2\theta u$$

is determined by the prefix $\mu(1), \ldots, \mu(\lfloor \theta u \rfloor)$. Consequently the intermediate block

$$(\theta u, u]$$

does not occur as a direct input in the divisor recursion determining the localized future window. As $\theta \downarrow \theta_*$, the omitted block has relative length tending to $1 - \theta_* = 0.3755\ldots$.

Proof. For $u < n \le 2\theta u$, every proper divisor $d \mid n$ satisfies $d \le n/2 \le \theta u$. Hence

$$\mu(n) = -\sum_{\substack{d \mid n \\ d < n}} \mu(d)$$

uses only $d \le \theta u$. □

Remark 8.41 (Direct dependency is not independence). Corollary 8.40 is an algebraic statement about which values occur as inputs of the divisor recursion. It does not assert statistical or arithmetic independence between the omitted block $(\theta u, u]$ and the earlier prefix.

In convolution form, (8.19) says that $(\mu - \mu_u) * 1 = \mu \mathbf{1}_{(u,\infty)}$ on $n \le 2u$, the order-one analogue of (8.31), which for $u < n \le u^2$ reads $\mu(n) = -(\mu_u * \mu_u * 1)(n)$; on $(u, 2u]$ both give $\mu(n)$, the bilinear form being the quadratic continuation of the linear one. Equivalently, (8.20) follows from $\sum_{d \le t} \mu(d) \lfloor t/d \rfloor = 1$, since $\lfloor t/d \rfloor = 1$ for $u < d \le t \le 2u$.

Proposition 8.42 (One-step recursion). *For real $x \ge 1$,*

$$\sum_{k \le x} M\Big(\frac{x}{k}\Big) = 1. \tag{8.22}$$

Hence, if $M(u) = 0$ and $t > u$,

$$M(t) = -\sum_{2 \le k \le t} \left[M\Big(\frac{t}{k}\Big) - M\Big(\frac{u}{k}\Big) \right], \tag{8.23}$$

with $M(y) = 0$ for $y < 1$. Every argument on the right is at most $t/2$; hence on a localized window $u < t \le \lambda u$ all arguments are at most $\lambda u/2$. For $\lambda = 5/4$ this is $5u/8$.

Proof. $\sum_{k \le x} M(x/k) = \sum_{k \le x} \sum_{d \le x/k} \mu(d) = \sum_{m \le x} \sum_{d \mid m} \mu(d) = 1$. Subtract the identity at u, where $M(u) = 0$, and note $M(u/k) = 0$ for $k > u$. □

Proposition 8.43 (Scale cascade in the localized excursion window). *Let $(u, v]$ be an excursion, let $\lambda_* < \lambda < 2$, and let $u < t \le \min\{v, \lambda u\}$. For $k \ge 2$ put*

$$\Delta_k(u,t) := M(t/k) - M(u/k),$$

and for an integer $K \ge 2$ put

$$\mathcal{R}_K(u,t) := -\sum_{K < k \le t} \Delta_k(u,t).$$

Then, exactly,

$$M(t) = -\sum_{k=2}^{K} \Delta_k(u,t) + \mathcal{R}_K(u,t), \tag{8.24}$$

and $\mathcal{R}_K(u,t)$ uses only values $M(q)$ with

$$q \le \frac{\lambda u}{K+1}.$$

Equivalently, all dependence on values above $\lambda u/(K+1)$ is concentrated in the $K-1$ increments $\Delta_2, \ldots, \Delta_K$. Moreover

$$\Delta_k(u,t) = \sum_{u/k < n \le t/k} \mu(n),$$

so the supporting interval has length $(t-u)/k$.

Proof. Equation (8.24) is Proposition 8.42 with the terms $2 \le k \le K$ separated. If $k > K$, then $t/k \le \lambda u/(K+1)$ and $u/k \le \lambda u/(K+1)$. The last identity is the definition of the summatory function on the interval $(u/k, t/k]$. □

Corollary 8.44 (First channels). *For $K = 2$,*

$$M(t) = -\big(M(t/2) - M(u/2)\big) + R_2(u,t),$$

where R_2 uses only $q \le \lambda u/3$. With the convenient localized choice $\lambda = 5/4$, the only channel above $5u/12$ is

$$\Delta_2(u, u+r) = \sum_{u/2 < n \le u/2 + r/2} \mu(n), \qquad 0 < r \le u/4,$$

whose supporting interval has length at most $u/8$. In the limiting sense $\lambda \downarrow \lambda_*$, the successive remainder cutoffs are

$$\frac{\lambda_*}{2}u = 0.6244\ldots u, \quad \frac{\lambda_*}{3}u = 0.4162\ldots u, \quad \frac{\lambda_*}{4}u = 0.3122\ldots u, \quad \frac{\lambda_*}{5}u = 0.2497\ldots u.$$

Lemma 8.45 (Channel persistence extracted from a high excursion). *Let $S_r = \sigma M(u+r) > 0$ on $1 \le r \le h$ with $S_0 = 0$ and $|S_r - S_{r-1}| \le 1$. Fix $K \ge 2$ and write, using Proposition 8.43,*

$$S_r = \sum_{k=2}^{K} X_k(r) + Y_K(r), \quad X_k(r) := -\sigma\Delta_k(u, u+r), \quad Y_K(r) := \sigma\mathcal{R}_K(u, u+r).$$

If $H = \max_{r \le h} S_r$ and τ is a point with $S_\tau = H$, then among the last $\lfloor H/2 \rfloor + 1$ indices

$$\tau - \lfloor H/2 \rfloor, \ldots, \tau$$

there is one fixed component among $X_2, \ldots, X_K, Y_K$ which is at least $H/(2K)$ at no fewer than $\lfloor H/(2K) \rfloor$ indices.

Proof. Since $S_0 = 0$ and the path has unit steps, $\tau \ge H$ and $S_{\tau - j} \ge H - j \ge H/2$ for $0 \le j \le \lfloor H/2 \rfloor$. At every one of these indices the sum of the K displayed components is at least $H/2$, hence some component is at least $H/(2K)$. Pigeonholing the indices among the K components gives the claim. □

Remark 8.46 (What the persistence lemma adds). Lemma 8.45 is not a bound for M. It converts a large value attained inside one excursion into a quantified persistence requirement: either one of the finitely many scaled interval sums $\Delta_2, \ldots, \Delta_K$ repeatedly carries a positive fraction of the height, or the responsibility lies repeatedly in the deeper remainder, which is supported entirely below $\lambda u/(K+1)$. This avoids the iteration defect of Remark 8.58: no new point is claimed to be the start-window point of another excursion.

Corollary 8.47 (Channel descent with explicit loss). *Let u be a zero of M, $t > u$, $K \ge 2$, and $0 < \alpha < 1$. Then either*

(i) *there are $2 \le k \le K$ and $q \in \{\lfloor u/k \rfloor, \lfloor t/k \rfloor\}$ with $|M(q)| \ge |M(t)|/(2K)$, hence*

$$\frac{|M(q)|}{q^\alpha} \ge \frac{k^\alpha}{2K} \cdot \frac{|M(t)|}{t^\alpha}, \qquad q \le \frac{t}{k};$$

or

(ii) *$|\mathcal{R}_K(u,t)| \ge |M(t)|/K$, where $\mathcal{R}_K$ depends only on $M(q)$ with $q \le t/(K+1)$.*

The statement holds at every $t > u$, not only in a localized window.

Proof. The proof of Proposition 8.43 applies verbatim to every $t > u$, with $\mathcal{R}_K$ supported on $q \le t/(K+1)$. By (8.24), one of the K terms $\Delta_2, \ldots, \Delta_K, \mathcal{R}_K$ has modulus at least $|M(t)|/K$. If it is $\Delta_k = M(\lfloor t/k \rfloor) - M(\lfloor u/k \rfloor)$, one of its two values has modulus at least $|M(t)|/(2K)$, and $q \le t/k$. □

Remark 8.48 (Channels do not iterate to a bound). Corollary 8.47 has no iteration gap: it applies at the new point q relative to the last zero of M before q. Its defect is the constant loss $k^\alpha/(2K) < 1$ per step. If alternative (i) held at every step, a chain of length $m \asymp \log t$ would give only $|M(t)| \le (2K)^m\, O(1) = t^{O(\log K)}$, weaker than the trivial bound. A closing argument must therefore show that alternative (ii), or alternative (i) with loss factor at least 1, occurs along the chain; both are statements about cancellation, not about dependency.

Remark 8.49 (What the recursion determines). Iterating (8.22) determines all of M from $M(1) = 1$, so determinacy alone carries no information; cf. Remark 10.2. The content of (8.23), after Corollary 8.39, is more precise: at the start u of an excursion, every value in an RH-equivalent window $u < t \le \lambda u$ is a signed sum of increments of M over scaled intervals $(u/k, t/k]$ lying below $\lambda u/2$. This is a descent of dependency only; no descent of normalized size follows from the identity.

Corollary 8.50 (RH in the localized linear window)**.** *RH holds if and only if, for every $\varepsilon > 0$,*

$$\left|\sum_{d\le 5u/8} \mu(d)\left(\left\lfloor \frac{t}{d}\right\rfloor - \left\lfloor \frac{u}{d}\right\rfloor\right)\right| = \left|\sum_{2\le k\le t}\left[M\left(\frac{t}{k}\right) - M\left(\frac{u}{k}\right)\right]\right| \ll_\varepsilon t^{1/2+\varepsilon}$$

for every excursion starting at u and every

$$u < t \le \min\left\{x_1, \frac{5}{4}u\right\}.$$

Equivalently, in intra-excursion coordinates $t = u + r$ it is enough to control

$$1 \le r \le \min\{\ell - 1, u/4\}.$$

Proof. Use Corollary 8.39 with $\theta = 5/8$, together with Proposition 8.42. □

Proposition 8.51 (Kernel form of the recursion)**.** *Let $M(u) = 0$ and let t be an integer with $u < t < 2u$. For $1 \le q \le t/2$ put*

$$c_q(t,u) = \left(\left\lfloor \frac{t}{q}\right\rfloor - \left\lfloor \frac{t}{q+1}\right\rfloor\right) - \left(\left\lfloor \frac{u}{q}\right\rfloor - \left\lfloor \frac{u}{q+1}\right\rfloor\right).$$

Then:

(i) $M(t) = -\sum_{1\le q\le t/2} c_q(t,u) M(q)$;

(ii) $c_q(t,u) = \dfrac{t-u}{q(q+1)} + e_q$ *with* $|e_q| < 2$; *for* $q > \sqrt{t}$ *one has* $c_q \in \{-1, 0, 1\}$, *and* $c_q \ne 0$ *only if* $q = \lfloor t/k\rfloor$ *or* $q = \lfloor u/k\rfloor$ *for some* $k < \sqrt{t}$;

(iii) $c_1(t,u) \ge \frac{1}{2}(t - u - 1)$;

(iv) $M(t) = -(t-u)\,\Upsilon(\sqrt{t}) + \mathrm{Err}(t,u)$, *where* Υ *is the coefficient* (11.10) *and*

$$|\mathrm{Err}(t,u)| \le 2\sum_{q\le\sqrt{t}} |M(q)| + \sum_{k<\sqrt{t}} \left(\left|M\left(\frac{t}{k}\right)\right| + \left|M\left(\frac{u}{k}\right)\right|\right).$$

Proof. (i) In (8.23) group the indices $k \ge 2$ by the value $q = \lfloor t/k\rfloor$, respectively $q = \lfloor u/k\rfloor$. The number of $k \ge 1$ with $\lfloor t/k\rfloor = q$ is $\lfloor t/q\rfloor - \lfloor t/(q+1)\rfloor$, and for $q \le t/2$ none of them is $k = 1$. The same holds for u, because $k = 1$ gives $q = u > t/2$. (ii) Each bracket equals $x/(q(q+1))$ minus a difference of two fractional parts, so it differs from $x/(q(q+1))$ by less than 1. For $q > \sqrt{t} \ge \sqrt{u}$ both $t/q - t/(q+1)$ and $u/q - u/(q+1)$ are < 1, so each bracket is 0 or 1, and it is 1 only if $q = \lfloor x/k\rfloor$ with $k \le x/q < \sqrt{t}$. (iii) $c_1 = (t-u) - (\lfloor t/2\rfloor - \lfloor u/2\rfloor)$ and $\lfloor t/2\rfloor - \lfloor u/2\rfloor \le \lceil (t-u)/2\rceil$. (iv) Insert (ii) into (i) for $q \le \sqrt{t}$; the smooth part is $(t-u)\sum_{q\le\sqrt{t}} M(q)/(q(q+1)) = (t-u)\Upsilon(\sqrt{t})$. For $q > \sqrt{t}$ use the last assertion of (ii). □

Corollary 8.52 (Loss of the pointwise induction)**.** *Let $0 < \alpha < 1$ and suppose $|M(q)| \le Cq^\alpha$ for all $q \le t/2$. Then, for $u < t < 2u$ with $M(u) = 0$,*

$$\big|M(t) + (t-u)\Upsilon(\sqrt{t})\big| \ll_\alpha C\, t^{(1+\alpha)/2}, \tag{8.25}$$

while the termwise bound of the full kernel gives only $|M(t)| \ll_\alpha C\, t$. For $\alpha = \frac{1}{2} + \varepsilon$ the pointwise induction therefore loses $t^{1/2-\varepsilon}$, and $t^{1/4-\varepsilon/2}$ after the smooth component $-(t-u)\Upsilon(\sqrt{t})$ is extracted.

Proof. By Proposition 8.51(iv), $|\mathrm{Err}| \le 2C\sum_{q\le\sqrt{t}} q^\alpha + 2Ct^\alpha \sum_{k<\sqrt{t}} k^{-\alpha} \ll_\alpha Ct^{(1+\alpha)/2}$. The full kernel satisfies $\sum_q |c_q| q^\alpha \ge c_1 \ge \frac{1}{2}(t-u-1)$ by (iii), and $\sum_q |c_q| q^\alpha \ll_\alpha t$ by (ii). □

The smooth component is not small for termwise reasons: by (11.12),

$$\Upsilon(\sqrt{t}) = -\sum_{n>\sqrt{t}} \frac{M(n)}{n(n+1)},$$

so its size is governed by the global identity $\sum_{n\ge1} M(n)/(n(n+1)) = 0$, i.e. by the pole of ζ at $s = 1$ (Remark 11.14), and involves values of M beyond $t/2$. Even under RH the tail gives only $(t-u)\Upsilon(\sqrt{t}) \ll t^{3/4+\varepsilon/2}$, the same order as the bound for $\mathrm{Err}(t,u)$; the RH scale for $M(t)$ is a statement about their sum.

Remark 8.53 (Localized intra-excursion system). Let $I = (u, x_1]$ be an excursion of length $\ell = x_1 - u$ and orientation σ, and put

$$S_r := \sigma M(u+r) > 0 \qquad (1 \le r < \ell), \qquad S_0 = S_\ell = 0.$$

For a fixed $\lambda \in (\lambda_*, 2)$ set

$$h_\lambda(u) := \min\{\ell - 1, (\lambda - 1)u\}.$$

Then the entire RH-relevant initial segment satisfies simultaneously

$$S_r = -\sigma \sum_{q \le \lambda u/2} c_q(u+r, u) M(q) > 0, \qquad 1 \le r \le h_\lambda(u),$$

and $|S_r - S_{r-1}| \le 1$. For $\lambda = 5/4$ this is the segment $r \le \min\{\ell - 1, u/4\}$ and all dependencies lie below $5u/8$.

Definition 8.54 (Relaxed divisor-sum window class)**.** Fix $u \ge 2$ and $1 \le h < u$. For $\delta = (\delta_d)_{d\le u} \in \{-1, 0, 1\}^u$ with $\delta_1 = 1$ and $\sum_{d\le u} \delta_d = 0$, put

$$D_\delta(n) := \sum_{\substack{d \mid n \\ d \le u}} \delta_d, \qquad u < n \le u + h.$$

We call δ window-admissible with orientation $\sigma \in \{-1, 1\}$ if $D_\delta(n) \in \{-1, 0, 1\}$ throughout the window and

$$\sigma \sum_{u < n \le u+r} D_\delta(n) < 0 \qquad (1 \le r \le h).$$

The induced oriented path is $S_r = -\sigma \sum_{u<n\le u+r} D_\delta(n) > 0$. For the genuine Möbius prefix, $\delta_d = \mu(d)$, these are exactly the increment and excursion conditions coming from $\mu * 1 = \delta_1$.

Proposition 8.55 (Ballistic counterexample to the relaxed window class)**.** *The relaxed class of Definition 8.54 admits ballistic paths. Namely, take*

$$\delta_1 = 1, \qquad \delta_u = -1, \qquad \delta_d = 0 \quad (1 < d < u).$$

Then for every $u < n < 2u$ one has $D_\delta(n) = 1$, and with $\sigma = -1$,

$$S_r = r \qquad (1 \le r < u).$$

If $m_q := \sum_{d\le q}\delta_d$, *then* $m_q = 1$ *for every* $q < u$. *Hence for any* $0 < \alpha < 1$, *any fixed* $1 < \lambda < 2$, *and* $h = \lfloor(\lambda-1)u\rfloor$,

$$\sup_{q\le\lambda u/2}\frac{|m_q|}{q^\alpha} = 1, \qquad \frac{S_h}{(u+h)^\alpha} \asymp_\lambda u^{1-\alpha} \longrightarrow \infty.$$

Thus a maximum principle comparing the normalized localized-window height to the normalized history is false on this relaxed class.

Proof. For $u < n < 2u$, the integer u does not divide n. Hence among the two nonzero coordinates of δ only $d = 1$ contributes to $D_\delta(n)$, so $D_\delta(n) = 1$. The remaining assertions are immediate. □

Remark 8.56 (What the relaxation loses). The counterexample violates the internal divisor identities of the Möbius prefix. Already for every prime $p < u$ the genuine identity requires $\delta_1+\delta_p = 0$, whereas the ballistic vector above has $\delta_p = 0$. Consequently any successful localized argument of this type must use some of the relations

$$\sum_{d|n}\delta_d = 0 \qquad (2 \le n \le u),$$

not merely the future-window constraints.

Remark 8.57 (The prefix correlation identity). The complete-excursion bridge identity must not be applied to a truncated localized window. For $1 \le h < \ell$, put $x_j = \sigma\mu(u+j)$, $S_r = \sum_{j\le r} x_j$, and $C_a^{[h]} = \sum_{j=1}^{h-a} x_j x_{j+a}$. Lemma 4.1 applied to the prefix of length h gives exactly

$$\sum_{r=1}^{h-1} S_r^2 + \sum_{a=1}^{h-1} aC_a^{[h]} = S_h \sum_{r=1}^{h-1} S_r.$$

The right-hand side vanishes only for a complete bridge. Thus $Q_\nu = -\sum_a aC_a^{(\nu)}$ cannot be used as an identity internal to a truncated horizon unless the excursion actually returns there. In particular, passing from a large localized value to the energy of the whole excursion mixes the retained window with the part deliberately discarded by the horizon theorem.

Remark 8.58 (The one-step iteration gap). Even a one-step implication sending a large value in the localized window to a large value at some $q \le \lambda u/2$ would not automatically iterate. The point q need not lie in the localized window of the excursion that contains q; it may lie beyond $\lambda u'$ relative to that excursion's start u'. Sparsity of the family of λ-long excursions does not repair a single descent chain. Iteration therefore requires either a principle valid for all t, or a mechanism that places the new point inside an admissible window at every step.

Proposition 8.59 (Termwise absolute-value contraction gives no nontrivial gain)**.** *Let* $(u, x_1]$ *be an excursion,* $1 \le h < \min\{x_1 - u, u\}$, *and let* $w_1, \dots, w_h \ge 0$, *not all zero. Put* $K_q = \sum_{r\le h} w_r c_q(u+r, u)$, *so that* $\sum_r w_r M(u+r) = -\sum_q K_q M(q)$. *If, for some* $0 < \alpha < 1$ *and* $C_\alpha > 0$,

$$\sum_q |K_q| q^\alpha \le C_\alpha \sum_{r\le h} w_r (u+r)^\alpha,$$

then

$$\sum_{r\le h} w_r |M(u+r)| \le (1 + 2C_\alpha) \sum_{r\le h} w_r (u+r)^\alpha.$$

Proof. Proposition 8.51(iii) gives $c_1(u+r, u) \ge (r-1)/2 \ge 0$, hence $\frac{1}{2}\sum_r w_r(r-1) \le K_1 \le C_\alpha \sum_r w_r (u+r)^\alpha$. Since $M(u) = 0$ and M has unit steps, $|M(u+r)| \le r$, and the stated inequality follows. □

Remark 8.60 (No strict coercivity from the first coefficient). Writing $t = u + r$ gives the exact formula

$$c_1(u+r,u) = \begin{cases} \lceil r/2 \rceil, & u \text{ even}, \\ \lfloor r/2 \rfloor, & u \text{ odd}. \end{cases}$$

In particular $c_1(u+1,u) = 0$ when u is odd. The preceding proposition therefore rules out only this particular termwise absolute-value mechanism; it does not rule out a global signed contraction or cancellation among the coefficients K_q.

Definition 8.61 (Multiplicative layers of the prefix)**.** Let $\Omega(n)$ denote the number of prime factors of n, counted with multiplicity. For $R \geq 2$ and $k \geq 1$, a vector $(\delta_d)_{d \leq R}$ satisfies the first k Möbius layers if $\delta_1 = 1$ and

$$\sum_{d \mid n} \delta_d = 0 \qquad (2 \leq n \leq R,\ \Omega(n) \leq k).$$

Thus layer 1 imposes the identities at primes, layer 2 adds prime squares and products of two primes, and so on.

Proposition 8.62 (One-layer propagation into the localized future)**.** *Suppose $(\delta_d)_{d \leq R}$ satisfies the first k Möbius layers. Then*

$$\delta_d = \mu(d) \qquad (d \leq R,\ \Omega(d) \leq k).$$

Moreover, for every integer n with

$$R < n \leq 2R, \qquad \Omega(n) \leq k+1,$$

one has

$$-\sum_{\substack{d \mid n \\ d \leq R}} \delta_d = \mu(n).$$

Thus the k-th multiplicative layer of the prefix determines exactly the $(k+1)$-st multiplicative layer of the future increments in a window of length at most R.

Proof. The first assertion follows recursively from $\delta_n = -\sum_{d \mid n,\ d<n} \delta_d$: every proper divisor of an integer with $\Omega(n) \leq k$ has smaller Ω. For the second, every proper divisor of $n \leq 2R$ is at most $n/2 \leq R$ and has $\Omega(d) \leq k$. Hence the displayed sum equals $-\sum_{d \mid n,\ d<n} \mu(d) = \mu(n)$. □

Question 8.63 (Multiplicative-layer depth)**.** Fix $\theta \in (\theta_*, 1)$ and the localized window $u < n \leq 2\theta u$. Let $k = k(u)$ be a depth up to which the internal Möbius divisor identities are imposed below $R = \theta u$. What growth of $k(u)$, if any, suffices to combine those identities with $M(u) = 0$ and the unit-step and excursion-sign constraints so as to force $\max_{u<t\leq 2\theta u} |M(t)| \ll_\varepsilon u^{1/2+\varepsilon}$? A bounded depth would give a compressed arithmetic mechanism; imposing all layers reconstructs the Möbius prefix. Proposition 8.62 makes the depth a quantifiable intermediate invariant; the results below treat only fixed k.

Proposition 8.64 (Layer sums in the localized window)**.** *Fix $1 < \lambda < 2$ and $j \geq 1$. As $u \to \infty$,*

$$\sum_{\substack{u < n \leq \lambda u \\ \Omega(n) = j}} \mu(n) = (-1)^j \,(1 + o(1))\, \frac{(\lambda - 1)\, u\, (\log\log u)^{j-1}}{(j-1)!\, \log u}. \tag{8.26}$$

Proof. If $\Omega(n) = j$ and $\mu(n) \neq 0$, then n is squarefree with exactly j prime factors and $\mu(n) = (-1)^j$. By Landau's theorem (see [14, Ch. 7]), the number $\pi_j^\flat(x)$ of such $n \leq x$ satisfies $\pi_j^\flat(x) \sim f_j(x) := x(\log\log x)^{j-1}/((j-1)!\log x)$. Since $f_j(x)/x$ is slowly varying, $f_j(\lambda u) - f_j(u) = (\lambda - 1 + o(1)) f_j(u)$. □

Corollary 8.65 (Bounded layer depth). *Fix $k \geq 1$, $\theta \in (\theta_*, 1)$, $R = \theta u$, and the window $u < n \leq 2\theta u$.*

(i) *The increments fixed by the first k layers (Proposition 8.62), namely those with $\Omega(n) \leq k+1$, form a set of $o(u)$ integers of the window.*

(ii) *Their summed contribution over the window is*

$$\sum_{\substack{u<n\leq 2\theta u \\ \Omega(n)\leq k+1}} \mu(n) = (-1)^{k+1}(1+o(1))\frac{(2\theta-1)\,u\,(\log\log u)^k}{k!\,\log u},$$

so it exceeds $u^{1-\varepsilon}$ in modulus and has a fixed sign.

(iii) *Let $\delta_d = \mu(d)$ for $\Omega(d) \leq k$ and $\delta_d = 0$ otherwise. For squarefree n in the window with $\omega(n) = w \geq k+2$,*

$$D_\delta(n) = \sum_{\substack{d\mid n \\ d\leq R}} \delta_d = \sum_{j=0}^{k} (-1)^j \binom{w}{j} = (-1)^k \binom{w-1}{k},$$

which has modulus at least $k+1 \geq 2$. Thus the zero extension of a k-layer prefix violates the unit-step constraint at every such n, i.e. at all but $o(u)$ squarefree integers of the window.

Proof. (i) and (ii) follow from Proposition 8.64 summed over $j \leq k+1$, the term $j = k+1$ dominating. For (iii), all divisors of n other than n are $\leq n/2 \leq R$, and n itself has $\Omega(n) = w > k$, so $D_\delta(n)$ is the truncated alternating binomial sum, evaluated by induction on k via $\binom{w-1}{k} = \binom{w}{k} - \binom{w-1}{k-1}$. The proportion of squarefree $n \leq x$ with $\omega(n) \leq k+1$ tends to 0 by Proposition 8.64. □

Remark 8.66 (What fixed layer depth proves). For every fixed k, Corollary 8.65 shows that the first k layers determine only a density-zero subset of the localized window, while the signed sum of the determined increments has modulus $(1+o(1))(2\theta-1)u(\log\log u)^k/(k!\log u)$, larger than $u^{1-\varepsilon}$ for all large u. Hence, if the full sum on the window is $O(u^{1/2+\varepsilon})$, substantial cancellation must occur between the increments determined by any fixed set of initial layers and the remaining ones. The results proved here do not determine how the required depth grows with u; in particular they do not show that $\asymp \log\log u$ layers (the typical size of Ω) are necessary or sufficient, which would require estimates uniform in $k = k(u)$. Part (iii) shows that the unit-step constraint is far from vacuous on a k-layer relaxation: it forbids the naive completion at almost every integer of the window, so the free coordinates δ_d with $\Omega(d) > k$ are strongly constrained.

8.6. **Conditional extreme-value consequence of excursion sparsity.** The counts of Section 8.1 bound the number of excursions in a family, not their heights. They acquire a second function when combined with a tail hypothesis on a normalized excursion statistic: they fix the scale of the family maximum, with no independence assumption.

Let $(\Omega, \mathcal{G}, \Pr)$ be a probability space carrying a random sequence of disjoint integer intervals $I_k = (x_{0,k}, x_{1,k}]$ with $x_{1,k} \leq x_{0,k+1}$, and real random variables Z_k (for instance a model of the Möbius walk with $Z_k = H_k/\sqrt{\ell_k}$). A family $\mathcal{F}$ is given by a rule selecting indices from $(x_{0,k}, x_{1,k})_k$, such as $\mathcal{F}_\flat$; let $k_\mathcal{F}(1) < k_\mathcal{F}(2) < \cdots$ enumerate its members, and put

$$Z_{\max,\mathcal{F}}(X) := \max\{Z_k : k \in \mathcal{F},\ x_{0,k} \leq X\} \qquad (\max\emptyset = -\infty).$$

The lemmas of Section 8.1 hold for every ω.

Hypothesis 8.67 (SG_ϖ). There are $C, \kappa_0, \varpi > 0$ such that for every $j \geq 1$ and $z \geq 0$

$$\Pr\big(k_\mathcal{F}(j) \text{ exists and } Z_{k_\mathcal{F}(j)} > z\big) \leq Ce^{-\kappa_0 z^\varpi}.$$

For $\varpi = 2$ this is a uniform sub-Gaussian tail, denoted (SG).

Theorem 8.68 (Extreme scale from a count)**.** *Assume* (SG_{ϖ}) *for* $\mathcal{F}$ *and suppose that* $E_{\mathcal{F}}(X) \le \Lambda(X)$ *for every* ω*, with* $\Lambda(X) \ge 1$ *deterministic. Then*

$$\Pr\big(Z_{\max,\mathcal{F}}(X) > z\big) \le \Lambda(X)\, C e^{-\kappa_0 z^{\varpi}}. \tag{8.27}$$

Consequently, for every $\delta > 0$*,*

$$\Pr\Big(Z_{\max,\mathcal{F}}(X) > \Big(\frac{1+\delta}{\kappa_0}\log\Lambda(X)\Big)^{1/\varpi}\Big) \le C\,\Lambda(X)^{-\delta}, \tag{8.28}$$

and $Z_{\max,\mathcal{F}}(X) = O_{\Pr}\big((\log\Lambda(X))^{1/\varpi}\big)$ *as* $\Lambda(X) \to \infty$*.*

Proof. The members of $\mathcal{F}$ starting at most X are the first $E_{\mathcal{F}}(X) \le \Lambda(X)$ members of the enumeration, so $\{Z_{\max,\mathcal{F}}(X) > z\} \subset \bigcup_{j \le \Lambda(X)} \{k_{\mathcal{F}}(j) \text{ exists},\ Z_{k_{\mathcal{F}}(j)} > z\}$, and (8.27) is the union bound. Substituting $\kappa_0 z^{\varpi} = (1+\delta)\log\Lambda(X)$ gives (8.28). □

Corollary 8.69 (Triple-logarithmic scale on the square family)**.** *Assume (SG) for the square-exceptional family* $\mathcal{F}_{x^2}$ *of the model. Then, for* $X \ge e^e$ *and every* $\delta > 0$*,*

$$\Pr\big(Z_{\max,\square}(X) > z\big) \le 4C(\log\log X)e^{-\kappa_0 z^2}, \tag{8.29}$$

$$\Pr\Big(Z_{\max,\square}(X) > \sqrt{\tfrac{1+\delta}{\kappa_0}\log\log\log X}\Big) \le 4C(\log\log X)^{-\delta}. \tag{8.30}$$

Thus $Z_{\max,\square}(X) = O_{\Pr}(\sqrt{\log\log\log X})$*. The same holds for the family* $\mathcal{F}_{cx^2}$ *of Lemma 8.2 with a constant depending on c.*

Proof. Use (8.2) in (8.27) and note that $e^{-\kappa_0 z^2} = (\log\log X)^{-(1+\delta)}$ at the stated z. □

The union bound gives convergence in probability; a Borel–Cantelli argument along a sequence adapted to Λ gives an almost sure bound with the same constant.

Theorem 8.70 (Almost sure extreme scale)**.** *Assume* (SG_{ϖ}) *for* $\mathcal{F}$ *and* $E_{\mathcal{F}}(X) \le \Lambda(X)$ *for every* ω*, where* Λ *is continuous, nondecreasing and unbounded. Then almost surely*

$$\limsup_{X\to\infty} \frac{Z_{\max,\mathcal{F}}(X)}{(\log\Lambda(X))^{1/\varpi}} \le \kappa_0^{-1/\varpi}.$$

Proof. Fix $\delta > 0$. For $j \ge j_0$ let X_j be the least X with $\Lambda(X) = 2^j$. Put $z_j = \big((1+\delta)(j+1)\log 2/\kappa_0\big)^{1/\varpi}$ and $A_j = \{Z_{\max,\mathcal{F}}(X_{j+1}) > z_j\}$. By (8.27), $\Pr(A_j) \le 2^{j+1}Ce^{-(1+\delta)(j+1)\log 2} = C2^{-(j+1)\delta}$, which is summable, so almost surely A_j occurs only finitely often. For $X_j \le X \le X_{j+1}$, monotonicity of $Z_{\max,\mathcal{F}}$ in X gives $Z_{\max,\mathcal{F}}(X) \le z_j$ outside A_j, while $\log\Lambda(X) \ge j\log 2$. Hence the lim sup is at most $\lim_j\big((1+\delta)(j+1)/(\kappa_0 j)\big)^{1/\varpi} = ((1+\delta)/\kappa_0)^{1/\varpi}$. Let $\delta \downarrow 0$ along a countable sequence. □

With $\Lambda_{\square}(X) = 2 + \log_2(\log X/\log 2)$, which is continuous and increasing, Theorem 8.70 gives under (SG)

$$\limsup_{X\to\infty} \frac{Z_{\max,\square}(X)}{\sqrt{\log\log\log X}} \le \kappa_0^{-1/2} \quad \text{almost surely.}$$

Applied to the three counts available for any sequence of disjoint intervals (all excursions: at most $X+1$; Lemma 8.4; Lemma 8.1), Theorems 8.68 and 8.70 give three levels:

family	members starting $\le X$	extreme scale under (SG)
all excursions	$\le X+1$	$\sqrt{\log X}$
$x_1 > \lambda x_0$	$O(\log X)$	$\sqrt{\log\log X}$
$x_1 > cx_0^2$ (or x_0^A)	$O(\log\log X)$	$\sqrt{\log\log\log X}$

Each level of lacunarity removes one logarithm from the extreme scale. Under (SG_{ϖ}) the last column reads $(\log X)^{1/\varpi}$, $(\log\log X)^{1/\varpi}$, $(\log\log\log X)^{1/\varpi}$.

Remark 8.71 (Scope). For M itself the square-exceptional family is finite by Corollary 8.8, so Theorems 8.68–8.70 concern stochastic models of the Möbius walk, in which the pathwise counts of Section 8.1 persist while no analogue of the Kaczorowski–Pintz localization is available. Hypothesis (SG) is an assumption on the model; the theorems state exactly what the counts contribute to its extremes.

8.7. **The square range as an exact bilinear form of the prefix.**

Proposition 8.72 (Bilinear square-range identity)**.** *Let $u \geq 1$ and $1 \leq t \leq u^2$. Then*

$$M(t) = 2M(\min(t,u)) - \sum_{d,j \leq u} \mu(d)\mu(j) \left\lfloor \frac{t}{dj} \right\rfloor. \tag{8.31}$$

In particular, if $M(u) = 0$ and $u < t \leq u^2$,

$$M(t) = -\mathcal{B}_u(t), \qquad \mathcal{B}_u(t) := \sum_{d,j \leq u} \mu(d)\mu(j) \left\lfloor \frac{t}{dj} \right\rfloor, \qquad \mathcal{B}_u(u) = 0. \tag{8.32}$$

Proof. Let $\mu_u = \mu \mathbf{1}_{[1,u]}$. Each factor of $(\mu - \mu_u) * (\mu - \mu_u)$ is supported on integers $\geq u+1$, so the convolution, and its convolution with 1, vanish on $n < (u+1)^2$, in particular on $n \leq u^2$. Expanding and using $\mu * 1 = \delta$ gives $\mu(n) = 2\mu_u(n) - (\mu_u * \mu_u * 1)(n)$ for $n \leq u^2$; sum over $n \leq t$. □

Remark 8.73 (Higher-order determinacy). For every $k \geq 1$ and $n \leq u^k$,

$$\mu(n) = \sum_{j=1}^{k} (-1)^{j-1} \binom{k}{j} \big(\mu_u^{*j} * 1^{*(j-1)}\big)(n), \tag{8.33}$$

because $(\mu - \mu_u)^{*k} * 1^{*(k-1)}$ vanishes on $n \leq u^k$; expand by the binomial theorem and use $\mu * 1 = \delta$. Thus determination by the prefix is not special to u^2. What is special to $t \leq u^2$ is the order-two form (8.32), with rank-one coefficients $\mu(d)\mu(j)$ and elementary kernel $\lfloor t/(dj) \rfloor$. This is a truncation identity of Heath-Brown/Vaughan type [4]; summed forms of this kind are used in the fast computation of M [5].

Question 8.74 (Square-excursion bridges)**.** Let $u = x_{0,\nu}$ and $x_{1,\nu} \leq u^2$. Then $\mathcal{B}_u(u) = \mathcal{B}_u(x_{1,\nu}) = 0$ and $-\sigma_\nu \mathcal{B}_u(t) > 0$ for $u < t < x_{1,\nu}$. Can the rank-one coefficients $\mu(d)\mu(j)$, the kernel $\lfloor t/(dj) \rfloor$ and these bridge and sign constraints be combined to prove

$$\max_{u < t \leq x_{1,\nu}} |\mathcal{B}_u(t)| \ll_\varepsilon x_{1,\nu}^{1/2+\varepsilon}$$

uniformly over square-ordinary excursions? By Corollary 8.12 this uniform bound is equivalent to RH. By Corollary 8.50 it suffices to consider $t \leq \min\{x_{1,\nu}, \frac{5}{4}u\}$, where $\mathcal{B}_u(t)$ reduces to the localized linear form and uses only $\mu(d)$ with $d \leq 5u/8$.

8.8. **Mellin mass of a complete excursion.**

Proposition 8.75 (Excursion blocks of $1/\zeta$)**.** *For a complete excursion $I = (x_0, x_1]$ put $D_I(s) = \sum_{x_0 < n \leq x_1} \mu(n) n^{-s}$. Then*

$$D_I(s) = \sum_{n=x_0+1}^{x_1-1} M(n)\big(n^{-s} - (n+1)^{-s}\big). \tag{8.34}$$

For real $\varsigma > 0$, $D_I(\varsigma)$ has the sign of the excursion and

$$|D_I(\varsigma)| \geq \varsigma \beta_I x_1^{-\varsigma-1} \geq \varsigma H_I^2 x_1^{-\varsigma-1}. \tag{8.35}$$

For $\Re s > 1$,

$$\frac{1}{\zeta(s)} = \sum_I D_I(s), \tag{8.36}$$

zero runs contributing nothing.

Proof. Summation by parts gives (8.34) because M vanishes at both endpoints. All interior $M(n)$ have the sign of the excursion, and $n^{-\varsigma} - (n+1)^{-\varsigma} \geq \varsigma x_1^{-\varsigma-1}$ for $n < x_1$. The bound $\beta_I \geq H_I^2$ is (4.14). If M vanishes at two consecutive integers the intervening $\mu(n)$ is zero, so every n with $\mu(n) \neq 0$ lies in an excursion; sum the disjoint blocks in the half-plane of absolute convergence. □

9. Two-point correlations and the energy of excursions

For integers $0 \leq u < X$ and $1 \leq h < X - u$ put

$$C_h^{(u,X]} := \sum_{u<n\leq X-h} \mu(n)\mu(n+h), \qquad C_h^{[X]} := C_h^{(0,X]}.$$

For an excursion $I_\nu = (x_{0,\nu}, x_{1,\nu}]$ one has $C_h^{(\nu)} = C_h^{(x_{0,\nu},x_{1,\nu}]}$, since $\sigma_\nu^2 = 1$. Chowla's conjecture asserts $C_h^{[X]} = o(X)$ for each fixed $h \geq 1$; it is open. Matomäki, Radziwiłł and Tao [11] proved the averaged form

$$\sum_{h\leq H} \left|C_h^{[X]}\right| = o(HX) \qquad (X \to \infty), \tag{9.1}$$

whenever $H = H(X) \leq X$ tends to infinity. It is stated there for the Liouville function and extended to bounded multiplicative functions in the form of an averaged (corrected) Elliott conjecture, which covers μ.

9.1. Signed lag averages are forced; weighted ones are not.

Remark 9.1 (Unweighted signed averages are degenerate). By Lemma 4.4, every excursion satisfies $\sum_h C_h^{(\nu)} = -q_\nu/2$, with $q_\nu \leq \ell_\nu$. Hence $\ell_\nu^{-2}\sum_h C_h^{(\nu)} = O(\ell_\nu^{-1})$ for purely algebraic reasons: a signed average over lags of the correlations inside a bridge is small without any pseudorandomness of μ. Statements of Chowla type inside an excursion must therefore carry absolute values inside the lag average, as in (9.1).

The weighted sum behaves differently. By (4.5), $Q_\nu = -\sum_h hC_h^{(\nu)}$: the weight h records at which lags the anomalous correlation of a high excursion sits. The same identity holds globally at every zero of M, and the cross terms between different excursions cancel exactly.

Proposition 9.2 (Weighted correlations at zeros of M)**.** *Let $v \geq 2$ be a zero of M. Then*

$$\sum_{n<v} M(n)^2 = -\sum_{h<v} h\, C_h^{[v]}, \qquad \sum_{h<v} h\, C_h^{[v]} = \sum_{\nu:\, x_{1,\nu}\leq v} \sum_h h\, C_h^{(\nu)}. \tag{9.2}$$

Thus, in the h-weighted sum, the correlations between increments lying in different completed excursions, or in zero runs, cancel exactly.

Proof. Apply Lemma 4.1 to the block $x_n = \mu(n)$, $1 \leq n \leq v$: then $S_t = M(t)$ and $S_v = M(v) = 0$, which gives the first identity. By (4.9), $\sum_{n\leq v} M(n)^2 = \sum_\nu Q_\nu$ over the completed excursions ending at or before v, and $Q_\nu = -\sum_h hC_h^{(\nu)}$ by (4.5). □

Theorem 9.3 (Weighted-correlation criterion)**.** *RH holds if and only if, for every $\varepsilon > 0$,*

$$-\sum_{h<v} h\, C_h^{[v]} \ll_\varepsilon v^{2+\varepsilon} \qquad \textit{for every zero } v \textit{ of } M.$$

Proof. If RH holds, Theorem 7.6 and (9.2) give the bound. Conversely, let x be large and let $v \geq x$ be the first zero of M at or after x. If x lies in an excursion starting at $u < x$, then $v \leq u^{1+\varepsilon} \leq x^{1+\varepsilon}$ by Corollary 8.8. Hence $\sum_{n\leq x} M(n)^2 \leq \sum_{n<v} M(n)^2 \ll_\varepsilon v^{2+\varepsilon} \leq x^{(1+\varepsilon)(2+\varepsilon)}$, and Theorem 7.6 applies. □

Remark 9.4 (Where the cancellation must occur). The trivial bound is $\sum_h h|C_h^{[v]}| \leq \sum_h h(v-h) \leq v^3/6$, and (9.1) with $H = v$ improves it only to $o(v^3)$. RH requires the signed weighted sum to be $v^{2+o(1)}$. Absolute values cannot give this at the natural scale: square-root cancellation $|C_h^{[v]}| \ll v^{1/2+o(1)}$ in every single lag would give only $\sum_h h|C_h^{[v]}| \ll v^{5/2+o(1)}$. Thus, in the formulation of Theorem 9.3, RH is a statement of cancellation *between lags* of the h-weighted correlations, beyond what any bound for $|C_h^{[v]}|$ lag by lag can provide. By Proposition 9.2 this cancellation may be localized excursion by excursion.

9.2. **Lag localization of the energy of an excursion.**

Proposition 9.5 (Energy lives at large lags)**.** *For every excursion I_ν of length $\ell = \ell_\nu$ and every $1 \leq h_0 < \ell$,*

$$\Big| Q_\nu + \sum_{h_0<h<\ell} h\, C_h^{(\nu)} \Big| \leq h_0 \sum_{h\leq h_0} \big|C_h^{(\nu)}\big|.$$

Consequently, if $\sum_{h\leq h_0} |C_h^{(\nu)}| \leq \delta h_0 \ell$, then

$$\sum_{h_0<h<\ell} h\, C_h^{(\nu)} \;\leq\; -Q_\nu + \delta h_0^2 \ell \;\leq\; -\frac{2H_\nu^3 + H_\nu}{3} + \delta h_0^2 \ell.$$

Proof. Split $Q_\nu = -\sum_{h<\ell} h C_h^{(\nu)}$ at h_0 and use Proposition 4.7. □

When Proposition 8.34 applies, Q_ν may be replaced by the larger lower bound $(\frac{2}{3} + o(1))H_\nu^3/\eta_u$.

Corollary 9.6 (Averaged Chowla on linear-scale intervals)**.** *Let $(u_j, u_j + L_j]$ be intervals with $L_j \to \infty$ and $u_j \leq BL_j$ for a fixed B, and let $h_j \to \infty$ with $h_j = o(L_j)$. Then*

$$\sum_{h\leq h_j} \big|C_h^{(u_j, u_j+L_j]}\big| = o(h_j L_j).$$

Proof. $C_h^{(u,u+L]}$ differs from $\sum_{n\leq u+L} \mu(n)\mu(n+h) - \sum_{n\leq u} \mu(n)\mu(n+h)$ by at most h terms. Summing over $h \leq h_j$, the boundary terms contribute $O(h_j^2) = o(h_j L_j)$. The first sum contributes $o(h_j(u_j + L_j)) = o(h_j L_j)$ by (9.1). For the second, given $\delta > 0$, the indices with $u_j \leq \delta L_j$ contribute at most $h_j u_j \leq \delta h_j L_j$, and along the others $u_j \to \infty$ and (9.1) gives $o(h_j u_j) = o(h_j L_j)$. □

The corollary does not use that the interval is an excursion; it is (9.1) transported to a shifted interval of comparable length. Its content for the present problem lies in the combination with Proposition 9.5.

Corollary 9.7 (Large-lag concentration in linear-scale excursions)**.** *Let I_j be excursions with $\ell_j \to \infty$, $x_{0,j} \leq B\ell_j$ and heights H_j with $H_j^3/\ell_j \to \infty$. For every $h_j \to \infty$ with $h_j^2 \ell_j = o(H_j^3)$,*

$$\sum_{h_j<h<\ell_j} h\, C_h^{(I_j)} = -(1 + o(1))\, Q_{I_j}.$$

In particular, if $\Theta > 1/2$ and such excursions carry Kaczorowski–Pintz witnesses, $H_j \geq \frac{1}{2} x_{0,j}^{\Theta-\eta}$, then all but a proportion $o(1)$ of their energy $Q_{I_j} \geq \frac{2}{3} H_j^3$ is carried by lags $h \gg x_{0,j}^{(3\Theta-1)/2-3\eta/2-o(1)}$.

Proof. By Corollary 9.6, $\sum_{h\le h_j}|C_h^{(I_j)}| = \delta_j h_j \ell_j$ with $\delta_j \to 0$, and Proposition 9.5 gives an error at most $\delta_j h_j^2 \ell_j = o(H_j^3) = o(Q_{I_j})$. For the witness case, $\ell_j \asymp x_{0,j}$ and $H_j^3/\ell_j \gg x_{0,j}^{3\Theta-3\eta-1} \to \infty$ since $\Theta > \frac12$; any $h_j = o(H_j^{3/2}\ell_j^{-1/2})$ is admissible. □

Remark 9.8 (The sublinear regime). Corollary 9.6 requires $L \gg u$; Section 9.3 removes this for $L \ge u^{3/5+\eta}$ at mesoscopic lags. The excursions of Corollary 8.14 satisfy $\ell \le (\lambda-1)x_0$ but may be much shorter; by Proposition 8.34, when $\Theta > 0.55$ they still satisfy $\ell \ge (2+o(1))H/\eta_{x_0}$, so that $\ell/H \to \infty$ and $\ell \ge x_0^{\Theta-\eta}$. For $L = o(u)$ a statement

$$\sum_{h\le h_0} \bigl|C_h^{(u,u+L]}\bigr| = o(h_0 L)$$

uniformly over intervals whose left endpoint is a zero of M is not a consequence of (9.1); results valid for almost all short intervals do not apply either, since the zeros of M form a sparse set selected by M itself. Such a local averaged Chowla estimate for excursions with $\ell \ge x_0^{1/2+\varepsilon}$, or down to $\ell \ge x_0^{\Theta-\varepsilon}$, would extend Corollary 9.7 to all Kaczorowski–Pintz witness excursions; only small lags are relevant for that purpose.

9.3. **Uniform two-point cancellation in sublinear intervals.** The linear-scale restriction of Corollary 9.6 can be removed, in a range of mesoscopic lags, using a Fourier-uniform input valid in *all* short intervals. Matomäki and Teräväinen [13, Theorem 1.5], improving the exponent 5/8 of Zhan, prove that for fixed $0 < A < 1/3$ and $\eta > 0$,

$$\sup_{\alpha\in\mathbb{R}}\Bigl|\sum_{x<n\le x+L}\mu(n)e(\alpha n)\Bigr| \ll_{A,\eta} \frac{L}{(\log x)^A} \qquad (x^{3/5+\eta} \le L \le x). \tag{9.3}$$

Theorem 9.9 (Uniform two-point Cesàro cancellation)**.** *Fix $0 < A < 1/3$ and $\eta > 0$. Uniformly for all integers u, L, R with $u^{3/5+\eta} \le L \le u$ and $1 \le R \le L/2$,*

$$\frac{1}{R}\sum_{h=1}^{R}\frac{|C_h^{(u,u+L]}|}{L-h} \ll_{A,\eta} \Bigl(\frac{L}{R}\Bigr)^{1/2}(\log u)^{-A}.$$

In particular, if $R \ge L(\log u)^{-2A+2\kappa}$ with $0 < \kappa < A$, the left side is $\ll (\log u)^{-\kappa}$; one may take $R \to \infty$ with $R = o(L)$.

Proof. Put $a_i = \mu(u+i)$ for $1 \le i \le L$ and $A(\alpha) = \sum_i a_i e(i\alpha)$. The numbers $C_{|h|}^{(u,u+L]}$, $|h| < L$ (with $C_0 = q = \sum_i a_i^2 \le L$), are the Fourier coefficients of $|A(\alpha)|^2$, so by Parseval

$$\sum_{|h|<L}\bigl|C_{|h|}^{(u,u+L]}\bigr|^2 = \int_0^1 |A(\alpha)|^4\,d\alpha \le \|A\|_\infty^2\int_0^1|A|^2 = \|A\|_\infty^2\, q \le L\|A\|_\infty^2.$$

Cauchy–Schwarz gives $\sum_{h\le R}|C_h| \le \sqrt{RL}\,\|A\|_\infty$, and $L - h \ge L/2$ for $h \le R$, so the left side is at most $2\|A\|_\infty/\sqrt{RL}$. Now apply (9.3) with $x = u$, since $\|A\|_\infty = \sup_\alpha|\sum_{u<n\le u+L}\mu(n)e(\alpha n)|$. □

Remark 9.10 (Status and scope). Theorem 9.9 is a short corollary of (9.3) by the Parseval argument standard in the Matomäki–Radziwiłł–Tao approach; it holds for every interval, hence for every excursion with $\ell \ge x_0^{3/5+\eta}$, and it does not use the excursion structure. Higher-order uniformity in all intervals of length $\ge x^{5/8+\varepsilon}$ is available in [12]; the Fourier input gives the two-point statement down to 3/5. If $\Theta > 3/5$, choosing η with $\Theta - \eta > 3/5 + \eta$, the theorem applies to all Kaczorowski–Pintz witness excursions, whose length is at least $2H \ge x_0^{\Theta-\eta}$. It does not localize their energy: Proposition 9.5 at $h_0 = R \ge L(\log u)^{-2A}$ requires $R^2L = o(H^3)$, which fails since $H \le L/2$. The range $1/2 < \Theta \le 3/5$ and lags $R \le L(\log u)^{-2/3}$ remain outside present technology.

Remark 9.11 (Excursion geometry alone does not give cancellation). For the tent excursion $a_i = +1$ $(i \le k)$, $a_i = -1$ $(k < i \le 2k)$, which satisfies every combinatorial constraint of a primitive excursion, $C_h = 2k - 3h$ for $1 \le h < k$. Hence for $R \to \infty$, $R = o(k)$,

$$\frac{1}{R}\sum_{h\le R}\frac{|C_h|}{2k-h} \longrightarrow 1, \qquad \sum_{h<R}(R-h)C_h \asymp kR^2,$$

whereas $RL \asymp kR$. Any cancellation of this type inside excursions must therefore come from arithmetic input, not from the bridge and sign constraints.

Lemma 9.12 (Fejér identity)**.** *For a real sequence $(a_i)_{1\le i\le L}$, extended by 0, $q = \sum a_i^2$, $C_h = \sum_i a_i a_{i+h}$ and $1 \le R \le L$,*

$$\sum_{m\in\mathbb{Z}}\Big(\sum_{j=1}^{R} a_{m+j}\Big)^2 = Rq + 2\sum_{h=1}^{R-1}(R-h)C_h.$$

Proof. Expand the square: the pair (j, j') contributes $\sum_m a_{m+j}a_{m+j'} = C_{|j-j'|}$, and the number of pairs with $|j - j'| = h \ge 1$ is $2(R - h)$. □

Thus the weighted target $\sum_{h<R}(R - h)C_h = o(RL)$, weaker than $\sum_{h\le R}|C_h| = o(RL)$ and adapted to $Q_\nu = -\sum_h hC_h$, is equivalent to a local variance statement: the R-block sums of μ along an excursion have mean square $q/L + o(1)$ times R, i.e. they behave diagonally. A bound $\sum_{j\le R}\mu(m + j) = o(R)$ for each block gives only $o(LR^2)$.

10. Trial-division horizon of an excursion

Lemma 10.1 (Square horizon)**.** *Let $u \ge 2$. Trial division by the primes $p \le u$ alone determines $\mu(n)$ for every $n \le u^2$.*

Proof. Remove the prime factors $p \le u$ of n; if one occurs twice, $\mu(n) = 0$. The cofactor $q \le u^2$ has all prime factors $> u$, so it is 1 or a single prime. □

Remark 10.2 (Determinacy is not information). Primes $> u$ are not logically new information, and determination by a finite prefix is not special to u^2: (8.33) propagates a prefix to every power range, and (8.22) determines M from $M(1)$. The ranges $(u, 2u]$ and $(u, u^2]$ are special only in the concrete sense that the exact summatory formula is of order one ((8.20)) or two ((8.32)) in the prefix, and that one-stage trial division by primes $\le u$ suffices on $(u, u^2]$.

Definition 10.3 (Smooth and rough parts)**.** For $y > 1$ write $n = s_y(n)\, r_y(n)$, where every prime factor of $s_y(n)$ is $\le y$ and every prime factor of $r_y(n)$ is $> y$. Let $P^+(n)$ be the largest prime factor of n, with $P^+(1) = 1$.

Lemma 10.4. *$\mu(n) = \mu(s_y(n))\,\mu(r_y(n))$. If $n < y^k$ then $\Omega(r_y(n)) \le k - 1$, and if $n \le y^2$ then $r_y(n)$ is 1 or a prime.*

Proof. The two parts are coprime. If $\Omega(r) \ge 1$ then $y^{\Omega(r)} < r \le n$. □

Remark 10.5 (The horizon u^2 is not the multiplicative threshold of an excursion). For a square-ordinary excursion $\sqrt{v_\nu} \le u_\nu$. Every $n \le v_\nu$ has at most one prime factor $> \sqrt{v_\nu}$, so the threshold that governs the multiplicative structure inside I_ν is $\sqrt{v_\nu}$, below u_ν. The regime $t > u_\nu^2$ of Section 11 lies beyond v_ν for a square-ordinary excursion; that section concerns $M(t)$ globally, and Section 12 gives the local version inside an excursion.

11. Large-prime decomposition at level U^2

The identities of this section are decompositions by large prime factors of Buchstab/Legendre type. The asymptotic statements use only the prime number theorem with classical error term and the Vinogradov–Korobov bound [14, 21].

Definition 11.1 (Smooth baseline and large-prime part)**.** For $U \geq 2$ and $t \geq 1$ put

$$M_U^{\mathrm{sm}}(t) := \sum_{\substack{n \leq t \\ P^+(n) \leq U^2}} \mu(n), \qquad M_U^{\mathrm{lp}}(t) := \sum_{\substack{n \leq t \\ P^+(n) > U^2}} \mu(n), \qquad M(t) = M_U^{\mathrm{sm}}(t) + M_U^{\mathrm{lp}}(t). \tag{11.1}$$

Proposition 11.2 (Single-prime copy)**.** *For a prime p and $t < p^2$,*

$$\Delta_p(t) := \sum_{a \leq t/p} \mu(ap) = -M(t/p). \tag{11.2}$$

Proof. $a \leq t/p < p$, so $p \nmid a$ and $\mu(ap) = -\mu(a)$. □

Proposition 11.3 (Sign content of one prime cone)**.** *Let $\mathcal{S}(y) = \sum_{a \leq y} \mu(a)^2$. Among the numbers ap with $a \leq y < p$, the numbers of values $+1, -1, 0$ of μ are*

$$N_+ = \frac{\mathcal{S}(y) - M(y)}{2}, \qquad N_- = \frac{\mathcal{S}(y) + M(y)}{2}, \qquad N_0 = \lfloor y \rfloor - \mathcal{S}(y), \tag{11.3}$$

with limiting proportions $3/\pi^2,\ 3/\pi^2,\ 1 - 6/\pi^2$.

Proof. $\mu(ap) = -\mu(a)$, and $\#\{a \leq y : \mu(a) = \mp 1\} = (\mathcal{S}(y) \mp M(y))/2$. □

Proposition 11.4 (First-order window)**.** *If $U^2 < t < U^4$, no $n \leq t$ has two prime factors $> U^2$ (with multiplicity), and*

$$M_U^{\mathrm{lp}}(t) = -\sum_{U^2 < p \leq t} M(t/p). \tag{11.4}$$

Proof. Two such factors would give $n > U^4 > t$. Hence $M_U^{\mathrm{lp}}(t) = \sum_{U^2 < p \leq t} \Delta_p(t)$, and $t < U^4 < p^2$; apply (11.2). □

Proposition 11.5 (Prime-count form)**.** *For $U^2 < t < U^4$ and $R := t/U^2$,*

$$M_U^{\mathrm{lp}}(t) = -\sum_{a \leq R} \mu(a)\big(\pi(t/a) - \pi(U^2)\big), \qquad M(t) = M_U^{\mathrm{sm}}(t) - \sum_{a \leq R} \mu(a)\pi(t/a) + M(R)\pi(U^2). \tag{11.5}$$

Proof. Expand (11.4) as $-\sum_p \sum_{a \leq t/p} \mu(a)$ and interchange: for fixed a the prime runs over $U^2 < p \leq t/a$, which is non-empty only if $a < R$. □

Thus the only input from primes $> U^2$ is the value of π at the points t/a, $a \leq R$, weighted by $\mu(a)$ at the small scale $R < U^2$.

Definition 11.6 (Copy multiplicities)**.** For $U^2 < t < U^4$, $R = t/U^2$ and $1 \leq y \leq \lfloor R \rfloor$ put

$$N_y(t, U) = \#\Big\{p :\ \max\big(U^2, \tfrac{t}{y+1}\big) < p \leq \tfrac{t}{y}\Big\}, \qquad \text{so that} \qquad M_U^{\mathrm{lp}}(t) = -\sum_{1 \leq y \leq \lfloor R \rfloor} N_y(t, U) M(y). \tag{11.6}$$

Remark 11.7 (Asymptotics of the multiplicities)*.* For fixed $R > 1$ and $U \to \infty$,

$$N_y \sim \frac{RU^2}{2 \log U} \cdot \frac{1}{y(y+1)} \quad (1 \leq y \leq \lfloor R \rfloor - 1), \qquad N_{\lfloor R \rfloor} \sim \frac{U^2}{2 \log U}\Big(\frac{R}{\lfloor R \rfloor} - 1\Big) \quad (R \notin \mathbb{Z}), \tag{11.7}$$

and $N_R = 0$ if $R \in \mathbb{Z}$. The top multiplicity is truncated because $t/(\lfloor R \rfloor + 1) < U^2$.

Definition 11.8 (Coherent coefficient)**.**

$$\mathcal{C}(R) := \int_1^R \frac{M(w)}{w^2}\,dw = \sum_{a\le R}\mu(a)\Big(\frac{1}{a}-\frac{1}{R}\Big) = \sum_{a\le R}\frac{\mu(a)}{a} - \frac{M(R)}{R}. \tag{11.8}$$

Proposition 11.9 (Fixed R)**.** *For fixed $R>1$, as $U\to\infty$,*

$$M_U^{\mathrm{lp}}(RU^2) = -\frac{RU^2}{2\log U}\big(\mathcal{C}(R)+o(1)\big). \tag{11.9}$$

Proof. In (11.5) the sum is finite; for $a<R$ the prime number theorem gives $\pi(RU^2/a)-\pi(U^2)\sim (R/a-1)U^2/(2\log U)$, and $\sum_{a\le R}\mu(a)(R/a-1)=R\mathcal{C}(R)$. □

Lemma 11.10 (The multiplicity-weighted coefficient)**.** *The coefficient*

$$\Upsilon(R) := \sum_{1\le y\le R}\frac{M(y)}{y(y+1)} = \sum_{n\le R}\frac{\mu(n)}{n} - \frac{M(R)}{\lfloor R\rfloor+1} \tag{11.10}$$

satisfies

$$\Upsilon(R) = \mathcal{C}\big(\lfloor R\rfloor+1\big), \qquad \mathcal{C}(R) = \Upsilon(R) - M(R)\Big(\frac{1}{R}-\frac{1}{\lfloor R\rfloor+1}\Big), \tag{11.11}$$

and

$$\mathcal{C}(R) = -\int_R^\infty \frac{M(w)}{w^2}\,dw, \qquad \Upsilon(R) = -\sum_{n>R}\frac{M(n)}{n(n+1)}. \tag{11.12}$$

Proof. M is constant on $[y,y+1)$ and $\int_y^{y+1}w^{-2}dw = 1/(y(y+1))$, which gives (11.11). For $s>1$, (7.1) gives $\int_1^\infty M(w)w^{-s-1}\,dw = 1/(s\zeta(s))\to 0$ as $s\downarrow 1$. For $s\in[1,1+\delta]$ the bound $M(w)\ll we^{-c\sqrt{\log w}}$ dominates the integrand by an integrable multiple of $e^{-c\sqrt{\log w}}/w$, so dominated convergence gives $\int_1^\infty M(w)w^{-2}dw=0$. □

Remark 11.11 ($\mathcal{C}$ versus Υ). Summing (11.7) against $M(y)$ suggests the coefficient $\Upsilon(R)$, but the truncated top multiplicity replaces it by $\mathcal{C}(R)$; by (11.11) the two agree exactly when $M(\lfloor R\rfloor)=0$. For $1<R<2$ one has $M_U^{\mathrm{lp}} = -(\pi(t)-\pi(U^2))\sim -(R-1)U^2/(2\log U)$, i.e. $\mathcal{C}(R)=1-1/R$, whereas $\Upsilon(R)=1/2$.

Corollary 11.12 (Decay of the coefficients)**.** *By* (11.12) *and* (6.2)*,*

$$\mathcal{C}(R),\ \Upsilon(R) \ll \exp\big(-c_1(\log R)^{3/5}(\log\log R)^{-1/5}\big). \tag{11.13}$$

Proposition 11.13 (Both parts are large)**.** (i) *For fixed $R>1$ with $\mathcal{C}(R)\neq 0$,*

$$M_U^{\mathrm{lp}}(RU^2) \sim -\frac{RU^2}{2\log U}\mathcal{C}(R), \qquad M_U^{\mathrm{sm}}(RU^2) \sim +\frac{RU^2}{2\log U}\mathcal{C}(R); \tag{11.14}$$

for $1<R<2$, $\mathcal{C}(R)=1-1/R>0$.

(ii) *There is $C>0$ such that, as $t\to\infty$, uniformly for $U^2<t<U^4$ with $R=t/U^2\ge \exp(C(\log\log t)^2)$,*

$$M_U^{\mathrm{lp}}(t) = (1+o(1))\frac{t}{(\log t)^2}, \qquad M_U^{\mathrm{sm}}(t) = -(1+o(1))\frac{t}{(\log t)^2}. \tag{11.15}$$

In particular, in both regimes $|M_U^{\mathrm{lp}}(t)|$ and $|M_U^{\mathrm{sm}}(t)|$ exceed $t^{1/2+\varepsilon}$ for every fixed $\varepsilon<1/2$ and all large t, and only their sum M can be compared with the scale $t^{1/2+\varepsilon}$.

Proof. We use the classical form [14]: for some $c > 0$,

$$\pi(y) = \mathrm{li}(y) + O\big(ye^{-c\sqrt{\log y}}\big), \quad M(y) \ll ye^{-c\sqrt{\log y}}, \quad \widetilde{m}(y) := \sum_{a\le y} \frac{\mu(a)}{a} \ll e^{-c\sqrt{\log y}}, \tag{11.16}$$

and $\sum_{n\ge1} \mu(n)\log n/n = -1$ as a convergent series (the value at $s=1$ of $-\frac{d}{ds}\zeta(s)^{-1}$, since $\zeta'/\zeta^2 \to -1$). The integral below converges absolutely by the bound on $\widetilde{m}$, and

$$\int_1^\infty \widetilde{m}(w)\frac{dw}{w} = \lim_{Y\to\infty}\Big(\widetilde{m}(Y)\log Y - \sum_{a\le Y}\frac{\mu(a)\log a}{a}\Big) = 1. \tag{11.17}$$

(i) Proposition 11.9, and $M(t) = o(t/\log t)$. For $1 < R < 2$, $M = 1$ on $[1, 2)$.

(ii) Since $t < U^4$, $R < \sqrt{t}$, so $\log(t/a) \ge \frac12 \log t$ for $a \le R$. By (11.5),

$$M_U^{\mathrm{lp}}(t) = -\sum_{a\le R}\mu(a)\,\mathrm{li}(t/a) - \sum_{a\le R}\mu(a)\big(\pi(t/a) - \mathrm{li}(t/a)\big) + M(R)\pi(U^2). \tag{11.18}$$

The second term is $\ll t(1+\log R)e^{-c\sqrt{\log t/2}} = o(t/\log^2 t)$. For the third, $\frac14\log t < \log U < \frac12 \log t$, so

$$|M(R)|\pi(U^2) \ll Re^{-c\sqrt{\log R}}\frac{U^2}{\log U} \ll e^{-c\sqrt{\log R}}\frac{t}{\log t} = o(t/\log^2 t)$$

once $c\sqrt{C} > 1$. For the first, with $F(w) = w\,\mathrm{li}(t/w)$,

$$\sum_{a\le R}\mu(a)\,\mathrm{li}(t/a) = \widetilde{m}(R)F(R) - \int_1^R \widetilde{m}(w)F'(w)\,dw, \tag{11.19}$$

$$F'(w) = \mathrm{li}(t/w) - \frac{t/w}{\log(t/w)} = \frac{t/w}{\log^2(t/w)}\Big(1 + O\Big(\frac{1}{\log t}\Big)\Big).$$

The boundary term is $\ll e^{-c\sqrt{\log R}}t/\log t = o(t/\log^2 t)$. The integral equals

$$\frac{t}{\log^2 t}\left(\int_1^R \widetilde{m}(w)g_t(w)\frac{dw}{w} + O(1/\log t)\right), \qquad g_t(w) = \frac{\log^2 t}{\log^2(t/w)} \in [1, 4].$$

Extend $g_t(w)\mathbf{1}_{\{w\le R\}}$ by zero to $[1,\infty)$. Since $R \ge \exp(C(\log\log t)^2) \to \infty$, this factor tends to 1 pointwise and is bounded by 4, while $|\widetilde{m}(w)|/w$ is integrable; dominated convergence gives $\int_1^R \widetilde{m}(w)g_t(w)\,dw/w = 1 + o(1)$, where the $o(1)$ depends only on t and tends to 0 as $t\to\infty$ uniformly over all pairs (U, R) in the stated range: indeed $g_t(w)\mathbf{1}_{\{w\le R\}} \to 1$ uniformly for w in compact sets, since $R \ge \exp(C(\log\log t)^2)$ and $w \le W$ give $|g_t(w) - 1| \ll \log W/\log t$. Hence $M_U^{\mathrm{lp}}(t) = (1 + o(1))t/\log^2 t$, and $M_U^{\mathrm{sm}} = M - M_U^{\mathrm{lp}}$ with $M(t) = o(t/\log^2 t)$. □

Remark 11.14 (Why $t/\log^2 t$). $\mathcal{C}(R)$ captures only the $1/\log t$ term, whose coefficient tends to zero by Corollary 11.12. The $1/\log^2 t$ term comes from $\zeta'(s)/\zeta(s)^2 \to -1$, that is, from the pole of ζ at $s = 1$, and does not decay in R.

Remark 11.15 (First copy transitions). For $1 < R < 2$ every copy has length 1 and $M_U^{\mathrm{lp}}(t) = -(\pi(t) - \pi(U^2))$. For $2 < R < 3$ copies of length 2 appear but $M(2) = 0$. For $3 < R < 4$ copies of length 3 enter with $M(3) = -1$, of opposite net sign.

12. Local large-prime decomposition of an excursion

Proposition 12.1 (Local decomposition)**.** *Let $I_\nu = (u, v]$ be a complete excursion and $y = \sqrt{v}$. For $u < t \le v$,*

$$M(t) = \sum_{\substack{u<n\le t\\ P^+(n)\le y}} \mu(n) - \sum_{a<y}\mu(a)\,\Pi_a(t), \qquad \Pi_a(t) := \#\Big\{p:\ \max\big(y, \tfrac{u}{a}\big) < p \le \tfrac{t}{a}\Big\}, \tag{12.1}$$

and the bridge condition $M(v) = 0$ reads

$$\sum_{\substack{u<n\le v\\ P^+(n)\le\sqrt{v}}} \mu(n) = \sum_{a<\sqrt{v}} \mu(a)\,\Pi_a(v). \tag{12.2}$$

Proof. $M(u) = 0$. An $n \le v = y^2$ has at most one prime factor $p > y$; then $n = ap$ with $a < v/y = y < p$, so $\mu(n) = -\mu(a)$, and $(a,p) \mapsto ap$ is a bijection onto these n. □

Remark 12.2 (Primes in short intervals). The decomposition holds for every excursion, with threshold $\sqrt{v_\nu}$. The large-prime input consists of prime counts in the intervals $(u/a, t/a]$, of length at most ℓ_ν/a at height about v_ν/a. Huxley proved the prime number theorem uniformly in intervals of length $h \ge x^{7/12+\varepsilon}$ [7]; the zero-density estimates of Guth–Maynard give intervals of length $x^{17/30+o(1)}$ [3]. For Möbius sums, Matomäki–Teräväinen prove $\sum_{x<n\le x+x^\theta} \mu(n) = o(x^\theta)$ for every $\theta > 0.55$ [13]. The cited results do not provide, below these ranges, an asymptotic for Π_a uniform in a.

Lemma 12.3 (Inverse logarithms)**.** *Let C be as in Proposition 11.13. Uniformly for*

$$\exp\bigl(C(\log\log x)^2\bigr) \le z \le \sqrt{x},$$

$$\sum_{a\le z} \frac{\mu(a)}{a\log(x/a)} = -\frac{1+o(1)}{(\log x)^2}. \tag{12.3}$$

Proof. Partial summation gives

$$\frac{\widetilde{m}(z)}{\log(x/z)} - \int_1^z \frac{\widetilde{m}(w)}{w\log^2(x/w)}\,dw.$$

Since $z \le \sqrt{x}$, the boundary term is $O(e^{-c\sqrt{\log z}}/\log x) = o(1/\log^2 x)$ uniformly. Write $\log^{-2}(x/w) = g_x(w)\log^{-2} x$ with $1 \le g_x(w) \le 4$, extend $g_x \mathbf{1}_{\{w\le z\}}$ by zero, and apply dominated convergence with (11.17). □

Heuristic 12.4 (Opposite drifts)**.** Let $u \ge \sqrt{v}\exp(C(\log\log v)^2)$. Near $t \in I_\nu$ the counts Π_a grow for $a < t/\sqrt{v}$, at rate $1/(a\log(t/a))$ if primes near t/a have their expected local density uniformly in a. Over $(t, t+h]$ the large-prime part then changes by about $-h\sum_{a<t/\sqrt{v}} \mu(a)/(a\log(t/a)) = (1+o(1))h/\log^2 t$ by Lemma 12.3. Since $M(v) - M(u) = 0$, the $\sqrt{v}$-smooth part drifts at rate about $-1/\log^2 v$, independently of σ_ν. As

$$\frac{\ell_\nu}{\log^2 v_\nu} \gg \sqrt{\ell_\nu} \iff \ell_\nu \gg \log^4 v_\nu,$$

for $\ell_\nu \gg (\log v_\nu)^4$ both drift contributions exceed $\sqrt{\ell_\nu}$, and (12.2) is a balance between two opposite drifts: the local analogue of Proposition 11.13.

13. A one-frequency model for the black curve

Heuristic 13.1 (Scale and period)**.** Motivated by (5.1), model $M(x) \approx \sqrt{x}\,F(\log x)$ with F almost periodic, and retain only the lowest zero γ_1 (RH does not imply that its coefficient dominates). Within an excursion of length $\asymp x$ this gives $H_\nu \asymp \sqrt{x}$ and $\sqrt{H_\nu}\ell_\nu \asymp x^{5/4}$, so the black curve oscillates at scale

$$|\mathcal{A}(x)| \asymp x^{5/4}, \tag{13.1}$$

the scale of Corollary 6.4, with period $2\pi/\gamma_1$ in $\log x$, i.e. a factor $e^{2\pi/\gamma_1} \approx 1.5597$ in x. In this model consecutive zeros of M are separated by the factor $e^{\pi/\gamma_1} = \lambda_*$, the threshold of Corollary 8.13(iii): a model excursion is exactly λ_*-long.

14. Numerical data

The following census covers $n \le N = 455\,052\,511$, with $Z(N) = 87\,385$ excursions, recording $(u_\nu, \tau_\nu, v_\nu, H_\nu, \sigma_\nu)$ and the curves $\mathcal{A}_+, \mathcal{A}_-, \mathcal{A}$:

$$\mathcal{A}_+(N) = 6.3449 \times 10^9, \quad \mathcal{A}_-(N) = -7.0877 \times 10^9, \quad \mathcal{A}(N) = -7.4279 \times 10^8, \tag{14.1}$$

$$\max_\nu \frac{H_\nu}{\sqrt{\ell_\nu}} = 2.0743, \qquad \max_\nu \frac{\beta_\nu}{\ell_\nu^{3/2}} = 1.0214. \tag{14.2}$$

Since $|M(x)| \le 0.571\sqrt{x}$ for $33 \le x \le 10^{16}$ [6], and $|M(n)| \le \sqrt{n}$ for $n < 33$ with equality only at $n = 1$, the constant $C_0 = 1$ is admissible in Lemma 6.1, which gives $\mathcal{A}_+(N) + |\mathcal{A}_-(N)| \le \frac{4}{5}N^{5/4} = 5.317 \times 10^{10}$ against the observed 1.3433×10^{10}, a ratio of 3.96. Heuristic 7.5 with $L = \sqrt{N}$ predicts $\#\{\ell_\nu > \sqrt{N}\} \asymp N^{1/4} \approx 146$. At this range $N^{0.05} \approx 2.7$, so factors x^ε are not separable from constants. Reproducing these values requires the excursion file, the area-integration convention of Proposition 3.2, and the terminal convention of Definition 2.6. These data are not used in any proof.

15. Concluding remarks

The main results can be summarized in four statements. First, the Kaczorowski–Pintz oscillation density, combined with elementary counts of lacunary families of excursions, localizes Littlewood's criterion to the initial part $x_0 < t \le \min\{x_1, \lambda x_0\}$ of each excursion for every $\lambda > \lambda_* = e^{\pi/\gamma_1}$, the exact threshold of the counting mechanism; in averaged form, the excursions carrying large values have mean logarithmic length at most $\log \lambda_*$. Second, on this window the future increments depend directly only on the Möbius prefix up to θu for every $\theta > \theta_* = \frac{1}{2}e^{\pi/\gamma_1}$, and the endpoints of late excursions satisfy $x_1 = x_0^{1+o(1)}$. Third, the remaining difficulty is not one of dependency: the one-step recursion, the relaxed divisor-sum cone and any bounded number of multiplicative layers provably fail to produce the RH scale by themselves. Fourth, in the language of two-point correlations, RH is equivalent to cancellation between lags of the h-weighted correlations at the zeros of M; uniform two-point Cesàro cancellation is available in all intervals of length $\ge u^{3/5+\eta}$ at mesoscopic lags, and the missing inputs are precisely a translated sublinear averaged Chowla estimate on blocks selected by the zeros of M for lengths between $u^{1/2}$ and $u^{3/5}$, and a local variance estimate of diagonal size in the sense of Lemma 9.12.

16. Open problems

(1) Extend Theorem 9.9 to excursions with $x_0^{1/2+\varepsilon} \le \ell < x_0^{3/5}$, and to lags $R \le \ell(\log x_0)^{-2/3}$; equivalently, prove for blocks starting at zeros of M the local variance statement $\sum_{h<R}(R-h)C_h^{(\nu)} = o(R\ell)$ of Lemma 9.12.

(2) In the localized form of Corollary 8.50, bound

$$\sum_{d \le 5u/8} \mu(d)\big(\lfloor t/d \rfloor - \lfloor u/d \rfloor\big)$$

by $t^{1/2+\varepsilon}$ on $u < t \le \min\{x_1, 5u/4\}$, using the full simultaneous intra-excursion constraints rather than pointwise estimates.

(3) Quantify the multiscale persistence forced by Lemma 8.45. By Corollary 8.47 and Remark 8.48, channel descent iterates but loses the factor $2K/k^\alpha$ per step; the problem is to show that along a descent chain from an RH-violating value the loss is eventually absent, or that the deep remainder $\mathcal{R}_K$ carries the value and can be bounded.

(4) Question 8.63 in the form left by Corollary 8.65: bounded depth fixes only $o(u)$ window increments, whose sum is of order $u(\log\log u)^k/\log u$; decide whether the unit-step and excursion constraints on the remaining free coordinates δ_d, $\Omega(d) > k$, already force the cancellation down to $u^{1/2+\varepsilon}$. The first case is $k = 1$ (primes fixed, $\delta_p = -1$).
(5) Can the threshold $\lambda_* = e^{\pi/\gamma_1}$ of Corollary 8.13 be lowered? By Remark 8.29 this requires information on the position of large values inside excursions, beyond the count (8.6).
(6) Whether (6.4) is equivalent to RH.
(7) Whether the pruned local condition (7.7) holds, and whether the bridge constraints (4.5), (4.6) admit a third independent relation involving the defect $\ell_\nu/2 - H_\nu \geq 0$.
(8) Establish (SG) for $Z = H/\sqrt{\ell}$ in the simple random walk model of the Möbius walk, and compute the extreme scale of Corollary 8.69 there; the Brownian-excursion limit suggests $c = 2$.
(9) Numerically: the number of excursions with $\ell_\nu > \sqrt{N}$ against $N^{1/4}$, the maxima of $H_\nu/\sqrt{\ell_\nu}$ and $\beta_\nu/\ell_\nu^{3/2}$ over them; the split (12.1) and the σ_ν-independence of Heuristic 12.4; the frequency of touching runs and of ballistic excursions (Heuristic 7.8); and the distribution of x_1/x_0, to be compared with λ_*.

References

[1] P. Carmona and N. Pétrélis, *Limit theorems for random walk excursion conditioned to enclose a typical area*, arXiv:1709.06448.
[2] D. Denisov, M. Kolb and V. Wachtel, *Local asymptotics for the area of random walk excursions*, J. London Math. Soc. **91** (2015), 495–513.
[3] L. Guth and J. Maynard, *New large value estimates for Dirichlet polynomials*, Ann. of Math. (2) **203** (2026), no. 2, 623–675, doi:10.4007/annals.2026.203.2.6.
[4] D. R. Heath-Brown, *Prime numbers in short intervals and a generalized Vaughan identity*, Canad. J. Math. **34** (1982), 1365–1377.
[5] H. A. Helfgott and L. Thompson, *Summing $\mu(n)$: a faster elementary algorithm*, Res. Number Theory **9** (2023), Paper No. 6.
[6] G. Hurst, *Computations of the Mertens function and improved bounds on the Mertens conjecture*, Math. Comp. **87** (2018), no. 310, 1013–1028.
[7] M. N. Huxley, *On the difference between consecutive primes*, Invent. Math. **15** (1972), 164–170.
[8] S. Janson, *Brownian excursion area, Wright's constants in graph enumeration, and other Brownian areas*, Probab. Surveys **4** (2007), 80–145.
[9] J. Kaczorowski and J. Pintz, *Oscillatory properties of arithmetical functions. I*, Acta Math. Hungar. **48** (1986), 173–185.
[10] T. Kotnik and H. te Riele, *The Mertens conjecture revisited*, in Algorithmic Number Theory (ANTS VII), Lecture Notes in Comput. Sci. **4076**, Springer, Berlin, 2006.
[11] K. Matomäki, M. Radziwiłł and T. Tao, *An averaged form of Chowla's conjecture*, Algebra Number Theory **9** (2015), no. 9, 2167–2196.
[12] K. Matomäki, X. Shao, T. Tao and J. Teräväinen, *Higher uniformity of arithmetic functions in short intervals I. All intervals*, Forum Math. Pi **11** (2023), Paper No. e29.
[13] K. Matomäki and J. Teräväinen, *On the Möbius function in all short intervals*, J. Eur. Math. Soc. **25** (2023), no. 4, 1207–1225.
[14] H. L. Montgomery and R. C. Vaughan, *Multiplicative Number Theory I: Classical Theory*, Cambridge Stud. Adv. Math. **97**, Cambridge University Press, 2007.
[15] N. Ng, *The distribution of the summatory function of the Möbius function*, Proc. London Math. Soc. (3) **89** (2004), no. 2, 361–389.
[16] M. Nguyên Thê, *Area and inertial moment of Dyck paths*, Combin. Probab. Comput. **13** (2004), 697–716.
[17] A. M. Odlyzko and H. J. J. te Riele, *Disproof of the Mertens conjecture*, J. Reine Angew. Math. **357** (1985), 138–160.
[18] C. Richard, *On q-functional equations and excursion moments*, arXiv:math/0503198.
[19] L. Takács, *A Bernoulli excursion and its various applications*, Adv. Appl. Probab. **23** (1991), 557–585.
[20] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2nd ed., revised by D. R. Heath-Brown, Clarendon Press, Oxford, 1986.

[21] A. Walfisz, *Weylsche Exponentialsummen in der neueren Zahlentheorie*, VEB Deutscher Verlag der Wissenschaften, Berlin, 1963.