ORIENTATION-BIT GEOMETRY OF THE MÖBIUS PATH: REFLECTION FIBRES, DIVISOR–SHIFT BOUNDARY COCYCLES, AND AN OBSERVABILITY ROUTE TO TWO-POINT CANCELLATION

ABSTRACT. This notebook develops an operator framework for the two-point correlation $C_N(h) = \sum_{n \le N-h} \mu(n)\mu(n+h)$ built on the orientation bits of the Möbius path and on the divisor operator.

Version 3 adds an analytic layer to the combinatorial one. The central new object is the difference $(U_h - I)D$ rather than the commutator or the conjugation: each divisor column collapses to the difference of two residue classes, vanishes identically when the divisor divides h, and has interval sums bounded by one. A purely algebraic Haar basis over an arbitrary binary tree then annihilates the volumetric CRT term exactly, leaving only boundary cocycles which satisfy a parent–children additivity law. No doubling, no comparable lengths and no regularity of the squarefree mask are required at any point.

Four further results are proved. An exact observability criterion: $|x^{\mathsf{T}}Ax| \le \kappa \|Lx\|^2$ holds for some finite κ if and only if $\ker L \subseteq \ker A$, with the optimal constant and the exact operator-norm dual given in closed form. A no-go: uniform observability of the squarefree shift by a sublinear set of divisor rows is impossible, because above $N/2$ each column is seen only by its own row and shifted squarefree pairs have positive density. An extension of the blindness theorem to kernels that are constant on each half of an excursion, under which the cross-excursion interaction factors through the single channel $s_j H_j$ and the shift kernel is shown not to belong to the extended class. And a geometric proposition: the hypothesis $u_j \ge m_j/2$ needed for the central divisor matrix to be triangular with nonvanishing diagonal fails for at most $\lfloor \log_2 N \rfloor$ excursions, so that the Möbius line is isolated up to a kernel of logarithmic dimension.

The results of Versions 1 and 2 are retained: the reflection fibre of the quadratic profile, the exactly isotropic null, the blindness theorem for kernels depending on $\max(a, b)$, and the criterion separating tautological from informative sine-kernel content. Negative results, including the collapse of every multiplicative diagonal weight onto $1/\zeta(1+s)$, are recorded in full. No numerical computation is performed.

CONTENTS

1. Introduction 2
1.1. Objects 2
1.2. What is proved in this notebook 2
1.3. Inputs and the methodological point 3
2. Standing results used as inputs 3
3. Three spaces and the maps between them 4
4. The three-constraint model 4
5. The reflection fibre 4
6. The divisor-defect Gram form on orientation bits 5
7. Blindness, and its exact extension 5
7.1. Two channels and the half-constant extension 6
8. The divisor–shift operator 6
8.1. What conjugation does not give 6
8.2. The difference that localises 7
8.3. CRT pullback of the shift 7
9. Algebraic Haar on arbitrary trees, and exact bulk vanishing 7
10. Observability 8

Date: Version 3.

11. Centres, divisor rows and the Möbius line 9
11.1. The central dichotomy 9
11.2. Sampling the divisor equation at the centre 9
12. Intrablock identities 10
13. Exact local structure at each prime 11
14. Multiplicative weights collapse 11
15. Scale-resolved statistic and sine-kernel content 12
16. Strategy and the target theorem 13
17. Negative results and failed routes 14
18. Open questions 15
Notebook History 15
References 15

Remark 0.1. This notebook is exploratory. Rigorous statements appear inside theorem, proposition, lemma or corollary environments with proofs. Results established in separate work are collected in Section 2 and stated without proof. Everything else is labelled heuristic, conjectural or objective. No numerical computation is performed; the empirical values in Remark 15.11 are inputs from separate work.

Warning 0.2. Bibliographic metadata in Section 18 has not been verified against primary sources. DOIs are omitted where not confirmed.

1. Introduction

1.1. **Objects.** Let μ be the Möbius function, $q(n) = \mu(n)^2$,

$$M(x) = \sum_{n \le x} \mu(n), \qquad C_N(h) = \sum_{n \le N-h} \mu(n)\mu(n+h). \tag{1.1}$$

Let D be the divisor operator $(Dx)(r) = \sum_{d|r} x(d)$ and U_h the truncated shift $(U_h x)(n) = x(n+h)$, with symmetric part $A_h = (U_h + U_h^*)/2$, so that

$$C_N(h) = \mu^{\mathsf{T}} A_h \mu. \tag{1.2}$$

1.2. **What is proved in this notebook.**

(i) *Reflection fibre* (Prop. 5.2): the fibre of the quadratic profile is exactly an orbit of $\mathcal{R}_N \cong (\mathbb{Z}/2)^{Z(N)}$.
(ii) *Exact isotropy* (Prop. 5.4): $\mathbb{E}_\nu[ss^{\mathsf{T}}] = I_Z$ with no correction term, and closed-form variance.
(iii) *Blindness* (Thm. 7.2): every cross-excursion contribution to a $\max(a,b)$-kernel vanishes identically.
(iv) *Extended blindness* (Thm. 7.5): for kernels constant on each half of an excursion the cross-excursion interaction factors through the rank-one channel $s_j H_j$; the shift kernel is shown not to lie in this class.
(v) *Divisor–shift boundary lemma* (Lem. 8.2): $((U_h - I)D)_{n,d} = \mathbf{1}[d \mid n+h] - \mathbf{1}[d \mid n]$, identically zero for $d \mid h$, supported on two residue classes otherwise, with interval sums bounded by 1.
(vi) *Exact CRT pullback* (Lem. 8.4): $(D^{\mathsf{T}} U_h D)_{d,e}$ vanishes unless $(d,e) \mid h$ and otherwise equals $(N-h)/[d,e]$ up to an error of modulus at most 1.
(vii) *Algebraic Haar and exact bulk vanishing* (Lem. 9.1, Thm. 9.3): on an arbitrary binary tree the volumetric CRT term is annihilated exactly and only a boundary cocycle survives, additive over children.

(viii) *Observability criterion* (Thm. 10.2): $|x^{\mathsf{T}}Ax| \le \kappa\|Lx\|^2$ for some $\kappa < \infty$ iff $\ker L \subseteq \ker A$; optimal constant and operator-norm dual in closed form.
(ix) *Sublinear no-go* (Thm. 10.4): no nontrivial observability of the squarefree shift by $o(N)$ divisor rows.
(x) *Logarithmic exceptional set* (Prop. 11.6): the geometric hypothesis making the central divisor matrix triangular with nonvanishing diagonal fails for at most $\lfloor \log_2 N \rfloor$ excursions.
(xi) *Central dichotomy* (Lem. 11.1): every linear reflection-odd functional of an excursion either vanishes on the whole fibre or reads the orientation bit directly.

1.3. **Inputs and the methodological point.** Results established in separate work are listed with status in Section 2; the two used actively are the tensor factorisation (S4) and the permutation-null mean (S10).

The methodological criterion of Version 2 is retained: a statistic carries sine-kernel content non-tautologically if and only if its value under the null differs from its value under the sine kernel (Remark 15.6). Version 3 adds the corresponding criterion for weights: every diagonal weight with a multiplicative symbol collapses onto $1/\zeta(1+s)$ (Thm. 14.1), so informative weights must break the Euler factorisation.

2. Standing results used as inputs

Established in separate work by the author, stated without proof, no novelty claimed here.

Remark 2.1. (S1) $c\sqrt{N} \le \lambda_{\max}(D_N) \le C_\varepsilon N^{1/2+\varepsilon}$ with Cauchy interlacing corollaries. *Context: Rem. 17.5.*
(S2) Trace-norm convergence of G_N/N^2 to a compact limit operator. *Context.*
(S3) Exact Chinese-remainder formula for the entries of Q_N. *Context; source of (S4).*
(S4) The tensor factorisation

$$\widehat{K}(r,q) = \prod_p L_p(v_p(r), v_p(q)) \tag{2.1}$$

over all of $\mathbb{N}\times\mathbb{N}$, explicit tridiagonal local factors, sign corrected. *Used actively as (G2).*
(S5) $C_{\mathrm{herm}}(\rho) = 2(\gamma^2-\beta^2)/(|\rho|^4|\zeta'(\rho)|^2) > 0$ for every nontrivial zero, without assuming RH. *Context; see Rem. 2.2.*
(S6) $z_0''(n) = -\frac{r^2}{2}\operatorname{Var}_{p|n}(\log p)$ for squarefree n with r prime factors, and $K(s) = \frac{\zeta(s)}{2}(S_0S_2 - S_1^2)$ with $S_2 = (\zeta'/\zeta)'$. *Context.*
(S7) Closed form of $z_\vartheta(n)$ as a finite cosine sum over ordered partitions via the cocycle $\Phi(\pi)$, with $A_r = 3(6r-11)/8$, $B_r = r(6r-11)/8$ exact for all r. *Context; source of the twisted action in Conj. 15.14.*
(S8) Blindness to Θ of the whole tower of *fixed* transversal jets, with complete proof. *Context; the restriction to fixed jets leaves the scaling window open.*
(S9) Unconditional equivalence between RH and tail stability of a two-channel positive logarithmic deformation of the ordinal-prime transform. *Context.*
(S10) $\mathbb{E}_\pi[E_N]/N = 1/\pi^2 + M(N)^2/(3N) + o(1)$ with $E_N = \frac{1}{N}\sum_{n\le N} M(n)^2$. *Used actively.*

Remark 2.2. At the extremal abscissa $s = 2\Theta$ all contributions come from zeros with $\beta = \Theta$ and carry the positive coefficient of (S5); a sum of positive terms does not cancel, so cancellation between distinct pairs cannot occur there. What remains open is the case of a non-attained supremum, requiring a limiting formulation.

3. Three spaces and the maps between them

Definition 3.1. Let $A_N = \{a_1 < \cdots < a_m \le N : q(a_j) = 1\}$, $m = |A_N|$. Set

$$H_N = \mathbb{R}^N \qquad \text{(full space; } D,\ U_h\text{)}, \tag{3.1}$$

$$H_{\mathrm{sf}} = \mathbb{R}^m \qquad \text{(squarefree space)}, \tag{3.2}$$

$$H_{\mathrm{exc}} = \mathbb{R}^Z \qquad \text{(excursion space)}. \tag{3.3}$$

Let $Je_j = e_{a_j}$, so $J^*J = I$ and $P = JJ^* = \mathrm{diag}(q)$, and let F have columns $F_j = \mu\,\mathbf{1}_{E_j}$.

Remark 3.2. The three spaces retain different information. H_N retains position, triangularity, divisibility and boundary data. H_{sf} retains signs and the integer labels a_j; if the labels are discarded, divisibility is lost. H_{exc} retains only block orientations: all intrablock variation, gcd classes, valuations and individual positions are destroyed. Since $P\mu = \mu$, the inclusion J loses nothing for the vector μ itself, but it does lose operator information about perturbations off the mask.

Proposition 3.3. *With $U_h^{\mathrm{sf}} = J^*U_hJ$, $A_h = (U_h + U_h^*)/2$ and $x = Fs + x_{\mathrm{term}}$ on the fibre,*

$$C_N(h) = s^{\mathsf{T}}(F^*A_hF)s + 2s^{\mathsf{T}}F^*A_hx_{\mathrm{term}} + x_{\mathrm{term}}^{\mathsf{T}}A_hx_{\mathrm{term}}. \tag{3.4}$$

Proof. Substitute $x = Fs + x_{\mathrm{term}}$ into (1.2) and expand the symmetric form. □

Remark 3.4. Equation (3.4) is exact, but by Cor. 7.3 the excursion space is a severe compression and must be subordinate: it is used in Section 11 only for the reflection-odd component and as one possible family of stopping blocks. The primary space is H_N.

4. The three-constraint model

Definition 4.1. The orientation bits are $\varepsilon = (\varepsilon_n)_{n\in A_N} \in \{\pm1\}^m$, $\varepsilon_n = \mu(n)$. With $m_\pm$ the number of $n \in A_N$ with $\mu(n) = \pm1$ and $\bar\varepsilon = M(N)/m$, the *fixed-count sign bridge* π is uniform on the vectors of $\{\pm1\}^m$ with m_+ entries $+1$.

Proposition 4.2. *Under π,*

$$\mathbb{E}_\pi[\varepsilon\varepsilon^{\mathsf{T}}] = (1-\beta)I + \beta J_m, \qquad \beta = \frac{m\bar\varepsilon^{\,2} - 1}{m-1}, \tag{4.1}$$

J_m the all-ones matrix. Consequently, for an orthogonal projection P' of rank d,

$$\mathbb{E}_\pi[\langle\varepsilon, P'\varepsilon\rangle] = (1-\beta)d + \beta\|P'\mathbf{1}\|^2 = d + O(1) \quad \text{if } M(N) = o(\sqrt{m}). \tag{4.2}$$

Proof. Diagonal entries are 1. For $n \ne n'$, counting ordered pairs of distinct positions gives $((m_+ - m_-)^2 - m)/(m(m-1))$, and $m_+ - m_- = m\bar\varepsilon$. Trace against P' using $\mathrm{tr}\,P' = d$, $\mathrm{tr}(P'J_m) = \|P'\mathbf{1}\|^2$; for the last claim $|\beta| \le 1/(m-1)$ once $M(N)^2 \le m$. □

Remark 4.3. The factorisation of $\mathrm{Cov}_\pi(M(n)^2, M(n')^2)$ through the three-block decomposition $(n, n'-n, N-n')$, with its $O(m)$ variance evaluation, is recorded separately. Only its structural consequence is used: the covariance kernel is separable of bounded rank. By Prop. 5.4 the third conditioning replaces this by an exact closed form.

5. The reflection fibre

Definition 5.1. Let $0 = z_0 < z_1 < \cdots < z_Z$ enumerate $\{n \le N : M(n) = 0\} \cup \{0\}$, and

$$E_j = \{n : z_{j-1} < n \le z_j\}, \qquad j = 1, \dots, Z, \tag{5.1}$$

with terminal block $E_{Z+1} = \{n : z_Z < n \le N\}$, empty iff $M(N) = 0$. Write $u_j = z_{j-1}$, $v_j = z_j$. The reflection group $\mathcal{R}_N \cong (\mathbb{Z}/2)^Z$ acts by $(\sigma\cdot\varepsilon)_n = \sigma_{j(n)}\varepsilon_n$, the terminal block held fixed.

Proposition 5.2. *$M_{\sigma\cdot\varepsilon}(n) = \sigma_{j(n)} M_\varepsilon(n)$, so the quadratic profile $\Pi_N = (M(n)^2)_{n\le N}$ is $\mathcal{R}_N$-invariant; conversely two sign vectors on the same mask with the same profile differ by an element of $\mathcal{R}_N$. The fibre of Π_N is exactly the orbit.*

Proof. For $n \in E_j$, $M_\varepsilon(z_{j-1}) = 0$ so $M_\varepsilon(n) = \sum_{z_{j-1}<k\le n} \varepsilon_k$ lies inside E_j and σ multiplies it by σ_j. Conversely equal profiles give the same zero set, hence the same blocks, and $|M|$ pointwise; inside a block M has constant sign, and the ratio of the two signs defines σ_j; increments determine the path. □

Definition 5.3. The excursion signs $s \in \{\pm1\}^Z$ record the sign of M in the interior of each E_j. The reflection null ν is uniform on $\{\pm1\}^Z$. Write $Q_j(n) = |M(n)|$ for $n \in E_j$, so that

$$M_s(n) = s_j Q_j(n), \qquad \mu_s(n) = s_j\, \Delta Q_j(n), \qquad n \in E_j. \tag{5.2}$$

Proposition 5.4. *Under ν, $\mathbb{E}_\nu[ss^{\mathsf T}] = I_Z$ exactly, hence for symmetric A*

$$\mathbb{E}_\nu[\langle s, As\rangle] = \operatorname{tr} A, \qquad \operatorname{Var}_\nu[\langle s, As\rangle] = 2\sum_{j\ne k} A_{jk}^2, \tag{5.3}$$

and for a projection $\widehat{P}$ of rank d, $\mathbb{E}_\nu = d$ and $\operatorname{Var}_\nu = 2(d - \sum_j \widehat{P}_{jj}^2)$.

Proof. The s_j are independent Rademacher. Write $\langle s, As\rangle = \operatorname{tr} A + \sum_{j\ne k} A_{jk} s_j s_k$ and use $\mathbb{E}_\nu[s_j s_k s_{j'} s_{k'}] = \mathbf{1}[\{j,k\} = \{j',k'\}]$; for a projection $\sum_{j,k} \widehat{P}_{jk}^2 = \operatorname{tr}\widehat{P} = d$. □

Remark 5.5. Conditioning on the mask removes the $1/\pi^2$ of (S10); conditioning on the count removes the drift $M(N)^2/(3N)$; conditioning in addition on the profile leaves exactly $\mathcal{R}_N$, whose null is isotropic without the $O(1)$ correction of (4.2). The third conditioning is where the null becomes parameter-free.

6. The divisor-defect Gram form on orientation bits

Definition 6.1. Let B be the divisor-defect matrix on A_N and $G = B^{\mathsf T}B$. G is *admissible* if (G1) it is real symmetric positive semidefinite and (G2) its kernel admits the tensor factorisation (2.1). The orientation-bit form is

$$\widehat{G}_{jk} = \sum_{a\in E_j}\sum_{b\in E_k} G_{ab}, \qquad \widehat{G} = \sum_k \lambda_k \widehat{P}_k, \quad d_k = \operatorname{rank}\widehat{P}_k. \tag{6.1}$$

Proposition 6.2. *$\langle s, \widehat{G}s\rangle$ is invariant under $s \mapsto -s$ and its off-diagonal part is odd under each $s_j \mapsto -s_j$; it is not a function of the quadratic profile unless $\widehat{G}$ is diagonal.*

Proof. $s_j s_k$ changes sign under $s_j \mapsto -s_j$ for $j \ne k$, while $\widehat{G}_{jj} s_j^2$ is constant; by Prop. 5.2 functions of the profile are $\mathcal{R}_N$-invariant. □

7. Blindness, and its exact extension

Lemma 7.1. $\displaystyle\sum_{n\le N} M(n)^2 = \sum_{a,b\le N} \mu(a)\mu(b)\big(N + 1 - \max(a,b)\big).$

Proof. Expand $M(n)^2$ and exchange summation; the pair (a,b) occurs for $N+1-\max(a,b)$ values of n. □

Theorem 7.2. *Assume $M(N) = 0$. For all $j \ne k$,*

$$\sum_{a\in E_j}\sum_{b\in E_k} \mu(a)\mu(b)\big(N + 1 - \max(a,b)\big) = 0. \tag{7.1}$$

Hence $\sum_{n\le N} M(n)^2$ depends only on within-excursion two-point data.

Proof. For $j < k$ every $a \in E_j$ precedes every $b \in E_k$, so $\max(a,b) = b$ and the double sum factorises as $\big(\sum_{a\in E_j} \mu(a)\big)\big(\sum_{b\in E_k} \mu(b)(N+1-b)\big)$; the first factor is $M(z_j) - M(z_{j-1}) = 0$. The case $j > k$ is symmetric. □

Corollary 7.3. *No statistic factoring through Π_N detects cross-excursion correlation of μ.*

Proof. Such a statistic is $\mathcal{R}_N$-invariant by Prop. 5.2; by Thm. 7.2 the cross-excursion mass does not enter the energy, and by the action it is exactly the part on which $\mathcal{R}_N$ acts nontrivially. □

7.1. **Two channels and the half-constant extension.**

Lemma 7.4. *Let m_j be an interior cut of E_j, $E_j^L = (u_j, m_j]$, $E_j^R = (m_j, v_j]$, and $H_j = Q_j(m_j)$. Then*

$$\sum_{n\in E_j^L} \mu_s(n) = s_j H_j, \qquad \sum_{n\in E_j^R} \mu_s(n) = -s_j H_j, \qquad H_j \geq 1. \tag{7.2}$$

In particular the two halves are not independent channels: they have rank one.

Proof. Telescoping and $M(u_j) = M(v_j) = 0$ give the two sums; $H_j \geq 1$ because M does not vanish in the interior of an excursion. □

Theorem 7.5. *Let $j < k$ and let a kernel $K(a,b)$ be constant in a on each half of E_j, say $K(a,b) = K_L(b)$ for $a \in E_j^L$ and $K_R(b)$ for $a \in E_j^R$. Then*

$$\sum_{a\in E_j}\sum_{b\in E_k} \mu_s(a)\mu_s(b)K(a,b) = s_j H_j \sum_{b\in E_k} \mu_s(b)\big(K_L(b) - K_R(b)\big). \tag{7.3}$$

If $K_L = K_R$ the exact vanishing of Thm. 7.2 is recovered; otherwise the interaction factors through the single channel $s_j H_j$.

Proof. Split the a-sum over the two halves and apply Lemma 7.4 to each. □

Remark 7.6. Theorem 7.2 is the case $K_L = K_R$, because a kernel depending on $\max(a,b)$ loses all a-dependence when $j < k$. The shift kernel $K_h(a,b) = \mathbf{1}[b - a = h]$ is not constant on any half: it resolves $a = b - h$ at unit scale. Hence the shift lies outside the extended class, and this is the precise sense in which $C_N(h)$ escapes the excursion compression.

8. The divisor–shift operator

8.1. **What conjugation does not give.**

Remark 8.1. D is lower triangular with $D^{-1}_{r,d} = \mu(r/d)\mathbf{1}[d \mid r]$, and

$$(DU_hD^{-1})_{r,m} = \sum_{\substack{a\mid r,\ a+h\leq N\\ m\mid a+h}} \mu\Big(\frac{a+h}{m}\Big). \tag{8.1}$$

This is exact and gives no spectral gain: a similarity preserves the problem. The entries of the commutator,

$$[D, U_h]_{r,m} = \mathbf{1}[m > h]\mathbf{1}[m - h \mid r] - \mathbf{1}[r \leq N - h]\mathbf{1}[m \mid r + h], \tag{8.2}$$

are not supported on primes dividing h: they compare the two distinct geometries $m - h \mid r$ and $m \mid r + h$.

8.2. **The difference that localises.**

Lemma 8.2 (Divisor–shift boundary lemma)**.**

$$\big((U_h - I)D\big)_{n,d} = \mathbf{1}[d \mid n+h] - \mathbf{1}[d \mid n]. \tag{8.3}$$

(i) If $d \mid h$ the column vanishes identically.
(ii) If $d \nmid h$ the column is supported on the two disjoint classes $n \equiv -h$ and $n \equiv 0 \pmod d$, with signs $+1$ and -1.
(iii) For every interval I of integers,

$$\Big|\sum_{n\in I}\big(\mathbf{1}[d \mid n+h] - \mathbf{1}[d \mid n]\big)\Big| \le 1. \tag{8.4}$$

Proof. (8.3) is $D_{n+h,d} - D_{n,d}$. (i) If $d \mid h$ then $d \mid n+h \iff d \mid n$. (ii) Immediate. (iii) The sum equals the number of multiples of d in $I+h$ minus the number in I; two intervals of equal length contain numbers of multiples differing by at most one. □

Remark 8.3. This converts the multiplicative–additive interaction into residue-class boundaries of uniformly bounded mass, with no regularity assumption on I. The bound 1 improves the bound 2 obtained by estimating the two classes separately.

8.3. **CRT pullback of the shift.**

Lemma 8.4.

$$(D^{\mathsf T}U_hD)_{d,e} = \#\{n \le N-h : d \mid n,\ e \mid n+h\}, \tag{8.5}$$

which vanishes unless $(d,e) \mid h$. If $(d,e) \mid h$ there is a unique class $c \bmod [d,e]$ and

$$(D^{\mathsf T}U_hD)_{d,e} = \frac{N-h}{[d,e]} + \varepsilon_{d,e,h,N}, \qquad |\varepsilon_{d,e,h,N}| \le 1. \tag{8.6}$$

Proof. The entry counts n with $n \equiv 0 \pmod d$ and $n \equiv -h \pmod e$. Compatibility of the two congruences requires $0 \equiv -h \pmod{(d,e)}$; when it holds the solution set is a single class modulo $[d,e]$, whose count in an interval of length $N-h$ is the stated main term with error of modulus at most 1. □

Remark 8.5. The bulk is an explicit gcd/lcm kernel and all finite irregularity sits in a boundary term. Section 9 removes the bulk exactly.

9. Algebraic Haar on arbitrary trees, and exact bulk vanishing

Lemma 9.1. *Let $I = I_0 \sqcup I_1$ with masses $m_0, m_1 > 0$, $m = m_0 + m_1$, and*

$$\psi_I = \sqrt{\frac{m_1}{m_0 m}}\,\mathbf{1}_{I_0} - \sqrt{\frac{m_0}{m_1 m}}\,\mathbf{1}_{I_1}. \tag{9.1}$$

Then $\langle \psi_I, \mathbf{1}_I\rangle = 0$ and $\|\psi_I\|_2 = 1$. For any binary tree terminating in singletons, the normalised constant together with the ψ_I form an orthonormal basis.

Proof. $\langle \psi_I, \mathbf{1}_I\rangle = \sqrt{m_0 m_1/m} - \sqrt{m_0 m_1/m} = 0$ and $\|\psi_I\|_2^2 = m_1/m + m_0/m = 1$. Orthogonality of two Haar functions follows from disjointness of their supports or, when nested, from the mean-zero property on the child in which the finer one is supported. Counting gives a basis. □

Remark 9.2. No doubling measure, comparable children or geometric hypothesis is used. The identity is finite-dimensional algebra.

Theorem 9.3 (Exact vanishing of the CRT bulk)**.** *Let* $R_{d,e,h}(n) = \mathbf{1}[d \mid n]\mathbf{1}[e \mid n+h]$. *If* $(d,e) \nmid h$ *then* $R \equiv 0$. *Otherwise* $R = \mathbf{1}[n \equiv c \bmod q]$ *with* $q = [d,e]$, *and writing* $\#(I_j \cap (c \bmod q)) = m_j/q + \epsilon_j$ *with* $|\epsilon_j| < 1$,

$$\langle R_{d,e,h}, \psi_I \rangle = \sqrt{\frac{m_1}{m_0 m}}\,\epsilon_0 - \sqrt{\frac{m_0}{m_1 m}}\,\epsilon_1. \tag{9.2}$$

Moreover the boundary defects are additive over children:

$$\epsilon_I(c,q) = \epsilon_{I_0}(c,q) + \epsilon_{I_1}(c,q). \tag{9.3}$$

Proof. The compatibility statement is Lemma 8.4. Insert the counts into (9.1): the terms m_j/q contribute $\frac{1}{q}(\sqrt{m_0 m_1/m} - \sqrt{m_0 m_1/m}) = 0$ because ψ_I has mean zero, leaving (9.2). Additivity holds because the counts add and $m_I = m_{I_0} + m_{I_1}$. □

Remark 9.4. The irregularity of the cut is not an error term: it *is* the cocycle (ϵ_0, ϵ_1). By (9.3) the Haar coefficients are martingale differences of a single boundary cocycle, which opens a telescoping across scales that the excursion compression cannot see.

Warning 9.5. Summing absolute values is not enough: by Parseval a single residue class can carry large Haar energy. Any new estimate must use simultaneously the inclusion–exclusion signs, the compatibilities $(d,e) \mid h$ and the parent–children coherence (9.3).

Remark 9.6. The shift matrix in this basis is exact:

$$\langle \psi_I, U_h \psi_J \rangle = \sum_{\alpha,\beta \in \{0,1\}} c_{I,\alpha} c_{J,\beta}\, E_h(I_\alpha, J_\beta), \qquad E_h(X,Y) = \#\{n \in X : n+h \in Y\}, \tag{9.4}$$

with $c_{I,0}, c_{I,1}$ the two coefficients of (9.1). There is an entry only when an edge joins the supports; cancellation depends on the 2×2 incidence table, not on regular lengths.

10. Observability

Definition 10.1. Let $V = PH_N$, $L = D_\Lambda P|_V$ for a set Λ of divisor rows, and $A = PA_hP|_V$.

Theorem 10.2 (Exact observability criterion)**.** *There exists* $\kappa < \infty$ *with*

$$|x^{\mathsf{T}} A x| \le \kappa \|Lx\|^2 \qquad \text{for all } x \in V \tag{10.1}$$

if and only if $\ker L \subseteq \ker A$. *In that case* $A = L^{\mathsf{T}} B L$ *with* $B = (L^+)^{\mathsf{T}} A L^+$ *and the best constant is*

$$\kappa_* = \big\| (L^{\mathsf{T}}L)^{\dagger/2} A (L^{\mathsf{T}}L)^{\dagger/2} \big\|_{\mathrm{op}}. \tag{10.2}$$

Proof. Sufficiency: write $x = x_0 + x_1$ with $x_0 \in \ker L$ and $x_1 \perp \ker L$. Then $Ax_0 = 0$ gives $x^{\mathsf{T}}Ax = x_1^{\mathsf{T}}Ax_1$, while $\|Lx\| = \|Lx_1\| \ge \sigma\|x_1\|$ with σ the least positive singular value, so (10.1) holds with $\kappa = \|A\|/\sigma^2$.

Necessity: let $z \in \ker L$ and $x = z + ty$. Then $\|Lx\|^2 = t^2\|Ly\|^2$ while $x^{\mathsf{T}}Ax = z^{\mathsf{T}}Az + 2t\, z^{\mathsf{T}}Ay + t^2 y^{\mathsf{T}}Ay$. Letting $t \to 0$ forces $z^{\mathsf{T}}Az = 0$, and then the linear term must be $O(t^2)$, forcing $z^{\mathsf{T}}Ay = 0$ for all y, i.e. $Az = 0$. Formula (10.2) is the induced norm on the range. □

Remark 10.3. Without the kernel condition the best Frobenius fit is $B_F = (L^+)^{\mathsf{T}} A L^+$ with residual $R_F = A - P_{(\ker L)^\perp} A P_{(\ker L)^\perp}$, and in operator norm the exact dual is

$$\inf_B \|A - L^{\mathsf{T}} B L\|_{\mathrm{op}} = \sup\{\,|\operatorname{tr}(AZ)| : Z = Z^{\mathsf{T}},\ \|Z\|_* \le 1,\ LZL^{\mathsf{T}} = 0\}, \tag{10.3}$$

the constraint $LZL^{\mathsf{T}} = 0$ being exactly annihilation of the subspace $\{L^{\mathsf{T}}BL\}$. For an error $\varepsilon\|x\|^2$ the optimal pair (κ, ε) solves the semidefinite programme $-\kappa L^{\mathsf{T}}L - \varepsilon I \preceq A \preceq \kappa L^{\mathsf{T}}L + \varepsilon I$.

Theorem 10.4 (No sublinear observability)**.** *Fix $h \neq 0$. If q and $q+h$ are squarefree with $q > N/2$ and neither row q nor row $q+h$ lies in Λ, then $x = e_q \pm e_{q+h} \in \ker L$ with $|x^{\mathsf{T}} A_h x| = 1$ and $\|x\|^2 = 2$; hence any inequality of the form $|x^{\mathsf{T}} A_h x| \leq \kappa \|Lx\|^2 + \varepsilon \|x\|^2$ forces $\varepsilon \geq 1/2$. Consequently, to achieve $\varepsilon < 1/2$ the set Λ must be a vertex cover of the shift graph on $(N/2, N]$, and since that graph has maximum degree 2 and, by Mirsky's theorem [9], a positive density of edges, $|\Lambda| = \Omega_h(N)$. No nontrivial observability is possible with $o(N)$ divisor rows.*

Proof. For $q > N/2$ the only multiple of q up to N is q itself, so column q of D is seen only by row q; the same for $q + h$. Hence $Lx = 0$ for the stated x, while $x^{\mathsf{T}} A_h x = \pm 2 \cdot \frac{1}{2} = \pm 1$ and $\|x\|^2 = 2$. The covering statement follows since each uncovered edge produces such an x; a graph of maximum degree 2 with $\Omega(N)$ edges needs $\Omega(N)$ vertices to cover. □

Remark 10.5. Centres, quarters and returns can therefore only work after restricting to a low-dimensional multiscale subspace; global identification is excluded outright.

11. Centres, divisor rows and the Möbius line

11.1. The central dichotomy.

Lemma 11.1 (Central dichotomy)**.** *Let L_j be any linear functional of the values or increments of M on E_j, fixed before the orientation is known. Then*

$$L_j(M_s) = s_j L_j(Q_j) = s_j \kappa_j, \tag{11.1}$$

so either $\kappa_j = 0$ and L_j vanishes on the whole fibre, or $\kappa_j \neq 0$ and $s_j = \operatorname{sgn}(L_j(M_s)/\kappa_j)$ is read off directly. There is no third possibility.

Proof. Immediate from (5.2). □

Remark 11.2. With m_j^{sf} the support centre, i.e. the vertex after ℓ_j of the $2\ell_j$ squarefree steps of E_j, one has $1 \leq Q_j(m_j^{\mathrm{sf}}) \leq \ell_j$ and $Q_j(m_j^{\mathrm{sf}}) \equiv \ell_j \pmod 2$, so the central path value never vanishes and always reads the bit. The support centre is the canonical choice: it is squarefree, so a divisor row evaluated there necessarily contains its own diagonal term, and it avoids spurious plateaux with $\mu(n) = 0$.

Warning 11.3. Augmenting the Gram form with central path readings gives $C_{\mathrm{path}} = -2\,\mathrm{diag}(h_1, \ldots, h_Z)$ and hence $G^+ = V^{\mathsf{T}} A V + C_{\mathrm{path}}^{\mathsf{T}} W C_{\mathrm{path}}$ with $\ker G^+ = \ker V \cap \ker C_{\mathrm{path}} = \{0\}$, since $h_j \neq 0$. The kernel disappears, but any J_h then admits $J_h = C_{\mathrm{path}}^{\mathsf{T}} (C_{\mathrm{path}}^{-\mathsf{T}} J_h C_{\mathrm{path}}^{-1}) C_{\mathrm{path}}$ with zero remainder. This is tautological: the whole difficulty has been moved inside a diagonal bit reader.

11.2. Sampling the divisor equation at the centre.

Definition 11.4. With $m_j = m_j^{\mathrm{sf}}$ and q_k the canonical positive increment of excursion k, set

$$C_{jk}^{\mathrm{div}} = \sum_{\substack{d \mid m_j \\ d \in E_k}} q_k(d). \tag{11.2}$$

Proposition 11.5. *$C_{jk}^{\mathrm{div}} = 0$ for $k > j$, so C^{div} is lower triangular. If moreover $u_j \geq m_j/2$, then $C_{jj}^{\mathrm{div}} = q_j(m_j) \in \{\pm 1\}$.*

Proof. For $k > j$ every element of E_k exceeds m_j, so no divisor of m_j lies in E_k. If $u_j \geq m_j/2$ then every proper divisor of m_j is $\leq m_j/2 \leq u_j$, hence outside $E_j = (u_j, v_j]$; only $d = m_j$ remains and m_j is squarefree. □

Proposition 11.6 (Logarithmic exceptional set)**.** *The hypothesis $u_j \geq m_j/2$ of Prop. 11.5 holds for all but at most $\lfloor \log_2 N \rfloor$ indices j.*

Proof. Since $m_j \le v_j$, the inequality $u_j \ge v_j/2$ implies $u_j \ge m_j/2$. So the hypothesis can fail only when $v_j > 2u_j$. The excursion endpoints satisfy $u_{j+1} = v_j$, so at each failing index the value of u more than doubles. Starting from $u_1 \ge 0$ and ending at $u \le N$, there can be at most $\lfloor \log_2 N \rfloor$ such indices. □

Theorem 11.7. *Let $\mathcal{J}$ be the set of indices satisfying $u_j \ge m_j/2$. In orientation coordinates, treating the row $n = 1$ affinely by $w(1) = 1$,*

$$C^{\mathrm{div}} s_\mu = 0, \qquad \operatorname{rank} C^{\mathrm{div}} \;\ge\; |\mathcal{J}| \;\ge\; Z - \lfloor \log_2 N \rfloor. \tag{11.3}$$

In particular $\dim \ker C^{\mathrm{div}} \le \lfloor \log_2 N \rfloor + 1$ *and s_μ lies in that kernel.*

Proof. Row j of $C^{\mathrm{div}} s_\mu$ equals $\sum_{d \mid m_j} \mu(d)$, which vanishes for $m_j > 1$; the affine treatment of $d = 1$ fixes the remaining normalisation. The rank bound follows from lower triangularity together with Prop. 11.5 on $\mathcal{J}$ and Prop. 11.6. □

Remark 11.8. This is a sparse orientational selector: roughly one divisor equation per excursion, chosen canonically at its centre, isolating the Möbius line up to a kernel of logarithmic dimension. For the full divisor operator these rows are already present, $C^{\mathrm{div}} = RV$ with R a row selector, so they do not reduce $\ker V$; for a truncation V_y they reduce the kernel exactly when $C^{\mathrm{div}}|_{\ker V_y} \ne 0$.

Warning 11.9. Isolating the Möbius line does not bound the shift on it. If the central rows reduce the kernel to $\operatorname{span}\{s_\mu\}$, then estimating J_h on that kernel is exactly estimating the cross-excursion Möbius correlation. Centres help as an arithmetic selector, not yet as an estimator.

12. Intrablock identities

Lemma 12.1. *Let $x_1, \dots, x_L$ have zero sum, $P_t = \sum_{r \le t} x_r$ extended by zero outside $[0, L]$, and $A_h = \sum_t P_t P_{t+h}$. Then*

$$C_h := \sum_t x_t x_{t+h} = 2A_h - A_{h-1} - A_{h+1}. \tag{12.1}$$

Moreover

$$C_0 + 2\sum_{h \ge 1} C_h = 0, \qquad \sum_{t=1}^{L-1} P_t^2 = -\sum_{h \ge 1} h C_h. \tag{12.2}$$

Proof. Expand $x_t = P_t - P_{t-1}$ and collect the four autocorrelations. The first identity in (12.2) is $\sum_{h \in \mathbb{Z}} C_h = (\sum_t x_t)^2 = 0$. For the second, apply (12.1) for all $h \in \mathbb{Z}$ and reindex: $\sum_h |h| C_h = \sum_h A_h(2|h| - |h+1| - |h-1|) = -2A_0$, since the bracket vanishes for $h \ne 0$ and equals -2 at $h = 0$; halve. □

Remark 12.2. Inside an excursion $P_t P_{t+h} = |P_t||P_{t+h}|$, so the intrablock autocorrelation is already determined by the absolute profile. The centre adds nothing there.

Remark 12.3 (The centre does not determine a lag). The two positive excursions of length 8

$$(+,+,+,-,+,-,-,-) \quad \text{and} \quad (+,+,-,+,-,+,-,-) \tag{12.3}$$

have paths $(1,2,3,2,3,2,1,0)$ and $(1,2,1,2,1,2,1,0)$, hence the same central value 2, but $C_1 = 1$ and $C_1 = -3$ respectively. Length, zero sum and central height do not control even the first lag.

Lemma 12.4 (Left–right splitting)**.** *With m the centre and $C_h = C_h^{LL} + C_h^{RR} + C_h^{\times}$, only indices $m - h < t \le m$ can cross, so $|C_h^{\times}| \le h$ and, summing over excursions, $|\sum_j C_{h,j}^{\times}| \le hZ$.*

Proof. A pair $(t, t+h)$ crosses m iff $t \le m < t + h$. □

Remark 12.5. Lemma 12.4 is $o(N)$ precisely when $Z = o(N)$, which is not proved anywhere in this project; see (Q5). The other two parts can be of the size of the excursion length: the centre isolates a layer of width h only.

Lemma 12.6 (Discrete coarea)**.**

$$\mu(n) = \sum_{\ell\in\mathbb{Z}} \Big(\mathbf{1}[M(n-1) \le \ell < M(n)] - \mathbf{1}[M(n) \le \ell < M(n-1)] \Big). \tag{12.4}$$

Consequently $C_N(h)$ is a sum of oriented intersections of level-set boundaries separated by h.

Proof. If $\mu(n) = +1$ the first indicator contributes once and the second is empty; if $\mu(n) = -1$ the roles are exchanged; if $\mu(n) = 0$ both are empty. □

13. Exact local structure at each prime

Proposition 13.1. *Fix a prime p and $t = v_p(h)$. Counting residues modulo p^2, the admissible states $(v_p(n), v_p(n+h)) \in \{0,1\}^2$ and their multiplicities are*

t	*states and multiplicities*	*local sign*
$t = 0$	$(0,0)$*:* $p^2 - 2p$*;* $(1,0)$*,* $(0,1)$*:* $p-1$ *each*	$+,-,-$
$t = 1$	$(0,0)$*:* $p^2 - p$*;* $(1,1)$*:* $p-2$	$+$
$t \ge 2$	$(0,0)$*:* $p^2 - p$*;* $(1,1)$*:* $p-1$	$+$

The local squarefree density of the pair is $1 - 2/p^2$ if $p^2 \nmid h$ and $1 - 1/p^2$ if $p^2 \mid h$, which is the Mirsky structure [9]*. For any finite set of primes P,*

$$\prod_{p\in P} (-1)^{v_p(n)+v_p(n+h)} \mathbf{1}[v_p(n), v_p(n+h) \le 1] \tag{13.1}$$

is an exact identity, not an independence hypothesis.

Proof. For $p \nmid h$ the classes $n \equiv 0$ and $n \equiv -h \pmod p$ are distinct, giving $(p-2)p$ residues with $(0,0)$ and $p-1$ each with $(1,0)$, $(0,1)$; the two excluded residues are $p^2 \mid n$ and $p^2 \mid n+h$. For $p \| h$, $p \mid n$ forces $p \mid n+h$, and among the p residues with $p \mid n$ one must delete $n \equiv 0$ and $n \equiv -h$, distinct modulo p^2, leaving $p-2$. For $p^2 \mid h$ the two deleted residues coincide, leaving $p-1$. Signs are $(-1)^{v_p(n)+v_p(n+h)}$; the densities follow by counting the excluded residues. CRT makes the joint count over distinct primes multiplicative. □

Remark 13.2. The shift does not break the tensor structure only at $p \mid h$: every prime participates, with a universal off-diagonal factor for $p \nmid h$ and a diagonal factor for $p \mid h$. What fails on globalising is the cut $n \le N$, the infinite product and uniform control of the tail of squares, not the local product.

14. Multiplicative weights collapse

Theorem 14.1. *Let $m_d(x) = \sum_{n\le x,\ (n,d)=1} \mu(n)/n$. For $p \nmid d$,*

$$m_d(x) = m_{dp}(x) - \frac{1}{p}\, m_{dp}(x/p), \qquad m_{dp}(x) = \sum_{k\ge 0} p^{-k} m_d(x/p^k), \tag{14.1}$$

the second sum being finite. Consequently

$$m_d(x) = \sum_{\substack{e\ge 1 \\ \mathrm{rad}(e)\mid d}} \frac{1}{e}\, m_1(x/e), \tag{14.2}$$

and with $G_{ab} = 2N/[a,b]$,

$$\frac{\langle \mu, G\mu\rangle}{2N} = \sum_{d \text{ squarefree}} \varphi(d) \left(\sum_{\mathrm{rad}(u)=d} \frac{m_1(N/u)}{u} \right)^2. \tag{14.3}$$

In Mellin variables the local operator has symbol $1 - p^{-1-s}$ and the full product has symbol $1/\zeta(1+s)$.

Proof. Split $\{(n,d)=1\}$ according to $p \mid n$; for $n = pn'$ squarefree with $(n', dp) = 1$ one has $\mu(n) = -\mu(n')$, giving the first identity. The Neumann series inverts it and terminates because $m_d(x/p^k) = 0$ for $p^k > x$. Composing over $p \mid d$ gives (14.2). For (14.3), write $\gcd(a,b) = \sum_{d|a,\ d|b} \varphi(d)$ to obtain $\langle \mu, G\mu\rangle/2N = \sum_d \varphi(d) d^{-2} \mu(d)^2 m_d(N/d)^2$; substituting (14.2) with $u = de$, the conditions $\mathrm{rad}(e) \mid d$, $d \mid u$ and d squarefree force $d = \mathrm{rad}(u)$, so each u lies in exactly one block and $1/e = d/u$. The Mellin symbol follows from $\sigma_p f(x) = f(x/p) \leftrightarrow p^{-s}$. □

Corollary 14.2. *Expanding G along its diagonals and using* $\gcd(a, a+h) = \gcd(a,h)$*,*

$$2\sum_{h\ge 1}\sum_{a} \mu(a)\mu(a+h)\frac{\gcd(a,h)}{a(a+h)} = \sum_d \varphi(d)\Big(\sum_{\mathrm{rad}(u)=d} \frac{m_1(N/u)}{u}\Big)^2 - \sum_{a\le N} \frac{\mu(a)^2}{a^2}, \tag{14.4}$$

so the gcd-weighted Chowla aggregate is determined by m_1, i.e. by $1/\zeta(1+s)$. The left side is in particular bounded below by $-\zeta(2)/\zeta(4)$ since the right side is a sum of squares.

Proof. Immediate from Theorem 14.1 and $G_{a,a+h} = 2N \gcd(a,h)/(a(a+h))$. □

Warning 14.3 (Dichotomy for diagonal weights). Any diagonal weight $\omega(a,h)$ with a multiplicative symbol collapses as in Theorem 14.1, because the divisor lattice closes; any weight without one decomposes into characters and turns $\sum_a \mu(a)\mu(a+h)\omega(a,h)$ into a twisted correlation, strictly harder than $C_N(h)$. There is no middle case. The proof of collapse uses $\mu(da') = \mu(d)\mu(a')$ and uniqueness of the radical, i.e. commutativity of the Euler product; a non-commutative deformation such as the cocycle $\Phi(\pi)$ of (S7) is the only place where the argument does not apply. That escape is a fixed jet in ϑ, and (S8) closes fixed jets; only the scaling window $\vartheta \asymp \vartheta_c(N)$ survives.

15. Scale-resolved statistic and sine-kernel content

Definition 15.1. $\mathcal{E}_k = \langle s, \widehat{P}_k s\rangle$, $\mathcal{R}_k = \mathcal{E}_k/d_k$.

Proposition 15.2. *Under ν, $\mathbb{E}_\nu[\mathcal{R}_k] = 1$ exactly and $\mathrm{Var}_\nu[\mathcal{R}_k] = 2d_k^{-2}(d_k - \sum_j \widehat{P}^2_{k,jj})$, with $\sum_k \mathcal{E}_k = Z = \sum_k d_k$.*

Proof. Prop. 5.4 with $A = \widehat{P}_k$, and $\sum_k \widehat{P}_k = I_Z$. □

Lemma 15.3. *The orthogonal projection onto a contiguous block of d characters of a cyclic group of order Z has Dirichlet kernel, whose scaling limit with d/Z fixed is* $\sin(\pi x)/(\pi x)$ [8]*. Equivalently*

$$1 - R_2(x) = \Big(\frac{\sin \pi x}{\pi x}\Big)^2 = \int_{-1}^{1} (1-|\tau|) e^{2\pi i \tau x}\, d\tau. \tag{15.1}$$

Warning 15.4. By Prop. 5.4 the reflection null has covariance I, which restricted to a band of characters is a sine kernel. The sine kernel is therefore the *null* of $\mathcal{R}_k$, not a consequence of it, and deriving it from flatness carries no evidential weight about the zeros of ζ.

Remark 15.5. Hypothesis (G2), i.e. (S4), makes the eigenbasis of G a character basis of the multiplicative structure, so the transported band gives a sine kernel in $\log n$, the variable in which $n^{i\gamma} = e^{i\gamma \log n}$ oscillates. This does not affect Warning 15.4.

Remark 15.6. A statistic carries sine-kernel content non-tautologically if and only if its null value is a number different from its sine-kernel value; the measured position between the two is the coefficient. The three quantities of Version 2 satisfying this are retained below.

Definition 15.7. For the superposition of p independent identically distributed spectra each of density $1/p$,

$$R_2^{(p)}(x) = 1 - \frac{1}{p}\Big(\frac{\sin(\pi x/p)}{\pi x/p}\Big)^2, \qquad Y_{jk}^{(p)} = \frac{1}{p}\Big(\frac{\sin(\pi(j-k)/p)}{\pi(j-k)/p}\Big)^2. \tag{15.2}$$

Proposition 15.8. *If $\tilde{r}$ is strictly monotone in p with $\tilde{r}(\infty) = 2\ln 2 - 1$ and $\tilde{r}(1)$ the unitary value, then a measured $\tilde{r} > 2\ln 2 - 1$ forces $p < \infty$ and hence a strictly positive sine-kernel fraction $1/p$.*

Proof. Strict monotonicity and the endpoint values. □

Lemma 15.9. $m_n = \displaystyle\int_0^1 (1-\tau)\tau^n \, d\tau = \frac{1}{(n+1)(n+2)}$, *and* $(1-\tau)_+$, *having compact support, is determined by* $\{m_n\}_{n\geq 0}$.

Proof. Elementary integration and determinacy of the Hausdorff moment problem. □

Lemma 15.10. *Under the Wigner surmises,* $\mathrm{Var}_{\mathrm{Poisson}}(s) = 1$, $\mathrm{Var}_{\mathrm{GOE}}(s) = 4/\pi - 1$, $\mathrm{Var}_{\mathrm{GUE}}(s) = 3\pi/8 - 1$.

Proof. For $p(s) = \frac{\pi}{2} s e^{-\pi s^2/4}$ and $p(s) = \frac{32}{\pi^2} s^2 e^{-4s^2/\pi}$, both normalised to $\langle s \rangle = 1$, ratios of Gaussian moments give $\langle s^2 \rangle = 4/\pi$ and $3\pi/8$; the Poisson case is the exponential law. □

Remark 15.11. Recorded in separate work, not computed here, and used only to indicate the scale of the effects: a spacing ratio $\tilde{r} \approx 0.42$ for the spectrum of the smooth arithmetic term; relative freeness defects of about 12% at order two and 32% at order three; an energy deficit ratio of about 0.137; a persistent bias of about 2.5%; and the systematic observation that $d = 1$ is the weakest anomaly. The reference values $2\ln 2 - 1$, ≈ 0.5307 and ≈ 0.5996 are recalled from [1] and not verified here.

Remark 15.12. The triangle moments give $m_3/m_2 = 0.6$, decreasing; Chebyshev-basis variance of linear statistics grows linearly [5], giving 1.5; the recorded defects give $32/12 \approx 2.67$. The observed growth is faster than linear where the triangle requires decay. The ratio is invariant under a global rescaling of the normalisation of relative defect, so the discrimination survives that convention.

Remark 15.13. $\mathcal{R}_N$ and $\widehat{G}$ are real, hence $\beta = 1$, whose form factor $2\tau - \tau\log(1+2\tau)$ on $\tau \leq 1$ [8] is not a band indicator; the conjectural class for the zeros is $\beta = 2$ [10, 11].

Conjecture 15.14. *Let $U(1)^Z$ act by $s_j \mapsto e^{i\vartheta_j} s_j$, extending the reflection action, with ϑ governed by the cocycle of (S7), and let $\widehat{G}^{(\vartheta)}$ be the transported Hermitian form. Its local statistics interpolate between the orthogonal and unitary classes, with crossover at $\vartheta_c \asymp 1/Z$* [13].

16. Strategy and the target theorem

Remark 16.1 (Ordered strategy).
(i) Difference, not commutator: build $(U_h - I)D$, split off the columns $d \mid h$ exactly by Lemma 8.2, and send the rest to adaptive Haar.
(ii) Approximate observability: work in $V/\ker D_\Lambda$, measure the optimal residual by pseudoinverse and the dual (10.3); do not attempt global identification, which Thm. 10.4 excludes.
(iii) Irregular tree: choose cuts by changes of divisor density, levels of M, centres and residue boundaries; exploit (9.3).
(iv) Local p-adic structure: couple Prop. 13.1 with the factors L_p of (S4), treating $p \nmid h$ as a universal factor and $p \mid h$ as diagonal defects.
(v) Excursions last: use them only for the reflection-odd component and as one family of stopping blocks, subject to (Q1).

Objective 16.2 (Adaptive boundary observability)**.** For fixed h, construct trees $\mathcal{T}_N$, multiscale subspaces $W_N \subset H_{\mathrm{sf}}$ and rows Λ_N adapted to CRT boundaries, not to the signs of μ, such that

$$\left\| P_{W_N} (A_h - L^{\mathsf{T}} B_{h,N} L) P_{W_N} \right\|_{\mathrm{op}} = o(1), \tag{16.1}$$

with $B_{h,N}$ the variational minimiser, and simultaneously

$$\left\| P_{W_N^{\perp}} \mu \right\|^2 = o(N). \tag{16.2}$$

Estimate (16.1) is to be proved by reducing all volumetric contributions to zero through Theorem 9.3 and controlling only the boundary cocycles; (16.2) must come from prefix, level and squarefree identities, not from probabilistic regularity. Together they would give $C_N(h) = o(N)$.

Objective 16.3 (Variant with central divisor rows)**.** With $F_{N,y} = \begin{bmatrix} V_y \\ C_N^{\mathrm{div}} \end{bmatrix}$, truncated divisor equations together with centre rows fixed independently of the orientations, prove

$$J_h^{\mathrm{off}} = F_{N,y}^{\mathsf{T}} B_{h,N} F_{N,y} + R_{h,N} \tag{16.3}$$

with $B_{h,N}(d,e) = 0$ unless $(d,e) \mid h$, prime-by-prime CRT structure, $\|B_{h,N}\| = N^{o(1)}$, and $\sup_{s\in\{\pm1\}^z} |s^{\mathsf{T}} R_{h,N} s| = o(N)$, no definition using s_μ or $C_N(h)$. Since $F_{N,y} s_\mu = 0$ by Theorem 11.7, this gives $s_\mu^{\mathsf{T}} J_h^{\mathrm{off}} s_\mu = o(N)$; combined with the independent intrablock objective $\sum_j \sum_{n,n+h\in E_j} \mu(n)\mu(n+h) = o(N)$ it gives $C_N(h) = o(N)$.

Warning 16.4. The ordinary correlation $C_N(h) = o(N)$ at fixed $h \neq 0$ is open; the logarithmically averaged version is known [15], building on [7]. Any inequality that yields $o(N)$ directly must therefore contain a genuinely new estimate, not a conjugation or a rewriting. Remark 8.1 and Warning 11.3 record two ways of failing this test.

17. Negative results and failed routes

Remark 17.1. The scalar quadratic energy at a single level carries no local information; the defect is the collapse to one number, not the weight.

Remark 17.2. Statistics built from M alone weight zeros by decaying factors in the ordinate and therefore mix heights; no operation on $\log N$ repairs this, since $\log N$ is conjugate to differences of ordinates, not to ordinates.

Remark 17.3. The excursion decomposition passes to continued fractions and hence to a Jacobi matrix [3], which is real symmetric and lands in the orthogonal class, as in Rem. 15.13.

Remark 17.4. A cell-complex description of M exists [2] but does not reach two-point cancellation: the complex is multiplicative while shifted correlations are additive, and no map relates the complex at n to that at $n + h$.

Remark 17.5. Any operator whose spectrum is the set of ordinates must reproduce $N(T) \sim (T/2\pi)\log(T/2\pi)$; the bounds (S1) place $\lambda_{\max}(D_N)$ at Wigner scale, whose integrated density is semicircular and fails by shape.

Remark 17.6. A dyadic tree of midpoints of $|M|$ satisfies Chapman–Kolmogorov, hence is Markov in the scale variable; a constraint at a coarse level therefore gives no information at finer levels beyond the endpoint values, and the degrees of freedom double at each refinement. The product structure that makes the tree null computable is exactly the property that forbids constraint propagation. This closes the route of propagating a short-interval input down the tree. The same applies to the Euler factorisation in the multiplicative direction: both natural indexings factorise, so neither transports a constraint across scales.

Remark 17.7. Any hierarchy of bounds on the Möbius walk must respect $M(x) = \Omega_\pm(\sqrt{x})$ with explicit constants [12]; no ε in such a hierarchy is removable.

Remark 17.8. Splitting an interval into two halves keeps an $O(1)$ CRT error in each half; it does not reduce it. A left–right difference can cancel the main term when the lengths agree, but leaves a difference of boundary errors of size $O(1)$ with no sign and no uniform cancellation. For halves defined in squarefree time the set is no longer an arithmetic interval and the mask must be inserted, which makes the CRT factorisation less clean, not more.

18. Open questions

(Q1) Does $\widehat{G}$ inherit the tensor factorisation (2.1) from G? The excursion blocks are defined by the zero set of M and respect no multiplicative structure, so clean inheritance is not expected; partial inheritance restricted to the local factors L_p with small p would be the interesting outcome. Everything in Sections 15–16 depends on it.

(Q2) Prove a norm estimate for the boundary CRT cocycle operator of Theorem 9.3, restricted to adaptive Haar subspaces and after extracting the exact local p-adic modes of Prop. 13.1, respecting Warning 9.5. This is the first theorem of Objective 16.2 and the identified bottleneck.

(Q3) Sharpen Prop. 11.6: is the exceptional set actually bounded, or can $\log_2 N$ be attained?

(Q4) Test the prediction that $1/p(1)$ is the minimum of the divisor-resolved sine-kernel fraction.

(Q5) Determine the growth of $Z(N)$. Lemma 12.4 and every statement with an $O(hZ)$ error depend on $Z = o(N)$, which is not proved here.

(Q6) Close the non-attained-supremum case of Rem. 2.2.

(Q7) Decide whether the non-commutative deformation of Warning 14.3 escapes (S8) in the scaling window $\vartheta \asymp \vartheta_c(N)$.

Notebook History

Version 1: Three-constraint model and exact bridge second moment; reflection fibre and the canonical parameter-free null; divisor-defect Gram form pushed to orientation bits; blindness theorem; scale-resolved statistic with exact mean and variance; negative results.

Version 2: Abstract added; introduction restructured into what is proved, what is input, and the methodological point; standing results listed with status; sine-kernel section rewritten around the non-tautology criterion, with three recoveries of the sine-kernel coefficient from measured quantities; extremal-abscissa gap narrowed.

Version 3: Operator layer added. Three spaces and their information ledger; divisor–shift boundary lemma with interval mass bound 1; exact CRT pullback; algebraic Haar on arbitrary trees with exact vanishing of the volumetric CRT term and parent–children cocycle additivity; exact observability criterion with optimal constant, operator-norm dual and semidefinite formulation; no-go for sublinear divisor rows; extension of the blindness theorem to half-constant kernels, with the shift shown to escape it; central dichotomy lemma; triangular central divisor matrix with the logarithmic exceptional set, isolating the Möbius line; intrablock second-difference and aggregate identities, discrete coarea, and a counterexample showing the centre does not determine a lag; exact local tables at each prime; collapse theorem for multiplicative diagonal weights and the resulting dichotomy; target objectives stated.

References

[1] Y. Y. Atas, E. Bogomolny, O. Giraud and G. Roux, *Distribution of the ratio of consecutive level spacings in random matrix ensembles*, Physical Review Letters **110** (2013), 084101.

[2] A. Björner, *A cell complex in number theory*, arXiv:1101.5704.

[3] P. Flajolet, *Combinatorial aspects of continued fractions*, Discrete Mathematics **32** (1980), 125–161.

[4] S. M. Gonek, *On negative moments of the Riemann zeta-function*, Mathematika **36** (1989), 71–88.

[5] K. Johansson, *On fluctuations of eigenvalues of random Hermitian matrices*, Duke Mathematical Journal **91** (1998), 151–204.

[6] J. C. Lagarias, *An elementary problem equivalent to the Riemann hypothesis*, American Mathematical Monthly **109** (2002), 534–543.

[7] K. Matomäki and M. Radziwiłł, *Multiplicative functions in short intervals*, Annals of Mathematics **183** (2016), 1015–1056.

[8] M. L. Mehta, *Random Matrices*, third edition, Elsevier, Amsterdam, 2004.

[9] L. Mirsky, *Arithmetical pattern problems relating to divisibility by rth powers*, Proceedings of the London Mathematical Society **50** (1949), 497–508.

[10] H. L. Montgomery, *The pair correlation of zeros of the zeta function*, in: Analytic Number Theory, Proceedings of Symposia in Pure Mathematics **24**, American Mathematical Society, 1973, 181–193.
[11] A. M. Odlyzko, *On the distribution of spacings between zeros of the zeta function*, Mathematics of Computation **48** (1987), 273–308.
[12] A. M. Odlyzko and H. J. J. te Riele, *Disproof of the Mertens conjecture*, Journal für die reine und angewandte Mathematik **357** (1985), 138–160.
[13] A. Pandey and M. L. Mehta, *Gaussian ensembles of random Hermitian matrices intermediate between orthogonal and unitary ones*, Communications in Mathematical Physics **87** (1983), 449–468.
[14] G. Robin, *Grandes valeurs de la fonction somme des diviseurs et hypothèse de Riemann*, Journal de Mathématiques Pures et Appliquées **63** (1984), 187–213.
[15] T. Tao, *The logarithmically averaged Chowla and Elliott conjectures for two-point correlations*, Forum of Mathematics, Pi **4** (2016), e8.
[16] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, second edition, revised by D. R. Heath-Brown, Oxford University Press, 1986.