CATALAN MERTENS, VERSION 5: THE EXCEPTIONAL CATALAN CLASS, CRITERION, AND MECHANISM

Contents

1. Introduction 2
2. The selection wall 3
3. Exact Catalan decomposition of the compressed Mertens walk 4
3.1. Compressed time 4
3.2. Return skeleton and factorization 4
4. Open Catalan sectors: meanders and pointwise control 5
5. Catalan free energy and a sufficient selection principle 5
6. Brownian excursion and meander as continuum Catalan mechanics 6
6.1. A variational action 6
7. Transfer matrices and the quantum Catalan picture 7
7.1. Height confinement as a spectral problem 7
7.2. Fredkin spin chains 7
8. Statistical mechanics of Dyck paths: pressure, area, and q-Catalan weights 8
9. Non-equilibrium transport and the Burgers route 8
10. The deterministic arithmetic selector 9
10.1. Björner's filtered complex 9
10.2. Subset-sum energy representation 9
11. Three possible selector mechanisms 10
11.1. Hyperplane-chamber entropy 10
11.2. Shadow inequalities and parity restoration 10
11.3. Persistent topology 11
12. Structural viability of the physical routes 11
13. Potential obstructions 11
13.1. The selector may remain unconstrained at the needed scale 11
13.2. Static topology may not control filtered dynamics 11
13.3. Parity barrier 11
13.4. Canonical collapse to a classical Dirichlet object 12
14. A hierarchy of possible sufficient principles 12
15. The prime-delay selector and its exact interference algebra 12
15.1. Simultaneous prime-shell identities 12
15.2. Threshold kernel and transfer recursion 13
15.3. A maximal-parity Walsh coefficient 14
15.4. Uniform prefix energy: a corrected sufficient principle 15
15.5. The exceptional Catalan class is exactly the RH obstruction 15
15.6. No-go audit: what the exact identities do not prove 17
16. An aggregate route that does not predict M pointwise 18
16.1. A hierarchy of moments equivalent to RH 18
16.2. Exact excursion decomposition of the moment budget 19
16.3. The arithmetic correlation target 20
16.4. Finite evidence and its exact limitation 20
16.5. A staged proof program 20

17. Research program 22
17.1. Exact finite combinatorics 22
17.2. Finite selector chambers 22
17.3. Dynamic shadows 22
17.4. Transfer operators 22
17.5. Hydrodynamic coarse graining 22
18. Relation to existing literature 22
19. Conclusions and open questions 23
Appendix A. Reproducible computations and archived outputs 23
Appendix B. Python source: moment hierarchy audit 24
Appendix C. Python source: exact selector identities 26
Appendix D. Python source: excursion heights and Catalan tails 28
Appendix E. Python source: finite τ-dephasing illustration 31
Appendix F. Python source: publication figures 32
Appendix G. Python source: Version 5 no-go audit 33
References 35

Remark 0.1. This document is an exploratory mathematical notebook. Its purpose is to isolate an exact Catalan structure inside the Mertens walk, to develop physical and probabilistic tools that act directly on that Catalan state space, and to identify a precise obstruction between ambient Catalan rarity and the deterministic arithmetic trajectory. Rigorous statements are labelled as such; proposed mechanisms are labelled as heuristics, conjectures, or questions.

1. Introduction

Let

$$M(x) = \sum_{n \le x} \mu(n) \tag{1.1}$$

be the Mertens function. The classical square-root target is

$$M(x) = O_\varepsilon\Big(x^{1/2+\varepsilon}\Big) \qquad (\varepsilon > 0), \tag{1.2}$$

which is equivalent to the Riemann hypothesis; see, for example, the standard analytic literature and the discussion in Björner's topological formulation [1]. The present notebook does not pursue an explicit-formula, zero-density, or Dirichlet-series proof strategy. The target (1.2) is retained as the benchmark against which a different mechanism is measured.

The starting observation is elementary but exact. Remove the zero values of the Möbius function. If

$$1 = q_1 < q_2 < q_3 < \cdots \tag{1.3}$$

are the squarefree positive integers and

$$X_m = \sum_{j \le m} \mu(q_j), \tag{1.4}$$

then every increment of X_m is exactly $+1$ or -1, and

$$X_m = M(q_m). \tag{1.5}$$

Consequently, each interval between consecutive zeros of X_m is a primitive lattice excursion. If such an excursion has length $2n$, then, after fixing its sign, the number of possible excursion shapes is exactly the Catalan number

$$C_{n-1} = \frac{1}{n}\binom{2n-2}{n-1}. \tag{1.6}$$

Thus the real Mertens excursion is one state inside a finite Catalan state space.

The usefulness of this observation depends entirely on a second question. Catalan combinatorics describes the ambient set of admissible excursion shapes, but the arithmetic trajectory is a deterministic selection from that set. A rare Catalan shape remains possible unless the arithmetic selector itself satisfies an additional restriction. The central problem of this notebook is therefore to identify a deterministic mechanism that controls the location of the Mertens trajectory inside Catalan state space.

The physical methods considered here are useful because they quantify geometry inside that state space. Brownian excursion theory identifies the diffusive $\sqrt{L}$ scale; transfer matrices convert height restrictions into spectral gaps; path-integral arguments assign an action of order H^2/L to an excursion of height H and duration L; area-weighted Dyck paths provide genuine partition functions and free energies; Fredkin spin chains realize uniform superpositions of Dyck paths as quantum ground states; and TASEP provides a route from excursion statistics to non-equilibrium hydrodynamics and the Burgers equation [2, 3, 5, 6, 7, 9, 10].

The arithmetic side also has a nontrivial exact structure. Björner associated to each cutoff x a simplicial complex of squarefree integers and proved that the Mertens function is its reduced Euler characteristic [1]. In that representation, the compressed walk is the one-dimensional trace of a filtered, shifted simplicial complex. This supplies a deterministic selector which is absent from a generic random walk.

The project therefore separates into three layers:

$$\text{Mertens filtration} \longrightarrow \text{Catalan excursion state} \longrightarrow \text{physical/entropic cost of its geometry.} \tag{1.7}$$

The difficult arrow is the first-to-second selection law: why should the filtration select a non-exceptional region of the Catalan space? The rest of the notebook develops several concrete formulations of this question.

2. The selection wall

This section states the main obstruction at the outset.

Definition 2.1. Let $\mathcal{D}_n^{\mathrm{prim}}$ denote the set of positive primitive excursions of length $2n$: paths $D = (D_j)_{0\le j\le 2n}$ satisfying

$$D_0 = D_{2n} = 0, \qquad D_j > 0 \quad (1 \le j \le 2n-1), \qquad D_{j+1} - D_j \in \{-1, +1\}. \tag{2.1}$$

Proposition 2.2. *For every $n \ge 1$,*

$$\left|\mathcal{D}_n^{\mathrm{prim}}\right| = C_{n-1}. \tag{2.2}$$

Proof. Delete the initial up-step and the final down-step and shift the remaining path down by one unit. This is a bijection with ordinary Dyck paths of length $2n-2$, counted by C_{n-1}. □

If a Mertens excursion has length $2n$, Proposition 2.2 places its shape in a set of size C_{n-1}. Under the uniform Catalan measure, high excursions occupy a small fraction of this set; the maximum of a uniform Dyck path is naturally of order $\sqrt{n}$, and exponential moments and moderate/large deviations are well studied [5, 6].

Warning 2.3. Ambient rarity is not an arithmetic restriction. A deterministic selector may choose a member of an exponentially small subset. Therefore a statement of the form

$$\frac{\#\{D \in \mathcal{D}_n^{\mathrm{prim}} : \max D \geq H\}}{C_{n-1}} \ll \exp\Bigl(-cH^2/n\Bigr) \tag{2.3}$$

becomes relevant to M only after one proves a property of the arithmetic selection rule.

This warning is the main logical checkpoint of the project. The physical theory can compute or estimate the cost of a region of Catalan space. The arithmetical task is to show that the real trajectory cannot systematically occupy regions whose cost is too large.

3. Exact Catalan decomposition of the compressed Mertens walk

3.1. **Compressed time.** Let

$$Q(x) = \#\{n \leq x : \mu(n) \neq 0\} \tag{3.1}$$

be the squarefree counting function. The standard asymptotic is

$$Q(x) = \frac{6}{\pi^2}x + O(\sqrt{x}), \tag{3.2}$$

so compressed time and ordinary integer time are linearly comparable.

Proposition 3.1. *For every $m \geq 1$,*

$$X_m = M(q_m), \tag{3.3}$$

and for every real $x \geq 1$,

$$M(x) = X_{Q(x)}. \tag{3.4}$$

Proof. The terms omitted from (1.4) are exactly those for which $\mu(n) = 0$. □

3.2. **Return skeleton and factorization.** Let

$$0 = z_0 < z_1 < z_2 < \cdots \tag{3.5}$$

be the successive return times of the compressed walk to zero, whenever they exist. Put

$$L_r = z_r - z_{r-1} = 2n_r. \tag{3.6}$$

Parity forces each complete excursion length to be even.

Proposition 3.2. *Fix a finite return skeleton $0 = z_0 < z_1 < \cdots < z_R$ and fix the sign of each excursion. The number of nearest-neighbour paths realizing exactly these complete excursions is*

$$\mathcal{N}(z_1, \ldots, z_R) = \prod_{r=1}^{R} C_{n_r-1}. \tag{3.7}$$

Proof. The excursion interiors are independent combinatorial choices once their durations and signs are specified. Proposition 2.2 applies to each factor. □

Equation (3.7) is an exact entropy associated with the zero-return skeleton:

$$S_{\mathrm{Cat}}(z_R) = \log \mathcal{N}(z_1, \ldots, z_R) = \sum_{r=1}^{R} \log C_{n_r-1}. \tag{3.8}$$

Using Stirling's formula,

$$\log C_n = n \log 4 - \frac{3}{2}\log n + O(1). \tag{3.9}$$

Thus each completed excursion carries an entropy deficit of order $\frac{3}{2}\log n$ relative to unrestricted ± 1 words of comparable length.

4. Open Catalan sectors: meanders and pointwise control

Complete excursions alone do not control a point lying inside a very long unfinished excursion. The natural pointwise object is therefore a Catalan prefix, or Bernoulli meander. This is also the correct continuum bridge to Brownian meander [3, 4].

Definition 4.1. For $t \geq 1$ and $h > 0$ with $t \equiv h \pmod 2$, let $\mathcal{M}_{t,h}$ be the set of paths $S_0 = 0$, $S_t = h$, with increments ± 1 and

$$S_j > 0 \qquad (1 \leq j \leq t). \tag{4.1}$$

Let $\mathcal{M}_t = \bigcup_h \mathcal{M}_{t,h}$.

Proposition 4.2. *The ballot theorem gives*

$$|\mathcal{M}_{t,h}| = \frac{h}{t}\binom{t}{(t+h)/2}, \tag{4.2}$$

and

$$|\mathcal{M}_t| = \binom{t-1}{\lfloor (t-1)/2 \rfloor}. \tag{4.3}$$

Remark 4.3. The meander is the open-boundary sector of the same Catalan geometry. Closing a positive meander by a first return produces a primitive Dyck excursion, while cutting a primitive Dyck excursion before its terminal return produces a positive meander. For pointwise estimates on M, the meander formulation avoids any dependence on a future return time.

For a compressed time m let

$$a(m) = \max\{z_r : z_r \leq m\}, \qquad t(m) = m - a(m), \qquad h(m) = |X_m|. \tag{4.4}$$

Whenever $h(m) > 0$, the segment from $a(m)$ to m is an element of $\mathcal{M}_{t(m),h(m)}$ after reflection if necessary.

Proposition 4.4. *There exist absolute constants $c, C > 0$ such that, under the uniform measure on $\mathcal{M}_t$,*

$$\mathbb{P}_t\{S_t \geq h\} \leq C\left(1 + \frac{h}{\sqrt{t}}\right)\exp\left(-c\frac{h^2}{t}\right) \tag{4.5}$$

for admissible t, h.

Proof. Equation (4.2) reduces the endpoint law to binomial coefficients. Standard Chernoff bounds for the symmetric binomial distribution give Gaussian decay in h^2/t, while (4.3) is of order $2^t/\sqrt{t}$. Summing the endpoint masses from h upward gives (4.5) after adjustment of absolute constants. □

This proposition supplies a rigorous ambient energy scale at the current time, not only at completed returns.

5. Catalan free energy and a sufficient selection principle

Definition 5.1. For the actual compressed Mertens path at a time m with $h(m) > 0$, define its open Catalan selection energy by

$$\mathcal{I}_{\mathrm{Cat}}(m) = -\log \mathbb{P}_{t(m)}\{S_{t(m)} \geq h(m)\}, \tag{5.1}$$

where $\mathbb{P}_t$ is the uniform measure on $\mathcal{M}_t$.

This is a deterministic number attached to the actual path: probability is used only to measure the volume of the ambient Catalan/meander sector containing an endpoint at least as extreme as the observed one.

Proposition 5.2. *There are constants $c_1, c_2, c_3 > 0$ such that*

$$\mathcal{I}_{\mathrm{Cat}}(m) \geq c_1 \frac{h(m)^2}{t(m)} - c_2 \log\left(2 + \frac{h(m)}{\sqrt{t(m)}}\right) - c_3. \tag{5.2}$$

Proof. Take minus logarithms in Proposition 4.4. □

Theorem 5.3. *Assume that for some constant $A > 0$ and all sufficiently large compressed times m,*

$$\mathcal{I}_{\mathrm{Cat}}(m) \leq A \log m. \tag{5.3}$$

Then

$$|X_m| = O\Big(\sqrt{m \log m}\Big), \tag{5.4}$$

and consequently

$$M(x) = O\Big(\sqrt{x \log x}\Big). \tag{5.5}$$

In particular, (1.2) *follows.*

Proof. By Proposition 5.2 and (5.3), the quantity $h(m)^2/t(m)$ is $O(\log m)$, up to a lower-order logarithmic correction. Since $t(m) \leq m$, this gives (5.4). Equation (3.4) and the linear comparability (3.2) then give (5.5). □

Warning 5.4. The hypothesis (5.3) is precisely the unresolved selector statement. Theorem 5.3 is useful because it isolates the target in a finite combinatorial language: the arithmetic path must be prevented from entering meander sectors whose relative Catalan volume is smaller than a power of the current scale.

6. Brownian excursion and meander as continuum Catalan mechanics

Kaigh proved an invariance principle for a recurrent random walk conditioned on a late first return to zero: after diffusive rescaling the path converges to Brownian excursion [2]. Durrett, Iglehart, and Miller established weak convergence results for Brownian meander and excursion in related conditional settings [3]. These results provide a continuum mechanics for the Catalan state space rather than an independent model imposed on the arithmetic data.

For an excursion of discrete duration L, the diffusive scaling is

$$y_L(s) = L^{-1/2} D_{\lfloor Ls \rfloor}, \qquad 0 \leq s \leq 1. \tag{6.1}$$

The limiting excursion has order-one height, so the natural discrete height scale is

$$H \asymp \sqrt{L}. \tag{6.2}$$

The exponential moments of the normalized Dyck maximum are uniformly controlled and converge to the corresponding Brownian-excursion quantity [5]. Kousha further developed large and moderate deviation principles for the Dyck maximum [6].

6.1. **A variational action.**

Heuristic 6.1. This is a physical large-deviation heuristic. For a continuum path $y : [0, L] \to \mathbb{R}_+$ with $y(0) = y(L) = 0$, assign the Euclidean action

$$\mathcal{A}[y] = \frac{\kappa}{2} \int_0^L \dot{y}(t)^2 \, dt. \tag{6.3}$$

Among paths that reach height H, the piecewise-linear triangular profile minimizes (6.3). Its action is

$$\mathcal{A}_{\min}(H, L) = 2\kappa \frac{H^2}{L}. \tag{6.4}$$

Thus the physical rate variable is H^2/L, matching the Gaussian scale already present in the exact ballot estimate (4.5).

The importance of (6.4) is conceptual: the physical energy scale is not being guessed from Mertens data. It is the continuum expression of a rate already visible in Catalan/ballot combinatorics.

7. Transfer matrices and the quantum Catalan picture

7.1. **Height confinement as a spectral problem.** Let A_h be the adjacency matrix of the path graph on heights $0, 1, \ldots, h$. The number of Dyck paths of length $2n$ confined to this height interval is

$$Z_n(h) = (A_h^{2n})_{00}. \tag{7.1}$$

The eigenvalues are

$$\lambda_k(h) = 2\cos\left(\frac{k\pi}{h+2}\right), \qquad 1 \le k \le h+1. \tag{7.2}$$

Consequently,

$$Z_n(h) = \frac{2}{h+2}\sum_{k=1}^{h+1} \sin^2\left(\frac{k\pi}{h+2}\right)\lambda_k(h)^{2n}. \tag{7.3}$$

This spectral representation is standard in the analysis of Dyck-path maxima and underlies related deviation results [6].

For large h,

$$\lambda_1(h) = 2 - \frac{\pi^2}{(h+2)^2} + O(h^{-4}), \tag{7.4}$$

so confinement below an atypically low ceiling has a free-energy scale of order

$$\Delta F_{\mathrm{conf}} \asymp \frac{n}{h^2}. \tag{7.5}$$

This is complementary to the high-excursion action H^2/n in (6.4). The two scales are the spectral and variational faces of the same diffusive geometry.

7.2. **Fredkin spin chains.** The Fredkin spin chain provides a genuine quantum realization of the Catalan ensemble. In the spin-1/2 model, the ground state can be written as an equal-weight superposition of Dyck paths; related models and their correlation structure are developed in [10, 11]. Abstractly, if $\mathcal{D}_n$ is the Dyck basis,

$$|\Psi_n\rangle = \frac{1}{\sqrt{C_n}}\sum_{D\in\mathcal{D}_n} |D\rangle. \tag{7.6}$$

For any subset $\mathcal{A} \subseteq \mathcal{D}_n$ with projector $\Pi_{\mathcal{A}}$,

$$\langle\Psi_n|\Pi_{\mathcal{A}}|\Psi_n\rangle = \frac{|\mathcal{A}|}{C_n}. \tag{7.7}$$

Thus a Catalan free energy

$$\mathcal{F}_{\mathrm{Cat}}(\mathcal{A}) = -\log\frac{|\mathcal{A}|}{C_n} \tag{7.8}$$

has a literal interpretation as the negative logarithm of a quantum-sector weight in an equal-amplitude Catalan state.

Question 7.1. Can the arithmetic selector be represented by a local or quasi-local deformation of a Catalan transfer matrix or Fredkin-type Hamiltonian whose ground-state or low-energy sector contains the Mertens excursion? A positive answer would turn the selector problem into a spectral-energy estimate rather than a statement of random sampling.

8. Statistical mechanics of Dyck paths: pressure, area, and q-Catalan weights

Dyck paths are established physical configuration spaces for directed polymers and vesicles. Owczarek and Prellberg study area-weighted Dyck paths as a two-dimensional vesicle model and define pressure through finite-size free-energy differences [7]. Let $A(D)$ be the area under a Dyck path and define

$$Z_n(q) = \sum_{D \in \mathcal{D}_n} q^{A(D)}. \tag{8.1}$$

At $q = 1$,

$$Z_n(1) = C_n, \tag{8.2}$$

while $q \neq 1$ biases the ensemble by area. The asymptotics of area-weighted Dyck paths exhibit Airy-type scaling and have been analyzed in detail [8].

For a Mertens excursion D_r, define its absolute area

$$A_r = \sum_{j=z_{r-1}}^{z_r} |X_j|. \tag{8.3}$$

The pair (H_r, A_r) separates a sharp spike from a sustained displacement. A physical route based on q-Catalan weights would seek an arithmetic inequality on the partition weight of the sector containing (H_r, A_r).

Heuristic 8.1. In a Catalan polymer interpretation, excluding a region of path space produces an entropic pressure equal to a free-energy difference. An arithmetic excursion that remains at large height for a long interval can therefore be assigned a pressure or free-energy cost determined entirely by the number of Catalan paths compatible with that geometric constraint. The open problem is to connect this cost to the deterministic arithmetic selector.

9. Non-equilibrium transport and the Burgers route

The asymmetric exclusion process provides a distinct physical use of Brownian excursions. Derrida, Enaud, and Lebowitz showed that in the maximal-current phase of open TASEP, the non-Gaussian part of density fluctuations can be described in terms of Brownian excursion statistics [9]. At hydrodynamic scale, TASEP is governed by a conservation law of Burgers type,

$$\partial_t \rho + \partial_x\big(\rho(1-\rho)\big) = 0. \tag{9.1}$$

This creates a rigorous physical chain linking constrained lattice paths, excursion fluctuations, current transport, and non-linear hydrodynamics.

The present project does not identify the Mertens walk with TASEP. The possible use is structural: construct a dynamics on Catalan states whose conserved current or entropy production encodes the arithmetic selector. If the selector can be expressed as a constrained transport process, tools from hydrodynamic limits, entropy solutions, shocks, and large-deviation functionals become available.

Question 9.1. Is there a local state variable on the filtered squarefree complex for which the parity imbalance X_m is a conserved or approximately conserved current, and for which the admissible Catalan updates induce a non-equilibrium transport law?

A positive answer could replace a direct estimate on $\mu(n)$ by an entropy-production inequality on the Catalan dynamics. This is speculative, but the TASEP/Brownian-excursion/Burgers chain shows that such a passage from discrete constrained paths to hydrodynamic free energies is a real mechanism in statistical physics, not merely a metaphor.

10. The deterministic arithmetic selector

10.1. **Björner's filtered complex.** For each squarefree integer k, let $P(k)$ be the set of its prime divisors and define

$$\Delta_x = \{P(k) : k \le x,\ k \text{ squarefree}\}. \tag{10.1}$$

This is an abstract simplicial complex because divisors of a squarefree integer correspond to subsets of its prime set. Björner observed and exploited the exact identity

$$M(x) = -\widetilde{\chi}(\Delta_x), \tag{10.2}$$

where $\widetilde{\chi}$ is the reduced Euler characteristic [1]. He further showed that these complexes are shifted and studied their Betti numbers, wedge-of-spheres homotopy type, and combinatorial shadow inequalities. Björner explicitly noted that these results did not by themselves yield new control of the growth of $M(x)$ [1].

The filtration

$$\Delta_1 \subseteq \Delta_2 \subseteq \Delta_3 \subseteq \cdots \tag{10.3}$$

adds a simplex precisely when the cutoff crosses a squarefree integer. The change in Euler characteristic is therefore the sign of the corresponding Möbius value. The compressed Mertens walk is exactly the Euler-characteristic trace of this filtration.

10.2. **Subset-sum energy representation.** A squarefree integer corresponds to a finite prime subset S, and its logarithm is additive:

$$E(S) = \log \prod_{p\in S} p = \sum_{p\in S} \log p. \tag{10.4}$$

The sign is the parity

$$\mu\left(\prod_{p\in S} p\right) = (-1)^{|S|}. \tag{10.5}$$

Thus the selector can be stated exactly as follows: order all finite prime subsets by the additive energy (10.4), then record the parity (10.5). Mertens is the cumulative parity imbalance.

For finitely many primes $p_1, \dots, p_d$, let

$$w = (w_1, \dots, w_d), \qquad w_j = \log p_j, \tag{10.6}$$

and

$$E_w(S) = \sum_{j\in S} w_j. \tag{10.7}$$

The relative order of two subsets changes on a hyperplane

$$H_{S,T} = \{w : \langle \mathbf{1}_S - \mathbf{1}_T, w\rangle = 0\}. \tag{10.8}$$

Unique factorization implies that the prime-log vector does not lie on any hyperplane $H_{S,T}$ with $S \neq T$.

Proposition 10.1. *If a and b are distinct positive integers with $a, b \le X$, then*

$$|\log a - \log b| \ge \log\left(1 + \frac{1}{X}\right). \tag{10.9}$$

Consequently, the ordering of any fixed finite family of squarefree products $\le X$ is separated from its subset-sum collision hyperplanes by a positive deterministic margin.

Proof. Assume $a > b$. Then $a \ge b + 1$ and $b \le X$, so

$$\log(a/b) \ge \log(1 + 1/b) \ge \log(1 + 1/X). \tag{10.10}$$

□

This hyperplane-arrangement formulation gives a concrete geometric meaning to "selection": a finite parity word, and hence a finite Catalan excursion, is stable on a chamber of weight space. The arithmetic vector $(\log p_j)$ lies in one such chamber.

Definition 10.2. For a finite family $\mathcal{S}$ of prime subsets, define the selector chamber of the prime-log vector to be the connected component containing $w = (\log p_j)$ of

$$\mathbb{R}^d_{>0} \setminus \bigcup_{S \ne T \in \mathcal{S}} H_{S,T}. \tag{10.11}$$

The chamber solid angle or normalized volume is a possible deterministic analogue of statistical weight. A useful theorem would compare the chamber volume of a Catalan-extreme sign word with that of ordinary sign words and then prove a lower bound for the chamber containing the actual prime-log vector.

11. Three possible selector mechanisms

11.1. **Hyperplane-chamber entropy.** The first route uses Definition 10.2. For a finite excursion involving a finite collection of subset energies, let Ω_r be the region of positive weight space that preserves the relevant ordering and hence the same parity word. Define a geometric selector energy

$$\mathcal{G}_r = -\log \nu(\Omega_r), \tag{11.1}$$

where ν is a normalized reference measure on an appropriate projective section of positive weight space.

Question 11.1. Does a high Catalan excursion necessarily correspond to a small union of ordering chambers, with a bound of the form

$$\nu\{w : \text{induced excursion has height} \ge H\} \le \exp(-\Psi(H, L)), \tag{11.2}$$

where $\Psi(H, L)$ is at least comparable to H^2/L in the relevant regime?

If so, the remaining arithmetic problem becomes a Diophantine non-concentration statement for the prime-log vector relative to those chambers. Proposition 10.1 gives a first, very weak, deterministic separation from the collision walls.

11.2. **Shadow inequalities and parity restoration.** Björner obtains number-theoretic inequalities from combinatorial shadow functions associated with the shifted complex [1]. A family of many high-dimensional faces forces a sufficiently large lower-dimensional shadow. Since adjacent dimensions contribute opposite signs to Euler characteristic, this is a plausible source of parity compensation.

Warning 11.2. A shadow inequality is primarily static: it says that certain lower-dimensional faces exist when a higher-dimensional face exists. The Mertens problem is temporal along the filtration. Many shadow faces may have entered much earlier. The useful statement must therefore be a filtered shadow inequality controlling how much opposite parity is forced to enter within a bounded future energy interval.

The corresponding research target is a dynamic inequality of the schematic form

$$\text{current parity imbalance } H \quad \Longrightarrow \quad \text{future opposite-parity influx} \geq \Phi(H, L), \tag{11.3}$$

with a scale strong enough to feed into the Catalan energy H^2/L.

11.3. **Persistent topology.** Equation (10.2) can be refined by the Euler–Poincaré formula. Writing $\beta_k(\Delta_x)$ for reduced Betti numbers,

$$M(x) = \sum_{k\geq 0} (-1)^{k-1} \beta_k(\Delta_x). \tag{11.4}$$

Björner studied the asymptotic behaviour of these Betti numbers and proved that Δ_x has the homotopy type of a wedge of spheres [1]. A large Mertens excursion is therefore a sustained imbalance between even- and odd-dimensional homological contributions.

The filtration suggests studying not only $\beta_k(\Delta_x)$ at a fixed x but the birth and death structure across x. Persistent-homology constraints could in principle control the duration for which one parity dominates. This route preserves information discarded by the scalar Euler characteristic.

Question 11.3. Can one prove an interlacing, pairing, or energy inequality between persistent homology classes of adjacent parity whose projection onto Euler characteristic yields a bound on Catalan excursion height?

12. Structural viability of the physical routes

The physical structures above do not by themselves prove a new estimate for M. They are nevertheless mathematically relevant for four specific reasons.

First, the Catalan state space is exact, not fitted. Proposition 2.2 follows from the deterministic zero-return geometry of the compressed Mertens walk.

Second, the physical rate H^2/L is already visible in exact combinatorics. Proposition 4.4 supplies a finite discrete Gaussian tail, while Brownian action and large-deviation theory explain the same scale in the continuum.

Third, the arithmetic selector has additional exact structure. Equations (10.2) and (10.4) show that the path is generated by a shifted simplicial filtration and an additive prime-log ordering. Therefore the selector problem is not an arbitrary deterministic choice from Catalan space.

Fourth, Theorem 5.3 converts a concrete selector-energy inequality into square-root control. This isolates a finite combinatorial target that can be attacked by geometry, topology, spectral theory, or statistical mechanics.

13. Potential obstructions

13.1. **The selector may remain unconstrained at the needed scale.** The strongest obstruction is that the known simplicial and subset-sum structure may fail to prevent the prime-log ordering from entering Catalan sectors with very large selection energy. Establishing such a prevention is the substantive new theorem required by the program.

13.2. **Static topology may not control filtered dynamics.** Euler characteristic and Betti-number identities can exhibit large cancellation only after many filtration steps. RH-scale control requires cancellation at every sufficiently large scale. Any topological approach must therefore retain filtration time and cannot rely solely on the terminal homotopy type.

13.3. **Parity barrier.** The selector records the parity $(-1)^{|S|}$ of prime subsets. Many coarse counting and sieve mechanisms are weak precisely when asked to distinguish even from odd numbers of prime factors. In the present language, a successful selector theorem must see parity at the level of the ordered subset-sum geometry rather than only the density of faces.

13.4. **Canonical collapse to a classical Dirichlet object.** A canonical partition function built directly from the signed energies (10.4) gives

$$\sum_S (-1)^{|S|} e^{-sE(S)} = \prod_p (1 - p^{-s}) = \frac{1}{\zeta(s)} \tag{13.1}$$

for $\Re s > 1$. Equation (13.1) is exact, but it collapses the excursion geometry into the familiar Euler product. The new information sought here lies in pathwise objects—return times, Catalan shapes, heights, areas, chamber geometry, and filtered topology—before such a transform removes the temporal structure.

Remark 13.1. Equation (13.1) is used as a diagnostic boundary: a proposed physical construction that immediately reduces to this canonical partition function has returned to the classical analytic representation and has not exploited the Catalan excursion structure.

14. A hierarchy of possible sufficient principles

The following principles are ordered from closest to the desired estimate to more structural formulations.

Conjecture 14.1 (Open Catalan selection principle)**.** *For the compressed Mertens walk,*

$$\mathcal{I}_{\mathrm{Cat}}(m) = O(\log m). \tag{14.1}$$

By Theorem 5.3, Conjecture 14.1 implies (5.5).

Conjecture 14.2 (Chamber non-concentration principle)**.** *For a natural finite-dimensional normalization of the prime-log selector, the chamber containing the arithmetic weight vector has geometric energy at most logarithmic in the cutoff, and Catalan-extreme endpoint sectors have chamber energy bounded below by a positive multiple of H^2/L.*

Conjecture 14.3 (Filtered parity-restoration principle)**.** *There is a deterministic filtered shadow or persistence functional $\mathcal{R}$ such that a parity imbalance H sustained for L compressed steps forces*

$$\mathcal{R} \geq c\frac{H^2}{L} - O(\log L), \tag{14.2}$$

and the arithmetic filtration satisfies $\mathcal{R} = O(\log m)$ up to compressed time m.

These conjectures are not independent claims of truth. They are candidate formulations of the same missing selector mechanism at different levels: direct Catalan volume, prime-log chamber geometry, and filtered topology.

15. The prime-delay selector and its exact interference algebra

This section makes the selector more explicit than the chamber description. The identities are exact; the passage from them to local Catalan moment bounds is not known and will be isolated separately.

15.1. **Simultaneous prime-shell identities.** For a prime p, define

$$A_p(x) = \sum_{\substack{n \leq x \\ p \nmid n}} \mu(n). \tag{15.1}$$

Terms divisible by p can be written uniquely as pm with $p \nmid m$; for these terms $\mu(pm) = -\mu(m)$. Therefore

$$M(x) = A_p(x) - A_p(x/p) = \sum_{\substack{x/p < n \leq x \\ p \nmid n}} \mu(n) \qquad (p \text{ prime}). \tag{15.2}$$

Thus a value $M(x)$ must be reproduced simultaneously as a filtered parity imbalance in every prime shell $(x/p, x]$. For $p = 2$ this is the shell underlying Björner's odd-squarefree Betti-number formula [1]. Identity (15.2) is elementary, but it is not a generic property of a Catalan path.

15.2. **Threshold kernel and transfer recursion.** Fix $y \geq 2$, let $P_y = \{p : p \leq y\}$, and put

$$K_y(t) = \sum_{\substack{S \subseteq P_y \\ \sum_{p \in S} \log p \leq t}} (-1)^{|S|}. \tag{15.3}$$

Unique factorization gives, for $0 \leq t \leq \log y$,

$$K_y(t) = M(e^t). \tag{15.4}$$

Here and below $M(u)$ for real u means $\sum_{n \leq u} \mu(n)$. The restriction $t \leq \log y$ is essential: outside it the finite prime set P_y no longer contains every prime that may occur below the threshold.

Enumerate the primes as $p_1 < p_2 < \cdots$ and define K_j by using only $p_1, \ldots, p_j$. Splitting subsets according to whether they contain p_j gives the exact recursion

$$K_j(t) = K_{j-1}(t) - K_{j-1}(t - \log p_j). \tag{15.5}$$

With $(T_a f)(t) = f(t - a)$ and $\mathcal{T}_a = I - T_a$,

$$K_j = \mathcal{T}_{\log p_j} K_{j-1}. \tag{15.6}$$

This is a deterministic delayed anti-copy operation: every new prime inserts a translated copy with the opposite sign.

Let dK_j denote the atomic jump measure. Distributionally,

$$dK_j = mathop{*}_{\ell \leq j} (\delta_0 - \delta_{\log p_\ell}), \qquad \widehat{dK_j}(\tau) = \prod_{\ell \leq j} (1 - e^{-i\tau \log p_\ell}), \tag{15.7}$$

and hence

$$|\widehat{dK_j}(\tau)| = 2^j \prod_{\ell \leq j} \left| \sin \frac{\tau \log p_\ell}{2} \right|. \tag{15.8}$$

Equations (15.5)–(15.8) are exact finite identities [DED].

For an integer $q \geq 1$, define the normalized frequency moment

$$\mathcal{D}_{j,q}(T) = \frac{1}{2T} \int_{-T}^{T} \left(2^{-j} |\widehat{dK_j}(\tau)| \right)^{2q} d\tau \tag{15.9}$$

and let $P_j = \prod_{\ell \leq j} p_\ell$.

Theorem 15.1 (Exact Cesàro dephasing law)**.** *For every fixed $j, q \geq 1$,*

$$\lim_{T \to \infty} \mathcal{D}_{j,q}(T) = \left(\frac{\binom{2q}{q}}{4^q} \right)^j. \tag{15.10}$$

Moreover,

$$\left| \mathcal{D}_{j,q}(T) - \left(\frac{\binom{2q}{q}}{4^q} \right)^j \right| \leq \min\left\{ 1, \frac{1}{T \log(1 + P_j^{-q})} \right\}. \tag{15.11}$$

Proof. The elementary Fourier expansion

$$\sin^{2q}(\theta/2) = 4^{-q} \sum_{m=-q}^{q} (-1)^m \binom{2q}{q+m} e^{im\theta} \tag{15.12}$$

has constant coefficient $\binom{2q}{q}/4^q$ and coefficient ℓ^1 norm equal to one. Multiplying these expansions for $\theta = \tau \log p_\ell$ produces frequencies

$$\lambda_m = \sum_{\ell \le j} m_\ell \log p_\ell = \log(A/B), \qquad |m_\ell| \le q. \tag{15.13}$$

Unique factorization makes every nonzero vector m a nonzero frequency. Its symmetric Cesàro average is $\sin(T\lambda_m)/(T\lambda_m)$, proving (15.10). After cancellation, $A, B \le P_j^q$; if $A > B$ then $\log(A/B) \ge \log(1 + P_j^{-q})$. Summing with the unit ℓ^1 norm gives (15.11). □

Warning 15.2 (What the τ theorem does and does not control). The same rational independence and Kronecker's theorem give

$$\sup_{\tau \in \mathbb{R}} 2^{-j} |\widehat{dK_j}(\tau)| = 1. \tag{15.14}$$

Thus Theorem 15.1 is an averaged statement, not pointwise decay. Its elementary finite-window error becomes useful only for very large T because P_j^q is enormous. More importantly, the product (15.7) contains all subsets of $p_1, \dots, p_j$, including products beyond the arithmetic cutoff y. Restriction to the causal region $t \le \log y$ destroys that product factorization. These two facts prevent the infinite-window average from being used directly as a proof of (15.20).

Figure 1 illustrates the theorem for the first few primes. The dashed levels are the exact $q = 1$ limits 2^{-j}; the curves are finite computations and have no asymptotic evidentiary role.




FIGURE 1. Normalized prime sine products and running Cesàro square means.

15.3. **A maximal-parity Walsh coefficient.** Put $j = \pi(y)$ and define the Boolean halfspace

$$f_{y,t}(\varepsilon) = \mathbf{1}\left\{ \sum_{\ell=1}^{j} \varepsilon_\ell \log p_\ell \le t \right\}, \qquad \varepsilon \in \{0,1\}^j. \tag{15.15}$$

Under the normalized Walsh convention,

$$\widehat{f}_{y,t}([j]) = 2^{-j} \sum_{\varepsilon \in \{0,1\}^j} (-1)^{\varepsilon_1 + \cdots + \varepsilon_j} f_{y,t}(\varepsilon), \tag{15.16}$$

we have

$$K_y(t) = 2^j \widehat{f}_{y,t}([j]), \qquad M(y) = 2^{\pi(y)} \widehat{f}_{y,\log y}([\pi(y)]). \tag{15.17}$$

Fourier analysis of Boolean halfspaces is well developed [15], but the coefficient in (15.17) has maximal degree and the threshold lies in a highly sparse tail of the cube. Standard low-degree and balanced-halfspace estimates do not directly give the required bound.

15.4. **Uniform prefix energy: a corrected sufficient principle.** Let $0 = z_0 < z_1 < \cdots$ be the return times of the compressed walk. On a complete excursion set

$$D_r(u) = |X_{z_r+u}|, \qquad 0 \le u \le L_r := z_{r+1} - z_r, \tag{15.18}$$

and for every prefix $1 \le v \le L_r$ define

$$\mathcal{A}_q(r; v) = \sum_{u=1}^{v} D_r(u)^{2q}. \tag{15.19}$$

The same definition is used for every prefix of the currently open excursion, even when its next return time is not yet known.

Conjecture 15.3 (Uniform prime-Catalan prefix bound)**.** *For every fixed integer $q \ge 1$ and every $\varepsilon > 0$,*

$$\mathcal{A}_q(r; v) \ll_{q,\varepsilon} v^{q+1}(z_r + v)^{\varepsilon} \tag{15.20}$$

uniformly over all completed excursions, all their prefixes, and all prefixes of the final open meander.

Theorem 15.4 (Uniform prefix moments imply RH)**.** *Conjecture 15.3 implies the Riemann hypothesis.*

Proof. Fix a compressed time N, let $z_* \le N$ be its last return time, and put $v = N - z_*$. If $v = 0$ there is nothing to prove. Otherwise the prefix containing N is included in Conjecture 15.3, so

$$|X_N|^{2q} \le \mathcal{A}_q(r; v) \ll_{q,\varepsilon} v^{q+1} N^{\varepsilon} \le N^{q+1+\varepsilon}. \tag{15.21}$$

Therefore

$$|X_N| \ll_{q,\varepsilon} N^{1/2+1/(2q)+\varepsilon/(2q)}. \tag{15.22}$$

Given $\delta > 0$, choose a fixed q sufficiently large and then choose $\varepsilon > 0$ sufficiently small. This gives $X_N = O_\delta(N^{1/2+\delta})$. Since the squarefree counting function $Q(x) \sim 6x/\pi^2$ and $M(x) = X_{Q(x)}$, the classical Mertens criterion for RH follows. □

Remark 15.5 (Logical strength). Theorem 15.4 is a rigorous implication, not an equivalence. The uniform prefix conjecture can be strictly stronger than RH. Its purpose is to expose a structured sufficient target that controls complete excursions, internal prefixes, and the open meander without predicting their individual signs.

15.5. **The exceptional Catalan class is exactly the RH obstruction.** Let $\mathfrak{C}$ denote the space of infinite nearest-neighbour paths, decomposed at their returns to zero into signed primitive Catalan excursions and, at every finite cutoff, a possibly open terminal meander. Define the RH-exceptional Catalan class by

$$\mathfrak{E}_{\mathrm{RH}} = \left\{ Y \in \mathfrak{C} : \exists \varepsilon_0 > 0 \text{ such that } \sup_{m \ge 1} \frac{|Y_m|}{m^{1/2+\varepsilon_0}} = \infty \right\}. \tag{15.23}$$

Equivalently, membership is tested over every internal excursion prefix and every open terminal meander. This last clause cannot be omitted.

Theorem 15.6 (Exceptional-class characterization)**.** *Let $X_m = \sum_{j \le m} \mu(q_j)$ be the squarefree-compressed Mertens walk. Then*

$$\boxed{X \notin \mathfrak{E}_{\mathrm{RH}} \quad \Longleftrightarrow \quad \mathrm{RH}.} \tag{15.24}$$

Thus, for the arithmetic path, not belonging to the exceptional Catalan class is exactly the Riemann hypothesis.

Proof. By squarefree counting,

$$Q(x) = \frac{6}{\pi^2}x + O(\sqrt{x}), \qquad q_m \asymp m, \tag{15.25}$$

and $M(x) = X_{Q(x)}$. Therefore

$$M(x) = O_\varepsilon(x^{1/2+\varepsilon}) \quad (\varepsilon > 0) \tag{15.26}$$

if and only if

$$X_m = O_\varepsilon(m^{1/2+\varepsilon}) \quad (\varepsilon > 0). \tag{15.27}$$

The latter statement is exactly $X \notin \mathfrak{E}_{\mathrm{RH}}$ by Equation (15.23); the former is the classical Mertens criterion for RH. □

Proposition 15.7 (Exact global class versus stronger local class)**.** *In excursion coordinates, the condition*

$$D_r(v) \ll_\varepsilon (z_r + v)^{1/2+\varepsilon} \qquad \textit{for every } r, v \textit{ and every } \varepsilon > 0 \tag{15.28}$$

including the open meander, is exactly equivalent to $X \notin \mathfrak{E}_{\mathrm{RH}}$ and hence to RH. It is neither stronger nor weaker. By contrast, a bound measured at the internal scale,

$$D_r(v) \ll_\varepsilon v^{1/2+\varepsilon}, \tag{15.29}$$

or the uniform local moment bound (15.20)*, is a strictly stronger property on the space of nearest-neighbour paths and is not known to follow from RH for the Mertens path.*

Proof. At the prefix (r, v) the current compressed time is $m = z_r + v$ and $D_r(v) = |X_m|$, which proves the first assertion point by point. To separate the two path properties, place at arbitrarily large times N a primitive mountain excursion of length $L \asymp \sqrt{N}$ and height $L/2$, separating successive mountains by unit excursions that return immediately to zero. The resulting nearest-neighbour path obeys a global bound $|Y_m| = O(\sqrt{m})$, but at the mountain peak it has height comparable to L, violating (15.29) and, for fixed q, the Brownian local moment scale. Thus local Brownian control contains information not present in the global RH-scale condition. □

Warning 15.8 (Criterion is not mechanism). Theorem 15.6 is an exact characterization and, by itself, proves no new estimate. Nevertheless, its equivalence to RH does not make every attempted proof circular. Circularity occurs only if exclusion from $\mathfrak{E}_{\mathrm{RH}}$ is assumed, renamed, or recovered by immediate substitution. A proof from an independent arithmetic theorem about the prime selector would be a non-circular proof of an equivalent criterion, just as for any other valid RH criterion.

There are therefore two distinct assertions:

$$\underbrace{X \notin \mathfrak{E}_{\mathrm{RH}} \iff \mathrm{RH}}_{\text{exact criterion [DED]}} \qquad \text{and} \qquad \underbrace{\text{prime selector} \Longrightarrow X \notin \mathfrak{E}_{\mathrm{RH}}}_{\text{open mechanism [ADAPT]}}. \tag{15.30}$$

The first statement does not close the Catalan programme. It identifies its target. The programme remains open precisely because the second statement is neither proved nor disproved.

Controlling only completed excursions by their future return time is not enough for the reverse implication: a return may lie far beyond the point being estimated, and an indefinitely open meander would escape the condition. The correct local formulation is the uniform prefix hierarchy (15.20), which acts at the present scale $z_r + v$.

15.6. **No-go audit: what the exact identities do not prove.** The exact Fourier and prime-shell identities above remain useful diagnostics, but Version 5 removes them from the proposed proof mechanism. The following statements isolate the reason.

Proposition 15.9 (Catalan universality obstruction)**.** *For every $n \geq 1$ there is a primitive Dyck excursion of length $2n$ and height n. More generally, every finite concatenation of primitive Dyck excursions followed by a strict meander is realized by a unique finite word of increments in $\{-1, +1\}$.*

Proof. The mountain word $+^n-^n$ is primitive and has height n. For the second statement, read the successive differences of the prescribed paths and concatenate them. The return skeleton and the final meander recover the decomposition uniquely. □

Corollary 15.10 (Catalan-only no-go theorem)**.** *No estimate that uses only the nearest-neighbour property, the return skeleton, primitive Catalan enumeration, or uniform Catalan large deviations can imply a sublinear bound for the selected path. An additional arithmetic statement excluding the mountain words is logically necessary.*

The corollary does not deny the exact Catalan structure. It says that ambient counting and typicality cannot select the Mertens word from that structure.

Proposition 15.11 (Redundant prime lift)**.** *Let $a = q_{z_r}$ be a return anchor, $x = q_{z_r+v}$, and*

$$A_p(y) = \sum_{\substack{n \leq y \\ p \nmid n}} \mu(n). \tag{15.31}$$

For every prime p,

$$\big(A_p(x) - A_p(a)\big) - \big(A_p(x/p) - A_p(a/p)\big) = X_{z_r+v}. \tag{15.32}$$

Consequently, averaging any power of the left-hand side over primes reproduces the corresponding Catalan prefix moment exactly and gives no estimate.

Proof. Subtract the exact shell identity $M(y) = A_p(y) - A_p(y/p)$ at $y = x$ and $y = a$, then use $M(a) = X_{z_r} = 0$. The final assertion follows because the difference is independent of p. □

This proposition invalidates the prime-dilation "alignment" proposal as an independent bridge: after definitions are expanded, its desired inequality is precisely Conjecture 15.3 again.

Proposition 15.12 (Stability floor for smoothed chambers)**.** *Fix $x \geq 2$ and the squarefree integers not exceeding $2x$. Perturb their prime weights by*

$$w_p(\theta, \xi) = \log p + \theta \xi_p, \qquad |\xi_p| \leq 1. \tag{15.33}$$

Put

$$\delta_x = \frac{\log(1 + 1/(2x))}{4 \log_2(2x)}. \tag{15.34}$$

For every choice of ξ, the complete order of those squarefree integers is unchanged whenever $|\theta| \leq \delta_x$. Hence any bad-excursion event true for the arithmetic order occupies at least the interval $[-\delta_x, \delta_x]$ in every perturbation direction.

Proof. Distinct integers $m, n \leq 2x$ satisfy

$$|\log m - \log n| \geq \log(1 + 1/(2x)). \tag{15.35}$$

The perturbation of their energy difference is at most

$$|\theta|(\omega(m) + \omega(n)) \leq 2|\theta| \log_2(2x). \tag{15.36}$$

At the radius (15.34) this is at most half the deterministic gap, so no pair reverses order. □

If θ is uniform on a fixed interval $[-\eta, \eta]$, an actual bad excursion therefore forces bad probability at least

$$\frac{\delta_x}{\eta} \asymp \frac{1}{x \log x}. \tag{15.37}$$

A smoothed Catalan tail smaller than this floor cannot be obtained merely by replacing the arithmetic order with a typical perturbed order: it must already control the rigid neighbourhood containing the arithmetic point. Version 5 therefore does not promote smoothed analysis to a proof route.

Definition 15.13 (Admissible Catalan–arithmetic bridge)**.** A future bridge is called admissible only if it satisfies all four tests:

(i) it is false for at least one primitive Dyck mountain word;
(ii) it uses a proved arithmetic relation not recoverable from the values of the single prefix path alone;
(iii) after all definitions are substituted, it is not the desired uniform prefix bound itself;
(iv) it is uniform over internal prefixes and the final open meander.

No admissible bridge is proved in this notebook. This is the precise boundary left after the audit. The exact target remains Conjecture 15.3; calling it a selector, alignment, dephasing, or free-energy estimate does not constitute progress unless an admissible arithmetic mechanism proves it.

16. An aggregate route that does not predict M pointwise

The pointwise selection principle in Conjecture 14.1 is deliberately strong. It asks for a restriction at every time and can therefore reproduce, in disguised form, the original difficulty of controlling $M(x)$. This section replaces that hypothesis by averaged positive quantities. A large value cannot hide from sufficiently high moments because the Mertens walk has increments of absolute value at most one.

Remark 16.1 (Provenance)*.* The use of moments to recover a point value from local regularity is a classical analytic mechanism. The specific time-moment criterion and its decomposition over Catalan excursions below are deductions in the present notebook [DED]. Verjovsky's local Fourier-moment criterion is a distinct recent realization of a related moment-to-point principle [14] [LIT]; it averages a Möbius Fourier polynomial over a short frequency arc, whereas the criterion below averages the Mertens walk over time. The weak Mertens conjecture is another classical averaged sufficient condition, but it is stronger than RH and also forces simplicity of the zeros [13] [LIT].

16.1. **A hierarchy of moments equivalent to RH.** For an integer $k \geq 1$, define

$$\mathcal{B}_k(X) = \sum_{n \leq X} |M(n)|^{2k}. \tag{16.1}$$

Theorem 16.2 (Aggregate moment criterion)**.** *The Riemann hypothesis is equivalent to the following family of estimates: for every fixed integer $k \geq 1$ and every $\varepsilon > 0$,*

$$\mathcal{B}_k(X) \ll_{k,\varepsilon} X^{k+1+\varepsilon}. \tag{16.2}$$

Proof. Assume RH. For every $\eta > 0$ the classical Mertens criterion gives $|M(n)| \ll_\eta n^{1/2+\eta}$. Given k and ε, take $\eta = \varepsilon/(2k)$. Summation gives (16.2), after an inessential adjustment of ε.

Conversely, suppose (16.2) holds for every fixed k. Fix N and write $H = |M(N)|$. Since

$$|M(n+1) - M(n)| \leq 1, \tag{16.3}$$

we have $|M(N+j)| \geq H/2$ for $0 \leq j \leq \lfloor H/2 \rfloor$. Moreover $H \leq N$, so these indices are at most $2N$. Hence, for $H \geq 4$,

$$\mathcal{B}_k(2N) \geq c_k H^{2k+1} \tag{16.4}$$

with an absolute admissible choice such as $c_k = 2^{-2k-2}$. It follows that

$$H \ll_{k,\varepsilon} N^{(k+1+\varepsilon)/(2k+1)}. \tag{16.5}$$

Given $\delta > 0$, choose k so large that $1/(4k+2) < \delta/2$, and then choose ε so that $\varepsilon/(2k+1) < \delta/2$. Equation (16.5) yields $M(N) = O_\delta(N^{1/2+\delta})$, which is equivalent to RH. □

The first RH-equivalent statement in the new route is therefore Theorem 16.2. No Catalan, physical, or topological statement preceding it proves RH. This is the dependency-graph checkpoint for the remainder of the program.

Corollary 16.3 (Finite-order milestones)**.** *For a single fixed k, estimate* (16.2) *implies*

$$M(X) \ll_{k,\varepsilon} X^{\frac{k+1}{2k+1}+\varepsilon}. \tag{16.6}$$

In particular, $k = 1, 2, 3$ give the successive exponents $2/3$, $3/5$, and $4/7$.

Warning 16.4. Even the case $k = 1$ would be a major unconditional advance: it would give a fixed zero-free half-plane $\Re s > 2/3$. The hierarchy avoids pointwise sign prediction, but it does not make the arithmetic cancellation automatic. Its advantage is that it supplies auditable intermediate targets rather than an all-or-nothing selector axiom.

16.2. **Exact excursion decomposition of the moment budget.** In compressed time put

$$\mathcal{B}_k^{\#}(Y) = \sum_{m \le Y} |X_m|^{2k}. \tag{16.7}$$

The proof of Theorem 16.2, together with $q_m \asymp m$, gives the analogous equivalence

$$\text{RH} \iff \mathcal{B}_k^{\#}(Y) ll_{k,\varepsilon} Y^{k+1+\varepsilon} \quad \text{for every fixed } k \text{ and every } \varepsilon > 0. \tag{16.8}$$

For an excursion D of length L, define its integrated $2k$-energy

$$\mathcal{E}_k(D) = \sum_{u=1}^{L-1} |D(u)|^{2k}. \tag{16.9}$$

At a cutoff Y, let $D_1, \dots, D_R$ be the completed excursions and let U_Y be the open terminal meander of length t_Y. Then the identity

$$\boxed{\mathcal{B}_k^{\#}(Y) = \sum_{r=1}^{R} \mathcal{E}_k(D_r) + \mathcal{E}_k(U_Y)} \tag{16.10}$$

is exact. It converts RH into a hierarchy of total geometric-energy bounds over the actual excursions, without requiring the prediction of any individual value of X_m.

Under the uniform Catalan law at fixed length L, Brownian scaling predicts

$$\mathbb{E}_{\mathrm{Cat},L} \mathcal{E}_k(D) \asymp_k L^{k+1}. \tag{16.11}$$

This motivates the length budget

$$\Lambda_k(Y) = \sum_{r=1}^{R} L_r^{k+1} + t_Y^{k+1}. \tag{16.12}$$

Since the segment lengths sum to Y,

$$\Lambda_k(Y) \le Y^{k+1}. \tag{16.13}$$

Conjecture 16.5 (Aggregate Catalan domination)**.** *For every fixed $k \ge 1$ and every $\varepsilon > 0$,*

$$\mathcal{B}_k^{\#}(Y) \ll_{k,\varepsilon} Y^{\varepsilon} \Lambda_k(Y). \tag{16.14}$$

Proposition 16.6. *Conjecture 16.5 for all fixed k implies RH.*

Proof. Combine (16.14) with (16.13) and then apply (16.8). □

Warning 16.7. Conjecture 16.5 is a sufficient selector principle and may be strictly stronger than RH. RH controls the total moment $\mathcal{B}_k^{\#}(Y)$ but does not by itself compare it with the potentially much smaller random length budget $\Lambda_k(Y)$. The conjecture is proposed as a structured attack target, not as an equivalent reformulation.

Unlike a pointwise percentile bound, aggregate Catalan domination permits exceptional excursions: one excursion may have energy far above its Catalan mean provided the total energy remains within the global length budget. This is the precise sense in which the route avoids predicting the real Mertens path excursion by excursion.

16.3. **The arithmetic correlation target.** Write $a_j = \mu(q_j) \in \{-1, +1\}$. Expanding the positive moment and changing the order of summation gives the exact identity

$$\mathcal{B}_k^{\#}(Y) = \sum_{1 \le i_1, \dots, i_{2k} \le Y} a_{i_1} \cdots a_{i_{2k}} \big(Y - \max(i_1, \dots, i_{2k}) + 1\big). \tag{16.15}$$

The configurations in which every index occurs with even multiplicity make a contribution of order at most $O_k(Y^{k+1})$. The remaining configurations form an explicit weighted off-diagonal correlation sum. Thus the analytic problem is not to predict a_j but to prove cancellation in the aggregate off-diagonal part of (16.15).

Definition 16.8 (Off-diagonal correlation budget)**.** Let $\mathcal{O}_k(Y)$ denote the part of (16.15) indexed by tuples for which at least one index has odd multiplicity.

A concrete sufficient target is

$$\mathcal{O}_k(Y) = O_{k,\varepsilon}(Y^{k+1+\varepsilon}) \tag{16.16}$$

for every fixed k. This target retains ordering and high-order sign correlations; it cannot be replaced by one-point equidistribution or by a fixed-shift Chowla statement without an additional uniform inversion theorem.

16.4. **Finite evidence and its exact limitation.**

Experiment 16.9. For the supplied file through ordinary cutoff 455052511, compressed time is $Y = 276638778$. Direct blockwise summation gave the following ratios. The second column divides $\mathcal{B}_k^{\#}(Y)$ by the leading simple-walk moment $((2k-1)!!/(k+1))Y^{k+1}$; the third divides it by $\Lambda_k(Y)$, including the open terminal meander.

k	simple-walk ratio	$\mathcal{B}_k^{\#}(Y)/\Lambda_k(Y)$
1	3.70956×10^{-2}	0.404561
2	8.92436×10^{-4}	0.251180
3	1.64775×10^{-5}	0.188051
4	2.48322×10^{-7}	0.154931
5	3.16760×10^{-9}	0.135848
6	3.50903×10^{-11}	0.125099

These are finite measured values, not estimates uniform in Y. They support the scale in Conjecture 16.5 on this one range but provide no asymptotic control.

16.5. **A staged proof program.** The route now has intermediate statements that can fail independently:

(M1) Prove (16.2) for $k = 1$. This yields the exponent $2/3 + \varepsilon$ and tests whether the method supplies any new arithmetic cancellation.

(M2) Prove the excursion-budget inequality (16.14) for $k = 1$ by a filtered parity-restoration or transfer-operator estimate.




FIGURE 2. Finite compressed-time moment ratios through $k = 6$. The logarithmic left panel compares with the leading simple-walk moment; the right panel shows $\mathcal{B}_k^{\#}(Y)/\Lambda_k(Y)$.




FIGURE 3. Observed excursion height and return-gap geometry through $N = 455{,}052{,}511$. Both panels are descriptive finite data only.

(M3) Extend the argument to each fixed k, with constants allowed to depend on k. No uniformity in growing k is required for Theorem 16.2.

(M4) Translate the resulting bounds into the off-diagonal formulation (16.16); identify which multiplicity patterns are controlled by known Möbius correlation theorems and which remain new.

(M5) Only after a genuine estimate is proved at M1 or M2 should one construct a pinning, transfer, or hydrodynamic model intended to explain the estimate. A physical model fitted to the finite data is not an arithmetic proof.

This hierarchy is a clear route around pointwise prediction, but it is not a shortcut around cancellation. Its first falsifiable analytic target is the quadratic energy

$$\sum_{m \le Y} X_m^2 \ll_\varepsilon Y^{2+\varepsilon}, \tag{16.17}$$

and its first RH-complete target is the same estimate at every fixed even moment.

17. Research program

The route can be advanced without assuming a probabilistic model for Möbius by attacking finite, exact questions.

17.1. **Exact finite combinatorics.** For each actual excursion or meander prefix, compute

$$(L, H, A, S_{\mathrm{Cat}}, \mathcal{I}_{\mathrm{Cat}}), \tag{17.1}$$

where L is duration, H is maximum height or current endpoint, A is area, S_{Cat} is Catalan entropy, and $\mathcal{I}_{\mathrm{Cat}}$ is the open-sector selection energy. The formulas for all ambient counts are exact or can be evaluated by stable dynamic programming/transfer matrices.

17.2. **Finite selector chambers.** For a finite set of primes and squarefree products, build the hyperplane arrangement (10.8). Determine which chambers generate Catalan excursions of a prescribed length and height. The first structural question is whether the total chamber volume of height-H words exhibits the same H^2/L rate as the uniform Catalan ensemble or a stronger one.

17.3. **Dynamic shadows.** Translate Björner's static shadow inequalities into statements on increments of face numbers over multiplicative windows

$$[x, xe^{\delta}]. \tag{17.2}$$

The target is a lower bound on opposite-parity arrivals conditional on a current Euler-characteristic imbalance.

17.4. **Transfer operators.** Construct a transfer operator whose states retain both Catalan height and a finite amount of arithmetic filtration data. The ordinary Dyck transfer matrix has spectral gap scale h^{-2}; an arithmetic deformation could reveal whether the prime-log selector adds a confining or restoring term.

17.5. **Hydrodynamic coarse graining.** If a local transport rule emerges, derive its conservation law and large-deviation functional. The TASEP example shows that Brownian-excursion fluctuations can coexist with Burgers hydrodynamics [9]. The corresponding Mertens question is whether the arithmetic selector generates an entropy condition that suppresses macroscopic parity shocks.

18. Relation to existing literature

The closest direct probabilistic work on Möbius located in this literature survey is the study of Mussardo and LeClair, which reorganizes the nonzero Möbius coefficients into a ± 1 walk and investigates Brownian/random-walk behaviour through statistical tests and block structures [12]. The present program differs in its primary invariant: the exact Catalan state space attached to each return excursion and its open meander sectors.

On the mathematical-probability side, conditioned random walks converging to Brownian excursion or meander are classical [2, 3, 4]; the maximum of a random Dyck path and its exponential moments and deviations are quantitatively understood [5, 6]. On the statistical-mechanics side, Dyck paths occur as vesicle/polymer configurations with genuine partition functions and free-energy differences [7, 8]. On the quantum side, Fredkin-type spin chains realize Dyck-path superpositions as physical states [10, 11]. On the non-equilibrium side, Brownian excursions arise in exact TASEP fluctuation calculations, with Burgers-type hydrodynamics at macroscopic scale [9].

On the arithmetic-topological side, Björner's complex gives the exact identity (10.2), shifted-complex structure, Betti asymptotics, and shadow inequalities, while explicitly leaving the growth problem for M unresolved [1]. The potential new intersection is therefore

$$\boxed{\text{filtered arithmetic selector} + \text{Catalan excursion state space} + \text{physical free-energy geometry}}. \tag{18.1}$$

The notebook treats the selector theorem as the decisive missing component rather than as an implicit randomness assumption.

19. Conclusions and open questions

The compressed Mertens function is an exact nearest-neighbour walk, and every complete zero-to-zero excursion is a primitive Dyck path counted by a Catalan number. This yields a canonical finite state space for each excursion and a natural open-boundary meander space for every point inside an excursion. Standard Catalan probability then supplies a quantitative geometric energy: large endpoint displacement or height is exponentially expensive on the scale H^2/L.

The central unresolved issue is selection. The arithmetic path is generated by a deterministic ordering of prime-subset energies and simultaneously by a filtered shifted simplicial complex whose Euler characteristic is Mertens. The project becomes viable only if one of these structures can be shown to bound the Catalan selection energy.

Version 5 performs the hidden-equivalence audit explicitly. The shell identity (15.2), transfer recursion (15.5), convolution product (15.7), and maximal Walsh coefficient (15.17) remain exact diagnostic identities. The Catalan-only no-go theorem, redundant-prime-lift proposition, and stability floor show why none of them currently supplies the missing cancellation.

The most concrete first targets are:

(i) prove nontrivial chamber-volume inequalities for parity words induced by positive additive subset weights;
(ii) derive filtered versions of Björner's shadow inequalities over multiplicative windows;
(iii) test whether persistent homology forces adjacent-parity compensation on the timescale of a Catalan excursion;
(iv) construct an arithmetic deformation of the Dyck transfer matrix and identify its spectral energy;
(v) determine whether a current/transport formulation leads to an entropy-production principle analogous to non-equilibrium lattice gases.
(vi) subject every proposed bridge to Definition 15.13 before treating it as a route toward the uniform Catalan prefix bound.

A successful result need not identify Möbius with a random walk. The new aggregate route separates two logically different targets. Theorem 16.2 is exactly equivalent to RH and asks only for all fixed even time moments; it does not predict the value or sign of any individual $M(n)$. The aggregate Catalan domination principle is a structured sufficient mechanism for those moment bounds, but is explicitly left as a possibly stronger conjecture. Thus the first falsifiable analytic milestone is the quadratic estimate $\sum_{m\le Y} X_m^2 \ll_\varepsilon Y^{2+\varepsilon}$, while the original pointwise selection-energy theorem remains a stronger alternative rather than the definition of the route.

The only logically valid conditional chain retained in Version 5 is

$$\begin{aligned} &\text{admissible arithmetic–Catalan bridge} \\ &\quad\Longrightarrow \text{uniform prefix moment hierarchy} \\ &\quad\Longrightarrow |M(x)| \ll_\varepsilon x^{1/2+\varepsilon} \Longrightarrow \mathrm{RH}. \end{aligned} \tag{19.1}$$

Theorem 15.4 supplies the second arrow. No claim is made that the first arrow has been proved or reduced to a standard estimate. This negative conclusion is deliberate: it prevents exact but tautological identities from being advertised as progress toward RH.

Appendix A. Reproducible computations and archived outputs

The accompanying archive contains the six Python programs printed below and their outputs. The binary input is not redistributed; it consists of $\mu(1),\dots,\mu(N)$ stored as signed bytes. For the supplied file with $N = 455{,}052{,}511$, the moment audit found 276,638,778 compressed steps,

53,078 returns, $M(N) = -5414$, and maximum observed absolute height 6695. The computed ratios $\mathcal{B}_k^{\#}/\Lambda_k$ for $k = 1, \dots, 6$ were respectively

$$0.4045606,\ 0.2511805,\ 0.1880514,\ 0.1549310,\ 0.1358477,\ 0.1250992. \tag{A.1}$$

These are finite measurements only and are not used as asymptotic premises.

The finite selector audit checked the prime-shell identity through 10^5 for $p = 2, 3, 5, 7, 11, 97$, the threshold-kernel identity through 30, the exact transfer factorization, and the Fourier product at four frequencies. All integer identities passed; the largest floating-point Fourier discrepancy was 9.54×10^{-14}.

Typical commands are

```
python3 code/mertens_moment_audit.py /path/to/moebius.bin \
  --output outputs/moment_audit.json
python3 code/prime_selector_identity_check.py \
  --output outputs/selector_check.json
python3 code/analiza_alturas_excursiones_mertens.py /path/to/moebius.bin \
  --output outputs/mertens_excursiones_alturas.csv
python3 code/tau_dephasing_audit.py \
  --output-png outputs/tau_dephasing.png \
  --output-json outputs/tau_dephasing.json
python3 code/plot_catalan_mertens_outputs.py \
  --moments outputs/moment_audit.json \
  --excursions outputs/mertens_excursiones_alturas.csv \
  --output-dir outputs
python3 code/catalan_no_go_audit_v5.py
```

Appendix B. Python source: moment hierarchy audit

```python
#!/usr/bin/env python3
"""Blockwise audit of compressed-time Mertens moments and excursion budgets.

Input format: one signed int8 value mu(n) for each n=1,...,N.
The program never materializes the entire Mertens walk in memory.
"""

from __future__ import annotations

import argparse
import json
import math
from pathlib import Path

import numpy as np


def odd_double_factorial(k: int) -> int:
    value = 1
    for j in range(1, k + 1):
        value *= 2 * j - 1
    return value


def parse_args() -> argparse.Namespace:
    parser = argparse.ArgumentParser()
    parser.add_argument("filename", type=Path)
    parser.add_argument("--block", type=int, default=5_000_000)
    parser.add_argument("--max-k", type=int, default=6)
    parser.add_argument("--output", type=Path, default=Path("moment_audit.json"))
    return parser.parse_args()


```

```python
def main() -> None:
    args = parse_args()
    if args.max_k < 1:
        raise ValueError("--max-k must be positive")

    size = args.filename.stat().st_size
    data = np.memmap(args.filename, dtype=np.int8, mode="r", shape=(size,))
    moments = np.zeros(args.max_k, dtype=np.float64)
    length_budgets = [0] * args.max_k

    mertens = 0
    compressed = 0
    previous_zero = 0
    returns = 0
    max_abs = 0

    for lo in range(0, size, args.block):
        hi = min(lo + args.block, size)
        block = np.asarray(data[lo:hi])
        nz = np.flatnonzero(block)
        if nz.size == 0:
            continue

        steps = block[nz].astype(np.int64, copy=False)
        walk = np.cumsum(steps, dtype=np.int64)
        walk += mertens
        absolute = np.abs(walk).astype(np.float64, copy=False)
        max_abs = max(max_abs, int(absolute.max(initial=0.0)))

        power = absolute * absolute
        for k in range(1, args.max_k + 1):
            if k > 1:
                power = power * absolute * absolute
            moments[k - 1] += float(np.sum(power, dtype=np.float64))

        hits = np.flatnonzero(walk == 0)
        for hit in hits:
            zero = compressed + int(hit) + 1
            gap = zero - previous_zero
            for k in range(1, args.max_k + 1):
                length_budgets[k - 1] += gap ** (k + 1)
            previous_zero = zero
            returns += 1

        mertens = int(walk[-1])
        compressed += int(nz.size)

    tail = compressed - previous_zero
    records = []
    for k in range(1, args.max_k + 1):
        budget = length_budgets[k - 1] + tail ** (k + 1)
        iid_leading = (
            odd_double_factorial(k) / (k + 1) * float(compressed) ** (k + 1)
        )
        records.append(
            {
                "k": k,
                "B_k": float(moments[k - 1]),
                "Lambda_k": str(budget),
                "B_k_over_Lambda_k": float(moments[k - 1] / budget),
                "B_k_over_iid_leading": float(moments[k - 1] / iid_leading),
            }
        )

    result = {
```

```
        "input": str(args.filename.resolve()),
        "N": size,
        "compressed_steps": compressed,
        "M_N": mertens,
        "returns_to_zero": returns,
        "last_zero_compressed": previous_zero,
        "open_tail_compressed": tail,
        "max_abs_M": max_abs,
        "moments": records,
        "interpretation": "finite computation only; no asymptotic inference",
    }
    args.output.parent.mkdir(parents=True, exist_ok=True)
    args.output.write_text(json.dumps(result, indent=2) + "\n", encoding="utf-8")
    print(json.dumps(result, indent=2))


if __name__ == "__main__":
    main()
```

Appendix C. Python source: exact selector identities

```
#!/usr/bin/env python3
"""Finite exact checks of the prime-shell, transfer, and Fourier identities."""

from __future__ import annotations

import argparse
import cmath
import json
import math
from pathlib import Path


def mobius_sieve(limit: int) -> list[int]:
    mu = [1] * (limit + 1)
    prime = [True] * (limit + 1)
    prime[0:2] = [False, False]
    primes: list[int] = []
    mu[0] = 0
    for n in range(2, limit + 1):
        if prime[n]:
            primes.append(n)
            mu[n] = -1
        for p in primes:
            if n * p > limit:
                break
            prime[n * p] = False
            if n % p == 0:
                mu[n * p] = 0
                break
            mu[n * p] = -mu[n]
    return mu


def primes_up_to(limit: int) -> list[int]:
    sieve = [True] * (limit + 1)
    sieve[0:2] = [False, False]
    for p in range(2, int(math.isqrt(limit)) + 1):
        if sieve[p]:
            for n in range(p * p, limit + 1, p):
                sieve[n] = False
    return [p for p in range(2, limit + 1) if sieve[p]]

```

```python

def parse_args() -> argparse.Namespace:
    parser = argparse.ArgumentParser()
    parser.add_argument("--limit", type=int, default=100_000)
    parser.add_argument("--kernel-x", type=int, default=30)
    parser.add_argument("--output", type=Path, default=Path("selector_check.json"))
    return parser.parse_args()


def main() -> None:
    args = parse_args()
    mu = mobius_sieve(args.limit)
    mertens = [0] * (args.limit + 1)
    for n in range(1, args.limit + 1):
        mertens[n] = mertens[n - 1] + mu[n]

    test_primes = [p for p in (2, 3, 5, 7, 11, 97) if p <= args.limit]
    shell_failures = []
    for p in test_primes:
        filtered = [0] * (args.limit + 1)
        for n in range(1, args.limit + 1):
            filtered[n] = filtered[n - 1] + (mu[n] if n % p else 0)
        for x in range(1, args.limit + 1):
            shell = filtered[x] - filtered[x // p]
            if shell != mertens[x]:
                shell_failures.append({"p": p, "x": x})
                break

    kernel_primes = primes_up_to(args.kernel_x)
    atoms = [(1, 1)]
    for p in kernel_primes:
        atoms += [(product * p, -sign) for product, sign in atoms]
    atoms.sort()
    kernel_failures = []
    for x in range(1, args.kernel_x + 1):
        value = sum(sign for product, sign in atoms if product <= x)
        if value != mertens[x]:
            kernel_failures.append({"x": x, "K": value, "M": mertens[x]})

    convolution = {1: 1}
    for p in kernel_primes:
        updated = dict(convolution)
        for product, sign in convolution.items():
            updated[product * p] = updated.get(product * p, 0) - sign
        convolution = updated
    transfer_ok = sorted(convolution.items()) == atoms

    fourier_errors = []
    for tau in (0.125, 0.5, 1.0, 2.75):
        direct = sum(sign * cmath.exp(-1j * tau * math.log(product)) for product, sign in atoms)
        product = 1.0 + 0.0j
        for p in kernel_primes:
            product *= 1.0 - cmath.exp(-1j * tau * math.log(p))
        fourier_errors.append(abs(direct - product))

    result = {
        "limit": args.limit,
        "tested_shell_primes": test_primes,
        "shell_failures": shell_failures,
        "kernel_x": args.kernel_x,
        "kernel_primes": kernel_primes,
        "kernel_failures": kernel_failures,
        "transfer_factorization_ok": transfer_ok,
        "max_fourier_factorization_error": max(fourier_errors),
        "status": "PASS" if not shell_failures and not kernel_failures and transfer_ok else "FAIL",
```

```
        "interpretation": "finite verification of exact identities, not an RH test",
    }
    args.output.parent.mkdir(parents=True, exist_ok=True)
    args.output.write_text(json.dumps(result, indent=2) + "\n", encoding="utf-8")
    print(json.dumps(result, indent=2))


if __name__ == "__main__":
    main()
```

Appendix D. Python source: excursion heights and Catalan tails

```
#!/usr/bin/env python3
"""Extrae longitudes, alturas y areas de las excursiones de Mertens.

El fichero de entrada contiene mu(1),...,mu(N) como int8. Las posiciones
mu(n)=0 se eliminan y el paseo comprimido tiene pasos +/-1.
"""

from __future__ import annotations

import argparse
import csv
import math
import time
from pathlib import Path

import numpy as np


def parse_args() -> argparse.Namespace:
    parser = argparse.ArgumentParser()
    parser.add_argument("filename", type=Path)
    parser.add_argument("--block", type=int, default=5_000_000)
    parser.add_argument(
        "--output",
        type=Path,
        default=Path("mertens_excursiones_alturas.csv"),
    )
    return parser.parse_args()


def catalan_height_upper_tail(length: int, height: int) -> float:
    """P(maximo >= height) en la excursion uniforme de longitud ``length``.

    Se quitan el primer paso ascendente y el ultimo descendente. Queda un
    Dyck path de longitud 2*n=length-2 y altura maxima height-1. El conteo
    acotado se obtiene del espectro del grafo camino de alturas 0,...,a.
    """
    if length == 2:
        return 1.0
    n = length // 2 - 1
    a = height - 2
    if a < 0:
        return 1.0
    j = np.arange(1, a + 2, dtype=np.float64)
    theta = math.pi * j / (a + 2)
    eigen_abs = np.abs(2.0 * np.cos(theta))
    with np.errstate(divide="ignore"):
        terms = (
            math.log(2.0 / (a + 2))
            + 2.0 * np.log(np.sin(theta))
            + 2.0 * n * np.log(eigen_abs)
```

```
        )
    peak = float(np.max(terms))
    log_bounded = peak + math.log(float(np.exp(terms - peak).sum()))
    log_catalan = (
        math.lgamma(2 * n + 1) - math.lgamma(n + 1) - math.lgamma(n + 2)
    )
    log_cdf = min(0.0, log_bounded - log_catalan)
    return min(1.0, max(0.0, -math.expm1(log_cdf)))


def main() -> None:
    args = parse_args()
    size = args.filename.stat().st_size
    data = np.memmap(args.filename, dtype=np.int8, mode="r", shape=(size,))

    rows: list[tuple[int, int, int, int, int, int, int, int, int, float]] = []
    mertens = 0
    compressed = 0
    previous_zero_n = 0
    previous_zero_m = 0
    open_height = 0
    open_area = 0
    open_sign = 0
    started = time.monotonic()

    for lo in range(0, size, args.block):
        hi = min(lo + args.block, size)
        block = np.asarray(data[lo:hi])
        nz = np.flatnonzero(block)
        if nz.size:
            steps = block[nz].astype(np.int64, copy=False)
            walk = np.cumsum(steps, dtype=np.int64)
            walk += mertens
            hits = np.flatnonzero(walk == 0)

            start = 0
            for hit in hits:
                segment = walk[start : hit + 1]
                if segment.size:
                    if open_sign == 0:
                        open_sign = int(np.sign(segment[0]))
                    open_height = max(open_height, int(np.max(np.abs(segment))))
                    open_area += int(np.sum(np.abs(segment), dtype=np.int64))

                end_m = compressed + int(hit) + 1
                end_n = lo + int(nz[hit]) + 1
                length = end_m - previous_zero_m
                rows.append(
                    (
                        len(rows) + 1,
                        previous_zero_n,
                        end_n,
                        previous_zero_m,
                        end_m,
                        length,
                        open_sign,
                        open_height,
                        open_area,
                        open_height / math.sqrt(length),
                    )
                )
                previous_zero_n = end_n
                previous_zero_m = end_m
                open_height = 0
                open_area = 0
```

```
                open_sign = 0
                start = int(hit) + 1

            if start < walk.size:
                segment = walk[start:]
                if open_sign == 0:
                    open_sign = int(np.sign(segment[0]))
                open_height = max(open_height, int(np.max(np.abs(segment))))
                open_area += int(np.sum(np.abs(segment), dtype=np.int64))

            mertens = int(walk[-1])
            compressed += int(nz.size)

        block_number = lo // args.block + 1
        if block_number % 10 == 0 or hi == size:
            elapsed = time.monotonic() - started
            print(
                f"bloque={block_number:3d} n={hi:12d} m={compressed:12d} "
                f"M={mertens:7d} retornos={len(rows):7d} t={elapsed:7.1f}s",
                flush=True,
            )

    args.output.parent.mkdir(parents=True, exist_ok=True)
    columns = (
        "Return",
        "Start_n",
        "End_n",
        "Start_m",
        "End_m",
        "Gap",
        "ExcursionSign",
        "MaxHeight",
        "Area",
        "HeightOverSqrtGap",
        "ReducedHeightOverSqrtLength",
        "CatalanHeightUpperTailExact",
        "HeightFreeEnergyNats",
        "HeightFreeEnergyBits",
    )
    with args.output.open("w", newline="") as handle:
        writer = csv.writer(handle)
        writer.writerow(columns)
        for row in rows:
            length = row[5]
            height = row[7]
            reduced_scaled = (
                (height - 1) / math.sqrt(length - 2) if length > 2 else 0.0
            )
            upper_tail = catalan_height_upper_tail(length, height)
            free_nats = -math.log(upper_tail)
            writer.writerow(
                row
                + (
                    reduced_scaled,
                    upper_tail,
                    free_nats,
                    free_nats / math.log(2.0),
                )
            )

    gaps = np.fromiter((row[5] for row in rows), dtype=np.int64)
    heights = np.fromiter((row[7] for row in rows), dtype=np.int64)
    scaled = heights / np.sqrt(gaps)
    print("\nRESUMEN")
    print(f"N={size}")
```

```
    print(f"m={compressed}")
    print(f"M(N)={mertens}")
    print(f"retornos={len(rows)}")
    print(f"ultimo_retorno_n={previous_zero_n}")
    print(f"ultimo_retorno_m={previous_zero_m}")
    print(f"cola_m={compressed-previous_zero_m}")
    print(f"gap_max={int(gaps.max())}")
    print(f"altura_max_absoluta={int(heights.max())}")
    print(f"max_altura_sobre_raiz_gap={float(scaled.max()):.12g}")
    for q in (0.5, 0.9, 0.95, 0.99, 0.999):
        print(f"q{q:g}_altura_sobre_raiz_gap={np.quantile(scaled, q):.12g}")
    print(f"csv={args.output.resolve()}")


if __name__ == "__main__":
    main()
```

Appendix E. Python source: finite τ-dephasing illustration

```
#!/usr/bin/env python3
"""Illustrate the exact Cesaro dephasing law for finite prime products.

The plot is numerical evidence only.  The limiting formula used for the
horizontal reference lines is proved algebraically in the accompanying paper.
"""

from __future__ import annotations

import argparse
import json
import math
from pathlib import Path

import matplotlib.pyplot as plt
import numpy as np


PRIMES = np.array([2, 3, 5, 7, 11, 13, 17, 19], dtype=np.float64)


def parse_args() -> argparse.Namespace:
    parser = argparse.ArgumentParser()
    parser.add_argument("--tau-max", type=float, default=10_000.0)
    parser.add_argument("--points", type=int, default=200_001)
    parser.add_argument("--output-png", type=Path, default=Path("tau_dephasing.png"))
    parser.add_argument("--output-json", type=Path, default=Path("tau_dephasing.json"))
    return parser.parse_args()


def running_cesaro(values: np.ndarray) -> np.ndarray:
    return np.cumsum(values, dtype=np.float64) / np.arange(1, values.size + 1)


def main() -> None:
    args = parse_args()
    tau = np.linspace(0.0, args.tau_max, args.points)
    fig, axes = plt.subplots(1, 2, figsize=(11.0, 4.2), constrained_layout=True)
    records = []

    for j, color in ((4, "#1967ad"), (6, "#c43c39"), (8, "#4b8f29")):
        normalized = np.prod(
            np.abs(np.sin(0.5 * tau[:, None] * np.log(PRIMES[:j])[None, :])),
            axis=1,
```

```
        )
        square_mean = running_cesaro(normalized * normalized)
        target = 2.0 ** (-j)
        axes[0].plot(tau[:4001], normalized[:4001], color=color, lw=0.8, label=f"j={j}")
        axes[1].plot(tau[1:], square_mean[1:], color=color, lw=1.0, label=f"j={j}")
        axes[1].axhline(target, color=color, ls="--", lw=0.8, alpha=0.8)
        records.append(
            {
                "j": j,
                "tau_max": args.tau_max,
                "sample_cesaro_square": float(square_mean[-1]),
                "exact_infinite_cesaro_square": target,
            }
        )

    axes[0].set_title("Normalized prime sine product")
    axes[0].set_xlabel(r"$\tau$ (initial window)")
    axes[0].set_ylabel(r"$2^{-j}|\widehat{dK_j}(\tau)|$")
    axes[0].legend(frameon=False)
    axes[0].grid(alpha=0.2)

    axes[1].set_title("Running Cesaro mean of the square")
    axes[1].set_xscale("log")
    axes[1].set_yscale("log")
    axes[1].set_xlabel(r"frequency cutoff $T$")
    axes[1].set_ylabel("running mean; dashed = exact limit")
    axes[1].legend(frameon=False)
    axes[1].grid(alpha=0.2, which="both")

    args.output_png.parent.mkdir(parents=True, exist_ok=True)
    fig.savefig(args.output_png, dpi=220)
    plt.close(fig)
    result = {
        "records": records,
        "interpretation": "finite illustration; the limit is an exact theorem",
    }
    args.output_json.parent.mkdir(parents=True, exist_ok=True)
    args.output_json.write_text(json.dumps(result, indent=2) + "\n", encoding="utf-8")
    print(json.dumps(result, indent=2))


if __name__ == "__main__":
    main()
```

Appendix F. Python source: publication figures

```
#!/usr/bin/env python3
"""Create publication PNGs from the archived moment and excursion outputs."""

from __future__ import annotations

import argparse
import csv
import json
from pathlib import Path

import matplotlib.pyplot as plt
import numpy as np


def parse_args() -> argparse.Namespace:
    parser = argparse.ArgumentParser()
    parser.add_argument("--moments", type=Path, required=True)
```

```
    parser.add_argument("--excursions", type=Path, required=True)
    parser.add_argument("--output-dir", type=Path, required=True)
    return parser.parse_args()


def main() -> None:
    args = parse_args()
    args.output_dir.mkdir(parents=True, exist_ok=True)
    moment_data = json.loads(args.moments.read_text(encoding="utf-8"))
    records = moment_data["moments"]
    k = np.array([row["k"] for row in records])
    iid = np.array([row["B_k_over_iid_leading"] for row in records])
    budget = np.array([row["B_k_over_Lambda_k"] for row in records])

    fig, axes = plt.subplots(1, 2, figsize=(10.5, 4.0), constrained_layout=True)
    axes[0].semilogy(k, iid, "o-", color="#1967ad")
    axes[0].set_xlabel("moment order k")
    axes[0].set_ylabel(r"$\mathcal{B}_k^{\#}$/simple-walk leading moment")
    axes[0].set_title("Compressed Mertens moment ratios")
    axes[0].grid(alpha=0.25, which="both")
    axes[1].plot(k, budget, "o-", color="#c43c39")
    axes[1].set_xlabel("moment order k")
    axes[1].set_ylabel(r"$\mathcal{B}_k^{\#}/\Lambda_k$")
    axes[1].set_ylim(0.0, max(0.45, 1.08 * budget.max()))
    axes[1].set_title("Observed excursion-budget ratio")
    axes[1].grid(alpha=0.25)
    fig.savefig(args.output_dir / "moment_ratios.png", dpi=220)
    plt.close(fig)

    gaps: list[int] = []
    scaled: list[float] = []
    with args.excursions.open(newline="") as handle:
        for row in csv.DictReader(handle):
            gaps.append(int(row["Gap"]))
            scaled.append(float(row["HeightOverSqrtGap"]))
    gap = np.asarray(gaps, dtype=np.float64)
    height_scaled = np.asarray(scaled, dtype=np.float64)
    ordered = np.sort(gap)
    ccdf = (ordered.size - np.arange(ordered.size)) / ordered.size

    fig, axes = plt.subplots(1, 2, figsize=(10.5, 4.0), constrained_layout=True)
    axes[0].scatter(gap, height_scaled, s=5, alpha=0.18, color="#1967ad", edgecolors="none")
    axes[0].set_xscale("log")
    axes[0].set_xlabel("compressed excursion length L")
    axes[0].set_ylabel(r"$H/\sqrt{L}$")
    axes[0].set_title("Height across excursion scales")
    axes[0].grid(alpha=0.2, which="both")
    axes[1].loglog(ordered, ccdf, color="#c43c39", lw=1.2)
    axes[1].set_xlabel("compressed gap L")
    axes[1].set_ylabel(r"empirical $\Pr(\mathrm{Gap}\geq L)$")
    axes[1].set_title("Heavy-tailed return gaps")
    axes[1].grid(alpha=0.2, which="both")
    fig.savefig(args.output_dir / "excursion_geometry.png", dpi=220)
    plt.close(fig)

    print(args.output_dir / "moment_ratios.png")
    print(args.output_dir / "excursion_geometry.png")


if __name__ == "__main__":
    main()
```

Appendix G. Python source: Version 5 no-go audit

```python
#!/usr/bin/env python3
"""Finite audit for the Catalan--Mertens v5 criterion/mechanism note."""

from __future__ import annotations

import json
import math
from pathlib import Path

def catalan(n: int) -> int:
    return math.comb(2 * n, n) // (n + 1)


def primitive_dyck_paths(semilength: int):
    length = 2 * semilength

    def rec(pos: int, height: int, word: list[int]):
        if pos == length:
            if height == 0:
                yield tuple(word)
            return
        for step in (1, -1):
            new_height = height + step
            if new_height < 0:
                continue
            if pos + 1 < length and new_height == 0:
                continue
            if pos + 1 == length and new_height != 0:
                continue
            word.append(step)
            yield from rec(pos + 1, new_height, word)
            word.pop()

    yield from rec(0, 0, [])


def heights(word):
    out = [0]
    for step in word:
        out.append(out[-1] + step)
    return out


def mobius_sieve(limit: int):
    mu = [1] * (limit + 1)
    prime = [True] * (limit + 1)
    prime[0:2] = [False, False]
    for p in range(2, limit + 1):
        if not prime[p]:
            continue
        for n in range(p, limit + 1, p):
            prime[n] = False if n != p else prime[n]
            mu[n] *= -1
        square = p * p
        for n in range(square, limit + 1, square):
            mu[n] = 0
    mu[0] = 0
    return mu


def prime_dilation_error(mu, limit: int):
    mertens = [0] * (limit + 1)
    for n in range(1, limit + 1):
        mertens[n] = mertens[n - 1] + mu[n]
```

```
    primes = [p for p in range(2, 50) if all(p % d for d in range(2, int(p**0.5) + 1))]
    maximum = 0
    for p in primes:
        ap = [0] * (limit + 1)
        for n in range(1, limit + 1):
            ap[n] = ap[n - 1] + (mu[n] if n % p else 0)
        for x in range(1, limit + 1):
            maximum = max(maximum, abs(mertens[x] - (ap[x] - ap[x // p])))
    return maximum


def main():
    root = Path(__file__).resolve().parents[1]
    output = root / "outputs"
    output.mkdir(parents=True, exist_ok=True)

    rows = []
    for n in range(1, 8):
        paths = list(primitive_dyck_paths(n))
        maxima = [max(heights(path)) for path in paths]
        rows.append(
            {
                "semilength": n,
                "enumerated_primitive_paths": len(paths),
                "expected_catalan": catalan(n - 1),
                "largest_possible_height": max(maxima),
            }
        )

    mu = mobius_sieve(5000)
    identity_error = prime_dilation_error(mu, 5000)
    report = {
        "purpose": "audit identities and demonstrate the Catalan-only obstruction",
        "primitive_excursions": rows,
        "prime_dilation_identity_max_abs_error": identity_error,
        "conclusion": (
            "Primitive Catalan structure permits mountain excursions of height L/2; "
            "the prime-dilation average reproduces M exactly and supplies no bound."
        ),
    }
    (output / "catalan_no_go_audit_v5.json").write_text(
        json.dumps(report, indent=2), encoding="utf-8"
    )

    print(json.dumps(report, indent=2))


if __name__ == "__main__":
    main()
```

References

[1] A. Björner, *A cell complex in number theory*, Adv. Appl. Math. **46** (2011), 71–85. DOI: 10.1016/j.aam.2010.09.007. arXiv:1101.5704.

[2] W. D. Kaigh, *An invariance principle for random walk conditioned by a late return to zero*, Ann. Probab. **4** (1976), 115–121. DOI: 10.1214/aop/1176996189.

[3] R. T. Durrett, D. L. Iglehart, and D. R. Miller, *Weak convergence to Brownian meander and Brownian excursion*, Ann. Probab. **5** (1977), 117–129. DOI: 10.1214/aop/1176995895.

[4] E. Csáki and S. G. Mohanty, *Excursion and meander in random walk*, Canad. J. Statist. **9** (1981), 57–70. DOI: 10.2307/3315296.

[5] O. Khorunzhiy and J.-F. Marckert, *Uniform bounds for exponential moment of maximum of a Dyck path*, Electron. Commun. Probab. **14** (2009), 327–333. DOI: 10.1214/ECP.v14-1486. arXiv:0902.1229.

[6] T. Kousha, *Asymptotic behavior and the moderate deviation principle for the maximum of a Dyck path*, Statist. Probab. Lett. **82** (2012), 340–347. DOI: 10.1016/j.spl.2011.10.015. arXiv:1008.0606.
[7] A. L. Owczarek and T. Prellberg, *Pressure exerted by a vesicle on a surface*, J. Phys. A: Math. Theor. **47** (2014), 215001. DOI: 10.1088/1751-8113/47/21/215001. arXiv:1311.2174.
[8] N. Haug and T. Prellberg, *Uniform asymptotics of area-weighted Dyck paths*, J. Math. Phys. **56** (2015), 043301. DOI: 10.1063/1.4917052.
[9] B. Derrida, C. Enaud, and J. L. Lebowitz, *The asymmetric exclusion process and Brownian excursions*, J. Stat. Phys. **115** (2004), 365–382. DOI: 10.1023/B:JOSS.0000019833.35328.b4. arXiv:cond-mat/0306078.
[10] L. Dell'Anna, O. Salberger, L. Barbiero, A. Trombettoni, and V. E. Korepin, *Violation of cluster decomposition and absence of light cones in local integer and half-integer spin chains*, Phys. Rev. B **94** (2016), 155140. DOI: 10.1103/PhysRevB.94.155140. arXiv:1604.08281.
[11] O. Salberger and V. Korepin, *Entangled spin chain*, Rev. Math. Phys. **29** (2017), 1750031. DOI: 10.1142/S0129055X17500313.
[12] G. Mussardo and A. LeClair, *Randomness of Möbius coefficients and Brownian motion: growth of the Mertens function and the Riemann hypothesis*, J. Stat. Mech. Theory Exp. (2021), 113106. DOI: 10.1088/1742-5468/ac22fb. arXiv:2101.10336.
[13] N. Ng, *The distribution of the summatory function of the Möbius function*, Proc. London Math. Soc. (3) **89** (2004), 361–389. DOI: 10.1112/S0024611504014721. arXiv:math/0310381.
[14] A. Verjovsky, *How random is the Möbius function? Smoothing, probability, and the Riemann hypothesis*, arXiv:2607.25002v2, 2026.
[15] K. Matulef, R. O'Donnell, R. Rubinfeld, and R. A. Servedio, *Testing halfspaces*, SIAM J. Comput. **39** (2010), 2004–2047. DOI: 10.1137/070707890.