HEISENBERG UNCERTAINTY AND TWISTED MÖBIUS DYNAMICS VIII: TWISTED MÖBIUS CURVATURE, EULER-LOCALITY DEFECT, AND RIGOROUS MÖBIUS FILTERING OF PRIME–ZETA REPLICAS

Abstract. We study an arithmetic specialization of the Weyl canonical commutation relations obtained from the map $n \mapsto (\ell \log n, \kappa\Omega(n))$. The resulting Weyl cocycle defines a twisted Dirichlet convolution and a deformed Möbius inverse z_ϑ. We prove an exact closed formula for its second deformation curvature, derive its Dirichlet generating function as a prime-log covariance determinant, and isolate the logarithmic branching inherited from the prime zeta function. We then compare the twisted inverse with a norm-convergent ordered Euler object. The odd infinitesimal Euler-locality defect has a Dirichlet series $\mathcal{O}(s)$ which, for $\Re s > 1$, admits both an absolutely convergent prime-pair expansion and a Fourier-distance integral.

The principal new analytic result of Version 8 is a domain-complete treatment of that integral. Absolute convergence, local uniform convergence, differentiation, and all sum–integral interchanges are justified in their stated domains. After continuation from $\Re s > 1$, every contribution arriving at a scaled zero $s_0 = \rho/r$ is included, including rational collisions. For the normalized energy $\mathcal{I}(s) = 2\pi\mathcal{O}(s)/\zeta(s)$, the singular–singular block attached to (ρ, r) has logarithmic-over-pole coefficient $-\pi m_\rho^2 \varphi(r) r^{-2} \sum_{d|r} \mu(d)$. Consequently this coefficient vanishes identically for every $r > 1$ and survives for $r = 1$. We prove that no other contour singularity can restore a term of the same order. At a pure replica the next coefficient is a strictly negative divisor energy β_r, for which we give both an integral and a finite divisor-sum formula, together with the exact radical law $\beta_r = (\operatorname{rad} r/r)\beta_{\operatorname{rad} r}$. We also record the exact distinction between the normalized energy $\mathcal{I}$ and $\mathcal{O}$ itself: multiplication by ζ lowers the singular order at an actual zeta zero. No new restriction on the location of the zeta zeros, and therefore no proof of the Riemann hypothesis, is claimed.

Contents

1. Introduction: the Version 8 structure 3
2. Notation, domains, and symbol index 4
2.1. Basic arithmetic and analytic notation 5
2.2. Operator and deformation notation 5
2.3. Euler-locality and replica notation 5
3. Exact canonical kinematics 5
3.1. Hilbert space, position, and momentum 5
3.2. The Heisenberg uncertainty principle and exact incompatibility 6
4. Weyl displacements and the arithmetic phase space 7
4.1. Weyl operators 7
4.2. Arithmetic embedding 8
5. Structural filter: linear Möbius states cannot escape second order 10
6. Arithmetic uncertainty generated by the CCR 10
6.1. The semiclassical and arithmetic-commutative limits are distinct 11
7. The Heisenberg-twisted Dirichlet algebra 12
7.1. Definition and associativity 12
7.2. Weighted Banach algebra and operator representation 12
8. Möbius inversion deformed by the CCR 13
8.1. The Möbius operator 13
8.2. Formal twisted inverse 13
8.3. A bounded inverse in a rigorous half-plane 14

9. The first Heisenberg arithmetic law 15
10. Curvature of the cocyclically deformed Möbius inverse 17
11. Multiplicative geometry of the curvature 19
12. Dirichlet summation of the curvature 20
13. Continuation of the curvature and detection of zeta zeros 22
14. Where the logarithmic branching comes from 23
15. Local monodromy and the Jordan-totient weight 23
16. Gauge invariance of the arithmetic cocycle 24
17. Cohomological obstruction to scalar Euler locality 25
18. Ordered Euler object and the global Euler-locality defect 25
19. The odd Euler-locality defect: existence and exact initial domain 26
19.1. Endpoint regularity and transverse derivatives 28
20. Continuation of the Fourier energy and complete replica accounting 29
20.1. Grouped log-zeta expansion and its normal tail 29
20.2. All contributions arriving at a scaled point 31
20.3. Pure replicas: the next coefficient and the complete local germ 33
20.4. Single-replica divisor energy 34
20.5. Actual zeta zeros, $\mathcal{O}$, and monodromy 36
20.6. Analytic checklist for the replica theorem 37
21. A two-variable Euler product and the covariance determinant 37
22. The positivity route and its exact obstruction 38
23. The Heisenberg-deformed zeta series 39
24. A genuine Möbius Hamiltonian 40
24.1. Definition and domain 40
24.2. Exact equations of motion for position and momentum 41
24.3. The CCR is visible in the dynamics 42
25. Exact non-localization inside the arithmetic dynamics 43
26. Literature boundary and novelty status in Version 8 43
26.1. Known framework 43
26.2. Notebook-specific formulas not located in the targeted search 44
26.3. What is specifically new mathematics inside this manuscript 44
27. Corrections and extensions carried into Version 8 44
28. Open problems 44
29. Notebook history 45
30. Literature status of the Version 4/5 formulas 45
References 46

Remark 0.1 (Version 8: mathematical status). The canonical Weyl calculus, Robertson uncertainty inequality, general 2-cocycle machinery, prime-zeta continuation, and distance-energy identities are standard and are cited to primary or standard references. The operator series based only on translations have direct precedents in the spectral operator literature; in particular, Herichi–Lapidus rigorously study $\zeta(\partial)$ and its Möbius inverse [12]. The notebook-specific content is the arithmetic specialization of the cocycle, the exact twisted Möbius curvature, the ordered Euler-locality defect, and the replica-filtering identities proved below.

Version 8 replaces the informal replica analysis of Version 7 by a complete analytic audit. Every theorem-like statement used in the main chain is accompanied by a proof. Domains of unbounded operators, domains of absolute convergence, differentiability, normal convergence, exchanges of

sums and integrals, boundary terms, branch choices, contour singularities, and possible rational collisions are stated explicitly. The phrase "principal singularity" always refers to the displayed asymptotic scale and never to an unstated global ordering of multivalued germs.

1. Introduction: the Version 8 structure

Version 1 placed a standard Heisenberg pair next to a zeta multiplier but did not couple the two structures. Version 2 corrected that defect: the exact canonical commutation relation deformed the arithmetic product itself through a nontrivial cocycle. The Version 2 audit nevertheless identified two remaining weaknesses. First, the actual uncertainty principle was not stated prominently. Second, several domain, inverse, and convergence claims required repair.

Version 8 starts from the self-adjoint position and momentum operators

$$(Q\psi)(q) = q\psi(q), \qquad (P\psi)(q) = -i\hbar\psi'(q), \tag{1.1}$$

with the canonical commutation relation

$$[Q, P] = i\hbar I \tag{1.2}$$

on the Schwartz core. For every normalized Schwartz state, the Heisenberg–Robertson uncertainty inequality is [1]

$$\boxed{\Delta_\psi Q \, \Delta_\psi P \geq \frac{\hbar}{2}.} \tag{1.3}$$

Here the uncertainty is a mathematical spread intrinsic to the state, not an experimental error.

The arithmetic specialization assigns to an integer n the phase-space increments

$$x_n = \ell \log n, \qquad y_n = \kappa\Omega(n), \tag{1.4}$$

and the self-adjoint generator

$$G_n = y_n Q - x_n P. \tag{1.5}$$

The arithmetic specialization of the standard Robertson inequality gives, for two integers,

$$\boxed{\Delta_\psi G_m \, \Delta_\psi G_n \geq \frac{\hbar\ell\kappa}{2} \left|\Omega(m)\log n - \log m\, \Omega(n)\right|.} \tag{1.6}$$

For distinct primes p, q, this becomes

$$\boxed{\Delta_\psi G_p \, \Delta_\psi G_q \geq \frac{\hbar\ell\kappa}{2} \left|\log \frac{q}{p}\right|.} \tag{1.7}$$

Thus the same arithmetic symplectic area that later appears in the twisted Möbius product already appears as a lower bound forced directly by the Heisenberg uncertainty principle.

The associated Weyl operators are

$$W_n = \exp\left(\frac{i}{\hbar} G_n\right), \tag{1.8}$$

and satisfy

$$W_m W_n = \chi_\vartheta(m, n) W_{mn}, \tag{1.9}$$

where

$$\chi_\vartheta(m, n) = \exp\left[\frac{i\vartheta}{2}\big(\Omega(m)\log n - \log m\, \Omega(n)\big)\right], \qquad \vartheta = \frac{\ell\kappa}{\hbar}. \tag{1.10}$$

The cocycle is proved below to be cohomologically nontrivial for every $\vartheta \neq 0$. It induces an associative noncommutative Dirichlet product. Möbius inversion in that product defines coefficients $z_\vartheta(n)$. For prime powers and distinct prime pairs one obtains

$$z_\vartheta(p^r) = 1, \tag{1.11}$$

$$z_\vartheta(pq) = 2\cos\left(\frac{\vartheta}{2}\log\frac{q}{p}\right) - 1. \tag{1.12}$$

The same determinant-like arithmetic quantity therefore controls both the uncertainty lower bound (1.6) and the phase responsible for the deformed inverse.

The deformation also differentiates the twisted inverse itself at the commuting point. For

$$n = \prod_{j=1}^{r} p_j^{a_j}, \qquad T(a) = \frac{a(a+1)}{2}, \tag{1.13}$$

one obtains

$$\boxed{-z_0''(n) = \frac{1}{2}\sum_{1\le i<j\le r} T(a_i)T(a_j)\log^2\frac{p_j}{p_i}.} \tag{1.14}$$

For square-free $n = p_1\cdots p_r$,

$$\boxed{-z_0''(n) = \frac{1}{2}\left[r\sum_{p\mid n}(\log p)^2 - (\log n)^2\right] = \frac{r^2}{2}\,\mathrm{Var}_{p\mid n}(\log p).} \tag{1.15}$$

Summing this curvature does not return a two-variable Möbius correlation. For $\Re s > 1$,

$$\mathcal{K}(s) = -\sum_{n\ge 1}\frac{z_0''(n)}{n^s} = \frac{\zeta(s)}{2}\big(S_0(s)S_2(s) - S_1(s)^2\big), \tag{1.16}$$

where

$$S_j(s) = \sum_p \frac{p^{-s}}{(1-p^{-s})^2}(\log p)^j. \tag{1.17}$$

Section 12 rewrites these functions through $\log\zeta(rs)$ and $\zeta'(rs)/\zeta(rs)$, producing, after a branch is fixed by continuation from $\Re s > 1$, a local continued germ in $\Re s > 1/2$ and an exact singularity criterion for nontrivial zeta zeros to the right of the critical line.

Finally, a bounded self-adjoint Möbius Hamiltonian is defined for $\sigma > 1$. Its Schrödinger evolution and the exact Heisenberg equations for $Q(t)$ and $P(t)$ are derived with domains stated explicitly. The paper does not claim an advance toward RH. It records exact reformulations and local singularity statements involving the classical zero set, but no new restriction on the location of those zeros is proved.

Remark 1.1 (Scope of Version 8). The operator deformation is developed in the half-planes stated for each operator. Version 8 additionally develops a direct local relation between the second deformation curvature and the classical zero set in Section 13.

2. Notation, domains, and symbol index

This section fixes the notation used throughout the notebook. Reusing a symbol with two unrelated meanings is avoided in Version 8.

2.1. **Basic arithmetic and analytic notation.**

$\mu(n)$:	the Möbius function.
$\Omega(n)$:	the completely additive number of prime factors of n, counted with multiplicity.
$\omega(n)$:	the number of distinct prime divisors of n.
$v_p(n)$:	the exponent of p in n.
$\mathrm{rad}(n)$:	the square-free kernel $\prod_{p\mid n} p$.
$\varphi(n)$:	Euler's totient and $J_2(n)$ the second Jordan totient.
$T(a)$:	the triangular weight $a(a+1)/2$.
$\zeta(s)$:	the Riemann zeta function and $L(s) = \zeta'(s)/\zeta(s)$ its logarithmic derivative.
$P(s)$:	the prime-zeta function $\sum_p p^{-s}$ in its initial half-plane $\Re s > 1$, followed by analytic continuation when explicitly stated.
ρ:	a nontrivial zero of ζ; m_ρ denotes its multiplicity.
s, t:	Write $s = \sigma + i\tau$. The real variable t is reserved for the Fourier/vertical shift in $B(s,t)$.

2.2. **Operator and deformation notation.**

$\mathcal{H}$:	$L^2(\mathbb{R}, dq)$.
Q, P:	position and momentum. Their commutator identity is asserted on the Schwartz core $\mathcal{S}(\mathbb{R})$, not on the whole intersection of maximal domains.
$W(x,y)$:	the Weyl displacement and W_n its arithmetic specialization.
ϑ:	the dimensionless deformation parameter $\ell\kappa/\hbar$.
$\chi_\vartheta(m,n)$:	the arithmetic Weyl 2-cocycle.
$\star_\vartheta$:	twisted Dirichlet convolution.
$z_\vartheta(n)$:	the coefficientwise twisted inverse of μ.
$\mathcal{C}(n)$:	the deformation curvature $-z_0''(n)$.
$\mathcal{K}(s)$:	the Dirichlet series of $\mathcal{C}(n)$.

2.3. **Euler-locality and replica notation.**

E_ϑ:	the increasing-prime ordered Euler object.
R_ϑ:	the Euler-locality defect $z_\vartheta - E_\vartheta$.
$\Phi_{\mathrm{ord}}(n)$:	the oriented prime-pair phase of the ordered Euler object.
$\mathcal{O}(s)$:	the odd Euler-locality Dirichlet defect, initially holomorphic for $\Re s > 1$.
$u_p(s)$:	$p^{-s}/(1-p^{-s}) = 1/(p^s-1)$.
$A(s)$:	$\sum_p u_p(s) = B(s,0)$, initially for $\Re s > 1$.
$B(s,t)$:	$\sum_p u_p(s)p^{it}$, initially for $\Re s > 1$.
$\mathcal{I}(s)$:	the normalized Fourier energy $2\pi\mathcal{O}(s)/\zeta(s)$ in $\Re s > 1$. Its continued germ is the object whose leading local term is naturally of $z^{-1}\log z$-type.
s_0:	a fixed point in the strip, and $z = s - s_0$ the local coordinate.
$\mathcal{R}(s_0)$:	the finite set of dilation indices $N \geq 1$ for which $\zeta(Ns_0) = 0$.
q_N:	the multiplicity of the zero Ns_0, for $N \in \mathcal{R}(s_0)$.
β_r:	the strictly negative divisor energy attached to a single pure replica of index $r > 1$.

Warning 2.1 (Normalized energy versus $\mathcal{O}$). The logarithmic-over-pole cancellation is a statement about $\mathcal{I} = 2\pi\mathcal{O}/\zeta$. At an actual zero $s = \rho$, the factor $\zeta(s)$ lowers the singular order of $\mathcal{O}$. Thus a formula containing $\log(s-\rho)/(s-\rho)$ must not be attributed to $\mathcal{O}$ itself at $r = 1$.

3. Exact canonical kinematics

3.1. **Hilbert space, position, and momentum.**

Definition 3.1 (Hilbert space)**.** Fix $\hbar > 0$ and set

$$\mathcal{H} = L^2(\mathbb{R}, dq) \tag{3.1}$$

with its standard complex Hilbert-space inner product.

Definition 3.2 (Position)**.** The position operator Q is

$$(Q\psi)(q) = q\psi(q) \tag{3.2}$$

with maximal domain

$$\mathcal{D}(Q) = \{\psi \in L^2(\mathbb{R}) : q\psi(q) \in L^2(\mathbb{R})\}. \tag{3.3}$$

Definition 3.3 (Momentum)**.** The momentum operator P is

$$(P\psi)(q) = -i\hbar\,\psi'(q) \tag{3.4}$$

with domain

$$\mathcal{D}(P) = H^1(\mathbb{R}). \tag{3.5}$$

Proposition 3.4 (Self-adjointness and common core)**.** *The operators Q and P are self-adjoint. The Schwartz space $\mathcal{S}(\mathbb{R})$ is a common invariant core for both.*

Proof. Q is the maximal multiplication operator by the real function q. Under the unitary Fourier transform with convention

$$(\mathcal{F}\psi)(p) = \frac{1}{\sqrt{2\pi\hbar}} \int_{\mathbb{R}} e^{-ipq/\hbar}\psi(q)\,dq, \tag{3.6}$$

P is unitarily equivalent to multiplication by the real variable p. The core statement follows by standard cutoff and mollification arguments. See, for example, [7]. □

Proposition 3.5 (Exact canonical commutation relation)**.** *For every $\psi \in \mathcal{S}(\mathbb{R})$,*

$$[Q, P]\psi = i\hbar\psi. \tag{3.7}$$

Proof. Direct differentiation gives

$$(QP\psi)(q) = -i\hbar q\psi'(q), \tag{3.8}$$

$$(PQ\psi)(q) = -i\hbar\psi(q) - i\hbar q\psi'(q). \tag{3.9}$$

Subtracting (3.9) from (3.8) proves (3.7). □

3.2. The Heisenberg uncertainty principle and exact incompatibility.

Definition 3.6 (Expectation and uncertainty)**.** Let A be self-adjoint and let $\psi \in \mathcal{D}(A)$ be normalized. Define

$$\langle A\rangle_\psi = \langle \psi, A\psi\rangle, \tag{3.10}$$

$$(\Delta_\psi A)^2 = \|(A - \langle A\rangle_\psi)\psi\|^2. \tag{3.11}$$

Theorem 3.7 (Heisenberg uncertainty principle)**.** *For every normalized $\psi \in \mathcal{S}(\mathbb{R})$,*

$$\boxed{\Delta_\psi Q\,\Delta_\psi P \geq \frac{\hbar}{2}.} \tag{3.12}$$

Proof. Set

$$\widetilde{Q} = Q - \langle Q\rangle_\psi I, \qquad \widetilde{P} = P - \langle P\rangle_\psi I. \tag{3.13}$$

By Cauchy–Schwarz,

$$\Delta_\psi Q\,\Delta_\psi P \geq \left|\langle \widetilde{Q}\psi, \widetilde{P}\psi\rangle\right|. \tag{3.14}$$

The modulus dominates the absolute value of the imaginary part, and on $\mathcal{S}(\mathbb{R})$

$$2i\ \mathrm{Im}\langle \widetilde{Q}\psi, \widetilde{P}\psi\rangle = \langle \psi, [Q,P]\psi\rangle \tag{3.15}$$

$$= i\hbar. \tag{3.16}$$

Combining (3.14)–(3.16) proves (3.12). □

Remark 3.8 (Meaning of the inequality). Equation (3.12) is the Heisenberg uncertainty principle used in this paper. The quantities $\Delta_\psi Q$ and $\Delta_\psi P$ are intrinsic standard deviations of a state. They are not measurement errors. Exact simultaneous localization would require both to vanish, which is prohibited by (3.12).

Theorem 3.9 (No simultaneous exact position and momentum)**.** *There is no nonzero state with simultaneous exact position and exact momentum. More precisely, both of the following are impossible for real q_0, p_0:*

(1) a nonzero vector ψ in a common invariant domain satisfying

$$Q\psi = q_0\psi, \qquad P\psi = p_0\psi; \tag{3.17}$$

(2) a nonzero tempered distribution $T \in \mathcal{S}'(\mathbb{R})$ satisfying

$$qT = q_0 T, \qquad -i\hbar T' = p_0 T. \tag{3.18}$$

Proof. For vectors, (3.17) gives $[Q,P]\psi = 0$, whereas the CCR gives $[Q,P]\psi = i\hbar\psi$; hence $\psi = 0$. For distributions define directly

$$QT = qT, \qquad PT = -i\hbar T'. \tag{3.19}$$

The distributional Leibniz rule gives

$$[Q,P]T = i\hbar T. \tag{3.20}$$

The simultaneous eigenvalue equations would make the left-hand side zero, so $T = 0$. □

4. Weyl displacements and the arithmetic phase space

4.1. **Weyl operators.**

Definition 4.1 (Weyl displacement)**.** For $(x,y) \in \mathbb{R}^2$, define

$$W(x,y) = \exp\left(\frac{i}{\hbar}(yQ - xP)\right). \tag{4.1}$$

This is a unitary operator on all of $\mathcal{H}$.

Proposition 4.2 (Coordinate action)**.** *For $\psi \in L^2(\mathbb{R})$,*

$$(W(x,y)\psi)(q) = \exp\left(\frac{i}{\hbar} y\left(q - \frac{x}{2}\right)\right)\psi(q-x) \tag{4.2}$$

almost everywhere.

Proof. Since $[iyQ/\hbar, -ixP/\hbar]$ is scalar, the Baker–Campbell–Hausdorff formula gives

$$W(x,y) = e^{-ixy/(2\hbar)} e^{iyQ/\hbar} e^{-ixP/\hbar}. \tag{4.3}$$

The last factor translates by x and the middle factor multiplies by $e^{iyq/\hbar}$, yielding (4.2). □

Proposition 4.3 (Weyl form of the CCR)**.** *Let $z = (x,y)$ and $z' = (x',y')$. Define*

$$\omega(z,z') = yx' - xy'. \tag{4.4}$$

Then

$$W(z)W(z') = \exp\left(\frac{i}{2\hbar}\omega(z,z')\right) W(z+z'). \tag{4.5}$$

Consequently,

$$[W(z), W(z')] = 2i \sin\left(\frac{\omega(z,z')}{2\hbar}\right) W(z+z'). \tag{4.6}$$

Proof. For

$$A_z = \frac{i}{\hbar}(yQ - xP), \tag{4.7}$$

the CCR gives

$$[A_z, A_{z'}] = \frac{i}{\hbar}\omega(z,z')I. \tag{4.8}$$

This commutator is central, so the Baker–Campbell–Hausdorff series stops after the commutator term and gives (4.5). Reversing z and z' changes the sign of ω, giving (4.6). The Weyl relations are standard; see [6]. □

Proposition 4.4 (Position and momentum displacement equations)**.** *On* $\mathcal{S}(\mathbb{R})$*,*

$$[Q, W(x,y)] = xW(x,y), \tag{4.9}$$

$$[P, W(x,y)] = yW(x,y). \tag{4.10}$$

Proof. Both identities follow by differentiating (4.2), or directly from (3.7) because the relevant commutators with the exponent are scalar. □

4.2. **Arithmetic embedding.**

Definition 4.5 (The additive prime-factor counter)**.** For

$$n = \prod_p p^{v_p(n)}, \tag{4.11}$$

define

$$\Omega(n) = \sum_p v_p(n). \tag{4.12}$$

Then

$$\Omega(mn) = \Omega(m) + \Omega(n). \tag{4.13}$$

Definition 4.6 (Arithmetic phase-space map)**.** Fix two positive scales $\ell > 0$ and $\kappa > 0$. For $n \in \mathbb{N}$, put

$$x_n = \ell \log n, \tag{4.14}$$

$$y_n = \kappa\Omega(n), \tag{4.15}$$

$$z_n = (x_n, y_n). \tag{4.16}$$

Define the dimensionless Heisenberg area parameter

$$\vartheta = \frac{\ell\kappa}{\hbar}. \tag{4.17}$$

Remark 4.7 (Why these two coordinates). The first coordinate is forced by multiplicative size because $\log(mn) = \log m + \log n$. The second coordinate is chosen as the simplest integer-valued completely additive function that directly controls the Möbius sign on square-free integers:

$$\mu(n) \neq 0 \quad \Longrightarrow \quad \mu(n) = (-1)^{\Omega(n)}. \tag{4.18}$$

The construction can be generalized by replacing Ω with another completely additive arithmetic function, but the present notebook fixes Ω throughout.

Definition 4.8 (Arithmetic Weyl operators)**.** For $n \in \mathbb{N}$, define

$$W_n = W(x_n, y_n). \tag{4.19}$$

Theorem 4.9 (Exact Heisenberg cocycle on the integers)**.** *For $m, n \in \mathbb{N}$,*

$$W_m W_n = \chi_\vartheta(m,n) W_{mn}, \tag{4.20}$$

where

$$\chi_\vartheta(m,n) = \exp\left[\frac{i\vartheta}{2}\big(\Omega(m)\log n - \log m\,\Omega(n)\big)\right]. \tag{4.21}$$

Moreover,

$$[W_m, W_n] = 2i\sin\left[\frac{\vartheta}{2}\big(\Omega(m)\log n - \log m\,\Omega(n)\big)\right] W_{mn}. \tag{4.22}$$

Proof. Complete additivity gives

$$z_{mn} = z_m + z_n. \tag{4.23}$$

Substitution of (4.14)–(4.17) into (4.5) gives (4.20); (4.22) follows from (4.6). □

Corollary 4.10 (Prime commutator)**.** *For distinct primes p, q,*

$$[W_p, W_q] = 2i\sin\left(\frac{\vartheta}{2}\log\frac{q}{p}\right) W_{pq}. \tag{4.24}$$

Thus the prime operators commute only at the resonant values

$$\vartheta \log\frac{q}{p} \in 2\pi\mathbb{Z}. \tag{4.25}$$

Proof. For primes p, q, one has $\Omega(p) = \Omega(q) = 1$. Substituting these values into the integer Weyl commutator formula of Theorem 4.9 gives (4.24). The commutator vanishes exactly when its sine factor vanishes, which is equivalent to (4.25). □

Remark 4.11 (Where Heisenberg enters nontrivially). Equation (4.20) is the first point where Version 2 ceases to be ordinary Dirichlet algebra in operator notation. If the canonical pair were commuting, the Weyl cocycle would be 1 and the arithmetic product below would reduce to ordinary Dirichlet convolution. For nonzero ϑ, the CCR creates order-dependent phases between different arithmetic factors.

Theorem 4.12 (The arithmetic cocycle is not gauge-trivial)**.** *Extend* log *and* Ω *additively to* $\mathbb{Q}_{>0}^\times$*. For every real $\vartheta \neq 0$, the cocycle χ_ϑ is not a scalar coboundary on $\mathbb{N}^\times$, and its extension is not a scalar coboundary on $\mathbb{Q}_{>0}^\times$.*

Proof. On a commutative semigroup, every scalar coboundary

$$(\delta a)(m,n) = a(m)a(n)a(mn)^{-1} \tag{4.26}$$

is symmetric in m, n. Hence a necessary condition for χ_ϑ to be a coboundary is

$$\frac{\chi_\vartheta(m,n)}{\chi_\vartheta(n,m)} = 1 \tag{4.27}$$

for all m, n. For distinct primes,

$$\frac{\chi_\vartheta(p,q)}{\chi_\vartheta(q,p)} = \exp\left(i\vartheta\log\frac{q}{p}\right). \tag{4.28}$$

If this were one simultaneously for $(p,q) = (2,3)$ and $(2,5)$, then there would be nonzero integers k, l such that

$$\vartheta\log(3/2) = 2\pi k, \qquad \vartheta\log(5/2) = 2\pi l. \tag{4.29}$$

Thus $\log(3/2)/\log(5/2) = k/l \in \mathbb{Q}$. Exponentiating a reduced rational relation would produce an equality between incompatible prime factorizations, contradicting unique factorization. Therefore (4.27) cannot hold for all primes. □

5. Structural filter: linear Möbius states cannot escape second order

Let $v_n \in \mathcal{H}$ be arbitrary vectors and define

$$\Psi_x = \sum_{n \le x} \mu(n) v_n. \tag{5.1}$$

Theorem 5.1 (Quadraticity of unnormalized sesquilinear moments)**.** *For every linear operator A for which the expression is defined,*

$$\langle \Psi_x, A\Psi_x \rangle = \sum_{m,n \le x} \mu(m)\mu(n) \langle v_m, A v_n \rangle. \tag{5.2}$$

Thus every unnormalized sesquilinear moment, including $\|\Psi_x\|^2$, is quadratic in the Möbius coefficients. Normalized expectations are quotients of such quadratic forms, and variances or Robertson expressions are rational combinations of them; they are not, in general, themselves quadratic forms.

Proof. Expand the inner product sesquilinearly and use that $\mu(n)$ is real. The assertions about normalized quantities follow directly by dividing by $\langle \Psi_x, \Psi_x \rangle$. □

Remark 5.2 (Scope of the linear-state filter)*.* Equation (5.2) is a structural warning, not a no-go theorem for every nonlinear observable constructed from a linear Möbius state. What is ruled out is only the claim that an unnormalized sesquilinear moment escapes second order in the coefficients.

6. Arithmetic uncertainty generated by the CCR

Definition 6.1 (Integer generator)**.** For $n \ge 1$, define on $\mathcal{S}(\mathbb{R})$ the symmetric operator

$$G_n^{(0)} = y_n Q - x_n P = \kappa \Omega(n) Q - \ell \log n \, P. \tag{6.1}$$

It is essentially self-adjoint on $\mathcal{S}(\mathbb{R})$; denote its self-adjoint closure by G_n. The corresponding unitary displacement is

$$W_n = \exp(i G_n/\hbar). \tag{6.2}$$

Proposition 6.2 (Arithmetic commutator)**.** *For all $m, n \ge 1$, the restrictions of G_m and G_n to $\mathcal{S}(\mathbb{R})$ satisfy*

$$[G_m, G_n] = i\hbar(x_m y_n - y_m x_n) I \tag{6.3}$$

$$= i\hbar\ell\kappa\big(\log m\, \Omega(n) - \Omega(m) \log n\big) I. \tag{6.4}$$

Proof. Expand the commutator and use $[Q, P] = i\hbar I$. Terms involving $[Q, Q]$ and $[P, P]$ vanish. □

Theorem 6.3 (Arithmetic Heisenberg uncertainty law)**.** *For every normalized $\psi \in \mathcal{S}(\mathbb{R})$ and all $m, n \ge 1$,*

$$\boxed{\Delta_\psi G_m \, \Delta_\psi G_n \ge \frac{\hbar\ell\kappa}{2} \left| \Omega(m) \log n - \log m \, \Omega(n) \right|.} \tag{6.5}$$

For distinct primes p, q,

$$\boxed{\Delta_\psi G_p \, \Delta_\psi G_q \ge \frac{\hbar\ell\kappa}{2} \left| \log \frac{q}{p} \right|.} \tag{6.6}$$

For two powers of the same prime the lower bound is zero.

Proof. Apply the Robertson inequality to the self-adjoint pair G_m, G_n:

$$\Delta_\psi G_m \, \Delta_\psi G_n \geq \frac{1}{2} \left| \langle \psi, [G_m, G_n]\psi \rangle \right|. \tag{6.7}$$

Substitution of (6.4) gives (6.5). If p, q are distinct primes then $\Omega(p) = \Omega(q) = 1$, which gives (6.6). If $m = p^a, n = p^b$, the determinant $\Omega(m)\log n - \log m\,\Omega(n)$ vanishes. □

Remark 6.4 (The common invariant). Define the arithmetic symplectic area

$$\mathcal{A}(m,n) = \Omega(m)\log n - \log m\,\Omega(n). \tag{6.8}$$

The uncertainty lower bound is proportional to $|\mathcal{A}(m,n)|$, while the twisted product below carries the phase $\exp(i\vartheta\mathcal{A}(m,n)/2)$. Thus uncertainty and twisted Möbius inversion are controlled by the same arithmetic invariant, although they are distinct consequences of the CCR.

6.1. **The semiclassical and arithmetic-commutative limits are distinct.** The parameter controlling the Weyl cocycle is

$$\vartheta(\hbar, \ell, \kappa) = \frac{\ell\kappa}{\hbar}. \tag{6.9}$$

The uncertainty scale and the deformation scale therefore depend on the same three parameters in different ways.

Proposition 6.5 (Opposite fixed-scale limits)**.** *Fix $\ell > 0$ and $\kappa > 0$. For every fixed pair m, n,*

$$\frac{\hbar\ell\kappa}{2}|\mathcal{A}(m,n)| \longrightarrow 0 \qquad (\hbar \to 0^+), \tag{6.10}$$

whereas

$$|\vartheta| = \frac{\ell\kappa}{\hbar} \longrightarrow \infty. \tag{6.11}$$

Consequently the ordinary semiclassical limit $\hbar \to 0^+$ with ℓ, κ fixed does not approach the untwisted Dirichlet product.

Proof. Equation (6.10) follows directly from Theorem 6.3, while (6.11) follows from (6.9). □

Proposition 6.6 (Joint semiclassical–commutative scaling)**.** *Let $\ell = \ell(\hbar) > 0$ and $\kappa = \kappa(\hbar) > 0$. Then*

$$\vartheta(\hbar) \longrightarrow 0 \qquad (\hbar \to 0^+) \tag{6.12}$$

if and only if

$$\ell(\hbar)\kappa(\hbar) = o(\hbar). \tag{6.13}$$

Under this scaling the twisted product converges coefficientwise to the ordinary Dirichlet product, and the arithmetic uncertainty lower bound obeys

$$\frac{\hbar\ell(\hbar)\kappa(\hbar)}{2}|\mathcal{A}(m,n)| = o(\hbar^2) \tag{6.14}$$

for every fixed m, n.

Proof. The equivalence of (6.12) and (6.13) is immediate from (6.9). For fixed m, n, $\chi_\vartheta(m,n) \to 1$ as $\vartheta \to 0$, so the finite divisor sum defining $f \star_\vartheta g$ converges to ordinary Dirichlet convolution coefficientwise. Multiplying $\ell(\hbar)\kappa(\hbar) = o(\hbar)$ by $\hbar/2$ gives (6.14). □

Remark 6.7 (Two independent notions of classicality). The limit $\hbar \to 0^+$ suppresses the Robertson–Heisenberg lower bound, whereas $\vartheta \to 0$ removes the arithmetic twist. They are not the same operation. With fixed ℓ, κ they move in opposite directions; simultaneous suppression requires the joint scaling (6.13).

7. The Heisenberg-twisted Dirichlet algebra

7.1. Definition and associativity.

Definition 7.1 (Heisenberg-twisted Dirichlet convolution)**.** For arithmetic functions $f, g : \mathbb{N} \to \mathbb{C}$, define

$$(f \star_\vartheta g)(k) = \sum_{d|k} f(d)g(k/d)\chi_\vartheta(d, k/d). \tag{7.1}$$

The sum is finite for every k.

Lemma 7.2 (Cocycle identity)**.** *For all $a, b, c \in \mathbb{N}$,*

$$\chi_\vartheta(a, b)\chi_\vartheta(ab, c) = \chi_\vartheta(b, c)\chi_\vartheta(a, bc). \tag{7.2}$$

Proof. The exponent in (4.21) is the alternating bilinear form obtained from the additive map $n \mapsto z_n$. Expanding both sides using $\log(ab) = \log a + \log b$ and $\Omega(ab) = \Omega(a) + \Omega(b)$ gives the same exponent. □

Proposition 7.3 (Associative noncommutative arithmetic algebra)**.** *The product $\star_\vartheta$ is associative and has identity δ_1, where*

$$\delta_1(1) = 1, \qquad \delta_1(n) = 0 \quad (n > 1). \tag{7.3}$$

For $\vartheta \neq 0$ it is generally noncommutative.

Proof. Associativity follows from Lemma 7.2 after grouping the finite divisor sums by triples $abc = k$. The identity property follows from $\chi_\vartheta(1, n) = \chi_\vartheta(n, 1) = 1$. Noncommutativity follows from Corollary 4.10; equivalently,

$$(\delta_p \star_\vartheta \delta_q)(pq) = \chi_\vartheta(p, q) \tag{7.4}$$

whereas

$$(\delta_q \star_\vartheta \delta_p)(pq) = \chi_\vartheta(q, p) = \overline{\chi_\vartheta(p, q)}. \tag{7.5}$$

□

7.2. Weighted Banach algebra and operator representation.

Definition 7.4 (Weighted arithmetic Banach space)**.** For $\sigma > 0$, set

$$\mathcal{A}_\sigma = \left\{ f : \mathbb{N} \to \mathbb{C} : \|f\|_\sigma := \sum_{n\geq 1} \frac{|f(n)|}{n^\sigma} < \infty \right\}. \tag{7.6}$$

Proposition 7.5 (Banach-algebra estimate)**.** *For $f, g \in \mathcal{A}_\sigma$,*

$$\|f \star_\vartheta g\|_\sigma \leq \|f\|_\sigma \|g\|_\sigma. \tag{7.7}$$

Hence $(\mathcal{A}_\sigma, \star_\vartheta)$ is a unital Banach algebra.

Proof. Since $|\chi_\vartheta| = 1$,

$$\|f \star_\vartheta g\|_\sigma \leq \sum_{k\geq 1} \frac{1}{k^\sigma} \sum_{d|k} |f(d)||g(k/d)| \tag{7.8}$$

$$= \sum_{d,e\geq 1} \frac{|f(d)|}{d^\sigma} \frac{|g(e)|}{e^\sigma} \tag{7.9}$$

$$= \|f\|_\sigma \|g\|_\sigma. \tag{7.10}$$

Completeness is the completeness of weighted ℓ^1. □

Definition 7.6 (Weyl–Dirichlet representation)**.** For $f \in \mathcal{A}_\sigma$, define

$$\Pi_{\sigma,\vartheta}(f) = \sum_{n\geq 1} \frac{f(n)}{n^\sigma} W_n, \tag{7.11}$$

where the series converges absolutely in operator norm.

Proposition 7.7 (Representation theorem)**.** *For $f, g \in \mathcal{A}_\sigma$,*

$$\|\Pi_{\sigma,\vartheta}(f)\| \leq \|f\|_\sigma, \tag{7.12}$$

$$\Pi_{\sigma,\vartheta}(f)\Pi_{\sigma,\vartheta}(g) = \Pi_{\sigma,\vartheta}(f \star_\vartheta g). \tag{7.13}$$

Proof. The norm bound follows from $\|W_n\| = 1$. Absolute norm convergence permits multiplication and regrouping of the double series. The coefficient of W_k is exactly (7.1) by Theorem 4.9. □

8. Möbius inversion deformed by the CCR

8.1. **The Möbius operator.**

Definition 8.1 (Möbius operator)**.** For $\sigma > 1$, define

$$\mathcal{M}_{\sigma,\vartheta} = \Pi_{\sigma,\vartheta}(\mu) = \sum_{n\geq 1} \frac{\mu(n)}{n^\sigma} W_n. \tag{8.1}$$

The series converges absolutely in operator norm, so $\mathcal{M}_{\sigma,\vartheta} \in \mathcal{B}(\mathcal{H})$ and

$$\mathcal{D}(\mathcal{M}_{\sigma,\vartheta}) = \mathcal{H}. \tag{8.2}$$

Remark 8.2 (Concrete coordinate action)*.* For $\psi \in L^2(\mathbb{R})$,

$$(\mathcal{M}_{\sigma,\vartheta}\psi)(q) = \sum_{n\geq 1} \frac{\mu(n)}{n^\sigma} \exp\left[\frac{i}{\hbar} y_n \left(q - \frac{x_n}{2}\right)\right] \psi(q - x_n), \tag{8.3}$$

with convergence in L^2. Thus the operator is a Möbius-weighted superposition of simultaneous translations in position and momentum; it is not a Fourier multiplier when $\kappa \neq 0$.

8.2. **Formal twisted inverse.**

Proposition 8.3 (Existence and uniqueness of arithmetic inverses)**.** *Let $f : \mathbb{N} \to \mathbb{C}$ satisfy $f(1) \neq 0$. Then there exists a unique arithmetic function g such that*

$$f \star_\vartheta g = g \star_\vartheta f = \delta_1. \tag{8.4}$$

Its coefficients are determined recursively from $g(1) = f(1)^{-1}$.

Proof. For $n > 1$, the equation $(f \star_\vartheta r)(n) = 0$ contains the term $f(1)r(n)$ and otherwise only values $r(d)$ with $d < n$. It therefore constructs a unique right inverse r satisfying $f \star_\vartheta r = \delta_1$. Separately, the equations $(\lambda \star_\vartheta f)(n) = 0$ construct a unique left inverse λ satisfying $\lambda \star_\vartheta f = \delta_1$. Associativity then gives

$$\lambda = \lambda \star_\vartheta (f \star_\vartheta r) = (\lambda \star_\vartheta f) \star_\vartheta r = r, \tag{8.5}$$

so the inverse is bilateral and unique. □

Definition 8.4 (Heisenberg-zeta coefficients)**.** Let $z_\vartheta : \mathbb{N} \to \mathbb{C}$ be the unique two-sided inverse of μ for $\star_\vartheta$:

$$\mu \star_\vartheta z_\vartheta = z_\vartheta \star_\vartheta \mu = \delta_1. \tag{8.6}$$

Thus

$$z_\vartheta(1) = 1, \tag{8.7}$$

and for $n > 1$,

$$z_\vartheta(n) = -\sum_{\substack{d|n \\ d>1}} \mu(d) z_\vartheta(n/d) \chi_\vartheta(d, n/d). \tag{8.8}$$

Proposition 8.5 (Commuting boundary)**.** *At $\vartheta = 0$,*

$$z_0(n) = 1 \tag{8.9}$$

for every $n \geq 1$.

Proof. When $\vartheta = 0$, $\chi_0 \equiv 1$ and $\star_\vartheta$ is ordinary Dirichlet convolution. Equation (8.6) becomes $\mu * z_0 = \delta_1$. Classical Möbius inversion gives $z_0 = \mathbf{1}$. See [10]. □

Proposition 8.6 (Ordered-factorization formula)**.** *For $n > 1$, the twisted inverse has the finite coefficient expansion*

$$\begin{aligned} z_\vartheta(n) = \sum_{r\geq 1} (-1)^r \sum_{\substack{n_1 \cdots n_r = n \\ n_j > 1}} \left(\prod_{j=1}^{r} \mu(n_j) \right) \\ \times \exp\left[\frac{i\vartheta}{2} \sum_{1 \leq i < j \leq r} \left(\Omega(n_i) \log n_j - \log n_i \, \Omega(n_j) \right) \right]. \end{aligned} \tag{8.10}$$

Only finitely many r and ordered factorizations contribute for fixed n.

Proof. Write $\mu = \delta_1 + h$ with $h(1) = 0$. Coefficientwise the formal Neumann expansion is

$$z_\vartheta = \sum_{r \geq 0} (-h)^{\star_\vartheta r}. \tag{8.11}$$

For fixed n, a product of more than $\log_2 n$ factors exceeding one cannot equal n, so the coefficient sum is finite. Iterating the bicharacter cocycle yields the pairwise phase in (8.10). □

8.3. **A bounded inverse in a rigorous half-plane.**

Definition 8.7 (Uniform Neumann threshold)**.** Let σ_* be the unique real number > 1 satisfying

$$\frac{\zeta(\sigma_*)}{\zeta(2\sigma_*)} = 2. \tag{8.12}$$

Numerically,

$$\sigma_* = 1.578017063034\ldots. \tag{8.13}$$

Theorem 8.8 (Uniform Neumann region)**.** *For every real ϑ and every $\sigma > \sigma_*$,*

$$z_\vartheta \in \mathcal{A}_\sigma. \tag{8.14}$$

Consequently

$$\mathcal{Z}_{\sigma,\vartheta} = \Pi_{\sigma,\vartheta}(z_\vartheta) = \sum_{n \geq 1} \frac{z_\vartheta(n)}{n^\sigma} W_n \tag{8.15}$$

converges absolutely in operator norm and satisfies

$$\mathcal{M}_{\sigma,\vartheta} \mathcal{Z}_{\sigma,\vartheta} = \mathcal{Z}_{\sigma,\vartheta} \mathcal{M}_{\sigma,\vartheta} = I. \tag{8.16}$$

Proof. Write $\mu = \delta_1 + h$, where $h(1) = 0$. For $\sigma > 1$,

$$\|h\|_\sigma = \sum_{n\geq 2} \frac{\mu(n)^2}{n^\sigma} \tag{8.17}$$

$$= \frac{\zeta(\sigma)}{\zeta(2\sigma)} - 1. \tag{8.18}$$

The right-hand side is continuous and strictly decreasing from $+\infty$ to 0 as σ runs from 1^+ to $+\infty$, so σ_* is unique and $\|h\|_\sigma < 1$ exactly when $\sigma > \sigma_*$. The Banach-algebra Neumann series

$$(\delta_1 + h)^{-1} = \sum_{r=0}^{\infty} (-h)^{\star_\vartheta r} \tag{8.19}$$

then converges in $\mathcal{A}_\sigma$. Uniqueness identifies it with z_ϑ, and the representation theorem gives (8.16). □

Warning 8.9 (Exact border of the proved inverse). The Möbius operator exists as a bounded operator for every $\sigma > 1$. The argument above proves a bounded inverse only for $\sigma > \sigma_*$. At $\sigma = \sigma_*$ the Neumann norm equals one and this argument is inconclusive. No claim is made here for $1 < \sigma \leq \sigma_*$.

9. The first Heisenberg arithmetic law

Theorem 9.1 (Prime rays are rigid; distinct primes interfere)**.** *Let p, q be distinct primes and let $r \geq 1$. Then*

$$z_\vartheta(p^r) = 1, \tag{9.1}$$

$$z_\vartheta(pq) = 2\cos\left(\frac{\vartheta}{2}\log\frac{q}{p}\right) - 1, \tag{9.2}$$

$$z_\vartheta(pq) - 1 = -4\sin^2\left(\frac{\vartheta}{4}\log\frac{q}{p}\right). \tag{9.3}$$

Proof. For p^r, only the divisors 1 and p contribute to the recursion because $\mu(p^j) = 0$ for $j \geq 2$. Moreover,

$$\Omega(p)\log(p^{r-1}) - \log p\,\Omega(p^{r-1}) = 0, \tag{9.4}$$

so $\chi_\vartheta(p, p^{r-1}) = 1$. Equation (8.8) gives

$$z_\vartheta(p^r) = z_\vartheta(p^{r-1}), \tag{9.5}$$

and (9.1) follows from $z_\vartheta(1) = 1$.

For pq, the four divisors $1, p, q, pq$ give

$$0 = z_\vartheta(pq) - \chi_\vartheta(p, q) - \chi_\vartheta(q, p) + 1. \tag{9.6}$$

Since

$$\chi_\vartheta(p, q) = \exp\left(\frac{i\vartheta}{2}\log\frac{q}{p}\right), \tag{9.7}$$

and $\chi_\vartheta(q, p)$ is its complex conjugate, (9.2) follows. The identity (9.3) is the elementary formula $2(\cos u - 1) = -4\sin^2(u/2)$. □

Corollary 9.2 (Loss of ordinary multiplicativity)**.** *Generically,*

$$z_\vartheta(pq) \neq z_\vartheta(p) z_\vartheta(q) = 1. \tag{9.8}$$

Thus the Heisenberg deformation preserves every one-prime ray but destroys ordinary multiplicativity through cross-prime interference.

Proof. The prime-pair formula gives $z_\vartheta(pq) = 2\cos(\frac{\vartheta}{2}\log(q/p)) - 1$. This equals one only on the stated resonance set; away from it the product of the two one-prime values, which is identically one, differs from the two-prime coefficient. □

Corollary 9.3 (Small-area expansion)**.** *For fixed distinct primes p, q,*

$$z_\vartheta(pq) = 1 - \frac{\vartheta^2}{4}\left(\log\frac{q}{p}\right)^2 + O(\vartheta^4) \qquad (\vartheta \to 0). \tag{9.9}$$

There is no linear correction.

Proof. Use $\cos x = 1 - x^2/2 + O(x^4)$ with $x = (\vartheta/2)\log(q/p)$ in the exact prime-pair formula. Multiplying out gives (9.9); evenness of the cosine excludes an odd term. □

Proposition 9.4 (Two-prime lattice recurrence)**.** *Let $p \neq q$ be primes, set*

$$Z_{a,b} = z_\vartheta(p^a q^b), \qquad \lambda = \frac{\vartheta}{2}\log\frac{q}{p}. \tag{9.10}$$

Then for $a, b \geq 1$,

$$Z_{a,b} = e^{ib\lambda}Z_{a-1,b} + e^{-ia\lambda}Z_{a,b-1} - e^{i(b-a)\lambda}Z_{a-1,b-1}, \tag{9.11}$$

with boundary values

$$Z_{a,0} = Z_{0,b} = 1. \tag{9.12}$$

In particular,

$$z_\vartheta(p^2 q) = 1 + 2\cos(2\lambda) - 2\cos\lambda. \tag{9.13}$$

Proof. Apply the right-inverse recursion (8.8) to $p^a q^b$. Since $\mu(d) \neq 0$ only when the exponents of p, q in d are at most one, only $d = p, q, pq$ contribute. Evaluation of the three cocycle factors gives (9.11); the boundary values are Theorem 9.1 on prime rays. □

Proposition 9.5 (Evenness of the full inverse)**.** *For every $n \geq 1$,*

$$z_{-\vartheta}(n) = z_\vartheta(n). \tag{9.14}$$

Proof. Changing ϑ to $-\vartheta$ reverses the cocycle:

$$(f \star_{-\vartheta} g)(n) = (g \star_\vartheta f)(n). \tag{9.15}$$

Since z_ϑ is a two-sided inverse of μ for $\star_\vartheta$, it is a two-sided inverse of μ for the opposite product $\star_{-\vartheta}$. Uniqueness of the inverse gives (9.14). □

Proposition 9.6 (Reality for real deformation)**.** *For real ϑ and every $n \geq 1$,*

$$z_\vartheta(n) \in \mathbb{R}. \tag{9.16}$$

Proof. Complex conjugation changes χ_ϑ to $\chi_{-\vartheta}$. Since μ is real, conjugating the inverse equation shows that $\overline{z_\vartheta}$ is the inverse for the $-\vartheta$ product. Proposition 9.5 therefore gives $\overline{z_\vartheta} = z_{-\vartheta} = z_\vartheta$. □

Remark 9.7 (Why this is the non-tautological step). At $\vartheta = 0$ the inverse coefficients are identically 1. At nonzero ϑ, equation (9.3) creates a new arithmetic coefficient depending on the logarithmic separation of two primes. This dependence is forced by the symplectic area in the Weyl product and disappears if the CCR is removed. It is therefore the first result in the notebook where Heisenberg noncommutativity changes the arithmetic inverse itself rather than merely constraining an unrelated state space.

10. Curvature of the cocyclically deformed Möbius inverse

Set

$$\mathcal{A}(d,e) = \Omega(d)\log e - \log d\,\Omega(e), \tag{10.1}$$

so that

$$\chi_\vartheta(d,e) = \exp\!\left(\frac{i\vartheta}{2}\mathcal{A}(d,e)\right). \tag{10.2}$$

Proposition 10.1 (Coefficientwise entire dependence)**.** *For each fixed $n \geq 1$, the map $\vartheta \mapsto z_\vartheta(n)$ is entire. It is even, and it is real-valued on the real axis.*

Proof. Formula (8.10) is a finite sum for fixed n, each term being an exponential $e^{ic\vartheta}$. Evenness and reality follow from Propositions 9.5 and 9.6. □

Proposition 10.2 (Second derivative of the inverse)**.** *Define*

$$g(k) = \frac{1}{4}\sum_{d|k}\mu(d)\mathcal{A}(d,k/d)^2. \tag{10.3}$$

Then

$$\boxed{z_0''(n) = \sum_{k|n} g(k).} \tag{10.4}$$

Proof. Differentiate the inverse equation

$$0 = \sum_{d|n}\mu(d) z_\vartheta(n/d)\exp\!\left(\frac{i\vartheta}{2}\mathcal{A}(d,n/d)\right) \tag{10.5}$$

twice at $\vartheta = 0$. Since $z_0 \equiv 1$ and $z_0' \equiv 0$,

$$(\mu * z_0'')(n) = \frac{1}{4}\sum_{d|n}\mu(d)\mathcal{A}(d,n/d)^2 = g(n). \tag{10.6}$$

Convolving with $\mathbf{1} = \mu^{-1}$ gives (10.4). □

Lemma 10.3 (Support collapse)**.** *Let $k = \prod_{j=1}^r p_j^{a_j}$ with distinct primes p_j. Then*

$$g(k) = 0 \qquad \text{for } r = 0,\ r = 1,\ \text{or } r \geq 3. \tag{10.7}$$

For two distinct primes,

$$\boxed{g(p^\alpha q^\beta) = -\frac{\alpha\beta}{2}\log^2\frac{q}{p}.} \tag{10.8}$$

Proof. A contributing divisor is square-free and therefore corresponds to a subset $S \subseteq \{1,\dots,r\}$. Put

$$R = \Omega(k) = \sum_j a_j, \qquad L = \log k, \qquad X_S = \sum_{j\in S}\log p_j. \tag{10.9}$$

Then

$$\mathcal{A}(d,k/d) = |S|L - RX_S. \tag{10.10}$$

Its square has Boolean degree at most two, so its full alternating difference vanishes when $r \geq 3$. For $r = 1$ the area is zero. For $r = 2$, direct evaluation of the four subsets gives

$$\sum_{d|p^\alpha q^\beta}\mu(d)\mathcal{A}(d,p^\alpha q^\beta/d)^2 = -2\alpha\beta\log^2(q/p), \tag{10.11}$$

which yields (10.8). □

Theorem 10.4 (Exact deformation curvature)**.** *If*

$$n = \prod_{j=1}^{r} p_j^{a_j}, \qquad T(a) = \frac{a(a+1)}{2}, \tag{10.12}$$

then

$$\boxed{z_0''(n) = -\frac{1}{2} \sum_{1 \le i < j \le r} T(a_i) T(a_j) \log^2 \frac{p_j}{p_i}.} \tag{10.13}$$

Thus $-z_0''(n) \ge 0$, with equality exactly for $n = 1$ or a prime power.

Proof. Only divisors supported on exactly two primes survive by Lemma 10.3. Summing (10.8) over $1 \le \alpha \le a_i$, $1 \le \beta \le a_j$ gives the factor $T(a_i)T(a_j)$. □

Corollary 10.5 (Square-free prime-log variance)**.** *If $n = p_1 \cdots p_r$ is square-free, then*

$$-z_0''(n) = \frac{1}{2} \sum_{i<j} (\log p_j - \log p_i)^2 \tag{10.14}$$

$$= \frac{1}{2} \left[r \sum_{p \mid n} (\log p)^2 - (\log n)^2 \right] \tag{10.15}$$

$$= \frac{r^2}{2} \operatorname{Var}_{p \mid n}(\log p). \tag{10.16}$$

Proof. In Theorem 10.4, square-freeness gives $T(v_p(n)) = T(1) = 1$ for every $p \mid n$. The first equality follows. Expanding the pairwise squares and counting each $(\log p)^2$ exactly $r - 1$ times gives $\sum_{i<j}(a_j - a_i)^2 = r \sum_i a_i^2 - (\sum_i a_i)^2$, which yields the second equality. Dividing by r identifies the usual empirical variance and gives the third. □

Corollary 10.6 (Infinitesimal symplectic size)**.** *For square-free $n = p_1 \cdots p_r$, let*

$$B_{ij} = \log p_j - \log p_i. \tag{10.17}$$

Then B has rank at most two. If $r \ge 2$, its two nonzero singular values are both

$$s(n) = \sqrt{r \sum_{p \mid n} (\log p)^2 - (\log n)^2}, \tag{10.18}$$

and

$$\boxed{-z_0''(n) = \frac{1}{2} s(n)^2.} \tag{10.19}$$

Proof. Write $B = \mathbf{1}a^T - a\mathbf{1}^T$, $a_i = \log p_i$. The two nonzero singular values of $uv^T - vu^T$ equal

$$\sqrt{\|u\|^2 \|v\|^2 - (u \cdot v)^2}. \tag{10.20}$$

Apply this with $u = \mathbf{1}$, $v = a$, and then use Corollary 10.5. □

Corollary 10.7 (Exact Taylor law for each integer)**.** *For every fixed n,*

$$z_\vartheta(n) = 1 + \frac{\vartheta^2}{2} z_0''(n) + O_n(\vartheta^4), \tag{10.21}$$

and the Taylor series has infinite radius of convergence. For square-free n,

$$\boxed{z_\vartheta(n) = 1 - \frac{\vartheta^2}{4} \left[\omega(n) \sum_{p \mid n} (\log p)^2 - (\log n)^2 \right] + O_n(\vartheta^4).} \tag{10.22}$$

Proof. Coefficientwise entire dependence was proved in Proposition 10.1. Evenness gives the Taylor expansion with no odd powers. The coefficient of ϑ^2 is $z_0''(n)/2$, proving the first formula. Substitution of the square-free formula from Corollary 10.5 yields the boxed identity. □

11. Multiplicative geometry of the curvature

Define

$$\mathcal{C}(n) := -z_0''(n) = \frac{1}{2}\sum_{p<q} T(v_p(n))T(v_q(n))\log^2\frac{q}{p}. \tag{11.1}$$

This section studies $\mathcal{C}$ before any summation over n.

Proposition 11.1 (Prime-ray rigidity)**.**

$$\boxed{\mathcal{C}(n) = 0 \iff n = 1 \text{ or } n = p^a.} \tag{11.2}$$

Proof. Every summand in (11.1) is nonnegative. It is strictly positive exactly when two distinct primes occur with positive valuation. □

Theorem 11.2 (Multiplicative superadditivity)**.** *For all positive integers m, n,*

$$\boxed{\mathcal{C}(mn) \geq \mathcal{C}(m) + \mathcal{C}(n).} \tag{11.3}$$

If $m, n > 1$, equality holds exactly when $m = p^a$ and $n = p^b$ for the same prime p.

Proof. Put $a_p = v_p(m)$, $b_p = v_p(n)$, $A_p = T(a_p)$, $B_p = T(b_p)$. Since $T(a+b) = T(a) + T(b) + ab$, subtraction of $\mathcal{C}(m) + \mathcal{C}(n)$ from $\mathcal{C}(mn)$ gives

$$\mathcal{C}(mn) - \mathcal{C}(m) - \mathcal{C}(n) = \frac{1}{2}\sum_{p<q} \Xi_{p,q}(m,n)\log^2\frac{q}{p}, \tag{11.4}$$

where

$$\Xi_{p,q}(m,n) = A_pB_q + B_pA_q + (A_p + B_p)a_qb_q + (A_q + B_q)a_pb_p + a_pb_pa_qb_q. \tag{11.5}$$

All terms are nonnegative. If $m, n > 1$, all cross terms vanish exactly when both supports consist of the same single prime. □

Corollary 11.3 (Monotonicity on the divisibility poset)**.** *If $m \mid n$, then*

$$\boxed{\mathcal{C}(m) \leq \mathcal{C}(n).} \tag{11.6}$$

Proof. Write $n = mq$. Theorem 11.2 gives $\mathcal{C}(n) \geq \mathcal{C}(m) + \mathcal{C}(q) \geq \mathcal{C}(m)$, because $\mathcal{C}(q) \geq 0$. This is exactly (11.6). □

Proposition 11.4 (Exact power law)**.** *If $n = \prod_p p^{a_p}$, then for every integer $k \geq 1$,*

$$\boxed{\mathcal{C}(n^k) = \frac{1}{8}\sum_{p<q}\Big[k^4a_p^2a_q^2 + k^3(a_p^2a_q + a_pa_q^2) + k^2a_pa_q\Big]\log^2\frac{q}{p}.} \tag{11.7}$$

Hence, if n contains at least two distinct primes,

$$\boxed{\lim_{k\to\infty}\frac{\mathcal{C}(n^k)}{k^4} = \frac{1}{8}\sum_{p<q} a_p^2a_q^2\log^2\frac{q}{p} > 0.} \tag{11.8}$$

Proof. For every prime exponent a_p,

$$T(ka_p) = \frac{k^2 a_p^2 + k a_p}{2}. \tag{11.9}$$

Thus for each pair $p < q$,

$$\frac{1}{2} T(ka_p) T(ka_q) = \frac{1}{8}\big[k^4 a_p^2 a_q^2 + k^3(a_p^2 a_q + a_p a_q^2) + k^2 a_p a_q\big]. \tag{11.10}$$

Substitution into (11.1) gives (11.7). Division by k^4 and termwise passage to the limit is legitimate because the sum contains only finitely many prime pairs. If the support has at least two primes, at least one coefficient $a_p^2 a_q^2 \log^2(q/p)$ is strictly positive, proving (11.8). □

Proposition 11.5 (Coprime block decomposition)**.** *Assume* $(m, n) = 1$. *Define*

$$W(m) = \sum_{p|m} T(v_p(m)), \tag{11.11}$$

$$\overline{L}(m) = \frac{1}{W(m)} \sum_{p|m} T(v_p(m)) \log p, \tag{11.12}$$

$$V(m) = \frac{1}{W(m)} \sum_{p|m} T(v_p(m))(\log p - \overline{L}(m))^2, \tag{11.13}$$

with the analogous definitions for n. *Then*

$$\boxed{\mathcal{C}(mn) = \mathcal{C}(m) + \mathcal{C}(n) + \frac{1}{2} W(m) W(n) \left[V(m) + V(n) + (\overline{L}(m) - \overline{L}(n))^2 \right].} \tag{11.14}$$

Proof. The cross-prime contribution is $\frac{1}{2} \sum_{p|m, q|n} A_p B_q (\log q - \log p)^2$. Expanding around the two weighted means gives (11.14). Abstractly this last algebraic identity is the standard weighted within/between variance decomposition; its use here is only to expose the multiplicative geometry of $\mathcal{C}$. □

12. Dirichlet summation of the curvature

Define

$$c(n) = -z_0''(n) \geq 0. \tag{12.1}$$

Theorem 12.1 (Exact curvature Dirichlet series)**.** *For* $\Re s > 1$,

$$\mathcal{K}(s) = \sum_{n \geq 1} \frac{c(n)}{n^s} \tag{12.2}$$

converges absolutely and

$$\boxed{\mathcal{K}(s) = \frac{\zeta(s)}{2} \left[S_0(s) S_2(s) - S_1(s)^2 \right],} \tag{12.3}$$

where

$$S_j(s) = \sum_p \frac{p^{-s}}{(1 - p^{-s})^2} (\log p)^j, \qquad j = 0, 1, 2. \tag{12.4}$$

Equivalently,

$$S_0 S_2 - S_1^2 = \sum_{p<q} \frac{p^{-s} q^{-s}}{(1 - p^{-s})^2 (1 - q^{-s})^2} \log^2 \frac{q}{p}. \tag{12.5}$$

Proof. Use Theorem 10.4 and

$$\sum_{a\geq 1} T(a)x^a = \frac{x}{(1-x)^3}. \tag{12.6}$$

Factoring the unrestricted Euler factors gives $\zeta(s)$, while a selected prime contributes

$$\frac{p^{-s}}{(1-p^{-s})^2}. \tag{12.7}$$

The pair sum is exactly the weighted variance determinant $S_0S_2 - S_1^2$. Absolute convergence follows for $\Re s > 1$. □

Corollary 12.2 (Prime-zeta reduction)**.** *Let $P(s) = \sum_p p^{-s}$ for $\Re s > 1$. Then*

$$S_0(s) = \sum_{m\geq 1} mP(ms), \tag{12.8}$$

$$S_1(s) = -\sum_{m\geq 1} mP'(ms), \tag{12.9}$$

$$S_2(s) = \sum_{m\geq 1} mP''(ms). \tag{12.10}$$

Using

$$P(s) = \sum_{k\geq 1} \frac{\mu(k)}{k} \log \zeta(ks), \tag{12.11}$$

gives

$$S_0(s) = \sum_{r\geq 1} \frac{J_2(r)}{r} \log \zeta(rs), \tag{12.12}$$

$$S_1(s) = -\sum_{r\geq 1} \varphi(r) \frac{\zeta'(rs)}{\zeta(rs)}, \tag{12.13}$$

$$S_2(s) = \left(\frac{\zeta'}{\zeta}\right)'(s), \tag{12.14}$$

where $J_2(r) = r^2 \prod_{p\mid r}(1-p^{-2})$.

Proof. Expand $x/(1-x)^2 = \sum_{m\geq 1} mx^m$, substitute (12.11), and group by $r = mk$. The needed divisor identities are

$$r\sum_{k\mid r} \frac{\mu(k)}{k^2} = \frac{J_2(r)}{r}, \tag{12.15}$$

$$r\sum_{k\mid r} \frac{\mu(k)}{k} = \varphi(r), \tag{12.16}$$

$$r\sum_{k\mid r} \mu(k) = \begin{cases} 1, & r = 1, \\ 0, & r > 1. \end{cases} \tag{12.17}$$

The last identity collapses S_2 to (12.14). □

Proposition 12.3 (Singularity at $s = 1$)**.** *As $s \to 1^+$,*

$$\boxed{\mathcal{K}(s) \sim \frac{1}{2(s-1)^3} \log \frac{1}{s-1}.} \tag{12.18}$$

Proof. From (12.11),

$$P(s) = \log \frac{1}{s-1} + O(1). \tag{12.19}$$

Hence

$$S_0(s) = \log \frac{1}{s-1} + O(1), \quad S_1(s) = \frac{1}{s-1} + O(1), \quad S_2(s) = \frac{1}{(s-1)^2} + O(1). \tag{12.20}$$

Combine with $\zeta(s) \sim (s-1)^{-1}$. □

13. Continuation of the curvature and detection of zeta zeros

Theorem 13.1 (Local continuation to $\Re s > 1/2$)**.** *Let $U \subset \{\Re s > 1/2\}$ be simply connected and contain no zero or pole of $\zeta(s)$. Choose a branch of $\log \zeta(s)$ on U. Then* (12.12)–(12.14)*, substituted into* (12.3)*, continue $\mathcal{K}$ holomorphically to U.*

Proof. For $r \geq 2$, $\Re(rs) > 1$, so the corresponding terms are holomorphic and converge normally on compact subsets. The $r = 1$ terms are holomorphic by the hypothesis on U. □

Theorem 13.2 (Zeta zeros are logarithmic curvature singularities)**.** *Let ρ be a zero of multiplicity $m \geq 1$ with $1/2 < \Re\rho < 1$, and write*

$$\zeta(s) = c_\rho (s-\rho)^m (1 + O(s-\rho)), \qquad c_\rho \neq 0. \tag{13.1}$$

Then

$$\boxed{\mathcal{K}(s) = -\frac{c_\rho m^2}{2}(s-\rho)^{m-2}\log(s-\rho) + O\Big((s-\rho)^{m-2}\Big).} \tag{13.2}$$

For a simple zero,

$$\boxed{\mathcal{K}(s) = -\frac{\zeta'(\rho)}{2}\frac{\log(s-\rho)}{s-\rho} + O\left(\frac{1}{s-\rho}\right).} \tag{13.3}$$

Proof. For $r \geq 2$, the terms in (12.12)–(12.13) are holomorphic near ρ. With $t = s - \rho$, the singular pieces are

$$S_0 = m \log t + O(1), \qquad S_1 = -\frac{m}{t} + O(1), \qquad S_2 = -\frac{m}{t^2} + O(1). \tag{13.4}$$

Thus

$$S_0 S_2 - S_1^2 = -\frac{m^2 \log t}{t^2} + O(t^{-2}). \tag{13.5}$$

Multiply by $\zeta(s)/2$. □

Corollary 13.3 (Curvature criterion equivalent to RH)**.** *RH is equivalent to the absence of logarithmic branch singularities of $\mathcal{K}$ in*

$$\boxed{\frac{1}{2} < \Re s < 1.} \tag{13.6}$$

Proof. Theorem 13.2 detects every zero in the right half of the critical strip, while Theorem 13.1 gives holomorphy on zero-free neighbourhoods. The functional equation makes the absence of zeros in this half-strip equivalent to RH. □

14. Where the logarithmic branching comes from

The logarithmic singularity in Theorem 13.2 does not come from a meromorphic function such as ζ, ζ', or ζ''. It comes from the prime-zeta component of S_0.

Proposition 14.1 (Exact source of the logarithm)**.** *For $\Re s > 1$,*

$$S_0(s) = \sum_{m\geq 1} mP(ms), \qquad P(s) = \sum_{r\geq 1} \frac{\mu(r)}{r} \log\zeta(rs). \tag{14.1}$$

Consequently

$$S_0(s) = \sum_{k\geq 1} \frac{J_2(k)}{k} \log\zeta(ks). \tag{14.2}$$

If ρ is a zero of ζ of multiplicity m and $1/2 < \Re\rho < 1$, then in a punctured simply connected neighbourhood of ρ,

$$S_0(s) = m\log(s-\rho) + O(1). \tag{14.3}$$

Proof. The first identity follows by expanding $p^{-s}(1-p^{-s})^{-2} = \sum_{m\geq 1} mp^{-ms}$ and summing over primes. The second follows from Möbius inversion for the prime zeta function and collecting the coefficient of $\log\zeta(ks)$, exactly as in Corollary 12.2. If $1/2 < \Re\rho < 1$, every term with $k \geq 2$ is holomorphic near ρ, because $\Re(k\rho) > 1$. The $k = 1$ term is $\log\zeta(s) = m\log(s-\rho) + O(1)$. □

Remark 14.2 (Global branch structure). For the classical prime-zeta function, only square-free dilation indices occur directly because of the factor $\mu(r)$. By contrast, S_0 carries the nonzero weight $J_2(r)/r$ for every r. The classical prime-zeta continuation generated by

$$P(s) = \sum_{r\geq 1} \frac{\mu(r)}{r} \log\zeta(rs) \tag{14.4}$$

has logarithmic branch points at scaled poles and zeros, including points $1/r$ and ρ/r, and the line $\Re s = 0$ is a natural boundary for P; see Fröberg [8]. This global phenomenon does not invalidate Corollary 13.3: in the open half-strip $1/2 < \Re s < 1$, scaled points ρ/r with $r \geq 2$ cannot occur, because $\Re(\rho/r) \leq 1/2$. Hence the only logarithmic branches there are the unscaled $r = 1$ branches attached to zeros of ζ itself.

15. Local monodromy and the Jordan-totient weight

The branch structure is used here only locally. No globally reordered double sum over zeta zeros is asserted.

Proposition 15.1 (Local monodromy at an isolated scaled zero)**.** *Let ρ be a zero of ζ of multiplicity m_ρ, let $r \geq 1$, and set $s_0 = \rho/r$. Suppose a sufficiently small punctured neighbourhood of s_0 contains no other branch point contributing to* (14.2)*. Analytic continuation once positively around s_0 changes S_0 by*

$$\boxed{\Delta_{\rho/r} S_0 = 2\pi i\, m_\rho \frac{J_2(r)}{r}.} \tag{15.1}$$

Since S_1 and S_2 are meromorphic and hence single-valued, the corresponding monodromy of $\mathcal{K}$ is

$$\boxed{\Delta_{\rho/r}\mathcal{K}(s) = \pi i\, m_\rho \frac{J_2(r)}{r}\, \zeta(s) \left(\frac{\zeta'}{\zeta}\right)'(s).} \tag{15.2}$$

If several scaled singularities collide at the same point, their local monodromy contributions are added before applying this formula.

Proof. In (14.2), the branch of $\log\zeta(rs)$ changes by $2\pi i m_\rho$ around $s_0 = \rho/r$. Multiplication by its coefficient $J_2(r)/r$ gives (15.1). The functions S_1 and S_2 are built from logarithmic derivatives of ζ, so they are meromorphic and carry no logarithmic monodromy. Substitution in $\mathcal{K} = \zeta(S_0S_2 - S_1^2)/2$ gives (15.2). □

Proposition 15.2 (Growth of the local branch weight)**.** *For every $r \geq 1$,*

$$\frac{r}{\zeta(2)} \leq \frac{J_2(r)}{r} = r\prod_{p\mid r}\left(1-\frac{1}{p^2}\right) \leq r. \tag{15.3}$$

Consequently the individual local weight is of linear size in r. In particular, the unregularized cumulative weight cannot be finite; more precisely,

$$\sum_{r\leq R}\frac{J_2(r)}{r} = \frac{R^2}{2\zeta(3)} + O(R\log R). \tag{15.4}$$

Proof. The product in (15.3) is bounded below by $\prod_p(1-p^{-2}) = 1/\zeta(2)$. For the average, use $J_2(r)/r = r\sum_{d\mid r}\mu(d)d^{-2}$, interchange the finite sums, and sum m up to R/d. The main term is $R^2\sum_{d\geq 1}\mu(d)d^{-3}/2 = R^2/(2\zeta(3))$; elementary estimation of the remaining terms gives the displayed error. □

Remark 15.3 (Why S_0 nevertheless converges away from its branches)*.* For fixed s with $\Re s > 0$, once $r\Re s > 1$, $\log\zeta(rs) = O(2^{-r\Re s})$. This exponential decay dominates the linear factor $J_2(r)/r$. Thus the growth of the local monodromy weight is compatible with convergence of (14.2) away from its singular points.

Warning 15.4 (No global zero-kernel identity is claimed)*.* A formal substitution of a Hadamard expansion for $\log\zeta$ into (14.2) produces nested sums over r and zeta zeros. Those sums require prescribed symmetric ordering in the zero variable and separate convergence control in r. Version 8 does not interchange them and does not claim a globally convergent identity of the form $\sum_{\rho,\rho'}\mathcal{K}_s(\rho,\rho')$. The local monodromy Proposition 15.1 does not require such a rearrangement.

16. Gauge invariance of the arithmetic cocycle

The previous sections use the completely additive coordinate $A(n) = \Omega(n)$. It is useful to record an exact invariance that holds for any completely additive A.

Proposition 16.1 (Exact gauge invariance)**.** *Let*

$$\mathcal{A}_A(m,n) = A(m)\log n - \log m\, A(n) \tag{16.1}$$

and let $\chi_{\vartheta,A} = \exp(i\vartheta\mathcal{A}_A/2)$. For every $c \in \mathbb{R}$,

$$\boxed{\mathcal{A}_{A+c\log}(m,n) = \mathcal{A}_A(m,n), \qquad \chi_{\vartheta,A+c\log} = \chi_{\vartheta,A}.} \tag{16.2}$$

Consequently every twisted product and twisted inverse constructed only from this cocycle depends on A only through its class modulo $\mathbb{R}\log$.

Proof. The additional contribution is

$$c\log m\log n - c\log m\log n = 0. \tag{16.3}$$

□

Remark 16.2. This is an obstruction as well as an invariance: the deformation cannot recover the component of an additive coordinate parallel to log from the cocycle alone. The statement concerns the chosen cocycle itself and should not be confused with invariance of the coefficient curvature under arbitrary replacement by a cohomologous cocycle.

17. Cohomological obstruction to scalar Euler locality

Write $e_n = \delta_n$ for the canonical arithmetic basis. For distinct primes $p < q$,

$$e_p \star_\vartheta e_q = e^{\,i\vartheta \log(q/p)/2} e_{pq}, \qquad e_q \star_\vartheta e_p = e^{-\,i\vartheta \log(q/p)/2} e_{pq}. \tag{17.1}$$

Theorem 17.1 (No scalar prime-local Euler factorization)**.** *Fix $\vartheta \in \mathbb{R}$. Suppose that there exist scalars $a_p(\vartheta)$ and a fixed increasing order of the primes such that the square-free coefficients of μ are produced by an ordered twisted product*

$$\mu = \overrightarrow{\prod_p}{}^{\star_\vartheta} (1 - a_p e_p). \tag{17.2}$$

Then $\vartheta = 0$. Hence, for every $\vartheta \neq 0$, the nontrivial arithmetic cocycle obstructs any scalar Euler factorization into independent one-prime factors of this form.

Proof. For primes $p < q$, comparison of the pq-coefficient gives

$$a_p a_q e^{\,i\vartheta \log(q/p)/2} = 1. \tag{17.3}$$

For $p < q < r$, the three corresponding equations imply

$$a_p^2 = e^{-i\vartheta \log(q/p)}. \tag{17.4}$$

The left side is independent of q. Taking two primes $q_1, q_2 > p$ therefore yields

$$e^{-i\vartheta \log(q_1/q_2)} = 1. \tag{17.5}$$

If $\vartheta \neq 0$, applying this simultaneously to independent prime ratios forces a nontrivial rational linear relation among logarithms of distinct primes, contradicting unique factorization. □

Remark 17.2 (What is and is not claimed). Nontrivial 2-cocycles and twisted algebras are classical. The theorem is a specific obstruction inside the present arithmetic deformation. It does not say that every nontrivial cocycle destroys every conceivable noncommutative or adelic Euler product; it rules out scalar independent one-prime factors compatible with the stated twisted product.

18. Ordered Euler object and the global Euler-locality defect

For $\sigma > 1$, let $\mathcal{A}_\sigma$ be the Banach space of arithmetic functions with

$$\|f\|_\sigma = \sum_{n \geq 1} |f(n)| n^{-\sigma}. \tag{18.1}$$

Since $|\chi_\vartheta| = 1$,

$$\|f \star_\vartheta g\|_\sigma \leq \|f\|_\sigma \|g\|_\sigma. \tag{18.2}$$

For a prime p, define

$$L_p = \sum_{a \geq 0} e_{p^a}. \tag{18.3}$$

The cocycle is trivial on a single prime ray, so

$$\boxed{L_p = (1 - e_p)^{-1}_{\star_\vartheta}.} \tag{18.4}$$

Moreover,

$$\sum_p \|L_p - 1\|_\sigma = \sum_p \frac{p^{-\sigma}}{1 - p^{-\sigma}} < \infty. \tag{18.5}$$

Proposition 18.1 (Norm-convergent ordered Euler object)**.** *For $\sigma > 1$, the increasing-prime product*

$$\boxed{E_\vartheta = \overrightarrow{\prod_p} \star_\vartheta L_p} \tag{18.6}$$

converges in $\mathcal{A}_\sigma$. If $n = \prod_{j=1}^r p_j^{a_j}$ with $p_1 < \cdots < p_r$, then

$$\boxed{E_\vartheta(n) = \exp\left[\frac{i\vartheta}{2}\sum_{i<j} a_i a_j \log\frac{p_j}{p_i}\right].} \tag{18.7}$$

Proof. The norm convergence follows from (18.5) and the standard convergence criterion for ordered infinite products in a Banach algebra. The coefficient formula follows by repeated use of the cocycle identity; all self-ray phases vanish. □

For $\sigma > \sigma_*$, the twisted Möbius inverse z_ϑ is already an element of $\mathcal{A}_\sigma$. Define

$$\boxed{R_\vartheta = z_\vartheta - E_\vartheta.} \tag{18.8}$$

Theorem 18.2 (Exact vanishing criterion for the Euler-locality defect)**.** *For real ϑ,*

$$\boxed{R_\vartheta \equiv 0 \iff \vartheta = 0.} \tag{18.9}$$

In fact, products of two primes already detect nonvanishing.

Proof. At $\vartheta = 0$, both coefficient systems equal 1. For $p < q$, put $\lambda = \frac{\vartheta}{2}\log(q/p)$. Then

$$R_\vartheta(pq) = 2\cos\lambda - 1 - e^{i\lambda} = e^{-i\lambda} - 1. \tag{18.10}$$

If this vanished for every prime pair, then $\vartheta\log(q/p) \in 4\pi\mathbb{Z}$ for every $p < q$, contradicting the same universal-resonance argument used in Theorem 17.1 unless $\vartheta = 0$. □

19. The odd Euler-locality defect: existence and exact initial domain

Because $z_\vartheta = z_{-\vartheta}$, one has $z_0'(n) = 0$. Define

$$\Phi_{\mathrm{ord}}(n) = \sum_{p<q} v_p(n) v_q(n) \log\frac{q}{p}. \tag{19.1}$$

Differentiating the coefficient formula for the ordered Euler object gives

$$R_0'(n) = -\frac{i}{2}\Phi_{\mathrm{ord}}(n). \tag{19.2}$$

Definition 19.1 (Odd Euler-locality Dirichlet defect)**.** For $\Re s > 1$, set

$$\mathcal{O}(s) = \frac{1}{2}\sum_{n\geq 1}\frac{\Phi_{\mathrm{ord}}(n)}{n^s}. \tag{19.3}$$

Whenever the differentiated defect series is used, it is understood through this absolutely convergent Dirichlet series; no interchange with ∂_ϑ is made outside the domain justified below.

Put

$$u_p(s) = \frac{p^{-s}}{1-p^{-s}} = \frac{1}{p^s - 1}, \qquad A(s) = \sum_p u_p(s), \qquad B(s,t) = \sum_p u_p(s) p^{it}. \tag{19.4}$$

Theorem 19.2 (Absolute convergence, holomorphy, and differentiation)**.** *The series* (19.3)*, $A(s)$, and $B(s,t)$ converge absolutely and locally uniformly for $\Re s > 1$, uniformly in real t on compact sets. The function $\mathcal{O}$ is holomorphic there. For every nonnegative integers a, b, the derivatives $\partial_s^a \partial_t^b B(s,t)$ are obtained by termwise differentiation on compact subsets of $\Re s > 1$ and bounded real t-intervals.*

Proof. Fix $\varepsilon > 0$ and restrict to $\Re s \geq 1 + \varepsilon$. Since $|1 - p^{-s}| \geq 1 - p^{-1-\varepsilon}$,

$$|u_p(s)| \leq \frac{p^{-1-\varepsilon}}{1 - p^{-1-\varepsilon}} \leq C_\varepsilon p^{-1-\varepsilon}, \tag{19.5}$$

with a constant independent of p. Therefore $\sum_p |u_p(s)|(\log p)^j$ converges uniformly on the stated compacta for every fixed $j \geq 0$, by comparison with $\sum_{n\geq 2} n^{-1-\varepsilon}(\log n)^j$. This proves normal convergence of A, B, and all the displayed termwise derivatives.

For $\mathcal{O}$, use

$$0 \leq \Phi_{\mathrm{ord}}(n) \leq \sum_{p<q} v_p(n)v_q(n)(\log p + \log q). \tag{19.6}$$

Summing first over n with fixed distinct p, q, absolute convergence of Euler products gives

$$\sum_{n\geq 1} \frac{v_p(n)v_q(n)}{n^s} = \zeta(s)u_p(s)u_q(s), \qquad \Re s > 1. \tag{19.7}$$

Consequently the absolute majorant for (19.3) is bounded by a constant multiple of

$$\left(\sum_p |u_p(s)|\right)\left(\sum_q |u_q(s)| \log q\right), \tag{19.8}$$

which is locally uniformly finite by (19.5). The Weierstrass theorem then gives holomorphy and termwise differentiation in s. □

Theorem 19.3 (Prime-pair and Fourier representations)**.** *For $\Re s > 1$,*

$$\mathcal{O}(s) = \frac{\zeta(s)}{2} \sum_{p<q} u_p(s)u_q(s) \log \frac{q}{p}, \tag{19.9}$$

$$= \frac{\zeta(s)}{2\pi} \int_0^\infty \frac{A(s)^2 - B(s,t)B(s,-t)}{t^2}\, dt. \tag{19.10}$$

The integral is absolutely convergent. Moreover,

$$B(s,t) = \sum_{k\geq 1} P(ks - it), \qquad \Re s > 1, \tag{19.11}$$

with absolute and locally uniform convergence for real t.

Proof. Identity (19.7) follows by multiplying the local geometric series at p and q:

$$\sum_{a\geq 0} ap^{-as} = \frac{p^{-s}}{(1 - p^{-s})^2}, \tag{19.12}$$

and division by the two ordinary local Euler factors leaves one factor $u_p(s)u_q(s)$. Inserting this identity into the absolutely convergent series of Theorem 19.2 proves (19.9).

Next, normal convergence permits multiplication of the two prime sums:

$$A(s)^2 - B(s,t)B(s,-t) = 2\sum_{p<q} u_p(s)u_q(s)\big[1 - \cos(t\log(q/p))\big]. \tag{19.13}$$

For real a, integration by parts gives

$$\int_0^\infty \frac{1 - \cos(at)}{t^2}\, dt = \frac{\pi}{2}|a|. \tag{19.14}$$

The boundary term vanishes because $(1-\cos(at))/t = O(t)$ at zero and is $O(t^{-1})$ at infinity; the remaining integral is the classical Dirichlet integral. Furthermore,

$$\sum_{p<q} |u_p(s)u_q(s)| \int_0^\infty \frac{|1-\cos(t\log(q/p))|}{t^2}\,dt$$

$$\le \frac{\pi}{2} \sum_{p<q} |u_p(s)u_q(s)|(\log p + \log q) < \infty, \tag{19.15}$$

by (19.8). Fubini's theorem is therefore applicable, and (19.10) follows.

Finally, $u_p(s) = \sum_{k\ge1} p^{-ks}$ absolutely for $\Re s > 1$. Thus

$$\sum_p \sum_{k\ge1} |p^{-ks}| \le \sum_p \frac{p^{-\sigma}}{1-p^{-\sigma}} < \infty, \tag{19.16}$$

so Tonelli–Fubini gives (19.11). The same majorant is independent of real t, proving the asserted local uniformity. □

Definition 19.4 (Normalized Fourier energy)**.** In $\Re s > 1$, define

$$\boxed{\mathcal{I}(s) := \int_0^\infty \frac{A(s)^2 - B(s,t)B(s,-t)}{t^2}\,dt = \frac{2\pi\mathcal{O}(s)}{\zeta(s)}.} \tag{19.17}$$

19.1. **Endpoint regularity and transverse derivatives.** For

$$M_j(s) = \sum_p u_p(s)(\log p)^j, \tag{19.18}$$

normal convergence from Theorem 19.2 yields

$$B(s,t) = A(s) + iM_1(s)t - \frac{1}{2}M_2(s)t^2 + O_K(t^3) \tag{19.19}$$

uniformly for s in every compact $K \subset \{\Re s > 1\}$. Hence

$$\lim_{t\to0} \frac{A(s)^2 - B(s,t)B(s,-t)}{t^2} = A(s)M_2(s) - M_1(s)^2. \tag{19.20}$$

At infinity, $|B(s,t)| \le \sum_p |u_p(s)|$, so the integrand is $O_K(t^{-2})$. These two estimates give a second direct proof of absolute integrability in (19.17).

Proposition 19.5 (Exact first transverse derivative)**.** *For $\Re s > 1$,*

$$\boxed{\partial_t B(s,0) = -i\frac{\zeta'}{\zeta}(s).} \tag{19.21}$$

For every integer $a \ge 0$,

$$\boxed{\partial_s^a \partial_t B(s,0) = -i\left(\frac{\zeta'}{\zeta}\right)^{(a)}(s).} \tag{19.22}$$

Both identities extend meromorphically wherever the right-hand side does.

Proof. Termwise differentiation is justified by Theorem 19.2; hence

$$\partial_t B(s,0) = i\sum_p \frac{\log p}{p^s - 1}. \tag{19.23}$$

The absolutely convergent Euler-product logarithmic derivative is

$$-\frac{\zeta'}{\zeta}(s) = \sum_p \sum_{a\ge1} (\log p)p^{-as} = \sum_p \frac{\log p}{p^s - 1}, \tag{19.24}$$

which proves (19.21). Differentiating in s termwise on compacta gives (19.22). The final statement follows from uniqueness of meromorphic continuation. □

Proposition 19.6 (Higher pure transverse derivatives)**.** *For every integer $j \geq 1$ and $\Re s > 1$,*

$$\partial_t^j B(s,0) = (-i)^j \sum_{n\geq 1} c_j(n) \left(\frac{\zeta'}{\zeta}\right)^{(j-1)}(ns), \tag{19.25}$$

where

$$c_j(n) = \sum_{d|n} \mu(d) d^{j-1} = \prod_{p|n} (1 - p^{j-1}). \tag{19.26}$$

In particular, $c_1(n) = \delta_{n1}$, explaining why only the first transverse derivative is a complete replica filter.

Proof. Insert the prime-zeta Möbius inversion

$$P(w) = \sum_{m\geq 1} \frac{\mu(m)}{m} \operatorname{Log} \zeta(mw), \qquad \Re w > 1, \tag{19.27}$$

into (19.11). Absolute convergence in $\Re s > 1$ permits termwise differentiation, giving

$$\partial_t^j B(s,0) = (-i)^j \sum_{k,m\geq 1} \mu(m) m^{j-1} L^{(j-1)}(mks), \tag{19.28}$$

with $L = \zeta'/\zeta$. Grouping by $n = mk$ is legitimate because the same differentiated series is absolutely convergent on compacta. The inner divisor sum is $c_j(n)$, proving (19.25). Since μ is supported on square-free integers, multiplicativity gives the Euler factor $1 - p^{j-1}$ in (19.26). For $j = 1$, the standard identity $\sum_{d|n} \mu(d) = \delta_{n1}$ completes the proof. □

20. Continuation of the Fourier energy and complete replica accounting

The prime-zeta identity (19.27) is classical [8]. What is needed here is not merely the location of its branch points, but a proof that the energy integral can be continued while retaining control of all sums, derivatives, contour singularities, and boundary terms.

20.1. **Grouped log-zeta expansion and its normal tail.** For $\Re s > 1$, regrouping the absolutely convergent double series gives

$$\boxed{B(s,t) = \sum_{n\geq 1} \sum_{d|n} \frac{\mu(d)}{d} \operatorname{Log} \zeta(ns - idt).} \tag{20.1}$$

Lemma 20.1 (Normal convergence of the large-dilation tail)**.** *Fix s_0 with $\sigma_0 := \Re s_0 > 0$. There are a neighbourhood U of s_0 and an integer N_0 such that the part of (20.1) with $n > N_0$ is single-valued and holomorphic in $s \in U$, is C^∞ in real t, and all mixed derivatives of fixed finite order converge normally on $U \times \mathbb{R}$ after multiplication by any fixed polynomially decaying weight in t.*

Proof. Choose U so that $\Re s \geq \sigma_0/2$ on U, and choose N_0 with $(N_0+1)\sigma_0/2 > 1+\delta$ for some $\delta > 0$. Then for $n > N_0$,

$$\Re(ns - idt) = n \Re s \geq 1 + \delta. \tag{20.2}$$

In that half-plane the Euler product defines the principal holomorphic logarithm and

$$|\operatorname{Log} \zeta(w)| \leq \sum_{m\geq 2} \frac{C_\delta}{m^{\Re w}} \leq C'_\delta 2^{-\Re w}. \tag{20.3}$$

Every fixed derivative of $\operatorname{Log}\zeta(w)$ satisfies the same estimate with an additional polynomial factor in $\Re w$, obtained by differentiating the absolutely convergent von Mangoldt series. The divisor coefficient satisfies

$$\sum_{d|n}\frac{1}{d}\le\tau(n), \tag{20.4}$$

and $\tau(n)$ has subpolynomial growth. Hence the resulting tail is majorized by a convergent series of the form $\sum_{n>N_0} n^C\tau(n)2^{-n\sigma_0/2}$, uniformly in real t. The Weierstrass theorem proves normal convergence and permits all asserted finite-order differentiations. □

Definition 20.2 (Continued branch of $\mathcal{I}$)**.** A branch $\mathcal{I}^\sharp$ near a point s_0 with $\Re s_0>0$ is obtained by analytic continuation from $\Re s>1$ of the finitely many head terms $n\le N_0$ in (20.1), while the normal tail of Lemma 20.1 is kept on its single-valued branch. Whenever a moving logarithmic singularity meets the real t-axis, the t-contour is deformed continuously around it. Different continuation paths may change lower-order branch constants; the principal coefficients computed below are independent of those choices.

Lemma 20.3 (Convergence of the continued contour integral)**.** *Every local branch from Definition 20.2 can be represented by a locally finite deformed t-contour on which the continued integrand is absolutely integrable at infinity and locally integrable at every logarithmic branch point. The contour may be chosen so that the contribution of all indentations above height T tends to zero as $T\to\infty$.*

Proof. Only the finite head $n\le N_0$ can contribute zeros or the pole of ζ to the t-contour. For each fixed head pair (n,d), those points are obtained from

$$ns-idt=\rho' \tag{20.5}$$

with ρ' a zero or pole of ζ. The classical zero-counting estimate

$$N_\zeta(T)=O(T\log T) \tag{20.6}$$

is unconditional; see [13]. Hence the number of contour singularities of height at most T is $O(T\log T)$.

Near a zero of multiplicity m, local factorization gives

$$\operatorname{Log}\zeta(\rho'+w)=m\operatorname{Log}w+h(w), \tag{20.7}$$

with h holomorphic. Thus each singularity is logarithmic and $|\operatorname{Log}|t-t_0||^2$ is locally integrable. Choose pairwise compatible indentations with radii no larger than $(1+|t_0|)^{-4}$, shrinking them further when necessary to avoid overlaps. On the complementary contour, standard bounds obtained from the Hadamard product for ξ imply

$$|\operatorname{Log}\zeta(\sigma+it)|\le C_K\log^2(2+|t|) \tag{20.8}$$

for the finitely many vertical strips involved, after the chosen disks have been removed; this standard estimate is sufficient here and follows from the zero count and local factorization, again as in [13]. Consequently

$$\frac{|A(s)^2-B(s,t)B(s,-t)|}{|t|^2}\le C_K\frac{1+\log^4(2+|t|)}{|t|^2} \tag{20.9}$$

away from the indentation arcs. The right-hand side is integrable.

On an indentation of radius r_0 around height t_0, the logarithmic factor is $O(|\log r_0|+\log(2+|t_0|))$, the arc length is $O(r_0)$, and the weight is $O(t_0^{-2})$ for $|t_0|\ge1$. The sum of arc contributions is therefore dominated by

$$\sum_{t_0}\frac{r_0\,[1+|\log r_0|+\log^2(2+|t_0|)]}{1+t_0^2}, \tag{20.10}$$

which converges because $r_0 \le (1+|t_0|)^{-4}$ and the number of t_0's in a unit interval is $O(\log(2+|t_0|))$ on average by (20.6). Local logarithmic integrability handles the finitely many points with $|t_0| < 1$. This proves absolute convergence and the vanishing of the indentation tail. □

Theorem 20.4 (Existence of the continued energy germ)**.** *For every s_0 with $\Re s_0 > 0$, every branch obtained from Definition 20.2 defines a sectorial analytic germ $\mathcal{I}^\sharp$ at s_0 away from its local logarithmic cuts. It coincides with $\mathcal{I}$ in the common domain $\Re s > 1$. No RH, zero-simplicity, or nonresonance assumption is used.*

Proof. Lemma 20.1 reduces the continuation to finitely many logarithms of meromorphic functions. On every compact subarc of the deformed contour not meeting an indentation, these logarithms are holomorphic in s, and their products are dominated by the integrable majorant (20.9). On each indentation, use the local factorization (20.7); after translating the local contour, the integral is a finite linear combination of parameter integrals of $\mathrm{Log}(a(s-s_*)+b\tau)$ and products of two such logarithms. These are holomorphic in a slit sector by differentiation under the finite local integral. The normal convergence of the sum of indentation contributions follows from (20.10). Therefore the sum of the compact-contour part, indentation parts, and tail is analytic in the slit sector. Continuation from $\Re s > 1$ fixes the branch on overlaps, so uniqueness of analytic continuation gives agreement with (19.17) in the initial half-plane. □

20.2. **All contributions arriving at a scaled point.** Fix s_0 with $0 < \Re s_0 < 1$, and define

$$\mathcal{R}(s_0) = \{N \in \mathbb{N} : \zeta(Ns_0) = 0\}, \qquad q_N = \mathrm{ord}_{w=Ns_0}\,\zeta(w). \tag{20.11}$$

The set $\mathcal{R}(s_0)$ is finite, because a nontrivial zero has real part in $(0,1)$, so $N\Re s_0 < 1$. Define

$$C(s_0) = \sum_{N\in\mathcal{R}(s_0)} q_N \frac{\varphi(N)}{N}, \tag{20.12}$$

$$D(s_0) = \sum_{N\in\mathcal{R}(s_0)} q_N \frac{\varphi(N)}{N}\log N, \tag{20.13}$$

and, for real u,

$$G_{s_0}(u) = \sum_{N\in\mathcal{R}(s_0)} q_N \sum_{d\mid N} \frac{\mu(d)}{d}\,\mathrm{Log}\left(1 - i\frac{d}{N}u\right). \tag{20.14}$$

Lemma 20.5 (Complete endpoint singular block)**.** *With $z = s - s_0$, the complete contribution of all log-zeta terms singular at $(s,t) = (s_0, 0)$ is*

$$S(z,t) = \sum_{N\in\mathcal{R}(s_0)} q_N \sum_{d\mid N} \frac{\mu(d)}{d}\,\mathrm{Log}(Nz - idt). \tag{20.15}$$

After subtracting S, the remainder $R(z,t) = B(s_0+z,t) - S(z,t)$ is holomorphic in z and twice continuously differentiable in real t in some product neighbourhood of $(0,0)$.

Proof. In the finite head of (20.1), a term indexed by (n,d) is singular at $(s_0,0)$ precisely when $\zeta(ns_0) = 0$. Thus $n = N \in \mathcal{R}(s_0)$, while every divisor $d \mid N$ contributes the same zero at $t = 0$. Local factorization (20.7) then yields exactly (20.15). Since the head is finite and all endpoint singular factors have been removed, discreteness of the zero set permits a small product neighbourhood containing no further head singularity. The normal tail is smooth by Lemma 20.1. Hence the remainder has the asserted regularity. □

Lemma 20.6 (Interior contour singularities are lower order)**.** *Let $t_0 > 0$ be any logarithmic singularity of a continued head term. Its local contribution to $\mathcal{I}^\sharp(s_0+z)$, including products of two logarithmic factors, is $O(1) + O(z\log^2 z)$ in a slit sector. In particular, no interior singularity can contribute either $z^{-1}\log z$ or z^{-1}.*

Proof. Write $t = t_0 + x$. The weight is analytic there:

$$\frac{1}{t^2} = \frac{1}{t_0^2} + O(x). \tag{20.16}$$

A local logarithmic factor has the form $\mathrm{Log}(az - ibx) + h(z, x)$, with h holomorphic. On a symmetric local contour of fixed radius, set $x = zu$ only in the central part $|x| \le c|z|$; the Jacobian contributes a factor z, while the weight has no inverse power of z. This region is therefore $O(z \log^2 z)$. On the complementary part $c|z| < |x| < \delta$, expand

$$\mathrm{Log}(az - ibx) = \mathrm{Log}(-ibx) + O\Big(\frac{|z|}{|x|}\Big) \tag{20.17}$$

uniformly on the chosen slit sector. The error integrates to $O(z|\log z|)$, and products of two logarithms to $O(z \log^2 z)$. The zeroth-order terms give a finite constant because $\mathrm{Log}\,|x|$ and $\mathrm{Log}^2\,|x|$ are locally integrable. Therefore no negative power of z is produced. □

Theorem 20.7 (Exact coefficient of the logarithmic-over-pole term)**.** *For every continued branch $\mathcal{I}^\sharp$ at s_0, the coefficient of $\log z/z$, $z = s - s_0$, is*

$$\boxed{[\log z/z]\,\mathcal{I}^\sharp(s_0 + z) = -\pi q_1 C(s_0),} \tag{20.18}$$

where $q_1 = 0$ when $1 \notin \mathcal{R}(s_0)$. Consequently,

$$\boxed{[\log z/z]\,\mathcal{I}^\sharp = 0 \iff \zeta(s_0) \neq 0.} \tag{20.19}$$

No singularity with $t_0 > 0$ can change this coefficient.

Proof. From (20.15), for $t = zu$ one has

$$S(z, zu) = C(s_0)\,\mathrm{Log}\,z + D(s_0) + G_{s_0}(u), \qquad S(z, 0) = C(s_0)\,\mathrm{Log}\,z + D(s_0). \tag{20.20}$$

The singular–singular numerator equals

$$\begin{aligned} &S(z,0)^2 - S(z,zu)S(z,-zu) \\ &= -\big(C\,\mathrm{Log}\,z + D\big)\big(G(u) + G(-u)\big) - G(u)G(-u). \end{aligned} \tag{20.21}$$

The contour rotation from the ray $u = t/z$ to the positive real axis is legitimate in every closed subsector because the branch points $u = \pm iN/d$ lie on the imaginary axis; the large circular arc contributes zero since the integrand is $O(\log^2 |u|/|u|^2)$. Therefore the coefficient of $\mathrm{Log}\,z/z$ is

$$-C(s_0) \int_0^\infty \frac{G(u) + G(-u)}{u^2}\, du. \tag{20.22}$$

For real u,

$$G(u) + G(-u) = \sum_{N \in \mathcal{R}(s_0)} q_N \sum_{d|N} \frac{\mu(d)}{d} \log\left(1 + \frac{d^2u^2}{N^2}\right). \tag{20.23}$$

Integration by parts gives, for $a > 0$,

$$\int_0^\infty \frac{\log(1 + a^2u^2)}{u^2}\, du = \pi a. \tag{20.24}$$

The boundary term vanishes at zero and infinity. Thus

$$\begin{aligned} \int_0^\infty \frac{G(u) + G(-u)}{u^2}\, du &= \pi \sum_{N \in \mathcal{R}(s_0)} \frac{q_N}{N} \sum_{d|N} \mu(d) \\ &= \pi q_1, \end{aligned} \tag{20.25}$$

because $\sum_{d|N}\mu(d)=\delta_{N1}$. This proves (20.18). Since all summands in $C(s_0)$ are positive, $C(s_0)>0$ whenever $\mathcal{R}(s_0)\neq\varnothing$, so the coefficient vanishes exactly when $q_1=0$, equivalently $\zeta(s_0)\neq 0$. Finally, Lemma 20.6 excludes any restoration of the same order by interior contour singularities. □

Corollary 20.8 (Blockwise Möbius cancellation)**.** *Let ρ be a zero of multiplicity m_ρ, let $r\geq 1$, and set $s_0=\rho/r$. The singular–singular block attached specifically to (ρ,r) contributes*

$$\boxed{-\pi m_\rho^2\frac{\varphi(r)}{r^2}\left(\sum_{d|r}\mu(d)\right)\frac{\operatorname{Log}(s-s_0)}{s-s_0}} \tag{20.26}$$

to the logarithmic-over-pole scale of $\mathcal{I}^\sharp$. Hence this block is zero for every $r>1$, whereas for $r=1$ it equals

$$-\pi m_\rho^2\frac{\operatorname{Log}(s-\rho)}{s-\rho}. \tag{20.27}$$

Proof. Restrict (20.15) to a single dilation index r. Then $C=m_\rho\varphi(r)/r$, while the integral (20.25) for that single block is $\pi m_\rho r^{-1}\sum_{d|r}\mu(d)$. Their negative product gives (20.26). The divisor identity $\sum_{d|r}\mu(d)=0$ for $r>1$ and equals one for $r=1$, proving both conclusions. □

20.3. **Pure replicas: the next coefficient and the complete local germ.** Assume now that $\mathcal{R}(s_0)\neq\varnothing$ but $\zeta(s_0)\neq 0$, so $q_1=0$. Define

$$\beta(s_0):=-\int_0^\infty\frac{|G_{s_0}(u)|^2}{u^2}\,du. \tag{20.28}$$

Lemma 20.9 (Convergence and strict negativity of the collision energy)**.** *Under $q_1=0$, the integral* (20.28) *converges and*

$$\boxed{\beta(s_0)<0.} \tag{20.29}$$

Proof. At zero,

$$G'_{s_0}(0)=-i\sum_{N\in\mathcal{R}(s_0)}\frac{q_N}{N}\sum_{d|N}\mu(d)=0, \tag{20.30}$$

because every $N\in\mathcal{R}(s_0)$ is greater than one. Since $G(0)=0$, it follows that $G(u)=O(u^2)$ and the integrand is $O(u^2)$ at zero. At infinity,

$$G(u)=C(s_0)\log u+O(1), \tag{20.31}$$

so the integrand is $O(\log^2 u/u^2)$. Thus the integral converges.

It remains to exclude $G\equiv 0$. Let $N_{\max}=\max\mathcal{R}(s_0)$. Differentiating (20.14) meromorphically in u, the $d=1$ term belonging to $N_{\max}$ has a simple pole at $u=-iN_{\max}$. Every pole from a term with $M<N_{\max}$ has modulus $M/d\leq M<N_{\max}$, and terms with $N=N_{\max}$, $d>1$, have modulus $N_{\max}/d<N_{\max}$. Hence that pole cannot cancel. Therefore $G\not\equiv 0$, its squared modulus is positive on a set of positive measure, and (20.29) follows. □

Theorem 20.10 (Complete local expansion at a pure replica point)**.** *Suppose $\mathcal{R}(s_0)\neq\varnothing$ and $\zeta(s_0)\neq 0$. Every continued branch satisfies, in every closed slit subsector,*

$$\boxed{\mathcal{I}^\sharp(s_0+z)=\frac{\beta(s_0)}{z}+\pi C(s_0)\frac{\zeta'}{\zeta}(s_0)\operatorname{Log}z+O(1).} \tag{20.32}$$

In particular the pole coefficient is nonzero and no term $\operatorname{Log}z/z$ is present.

Proof. Decompose $B = S + R$ as in Lemma 20.5. The singular–singular part is given by (20.21). Since $q_1 = 0$, equation (20.25) makes the entire term linear in $C \operatorname{Log} z + D$ vanish. Therefore its negative-power contribution is exactly

$$\frac{1}{z}\left(-\int_0^\infty \frac{G(u)G(-u)}{u^2}\,du\right) = \frac{\beta(s_0)}{z}, \tag{20.33}$$

because on the positive real axis $G(-u) = \overline{G(u)}$. The contour rotation used in Theorem 20.7 shows the same coefficient in every slit subsector.

Near $(0,0)$, write

$$R(z, \pm t) = R_0(z) \pm h_1 t + O(t^2 + |z|t), \tag{20.34}$$

with $h_1 = \partial_t R(0,0)$. The constant cross term is proportional to

$$\int_0^\infty \frac{G(u) + G(-u)}{u^2}\,du = 0, \tag{20.35}$$

so it produces neither z^{-1} nor $z^{-1}\operatorname{Log} z$. For the odd linear term, one obtains

$$h_1 \int \frac{S(z,t) - S(z,-t)}{t}\,dt. \tag{20.36}$$

As $u \to +\infty$,

$$G(u) - G(-u) = -i\pi C(s_0) + O(u^{-1}), \tag{20.37}$$

so integration from $u = 1$ to $u = \delta/|z|$ gives

$$i\pi C(s_0) h_1 \operatorname{Log} z + O(1). \tag{20.38}$$

All Taylor terms of order t^2 or higher are integrable at the endpoint and contribute $O(1)$; terms involving z are also $O(1)$. By Lemma 20.6, every interior contour singularity is bounded up to $O(z \log^2 z)$, and the contour tail is normally convergent by Lemma 20.3. Thus no additional unbounded term of order z^{-1}, $z^{-1}\operatorname{Log} z$, or $\operatorname{Log} z$ is present.

Finally, differentiating S at $t = 0$ gives

$$\partial_t S(z,0) = -\frac{i}{z} \sum_{N \in \mathcal{R}(s_0)} \frac{q_N}{N} \sum_{d \mid N} \mu(d) = 0, \tag{20.39}$$

so $h_1 = \partial_t B(s_0, 0)$. Proposition 19.5 and meromorphic continuation give

$$h_1 = -i\frac{\zeta'}{\zeta}(s_0), \tag{20.40}$$

because $\zeta(s_0) \neq 0$. Substitution into (20.38) proves (20.32). □

20.4. **Single-replica divisor energy.** For a single zero ρ of multiplicity m_ρ and one dilation index $r > 1$, define

$$G_r(u) = \sum_{d \mid r} \frac{\mu(d)}{d} \operatorname{Log}\left(1 - i\frac{d}{r}u\right). \tag{20.41}$$

Proposition 20.11 (Replica energy, finite formula, and sign)**.** *Define*

$$\boxed{\beta_r = -\int_0^\infty \frac{|G_r(u)|^2}{u^2}\,du.} \tag{20.42}$$

Then $\beta_r < 0$ for every $r > 1$, and

$$\boxed{\beta_r = -\frac{\pi}{r} \sum_{d,e \mid r} \frac{\mu(d)\mu(e)}{de}\Big[(d+e)\log(d+e) - d\log d - e\log e\Big].} \tag{20.43}$$

For an isolated pure replica point $s_0 = \rho/r$, the pole coefficient in (20.32) *is $m_\rho^2 \beta_r$.*

Proof. The convergence and strict sign follow from the proof of Lemma 20.9; for a single $r > 1$, $G_r(0) = G_r'(0) = 0$ and $G_r \not\equiv 0$.

For the finite formula, set

$$F(a,b) = \Re \int_0^\infty \frac{\operatorname{Log}(1 - iau)\operatorname{Log}(1 + ibu)}{u^2}\, du, \qquad a, b > 0. \tag{20.44}$$

The integral converges absolutely after taking the real part of the paired expression. Differentiate first with respect to a and b, justified on compact parameter rectangles by dominated convergence after splitting $(0,1) \cup (1,\infty)$. One obtains

$$\frac{\partial^2 F}{\partial a \partial b} = \Re \int_0^\infty \frac{du}{(1 - iau)(1 + ibu)} = \frac{\pi}{a + b}. \tag{20.45}$$

The last equality follows from partial fractions and evaluation of the logarithms at a finite cutoff before taking the limit. Since $F(a,0) = F(0,b) = 0$, two integrations give

$$F(a,b) = \pi\big[(a+b)\log(a+b) - a\log a - b\log b\big]. \tag{20.46}$$

Expanding $|G_r(u)|^2$, using $a = d/r$, $b = e/r$, and the homogeneity of (20.46), gives (20.43). □

Corollary 20.12 (Radical law)**.** *For every $r > 1$,*

$$\boxed{\beta_r = \frac{\operatorname{rad}(r)}{r}\beta_{\operatorname{rad}(r)}.} \tag{20.47}$$

Equivalently, $r\beta_r$ depends only on the square-free kernel of r.

Proof. Put $q = \operatorname{rad}(r)$. In the rescaled form of (20.42), let $u = rx$. Then

$$\beta_r = -\frac{1}{r}\int_0^\infty \frac{1}{x^2}\left|\sum_{d \mid r} \frac{\mu(d)}{d}\operatorname{Log}(1 - idx)\right|^2 dx. \tag{20.48}$$

Only square-free divisors contribute, and those are exactly the divisors of q. The integral in (20.48) is therefore the same as the corresponding integral for q, while the exterior factors are $1/r$ and $1/q$. Hence $r\beta_r = q\beta_q$, which is (20.47). □

Corollary 20.13 (Prime and prime-power replica asymptotics)**.** *For a prime p,*

$$\boxed{\beta_p = -\frac{2\pi}{p^2}\left[p\log p - (p+1)\log(p+1) + (p+1)\log 2\right].} \tag{20.49}$$

As $p \to \infty$,

$$\beta_p = -\frac{2\pi\log 2}{p} + O\Big(\frac{\log p}{p^2}\Big). \tag{20.50}$$

For every integer $a \geq 1$,

$$\boxed{\beta_{p^a} = p^{1-a}\beta_p.} \tag{20.51}$$

Proof. In (20.43), for $r = p$ the only divisors are $1, p$. Writing out the four terms and collecting logarithms gives (20.49). Expanding $(p+1)\log(p+1) = p\log p + \log p + 1 + O(p^{-1})$ yields (20.50); the weaker displayed error is sufficient and uniform. Finally, $\operatorname{rad}(p^a) = p$, so Corollary 20.12 gives (20.51). □

20.5. **Actual zeta zeros, $\mathcal{O}$, and monodromy.**

Theorem 20.14 (Survival at a genuine zero)**.** *Let ρ be a zero of multiplicity m_ρ. For every continued branch,*

$$\mathcal{I}^\sharp(\rho+z) = -\pi m_\rho C(\rho)\frac{\operatorname{Log} z}{z} + O\!\left(\frac{1}{z}\right), \tag{20.52}$$

where

$$C(\rho) = \sum_{N\in\mathcal{R}(\rho)} q_N \frac{\varphi(N)}{N} \geq m_\rho. \tag{20.53}$$

If no scaled zero other than $N=1$ collides at ρ, then

$$\boxed{\mathcal{I}^\sharp(\rho+z) = -\pi m_\rho^2 \frac{\operatorname{Log} z}{z} + O(z^{-1}).} \tag{20.54}$$

Proof. At $s_0=\rho$, one has $q_1 = m_\rho > 0$. Theorem 20.7 gives the coefficient $-\pi q_1 C(\rho)$, proving (20.52). All other terms are at most $O(z^{-1})$: the singular–singular expansion (20.21) has no stronger scale, cross terms contain at most one singular logarithm, and interior singularities are lower order by Lemma 20.6. In the absence of collisions, $C(\rho)=m_\rho$, yielding (20.54). □

Corollary 20.15 (Correct singular orders for $\mathcal{O}$)**.** *Let $\zeta(\rho+z) = a_m z^m + O(z^{m+1})$, $a_m \neq 0$, with $m = m_\rho$. In the noncolliding case,*

$$\boxed{\mathcal{O}(\rho+z) = -\frac{a_m m^2}{2} z^{m-1}\operatorname{Log} z + O(z^{m-1}).} \tag{20.55}$$

For a simple zero,

$$\boxed{\mathcal{O}(\rho+z) = -\frac{\zeta'(\rho)}{2}\operatorname{Log} z + O(1).} \tag{20.56}$$

At a pure replica $s_0 = \rho/r$, $r>1$, with no collision with an actual zero at s_0,

$$\boxed{\mathcal{O}(s_0+z) = \frac{\zeta(s_0)\,\beta(s_0)}{2\pi z} + \frac{C(s_0)\zeta'(s_0)}{2}\operatorname{Log} z + O(1).} \tag{20.57}$$

Proof. Use the identity $\mathcal{O} = \zeta\mathcal{I}/(2\pi)$, initially valid for $\Re s > 1$ and then on every continued branch by analytic continuation. Multiplying (20.54) by $a_m z^m + O(z^{m+1})$ gives (20.55); the case $m=1$ is (20.56). At a pure replica, $\zeta(s_0)\neq 0$, so multiplication of (20.32) gives (20.57). □

Warning 20.16 (The requested $\log z/z$ formula belongs to $\mathcal{I}$). A formula of the form

$$C_{\rho,r}\left(\sum_{d\mid r}\mu(d)\right)\frac{\operatorname{Log}(s-\rho/r)}{s-\rho/r} \tag{20.58}$$

is correct for the normalized energy $\mathcal{I} = 2\pi\mathcal{O}/\zeta$, with

$$C_{\rho,r} = -\pi m_\rho^2 \frac{\varphi(r)}{r^2} \tag{20.59}$$

for the individual singular–singular block. It is not the correct local order of $\mathcal{O}$ itself at $r=1$, because ζ vanishes there and cancels the pole. Equations (20.55) and (20.57) are the correct statements for $\mathcal{O}$.

Corollary 20.17 (Monodromy detector for genuine zeros)**.** *Let $\Delta\mathcal{I}^\sharp$ denote one positive turn of the local branch. Then*

$$\operatorname*{Res}_{s=s_0} \Delta\mathcal{I}^\sharp(s) = -2\pi^2 i\, q_1 C(s_0). \tag{20.60}$$

Consequently

$$\boxed{\zeta(s_0) = 0 \iff \Delta\mathcal{I}^\sharp \text{ has a pole at } s_0.} \tag{20.61}$$

Proof. Only the term $-\pi q_1 C(s_0)\operatorname{Log} z/z$ changes by a term of order z^{-1} after $\operatorname{Log} z \mapsto \operatorname{Log} z + 2\pi i$. Its change is $-2\pi^2 i q_1 C(s_0)/z$. All pole terms without a logarithm are single-valued, and all lower logarithmic terms have bounded monodromy. Since $C(s_0) > 0$, the residue is nonzero exactly when $q_1 > 0$, i.e. when $\zeta(s_0) = 0$. □

20.6. **Analytic checklist for the replica theorem.**

Proposition 20.18 (Eight-point analytic closure)**.** *The replica analysis above establishes the following items without RH, zero-simplicity, or a nonresonance assumption:*

1. *$\mathcal{O}(s)$ and $\mathcal{I}(s)$ exist absolutely for $\Re s > 1$ and are holomorphic there.*
2. *Every local branch required for the replica analysis is obtained by continuation from that half-plane, with a normally convergent large-dilation tail.*
3. *All exchanges among prime sums, dilation sums, the t-integral, and the differentiations used in the initial half-plane are justified by the majorants displayed above.*
4. *The complete singular expansion through the unbounded terms is* (20.32) *at pure replica points and* (20.52) *at actual zeros.*
5. *Every contribution reaching s_0 at the endpoint is included by the finite set $\mathcal{R}(s_0)$; every interior contour contribution is covered by Lemma 20.6.*
6. *For each individual index $r > 1$, the coefficient of $z^{-1}\operatorname{Log} z$ vanishes by $\sum_{d|r}\mu(d) = 0$.*
7. *No other endpoint or interior term can restore a $z^{-1}\operatorname{Log} z$ contribution when $\zeta(s_0) \neq 0$.*
8. *The corresponding $r = 1$ term survives with nonzero coefficient $-\pi m_\rho^2$ in the noncolliding case, and more generally with $-\pi q_1 C(s_0)$.*

Proof. Items (1)–(3) are Theorems 19.2, 19.3, Lemmas 20.1 and 20.3, and Theorem 20.4. Item (4) is Theorems 20.10 and 20.14. Item (5) is Lemmas 20.5 and 20.6. Item (6) is Corollary 20.8. Item (7) follows from Theorem 20.7 together with Lemma 20.6. Item (8) is Theorem 20.14. Thus every item is a consequence of a proved statement in the preceding subsections. □

21. A two-variable Euler product and the covariance determinant

The determinant in (12.3) has an intrinsic interpretation that does not involve a linearly Möbius-weighted quantum state. For $\Re s > 1$ and u in a neighbourhood of zero define

$$\mathfrak{Z}(s,u) = \prod_p \left(1 - e^u p^{-s}\right)^{-1}, \qquad F(s,u) = \log \mathfrak{Z}(s,u). \tag{21.1}$$

Then $\mathfrak{Z}(s,0) = \zeta(s)$.

Theorem 21.1 (Hessian form of the curvature)**.** *For $\Re s > 1$,*

$$F_{uu}(s,0) = S_0(s), \tag{21.2}$$

$$-F_{su}(s,0) = S_1(s), \tag{21.3}$$

$$F_{ss}(s,0) = S_2(s). \tag{21.4}$$

Consequently

$$\boxed{\frac{2\mathcal{K}(s)}{\zeta(s)} = \det\begin{pmatrix} F_{uu}(s,0) & F_{us}(s,0) \\ F_{su}(s,0) & F_{ss}(s,0) \end{pmatrix}.} \tag{21.5}$$

Proof. Differentiate

$$F(s,u) = -\sum_p \log(1 - e^u p^{-s}) \tag{21.6}$$

termwise in a neighbourhood of $u = 0$, justified by normal convergence for $\Re s > 1$. The three derivatives are exactly (12.4). Equation (21.5) is then Theorem 12.1. □

Theorem 21.2 (Zeta-distribution covariance identity)**.** *Let $\sigma > 1$ be real and let N_σ be the zeta-distributed integer*

$$\mathbb{P}_\sigma(N_\sigma = n) = \frac{n^{-\sigma}}{\zeta(\sigma)}. \tag{21.7}$$

Then

$$\mathrm{Var}_\sigma(\Omega(N_\sigma)) = S_0(\sigma), \tag{21.8}$$

$$\mathrm{Cov}_\sigma(\Omega(N_\sigma), \log N_\sigma) = S_1(\sigma), \tag{21.9}$$

$$\mathrm{Var}_\sigma(\log N_\sigma) = S_2(\sigma). \tag{21.10}$$

Therefore

$$\boxed{\frac{2\mathcal{K}(\sigma)}{\zeta(\sigma)} = \det \mathrm{Cov}_\sigma \begin{pmatrix} \Omega(N_\sigma) \\ \log N_\sigma \end{pmatrix} > 0.} \tag{21.11}$$

Proof. Under (21.7), the prime valuations $v_p(N_\sigma)$ are independent geometric random variables with

$$\mathrm{Var}(v_p) = \frac{p^{-\sigma}}{(1 - p^{-\sigma})^2}. \tag{21.12}$$

Since

$$\Omega(N_\sigma) = \sum_p v_p(N_\sigma), \qquad \log N_\sigma = \sum_p v_p(N_\sigma) \log p, \tag{21.13}$$

independence gives (21.8)–(21.10). Strict positivity follows from

$$S_0 S_2 - S_1^2 = \sum_{p<q} w_p w_q \log^2(q/p), \qquad w_p = \frac{p^{-\sigma}}{(1 - p^{-\sigma})^2} > 0. \tag{21.14}$$

□

22. The positivity route and its exact obstruction

The coefficients $c(n) = -z_0''(n)$ are nonnegative. It is therefore natural to ask whether positivity itself forces the absence of the logarithmic branches in $1/2 < \Re s < 1$. It does not.

Proposition 22.1 (Complete monotonicity on the real axis)**.** *For every real $\sigma > 1$ and every integer $r \geq 0$,*

$$\boxed{(-1)^r \mathcal{K}^{(r)}(\sigma) = \sum_{n \geq 1} \frac{c(n)(\log n)^r}{n^\sigma} \geq 0.} \tag{22.1}$$

In particular $\mathcal{K}$ is completely monotone on $(1, \infty)$.

Proof. Differentiate the absolutely convergent Dirichlet series termwise. □

Proposition 22.2 (Positivity does not remove complex logarithmic branches)**.** *Nonnegative Dirichlet coefficients, complete monotonicity on the real axis, and the strict covariance inequality* (21.11) *do not imply a branch-free continuation of $\mathcal{K}$ to $1/2 < \Re s < 1$.*

Proof. The source term $P(s) = \sum_p p^{-s}$ itself has nonnegative Dirichlet coefficients for $\Re s > 1$, yet its continuation is expressed through $\log \zeta(rs)$ and has logarithmic branch points at scaled zeros of ζ; see (14.4). More directly, the positive-coefficient function S_0 in (14.2) is exactly the factor that contributes $m \log(s - \rho)$ at a hypothetical zero ρ with $1/2 < \Re\rho < 1$. Thus real-axis positivity and complete monotonicity do not control the monodromy of the analytic continuation. □

Corollary 22.3 (What would still be needed to prove RH)**.** *To deduce Corollary 13.3 from the curvature side, one needs an additional property that controls the complex continuation of $\mathcal{K}$, not merely the signs of its Dirichlet coefficients or the positive-semidefiniteness of the real covariance matrix in* (21.11)*.*

Proof. Proposition 22.1 and Theorem 21.2 control only real-axis signs and positive semidefiniteness. Proposition 22.2 exhibits logarithmic branches compatible with those same properties. Therefore any proof using this route must add information that controls the complex continuation of $\mathcal{K}$, as asserted. □

Remark 22.4 (Status of the attempted step)*.* The expected coefficient-positivity step has therefore been carried out and closed: $c(n) \geq 0$ gives complete monotonicity and a positive covariance Hessian, but those properties do not forbid the logarithmic branches whose absence is equivalent to RH. The obstruction is visible inside the same formula: S_0 has positive coefficients and is simultaneously the carrier of the prime-zeta logarithmic monodromy.

23. The Heisenberg-deformed zeta series

Definition 23.1 (Heisenberg-deformed zeta series)**.** For $\Re s > \sigma_*$, define

$$\zeta_{\mathrm{H}}(s; \vartheta) = \sum_{n \geq 1} \frac{z_\vartheta(n)}{n^s}. \tag{23.1}$$

Theorem 8.8 guarantees absolute convergence for every $\Re s > \sigma_*$.

Proposition 23.2 (Classical boundary)**.** *For $\Re s > 1$,*

$$\zeta_{\mathrm{H}}(s; 0) = \zeta(s). \tag{23.2}$$

Proof. By Proposition 8.5, $z_0(n) = 1$ for all n. Substitution into (23.1) gives the ordinary Dirichlet series for $\zeta(s)$. □

Proposition 23.3 (Even Heisenberg deformation)**.** *For $\Re s > \sigma_*$,*

$$\zeta_{\mathrm{H}}(s; -\vartheta) = \zeta_{\mathrm{H}}(s; \vartheta). \tag{23.3}$$

Proof. This follows termwise from Proposition 9.5 and absolute convergence. □

Warning 23.4 (No analytic continuation is asserted)*.* The series (23.1) is absolutely convergent and holomorphic on the open half-plane $\Re s > \sigma_*$. No meromorphic continuation, functional equation, scalar Euler product, or critical-strip theory for $\zeta_{\mathrm{H}}(s; \vartheta)$ is asserted here.

Proposition 23.5 (Conditional first-order rigidity of simple zeros)**.** *Assume that, for some simple zero ρ of the ordinary zeta function, there exists a function $F(s, \vartheta)$ analytic in a neighbourhood of $(\rho, 0)$ such that*

$$F(s, 0) = \zeta(s), \tag{23.4}$$

$$F(s, -\vartheta) = F(s, \vartheta), \tag{23.5}$$

and that F agrees with ζ_{H} wherever both are defined. Then the local zero branch $\rho(\vartheta)$ through ρ satisfies

$$\rho'(0) = 0. \tag{23.6}$$

Proof. The implicit-function theorem applies because $\zeta'(\rho) \neq 0$. Evenness (23.5) makes the unique local zero branch even, so its first derivative at the origin vanishes. □

Warning 23.6. Proposition 23.5 is conditional on an analytic continuation in both variables and does not supply such a continuation. It is included because it identifies a precise way in which the exact $\vartheta \mapsto -\vartheta$ symmetry would constrain the motion of classical Riemann zeros under the Heisenberg deformation if that continuation exists.

24. A genuine Möbius Hamiltonian

24.1. **Definition and domain.**

Definition 24.1 (Cosine and sine Weyl quadratures)**.** For $n \geq 1$, set

$$C_n = \frac{W_n + W_n^*}{2}, \tag{24.1}$$

$$J_n = \frac{W_n - W_n^*}{2i}. \tag{24.2}$$

Both are bounded self-adjoint operators with norm at most 1.

Definition 24.2 (Möbius Hamiltonian)**.** Fix an energy scale $E_0 > 0$ and $\sigma > 1$. Define

$$H_{\mu,\sigma,\vartheta} = E_0 \left(\operatorname{Re} \mathcal{M}_{\sigma,\vartheta} - I\right) = E_0 \sum_{n\geq 2} \frac{\mu(n)}{n^\sigma} C_n. \tag{24.3}$$

The series converges absolutely in operator norm.

Proposition 24.3 (Self-adjoint bounded dynamics)**.** *For $\sigma > 1$, $H_{\mu,\sigma,\vartheta}$ is bounded and self-adjoint on all of $\mathcal{H}$:*

$$\mathcal{D}(H_{\mu,\sigma,\vartheta}) = \mathcal{H}. \tag{24.4}$$

It therefore generates the unitary group

$$U(t) = \exp\left(-\frac{i}{\hbar} t H_{\mu,\sigma,\vartheta}\right). \tag{24.5}$$

Proof. The norm series in (24.3) is dominated by $E_0 \sum_{n\geq 2} |\mu(n)| n^{-\sigma}$, which converges for $\sigma > 1$. Every C_n is self-adjoint, hence so is the norm limit. The spectral theorem then defines (24.5). □

Proposition 24.4 (Explicit Schrödinger equation)**.** *For $\psi_0 \in \mathcal{H}$, the evolution $\psi(t) = U(t)\psi_0$ satisfies*

$$i\hbar\, \partial_t \psi(t,q) = (H_{\mu,\sigma,\vartheta}\psi(t))(q) \tag{24.6}$$

in the strong L^2 sense, where

$$(H_{\mu,\sigma,\vartheta}\psi)(q) = \frac{E_0}{2} \sum_{n\geq 2} \frac{\mu(n)}{n^\sigma} \Big[e^{\frac{i}{\hbar} y_n (q - x_n/2)} \psi(q - x_n) + e^{-\frac{i}{\hbar} y_n (q + x_n/2)} \psi(q + x_n) \Big]. \tag{24.7}$$

Proof. Equation (24.7) follows from Proposition 4.2 applied to W_n and W_n^*. The series converges in operator norm, so the bounded Hamiltonian gives a strong Schrödinger evolution for every initial vector. □

24.2. **Exact equations of motion for position and momentum.**

Lemma 24.5 (Bounded first commutators)**.** *For $\sigma > 1$, the series*

$$\sum_{n\geq 2} \frac{|\mu(n)|x_n}{n^\sigma}, \tag{24.8}$$

$$\sum_{n\geq 2} \frac{|\mu(n)|y_n}{n^\sigma} \tag{24.9}$$

converge. Hence $[H_{\mu,\sigma,\vartheta}, Q]$ and $[H_{\mu,\sigma,\vartheta}, P]$, initially defined on $\mathcal{S}(\mathbb{R})$, extend to bounded operators.

Proof. The first series is bounded by a constant multiple of $\sum n^{-\sigma}\log n$. Since

$$\Omega(n) \leq \frac{\log n}{\log 2}, \tag{24.10}$$

the second is bounded by another constant multiple of the same convergent series. □

Theorem 24.6 (Exact Heisenberg equations of motion)**.** *Let*

$$Q(t) = U(t)^* Q U(t), \tag{24.11}$$

$$P(t) = U(t)^* P U(t). \tag{24.12}$$

The bounded commutators in Lemma 24.5 imply invariance of $\mathcal{D}(Q)$ and $\mathcal{D}(P)$ under $U(t)$. Indeed, for $A = Q$ or P and $\psi \in \mathcal{D}(A)$, Duhamel's formula first on a core and then by graph-norm closure gives

$$AU(t)\psi = U(t)A\psi + \frac{i}{\hbar}\int_0^t U(t-\tau)[H, A]U(\tau)\psi\, d\tau. \tag{24.13}$$

whose right-hand side lies in L^2 because $[H, A]$ is bounded. On these domains, and equivalently as quadratic-form identities, the Heisenberg equations are

$$\dot{Q}(t) = \frac{E_0}{\hbar} U(t)^* \left[\sum_{n\geq 2} \frac{\mu(n)x_n}{n^\sigma} J_n\right] U(t), \tag{24.14}$$

$$\dot{P}(t) = \frac{E_0}{\hbar} U(t)^* \left[\sum_{n\geq 2} \frac{\mu(n)y_n}{n^\sigma} J_n\right] U(t). \tag{24.15}$$

At $t = 0$ this becomes

$$\dot{Q}(0) = \frac{E_0 \ell}{\hbar} \sum_{n\geq 2} \frac{\mu(n)\log n}{n^\sigma} J_n, \tag{24.16}$$

$$\dot{P}(0) = \frac{E_0 \kappa}{\hbar} \sum_{n\geq 2} \frac{\mu(n)\Omega(n)}{n^\sigma} J_n. \tag{24.17}$$

Proof. For $A = Q$ or P, the bounded-commutator theorem for unitary dynamics gives

$$\dot{A}(t) = \frac{i}{\hbar} U(t)^* [H_{\mu,\sigma,\vartheta}, A] U(t). \tag{24.18}$$

From Proposition 4.4,

$$[Q, C_n] = i x_n J_n, \tag{24.19}$$

$$[P, C_n] = i y_n J_n. \tag{24.20}$$

Therefore

$$[H_{\mu,\sigma,\vartheta}, Q] = -iE_0 \sum_{n\geq 2} \frac{\mu(n)x_n}{n^\sigma} J_n, \tag{24.21}$$

$$[H_{\mu,\sigma,\vartheta}, P] = -iE_0 \sum_{n\geq 2} \frac{\mu(n)y_n}{n^\sigma} J_n. \tag{24.22}$$

Lemma 24.5 justifies the sums as bounded operators. Substitution into (24.18) gives (24.14) and (24.15). □

Remark 24.7 (Physical and arithmetic meaning of the two equations). The position variable is Q, the logarithmic-scale coordinate. The momentum variable is $P = -i\hbar\partial_q$, the generator conjugate to that coordinate. Multiplication by an integer n is represented by a phase-space displacement of size $\ell \log n$ in Q and $\kappa\Omega(n)$ in P. Equations (24.16) and (24.17) therefore say that the Möbius Hamiltonian drives position by logarithmic size and momentum by prime-factor count, with the same quantum current operators J_n.

24.3. **The CCR is visible in the dynamics.**

Proposition 24.8 (Pairwise dynamical noncommutativity). *Extend* log *and* Ω *additively to positive rationals and define* W_r *by* (4.19) *for* $r \in \mathbb{Q}_{>0}$. *For* $m, n \in \mathbb{N}$, *put*

$$\Phi_{m,n} = \vartheta(\Omega(m)\log n - \log m\,\Omega(n)). \tag{24.23}$$

Then

$$[C_m, J_n] = \frac{1}{2}\sin\Big(\frac{\Phi_{m,n}}{2}\Big)\Big(W_{mn} + W_{m/n} - W_{n/m} - W_{1/(mn)}\Big). \tag{24.24}$$

Proof. Expand C_m and J_n using (24.1)–(24.2). Apply (4.6) to the four pairs (z_m, z_n), $(z_m, -z_n)$, $(-z_m, z_n)$, and $(-z_m, -z_n)$. □

Corollary 24.9 (Second-order arithmetic commutator). *At* $t = 0$,

$$\ddot{Q}(0) = \frac{iE_0^2}{2\hbar^2} \sum_{m,n\geq 2} \frac{\mu(m)\mu(n)x_n}{(mn)^\sigma} \sin\Big(\frac{\Phi_{m,n}}{2}\Big) \times \Big(W_{mn} + W_{m/n} - W_{n/m} - W_{1/(mn)}\Big), \tag{24.25}$$

and the analogous momentum equation is

$$\ddot{P}(0) = \frac{iE_0^2}{2\hbar^2} \sum_{m,n\geq 2} \frac{\mu(m)\mu(n)y_n}{(mn)^\sigma} \sin\Big(\frac{\Phi_{m,n}}{2}\Big) \times \Big(W_{mn} + W_{m/n} - W_{n/m} - W_{1/(mn)}\Big). \tag{24.26}$$

The double sums converge in operator norm for $\sigma > 1$.

Proof. Differentiate (24.14) and (24.15) once more and use Proposition 24.8. Polynomial logarithmic weights are summable against $(mn)^{-\sigma}$ for $\sigma > 1$. □

Corollary 24.10 (The interaction disappears with the Weyl cocycle). *If* $\vartheta = 0$, *then*

$$\Phi_{m,n} = 0 \tag{24.27}$$

for all m, n, *and all pairwise terms in* (24.25)–(24.26) *vanish. Thus the second-order cross-arithmetic second-order term displayed above is generated entirely by the exact Heisenberg cocycle.*

Proof. If $\vartheta = 0$, the phase $\Phi_{m,n}$ in Proposition 24.8 is zero for every pair, so every summand in the displayed second-order cross terms contains a factor $\sin(\Phi_{m,n}/2) = 0$ or the equivalent commutator factor. The pairwise contributions therefore vanish identically. □

25. Exact non-localization inside the arithmetic dynamics

Proposition 25.1 (Schwartz action and distributional transpose)**.** *For $\sigma > 1$, the Hamiltonian $H_{\mu,\sigma,\vartheta}$ maps $\mathcal{S}(\mathbb{R})$ continuously into itself. Its continuous transpose $H^{\times} : \mathcal{S}'(\mathbb{R}) \to \mathcal{S}'(\mathbb{R})$ is defined by*

$$\langle H^{\times} T, \phi \rangle = \langle T, H_{\mu,\sigma,\vartheta} \phi \rangle, \qquad T \in \mathcal{S}'(\mathbb{R}), \quad \phi \in \mathcal{S}(\mathbb{R}). \tag{25.1}$$

Then $H^{\times}$ is continuous on $\mathcal{S}'(\mathbb{R})$.

Proof. For every Schwartz seminorm $p_{a,b}(\phi) = \sup_q |q^a \phi^{(b)}(q)|$, the coordinate action of a Weyl operator gives

$$p_{a,b}(W_n \phi) \le C_{a,b}(1 + |x_n| + |y_n|)^{N_{a,b}} \sum_{j \le b} p_{a+j,b-j}(\phi) \tag{25.2}$$

for suitable constants. Since $x_n, y_n = O(\log n)$ and

$$\sum_{n \ge 2} n^{-\sigma} (1 + \log n)^N < \infty \qquad (\sigma > 1) \tag{25.3}$$

for every fixed N, the Hamiltonian series converges in every Schwartz seminorm and defines a continuous map $\mathcal{S} \to \mathcal{S}$. Therefore (25.1) defines the standard continuous dual action on $\mathcal{S}'$. No termwise extension to all tempered distributions is assumed without this transpose construction. □

Proposition 25.2 (No exact position point is a stationary generalized mode)**.** *Let $\sigma > 1$, $\ell > 0$, and $E_0 > 0$. For no $q_0 \in \mathbb{R}$ and no scalar $E \in \mathbb{C}$ does $\delta_{q_0} \in \mathcal{S}'(\mathbb{R})$ satisfy*

$$H^{\times}_{\mu,\sigma,\vartheta} \delta_{q_0} = E \delta_{q_0}. \tag{25.4}$$

Proof. The transpose action of a Weyl displacement on a point mass is

$$W(x,y)^{\times} \delta_{q_0} = \exp\left(\frac{i}{\hbar} y(q_0 - x/2) \right) \delta_{q_0 - x}, \tag{25.5}$$

$$(W(x,y)^*)^{\times} \delta_{q_0} = \exp\left(-\frac{i}{\hbar} y(q_0 + x/2) \right) \delta_{q_0 + x}. \tag{25.6}$$

The $n = 2$ term therefore contributes nonzero masses at $q_0 \pm \ell \log 2$. No other term produces the same support point as the unique $n = 2$ contribution at either $q_0 - \ell \log 2$ or $q_0 + \ell \log 2$. Since $\mu(2) = -1$, the corresponding coefficient cannot cancel. Hence the support of the left side of (25.4) contains a point distinct from q_0, while the right side is supported only at q_0. □

Remark 25.3 (Relation to exact Heisenberg impossibility)*.* Theorem 3.9 is a universal consequence of the CCR. Proposition 25.2 is more specific: the Möbius Hamiltonian itself does not admit an exactly position-localized stationary mode. The reason is not a measurement uncertainty but the exact translation content of the arithmetic Weyl operators.

26. Literature boundary and novelty status in Version 8

The literature boundary is stated component by component rather than by assigning novelty to the entire package.

26.1. **Known framework.** The use of translations and functional calculus to form zeta operators has a direct literature precedent: Herichi–Lapidus construct spectral operators of the form $\zeta(\partial)$, together with truncated shifts and operator criteria related to RH [12]. Version 8 therefore does not claim novelty for translation representations or Möbius inversion at the untwisted operator level.

The following are classical or direct specializations of classical theory:

(1) the Schrödinger pair, CCR, Weyl operators, and the Robertson inequality for arbitrary observables [1, 6];

(2) projective representations and multipliers of abelian groups, including classification through alternating commutator bicharacters [2, 3];
(3) twisted group and semigroup algebras [9, 4];
(4) classical Möbius inversion, Jordan totients, and additive functions such as Ω [10];
(5) the prime-zeta identity $P(s) = \sum_{r\geq 1} \mu(r) \log \zeta(rs)/r$, its scaled logarithmic singularities, and its natural boundary at $\Re s = 0$ [8];
(6) the weighted variance decomposition used in (11.14).

In particular, Theorem 6.3 is not a new uncertainty principle: it is Robertson's inequality applied to the specific quadratures G_m, G_n.

26.2. **Notebook-specific formulas not located in the targeted search.** The targeted literature search carried out for Version 8 did not locate the following exact formulas as previously published results:

(1) the arithmetic embedding $n \mapsto (\ell \log n, \kappa\Omega(n))$ combined with twisted Möbius inversion;
(2) the resulting coefficients $z_\vartheta(n)$, including the prime-pair and two-prime recurrences;
(3) the exact curvature formula (10.13) and its prime-log variance form;
(4) the curvature Dirichlet determinant $\mathcal{K} = \zeta(S_0 S_2 - S_1^2)/2$;
(5) the local monodromy formula (15.2) with weight $m_\rho J_2(r)/r$;
(6) the multiplicative superadditivity, prime-ray rigidity, and power laws (11.3)–(11.8).

This is a priority statement only in the limited sense "not located in the search performed"; it is not a proof of bibliographic novelty.

26.3. **What is specifically new mathematics inside this manuscript.** Independently of priority, the deductions that are new relative to the standard ingredients listed above are the exact consequences proved here for the chosen arithmetic cocycle: the twisted inverse, its coefficientwise curvature, the closed prime-pair geometry, the determinant Dirichlet series, its local branch weight, and the multiplicative laws of $\mathcal{C}(n)$. By contrast, the ambient Weyl/cohomology machinery and the abstract variance identities are not new.

27. Corrections and extensions carried into Version 8

Remark 27.1. Version 8 incorporates the following mathematical corrections and clarifications: the cocycle is proved nontrivial rather than merely described as noncommutative; no faithfulness of $\Pi_{\sigma,\vartheta}$ is claimed; left and right inverses are constructed separately before being identified; the uniform Neumann half-plane is sharpened to $\sigma > \sigma_*$; the ordered-factorization formula and a two-prime recurrence are included; real-valuedness of $z_\vartheta(n)$ for real ϑ is proved; the deformed Dirichlet series is asserted holomorphic only on its proved open half-plane; the unitary evolution is not claimed to preserve Schwartz space; and the action on tempered distributions is defined by continuous transposition; the signs and supports of the two point-mass transpose formulas are corrected explicitly; and no global rearrangement of conditionally convergent sums over zeta zeros is asserted.

28. Open problems

Question 28.1 (Analytic continuation)**.** Does $\zeta_{\mathrm{H}}(s;\vartheta)$ admit a natural meromorphic continuation from $\Re s \geq 2$ to a larger domain for fixed nonzero ϑ?

Question 28.2 (Functional symmetry)**.** Is there a completion of $\zeta_{\mathrm{H}}(s;\vartheta)$ with a symmetry that reduces at $\vartheta = 0$ to the classical reflection $s \mapsto 1 - s$? A nontrivial answer would be required before using the deformation to say anything substantive about the critical line.

Question 28.3 (Spectral zeros)**.** Can the zero set of a continued ζ_{H} be realized as a spectral or resonance set of an operator built intrinsically from $H_{\mu,\sigma,\vartheta}$, rather than as zeros inserted into a multiplier?

Question 28.4 (Classical-zero rigidity)**.** If an analytic two-variable continuation through $\vartheta = 0$ exists, does the exact evenness in ϑ impose any second-order constraint on the motion of the classical Riemann zeros beyond Proposition 23.5?

Question 28.5 (Choice of the second additive coordinate)**.** Which completely additive arithmetic functions $A(n)$, replacing $\Omega(n)$ in $y_n = \kappa A(n)$, produce inequivalent deformations? The choice $A = \Omega$ is distinguished here by its direct relation to the sign of μ on square-free integers, but a classification is open.

Question 28.6 (Higher deformation curvatures)**.** Determine closed formulas for

$$\frac{d^{2k}}{d\vartheta^{2k}} z_\vartheta(n)\Big|_{\vartheta=0}, \qquad k \geq 2, \tag{28.1}$$

and for their Dirichlet generating series. The fourth-order case is the first test of whether the collapse to finitely many one-prime moments persists or irreducible higher prime geometry appears.

29. Notebook history

Version 8. Added the full analytic audit of $\mathcal{O}$ and $\mathcal{I}$: absolute convergence, differentiability, sum–integral interchange, grouped prime-zeta continuation, all endpoint collisions, interior contour singularities, exact blockwise Möbius cancellation, strict replica energy, radical law, correct distinction between $\mathcal{I}$ and $\mathcal{O}$, and the monodromy detector. Added an abstract and a symbol index, and audited the bibliography and theorem proofs.

Remark 29.1. Version 1 used Möbius-weighted logarithmic translations, Fourier multipliers, an eta multiplier in the critical strip, generalized plane-wave null modes, and a standard CCR non-localization statement. Its main defect was that the arithmetic multiplier and the Heisenberg pair were juxtaposed rather than coupled.

Remark 29.2. Version 2 removes the critical-strip multiplier as the central mechanism and replaces ordinary Dirichlet convolution by the Weyl-cocycle product (7.1). Heisenberg noncommutativity now changes the Möbius inverse coefficients themselves and enters the exact equations of motion of the Möbius Hamiltonian.

Remark 29.3 (Version 8). Version 8 corrects the Version 8 audit issues and adds the gauge-invariance obstruction, the scalar Euler-locality obstruction, the norm-convergent ordered Euler object, its exact defect R_ϑ, the odd Dirichlet defect $\mathcal{O}(s)$, and the Möbius filtering of the leading logarithmic-over-pole term at rational prime-zeta replicas.

30. Literature status of the Version 4/5 formulas

The general ingredients are established mathematics. Twisted products by 2-cocycles and projective representations belong to the standard theory of twisted group algebras; see, for example, Packer–Raeburn [9]. The prime-zeta identity

$$P(s) = \sum_{r\geq 1} \frac{\mu(r)}{r} \log \zeta(rs) \tag{30.1}$$

and the associated logarithmic branch structure and natural boundary are classical; see Fröberg [8].

The specific chain developed here—the Heisenberg arithmetic cocycle based on $(\log n, \Omega(n))$, its twisted Möbius inverse $z_\vartheta(n)$, the closed coefficient curvature (10.13), and its Dirichlet summation (12.3)—is not identified here with a prior theorem. The classical prime-zeta formulas explain the

continuation of the resulting S_j, but they do not by themselves produce the curvature formula or the Hessian identity (21.5).

Remark 30.1. Version 8 replaces the overstrong linear-state no-go claim by the exact sesquilinear statement and adds the exact prime-zeta branching source, Hessian/covariance representation, and positivity obstruction and moves the analysis to the cocycle itself. Differentiating the twisted Möbius inverse at $\vartheta = 0$ produces the prime-factor curvature (10.13). Summing it yields (12.3), whose local continuation detects every zeta zero in $\Re s > 1/2$ through (13.2).

References

[1] H. P. Robertson, *The uncertainty principle*, Physical Review **34** (1929), 163–164. doi:10.1103/PhysRev.34.163.

[2] A. Kleppner, *Multipliers on Abelian groups*, Mathematische Annalen **158** (1965), 11–34.

[3] L. Baggett and A. Kleppner, *Multiplier representations of Abelian groups*, Journal of Functional Analysis **14** (1973), 299–324. doi:10.1016/0022-1236(73)90075-X.

[4] L. Rigal and P. Zadunaisky, *Twisted semigroup algebras*, Algebras and Representation Theory **18** (2015), no. 5, 1155–1186. doi:10.1007/s10468-015-9525-z; arXiv:1406.2985.

[5] W. Heisenberg, *Über den anschaulichen Inhalt der quantentheoretischen Kinematik und Mechanik*, Zeitschrift für Physik **43** (1927), 172–198. doi:10.1007/BF01397280.

[6] G. B. Folland, *Harmonic Analysis in Phase Space*, Annals of Mathematics Studies 122, Princeton University Press, Princeton, 1989.

[7] M. Reed and B. Simon, *Methods of Modern Mathematical Physics, Vol. I: Functional Analysis*, revised and enlarged ed., Academic Press, 1980.

[8] C.-E. Fröberg, *On the prime zeta function*, BIT Numerical Mathematics **8** (1968), 187–202. doi:10.1007/BF01933420.

[9] J. A. Packer and I. Raeburn, *On the structure of twisted group C^*-algebras*, Transactions of the American Mathematical Society **334** (1992), 685–718. doi:10.2307/2154478.

[10] T. M. Apostol, *Introduction to Analytic Number Theory*, Springer, New York, 1976. doi:10.1007/978-1-4757-5579-4.

[11] G. J. Székely and M. L. Rizzo, *Energy statistics: A class of statistics based on distances*, Journal of Statistical Planning and Inference **143** (2013), no. 8, 1249–1272. doi:10.1016/j.jspi.2013.03.018.

[12] H. Herichi and M. L. Lapidus, *Truncated infinitesimal shifts, spectral operators and quantized universality of the Riemann zeta function*, Annales de la Faculté des Sciences de Toulouse, Série 6, **23** (2014), no. 3, 621–664. doi:10.5802/afst.1419.

[13] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2nd ed., revised by D. R. Heath-Brown, Oxford University Press, Oxford, 1986.