

CS 247 – Scientific Visualization

Lecture 24: Vector / Flow Visualization, Pt. 3

Markus Hadwiger, KAUST

Reading Assignment #13 (until Apr 25)



Read (required):

- Data Visualization book
 - Chapter 6.1 (Divergence and Vorticity)
- Diffeomorphisms / smooth deformations

<https://en.wikipedia.org/wiki/Diffeomorphism>

- Integral curves: Stream lines, path lines, streak lines

https://en.wikipedia.org/wiki/Integral_curve

https://en.wikipedia.org/wiki/Streamlines,_streaklines,_and_pathlines

- Paper:

Bruno Jobard and Wilfrid Lefer

Creating Evenly-Spaced Streamlines of Arbitrary Density,

<http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.29.9498>



Quiz #3: Apr 25

Organization

- First 30 min of lecture
- No material (book, notes, ...) allowed

Content of questions


- Lectures (both actual lectures and slides)
- Reading assignments (except optional ones)
- Programming assignments (algorithms, methods)
- Solve short practical examples

Vector Fields




Each vector is usually thought of as a velocity vector

- Example for actual velocity: fluid flow
- But also force fields, etc. (e.g., electrostatic field)



vectors given at grid points



vectors given at particle positions



Vector Fields

Each vector is usually thought of as a velocity vector

- Example for actual velocity: fluid flow
- But also force fields, etc. (e.g., electrostatic field)

Each vector in a vector field
lives in the **tangent space**
of the manifold at that point:

Each vector is a **tangent vector**





image from wikipedia

Vector Fields



Vector fields on general manifolds M (not just Euclidean space)

Tangent space at a point $x \in M$:

$$T_x M$$

Tangent bundle: Manifold of all tangent spaces over base manifold

$$\pi: TM \rightarrow M$$

Vector field: *Section of tangent bundle*

$$s: M \rightarrow TM,$$

$$x \mapsto s(x). \quad \pi(s(x)) = x$$





image from wikipedia

Vector Fields



Vector fields on general manifolds M (not just Euclidean space)

Tangent space at a point $x \in M$:

$$T_x M$$

Tangent bundle: Manifold of all tangent spaces over base manifold

$$\pi: TM \rightarrow M$$

Vector field: *Section of tangent bundle*

$$\mathbf{v}: M \rightarrow TM,$$

$$x \mapsto \mathbf{v}(x).$$

$$\mathbf{v}(x) \in T_x M$$





image from wikipedia

Interlude: Coordinate Charts




Interlude: Coordinate Charts

Coordinate chart

$$\phi: U \subset M \rightarrow \mathbb{R}^n,$$

$$x \mapsto (x^1, x^2, \dots, x^n).$$






Interlude: Coordinate Charts

Coordinate chart

$$\begin{aligned}\phi: U \subset M &\rightarrow \mathbb{R}^n, \\ x &\mapsto (x^1, x^2, \dots, x^n).\end{aligned}$$

Coordinate functions

$$\begin{aligned}x^i: U \subset M &\rightarrow \mathbb{R}, \\ x &\mapsto x^i(x).\end{aligned}$$






Interlude: Coordinate Charts

Coordinate charts

$$\begin{aligned}\phi_\alpha: U_\alpha \subset M &\rightarrow \mathbb{R}^n, \\ x &\mapsto (x^1, x^2, \dots, x^n).\end{aligned}$$

Atlas

$$\{(U_\alpha, \phi_\alpha)\}_{\alpha \in I}$$





Interlude: Coordinate Charts


Coordinate charts

$$\begin{aligned}\phi_\alpha : U_\alpha \subset M &\rightarrow \mathbb{R}^n, \\ x &\mapsto (x^1, x^2, \dots, x^n).\end{aligned}$$

Atlas

$$\{(U_\alpha, \phi_\alpha)\}_{\alpha \in I}$$

$$\begin{aligned}\phi_\alpha : U_\alpha \subset M &\rightarrow \mathbb{R}^n, \\ x &\mapsto (x^1(x), x^2(x), \dots, x^n(x)).\end{aligned}$$





Vector Fields vs. Vectors in Components

Because Euclidean space is most common, often slightly sloppy notation

$$\mathbf{v}: U \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2,$$

$$(x, y) \mapsto \begin{bmatrix} u \\ v \end{bmatrix}.$$



Vector Fields vs. Vectors in Components

Because Euclidean space is most common, often slightly sloppy notation

$$\mathbf{v}: U \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2,$$

$$(x, y) \mapsto \begin{bmatrix} u \\ v \end{bmatrix}.$$

$$\mathbf{v}: U \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2,$$

$$(x, y) \mapsto \begin{pmatrix} u(x, y) \\ v(x, y) \end{pmatrix}.$$



Vector Fields vs. Vectors in Components

Because Euclidean space is most common, often slightly sloppy notation

$$\mathbf{v}: U \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2,$$

$$(x, y) \mapsto \begin{bmatrix} u \\ v \end{bmatrix}.$$

$$\mathbf{v}: U \subset \mathbb{R}^3 \rightarrow \mathbb{R}^3,$$

$$(x, y, z) \mapsto \begin{bmatrix} u \\ v \\ w \end{bmatrix}.$$

$$\mathbf{v}: U \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2,$$

$$(x, y) \mapsto \begin{pmatrix} u(x, y) \\ v(x, y) \end{pmatrix}.$$

$$\mathbf{v}: U \subset \mathbb{R}^3 \rightarrow \mathbb{R}^3,$$

$$(x, y, z) \mapsto \begin{pmatrix} u(x, y, z) \\ v(x, y, z) \\ w(x, y, z) \end{pmatrix}.$$



Vector Fields vs. Vectors in Components

$\mathbf{v}: U \subset \mathbb{R}^n \rightarrow \mathbb{R}^n,$

$$(x^1, x^2, \dots, x^n) \mapsto \begin{bmatrix} v^1 \\ v^2 \\ \vdots \\ v^n \end{bmatrix}.$$


$\mathbf{v}: U \subset \mathbb{R}^n \rightarrow \mathbb{R}^n,$

$$(x^1, x^2, \dots, x^n) \mapsto \begin{pmatrix} v^1(x^1, x^2, \dots, x^n) \\ v^2(x^1, x^2, \dots, x^n) \\ \vdots \\ v^n(x^1, x^2, \dots, x^n) \end{pmatrix}.$$



Vector Fields vs. Vectors in Components

$$\mathbf{v}: U \subset \mathbb{R}^n \rightarrow \mathbb{R}^n,$$
$$(x^1, x^2, \dots, x^n) \mapsto \begin{bmatrix} v^1 \\ v^2 \\ \vdots \\ v^n \end{bmatrix}.$$



$$\mathbf{v}: U \subset \mathbb{R}^n \rightarrow \mathbb{R}^n,$$
$$(x^1, x^2, \dots, x^n) \mapsto \begin{pmatrix} v^1(x^1, x^2, \dots, x^n) \\ v^2(x^1, x^2, \dots, x^n) \\ \vdots \\ v^n(x^1, x^2, \dots, x^n) \end{pmatrix}.$$

$$\mathbf{v}|_U: \phi(U) \subset \mathbb{R}^n \rightarrow \mathbb{R}^n,$$
$$(x^1, x^2, \dots, x^n) \mapsto \begin{bmatrix} v^1 \\ v^2 \\ \vdots \\ v^n \end{bmatrix}.$$



Vector Fields vs. Vectors in Components

Need basis vector fields

$$\begin{aligned}\mathbf{e}_i : U \subset M &\rightarrow TM, \\ x &\mapsto \mathbf{e}_i(x)\end{aligned}\quad \left\{\mathbf{e}_i(x)\right\}_{i=1}^n \text{ basis for } T_x M$$



Vector Fields vs. Vectors in Components

Need basis vector fields

$$\begin{aligned}\mathbf{e}_i : U \subset M &\rightarrow TM, \\ x &\mapsto \mathbf{e}_i(x)\end{aligned}\quad \left\{\mathbf{e}_i(x)\right\}_{i=1}^n \text{ basis for } T_x M$$

$$\begin{aligned}\mathbf{v} : U \subset M &\rightarrow TM, \\ x &\mapsto v^1 \mathbf{e}_1 + v^2 \mathbf{e}_2 + \dots + v^n \mathbf{e}_n.\end{aligned}$$

$$\begin{aligned}\mathbf{v} : U \subset M &\rightarrow TM, \\ x &\mapsto v^1(x) \mathbf{e}_1(x) + v^2(x) \mathbf{e}_2(x) + \dots + v^n(x) \mathbf{e}_n(x).\end{aligned}$$



Vector Fields vs. Vectors in Components

Need basis vector fields

$$\mathbf{e}_i : U \subset M \rightarrow TM, \quad x \mapsto \mathbf{e}_i(x) \quad \{\mathbf{e}_i(x)\}_{i=1}^n \text{ basis for } T_x M$$

Coordinate basis:

$$\mathbf{e}_i := \frac{\partial}{\partial x^i}$$

$$\mathbf{v} : U \subset M \rightarrow TM, \quad x \mapsto v^1 \mathbf{e}_1 + v^2 \mathbf{e}_2 + \dots + v^n \mathbf{e}_n.$$

$$\mathbf{v} : U \subset M \rightarrow TM, \quad x \mapsto v^1(x) \mathbf{e}_1(x) + v^2(x) \mathbf{e}_2(x) + \dots + v^n(x) \mathbf{e}_n(x).$$

Examples of Coordinate Curves and Bases



Coordinate functions, coordinate curves, bases

- Coordinate functions are real-valued (“scalar”) functions on the domain
- On each coordinate curve, *one* coordinate changes, *all others stay constant*
- Basis: n linearly independent vectors at each point of domain


Cartesian coordinates

$$x^1 = x$$

$$x^2 = y$$

$$\mathbf{e}_1 = \frac{\partial}{\partial x} = \mathbf{i}$$

$$\mathbf{e}_2 = \frac{\partial}{\partial y} = \mathbf{j}$$




polar coordinates

$$x^1 = r$$

$$x^2 = \theta$$

$$\mathbf{e}_1 = \frac{\partial}{\partial r} = \mathbf{e}_r$$

$$\mathbf{e}_2 = \frac{\partial}{\partial \theta} = \mathbf{e}_\theta$$







Flow Field Example (1)

Potential flow around a circular cylinder

https://en.wikipedia.org/wiki/Potential_flow_around_a_circular_cylinder

Inviscid, incompressible flow that is irrotational (curl-free) and can be modeled as the gradient of a scalar function called the (scalar) velocity potential



images from wikipedia






Flow Field Example (2)

Depending on Reynolds number, turbulence will develop

Example: von Kármán vortex street: vortex shedding

https://en.wikipedia.org/wiki/Karman_vortex_street



images from wikipedia



Steady vs. Unsteady Flow

- Steady flow: time-independent
 - Flow itself is static over time: $\mathbf{v}(\mathbf{x})$ $\mathbf{v}: \mathbb{R}^n \rightarrow \mathbb{R}^n,$
 - Example: laminar flows $x \mapsto \mathbf{v}(x).$
- Unsteady flow: time-dependent
 - Flow itself changes over time: $\mathbf{v}(\mathbf{x}, t)$ $\mathbf{v}: \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n,$
 - Example: turbulent flows $x \mapsto \mathbf{v}(x, t).$

(here just for Euclidean domain; analogous on general manifolds)




Direct vs. Indirect Flow Visualization

- Direct flow visualization
 - Overview of current flow state
 - Visualization of vectors: arrow plots (“hedgehog” plots)
- Indirect flow visualization
 - Use intermediate representation: vector field integration over time
 - Visualization of temporal evolution
 - Integral curves: streamlines, pathlines, streaklines, timelines
 - Integral surfaces: streamsurfaces, pathsurfaces, streaksurfaces



Direct vs. Indirect Flow Visualization




Integral Curves: Intro

Integral Curves / Stream Objects



Integrating velocity over time yields spatial motion




Particle Trajectories



Courtesy Jens Krüger




Particle Trajectories



Courtesy Jens Krüger




Particle Trajectories




Courtesy Jens Krüger

Particle Trajectories



Courtesy Jens Krüger

Integral Curves



Streamline

- Curve parallel to the vector field in each point for a fixed time

Pathline

- Describes motion of a massless particle over time

Streakline

- Location of all particles released at a *fixed position* over time

Timeline


- Location of all particles released along a line at a *fixed time*



Streamlines Over Time

Defined only for steady flow or for a fixed time step (of unsteady flow)

Different tangent curves in every time step for time-dependent vector fields (unsteady flow)




Stream Lines vs. Path Lines Viewed Over Time




Plotted with time as third dimension

- Tangent curves to a $(n + 1)$ -dimensional vector field




Stream Lines



Path Lines

Numerical Integration

- Numerical integration of stream lines:
- approximate streamline by polygon \mathbf{x}_i
- Testing example:
 - $\mathbf{v}(x,y) = (-y, x/2)^T$
 - exact solution: ellipses
 - starting integration from $(0,-1)$





Streamlines – Practice

■ Basic approach:

- theory: $\mathbf{s}(t) = \mathbf{s}_0 + \int_{0 \leq u \leq t} \mathbf{v}(\mathbf{s}(u)) du$
- practice: numerical integration
- idea:
(very) locally, the solution is (approx.) linear
- Euler integration:
follow the current flow vector $\mathbf{v}(\mathbf{s}_i)$ from the current streamline point \mathbf{s}_i for a very small time (dt) and therefore distance
- Euler integration: $\mathbf{s}_{i+1} = \mathbf{s}_i + dt \cdot \mathbf{v}(\mathbf{s}_i)$,
integration of small steps (dt very small)


Euler Integration – Example

- 2D model data:

$$v_x = dx/dt = -y$$
$$v_y = dy/dt = x/2$$


- Sample arrows:

- True solution: ellipses!




Euler Integration – Example

- Seed point $s_0 = (0|-1)^T$;
current flow vector $v(s_0) = (1|0)^T$;
 $dt = 1/2$




Euler Integration – Example

- New point $\mathbf{s}_1 = \mathbf{s}_0 + \mathbf{v}(\mathbf{s}_0) \cdot dt = (1/2 | -1)^T$;
 current flow vector $\mathbf{v}(\mathbf{s}_1) = (1 | 1/4)^T$;




Euler Integration – Example

- New point $\mathbf{s}_2 = \mathbf{s}_1 + \mathbf{v}(\mathbf{s}_1) \cdot dt = (1 | -7/8)^T$;
current flow vector $\mathbf{v}(\mathbf{s}_2) = (7/8 | 1/2)^T$;




Euler Integration – Example

- $\mathbf{s}_3 = (23/16 | -5/8)^T \approx (1.44 | -0.63)^T;$
- $\mathbf{v}(\mathbf{s}_3) = (5/8 | 23/32)^T \approx (0.63 | 0.72)^T;$




Euler Integration – Example

- $s_4 = (7/4 \mid -17/64)^T \approx (1.75 \mid -0.27)^T;$
- $v(s_4) = (17/64 \mid 7/8)^T \approx (0.27 \mid 0.88)^T;$




Euler Integration – Example

- $s_9 \approx (0.20 | 1.69)^T;$
 $v(s_9) \approx (-1.69 | 0.10)^T;$




Euler Integration – Example

- $s_{14} \approx (-3.22 | -0.10)^T;$
 $v(s_{14}) \approx (0.10 | -1.61)^T;$




Euler Integration – Example

- $\mathbf{s}_{19} \approx (0.75|-3.02)^T$; $\mathbf{v}(\mathbf{s}_{19}) \approx (3.02|0.37)^T$;
clearly: large integration error, dt too large!
19 steps




Euler Integration – Example

- dt smaller ($1/4$): more steps, more exact!
 $\mathbf{s}_{36} \approx (0.04|-1.74)^T$; $\mathbf{v}(\mathbf{s}_{36}) \approx (1.74|0.02)^T$;
- 36 steps



Comparison Euler, Step Sizes

Euler
is getting
better
proportionally
to dt





Better than Euler Integr.: RK

■ Runge-Kutta Approach:

- theory: $\mathbf{s}(t) = \mathbf{s}_0 + \int_{0 \leq u \leq t} \mathbf{v}(\mathbf{s}(u)) du$

- Euler: $\mathbf{s}_i = \mathbf{s}_0 + \sum_{0 \leq u < i} \mathbf{v}(\mathbf{s}_u) \cdot dt$

- Runge-Kutta integration:

 - idea: cut short the curve arc

 - RK-2 (second order RK):

 - 1.: do half a Euler step

 - 2.: evaluate flow vector there


 - 3.: use it in the origin

 - RK-2 (two evaluations of \mathbf{v} per step):

$$\mathbf{s}_{i+1} = \mathbf{s}_i + \mathbf{v}(\mathbf{s}_i + \mathbf{v}(\mathbf{s}_i) \cdot dt/2) \cdot dt$$


RK-2 Integration – One Step

- Seed point $s_0 = (0|-2)^T$;
 current flow vector $v(s_0) = (2|0)^T$;
 preview vector $v(s_0 + v(s_0) \cdot dt/2) = (2|0.5)^T$;
 $dt = 1$




RK-2 – One more step

- Seed point $s_1 = (2|-1.5)^T$;
 current flow vector $v(s_1) = (1.5|1)^T$;
 preview vector $v(s_1 + v(s_1) \cdot dt/2) \approx (1|1.4)^T$;
 $dt = 1$



RK-2 – A Quick Round

- RK-2: even with $dt=1$ (9 steps)
better
than Euler
with $dt=1/8$
(72 steps)






RK-4 vs. Euler, RK-2

- Even better: fourth order RK:
 - four vectors \mathbf{a} , \mathbf{b} , \mathbf{c} , \mathbf{d}
 - one step is a convex combination:
$$\mathbf{s}_{i+1} = \mathbf{s}_i + (\mathbf{a} + 2\cdot\mathbf{b} + 2\cdot\mathbf{c} + \mathbf{d})/6$$
 - vectors:
 - $\mathbf{a} = dt \cdot \mathbf{v}(\mathbf{s}_i)$... original vector
 - $\mathbf{b} = dt \cdot \mathbf{v}(\mathbf{s}_i + \mathbf{a}/2)$... RK-2 vector
 - $\mathbf{c} = dt \cdot \mathbf{v}(\mathbf{s}_i + \mathbf{b}/2)$... use RK-2 ...
 - $\mathbf{d} = dt \cdot \mathbf{v}(\mathbf{s}_i + \mathbf{c})$... and again!

Euler vs. Runge-Kutta

- RK-4: pays off only with complex flows
- Here approx. like RK-2





■ Summary:

- analytic determination of streamlines
usually not possible
- hence: numerical integration
- several methods available
(Euler, Runge-Kutta, etc.)
- Euler: simple, imprecise, esp. with small dt
- RK: more accurate in higher orders
- furthermore: adaptive methods, implicit methods, etc.



Bonus Slides: Vectors as Derivative Operators



Vectors as Derivative Operators

A vector applied to a (real) function on the manifold gives the *directional derivative* in that direction

- From this viewpoint, the vector is a derivative operator (actually, a *derivation*)
- Can be used as *definition* of a vector (must fulfill props. of a derivation; esp. Leibniz rule)

$$f: M \rightarrow \mathbb{R}, \quad \mathbf{v} f \\ x \mapsto f(x).$$



Vectors as Derivative Operators

A vector applied to a (real) function on the manifold gives the *directional derivative* in that direction

- From this viewpoint, the vector is a derivative operator (actually, a *derivation*)
- Can be used as *definition* of a vector (must fulfill props. of a derivation; esp. Leibniz rule)

$$f: M \rightarrow \mathbb{R}, \quad \mathbf{v} f := df(\mathbf{v}) \\ x \mapsto f(x).$$



Vectors as Derivative Operators

A vector applied to a (real) function on the manifold gives the *directional derivative* in that direction

- From this viewpoint, the vector is a derivative operator (actually, a *derivation*)
- Can be used as *definition* of a vector (must fulfill props. of a derivation; esp. Leibniz rule)

$$f: M \rightarrow \mathbb{R}, \quad \mathbf{v} f := df(\mathbf{v}) \quad \mathbf{e}_i f := df(\mathbf{e}_i)$$
$$x \mapsto f(x).$$



Vectors as Derivative Operators

A vector applied to a (real) function on the manifold gives the *directional derivative* in that direction

- From this viewpoint, the vector is a derivative operator (actually, a *derivation*)
- Can be used as *definition* of a vector (must fulfill props. of a derivation; esp. Leibniz rule)

$$f: M \rightarrow \mathbb{R}, \quad \mathbf{v}f := df(\mathbf{v}) \quad \mathbf{e}_i f := df(\mathbf{e}_i)$$

$$x \mapsto f(x).$$

$$\frac{\partial}{\partial x^i} f = df \left(\frac{\partial}{\partial x^i} \right) = \frac{\partial f}{\partial x^i}$$



Vectors as Derivative Operators

A vector applied to a (real) function on the manifold gives the *directional derivative* in that direction

- From this viewpoint, the vector is a derivative operator (actually, a *derivation*)
- Can be used as *definition* of a vector (must fulfill props. of a derivation; esp. Leibniz rule)

$$f: M \rightarrow \mathbb{R}, \quad x \mapsto f(x).$$

$$\mathbf{v}f := df(\mathbf{v})$$

$$\mathbf{e}_i f := df(\mathbf{e}_i)$$

$$\frac{\partial}{\partial x^i} f = df \left(\frac{\partial}{\partial x^i} \right) = \frac{\partial f}{\partial x^i}$$

$$\frac{\partial}{\partial x^i} x^j = dx^j \left(\frac{\partial}{\partial x^i} \right) = \delta_i^j$$

Kronecker delta
("identity matrix")




Vectors as Derivative Operators

A vector applied to a (real) function on the manifold gives the *directional derivative* in that direction

- From this viewpoint, the vector is a derivative operator (actually, a *derivation*)
- Can be used as *definition* of a vector (must fulfill props. of a derivation; esp. Leibniz rule)

$$f: M \rightarrow \mathbb{R}, \quad x \mapsto f(x).$$

$$\mathbf{v}f := df(\mathbf{v})$$

$$\mathbf{e}_i f := df(\mathbf{e}_i)$$

$$\frac{\partial}{\partial x^i} f = df \left(\frac{\partial}{\partial x^i} \right) = \frac{\partial f}{\partial x^i} \quad \frac{\partial}{\partial x^i} x^j = dx^j \left(\frac{\partial}{\partial x^i} \right) = \delta_i^j$$

For vector field: obtain directional derivative at each point

Kronecker delta
("identity matrix")

$$\mathbf{v}f: M \rightarrow \mathbb{R},$$

$$x \mapsto \mathbf{v}(x) f = df(\mathbf{v}(x)).$$

(remember that this just looks scary (maybe) ...)

Thank you.

Thanks for material

- Helwig Hauser
- Eduard Gröller
- Daniel Weiskopf
- Torsten Möller
- Ronny Peikert
- Philipp Muigg
- Christof Rezk-Salama