

# Univariate Time Series Data and Model Card

Generated by [Cardtale](#) - Automated Model and Data Card Generator

This report provides an automated, comprehensive analysis of univariate time series data. Generated by Cardtale, it explores basic aspects and potential challenges in your data to support informed decision-making and modeling choices.

**Generated:** 2024-11-01 09:39

**Series Name:** M100

## Table of Contents

1	<b>Data Overview</b>	Time series fundamental characteristics and statistical properties
2	<b>Trend</b>	Long-term time series growth and dynamics. Analysis of level stabilization methods.
3	<b>Seasonality</b>	Analysing recurring patterns in the time series. Assessing the impact of different seasonality modeling strategies
4	<b>Variance</b>	Exploring the variability of values over time. Assessing the impact of variance stabilization methods
5	<b>Change Detection</b>	Change detection in the time series distribution

Other aspects were explored but omitted from the final report:

## Data Overview

This section examines the core characteristics and statistical properties of the time series. Understanding these attributes is important for assessing data quality and

gaining a preliminary context. We explore the temporal structure, summary statistics, and distribution patterns to create a baseline understanding of your data.

## Time Series Plot




Figure 1: Time series line plot.

- A total of 69 month observations which span from January 1990 to September 1995.
- The mean value of the series is 6500.43 (median equal to 6230), with a standard deviation of 1294.05. The data ranges from a minimum of 4360 to a maximum of 11700.

## Data Distribution




Figure 2: Distribution of the time series using an histogram (left) and a boxplot (right).

- The Kolmogorov-Smirnov test rejects the hypothesis that the series is distributed according to the following distributions: Exponential, Pareto, Power-law, and Chi-squared
- The distribution with largest p-value is Gamma (p-value equal to 0.91). But, we cannot reject the hypothesis that the data follows the following distributions

(ordered decreasingly by p-value): Log-Normal, Exponentially Modified Gaussian distribution, Logistic, Gaussian, and Cauchy.

- There are 1 outliers in the data, all of which are upper outliers. The outliers represent 1.45% of the complete data set.
- The skewness is equal to 1.14, indicates that the right tail is long relative to the left tail.
- The excess kurtosis is equal to 2.24. This indicates that the data has a heavy tailed distribution.

## Trend and Seasonality




Figure 3: Seasonal and Trend components after decomposition using the STL (Season-Trend decomposition using LOESS) method.

- All hypothesis tests carried out (Augmented Dickey-Fuller and Philips-Perron) indicate that the time series is stationary in trend/level.
- All hypothesis tests carried out (Wang-Smith-Hyndman and OCSB) indicate that the time series is not stationary in seasonality for the specified period.

## Auto-Correlation




Figure 4: Auto-correlation plot up to 24 lags.

- The following lags show significant autocorrelation: t-1, t-2, t-3, t-4, t-5, t-6, t-7, t-8, t-9, t-10, t-11, and t-12. The autocorrelation is positive for all lags with a significant value.
- None of the lags relative to the seasonal period (t-12 and t-24) show any significant autocorrelation.

## Partial Auto-Correlation




Figure 5: Partial Auto-correlation plot up to 24 lags. At each lag, the partial auto-correlation takes into account the previous correlations.

- The following lags show significant partial autocorrelation: t-1, t-2, t-13, and t-23.
- None of the lags relative to the seasonal period (t-12 and t-24) show any significant partial autocorrelation.

## Trend

Trend refers to the long-term change in the mean level of a time series. It reflects systematic and gradual changes in the data over time. Understanding the trend is important for identifying long-term growth or decline, structural changes, and making informed modeling decisions. This section examines the characteristics of the trend of the time series.

## Trend Line Plot




Figure 6: Time series trend plot.

- The time series has non-stationary trend according to the statistical test(s): . There is a strong downward trend. On the other hand, the method's Augmented Dickey-Fuller and Philips-Perron did not find evidence for the presence of trend.
- The same tests were applied to analyse whether the time series is stationary around a constant level. The method(s) Augmented Dickey-Fuller reject this hypothesis. But, the test(s) Philips-Perron fail to reject.
- Including a trend explanatory variable which denotes the position of each observation does not improve forecasting performance.

## Distribution of Differences




Figure 7: Distribution of percentage changes (left), and a Lag-plot (right). These plots help to understand how the data changes over consecutive observations. The histogram shown the distribution of these changes. The lag-plot depicts the randomness in the data. The time series shows greater randomness as the points deviate from the dotted line.

- The Kolmogorov-Smirnov test rejects the hypothesis that the differenced series is distributed according to the following distributions: Power-law, Exponential, Pareto, Log-Normal, and Chi-squared.

- The distribution with largest p-value is Logistic (p-value equal to 0.98). But, we cannot reject the hypothesis that the differenced series follows the following distributions (ordered decreasingly by p-value): Cauchy, Exponentially Modified Gaussian distribution, Gaussian, and Gamma.
- The skewness of the differenced series is equal to -0.4, which is close to zero. This indicates a symmetric distribution, though there is a slight left skewness.
- The excess kurtosis of the differenced series is equal to 1.76. This indicates that the data has a heavy tailed distribution.
- **\*\*Forecasting experiments\*\***: Taking first differences improves forecasting performance.

## Seasonality

Seasonality represents recurring patterns or cycles that appear at regular intervals in time series data. These are predictable fluctuations that reflect periodic influences such as monthly, quarterly, or yearly cycles. Understanding seasonal patterns is crucial for forecasting, trend analysis, and identifying anomalies. This section examines the presence, strength, and characteristics of seasonal components in the input time series.

### Seasonal Line Plot (Monthly)




Figure 8: Seasonal plot of monthly values grouped by year.

- All hypothesis tests carried out (Wang-Smith-Hyndman and OCSB) indicate that the time series is not stationary in seasonality for a yearly period.
- **\*\*Forecasting experiments\*\***: Including monthly information in the predictive model decreases forecasting performance. This information was included as Fourier terms and repeating basis function terms in the explanatory variables.

## Seasonal Sub-series Plot (Monthly)




Figure 9: Monthly seasonal sub-series. This plot helps to understand how the data varies within and across monthly groups.

- Statistical tests were carried out to check for differences among means and variances across months. No significant differences were found.
- Overall, there is a strong evidence that the time series is not stationary around a constant level. But, the data is constant around a level in each Month.
- **\*\*Forecasting experiments\*\***: There is evidence for a yearly seasonal pattern from statistical tests. Yet, including information about this period in the forecasting model decreased its performance.

## Seasonal Sub-series Plot (Quarterly)




Figure 10: Quarterly seasonal sub-series. This plot helps to understand how the data varies within and across quarterly groups.

- Statistical tests were carried out to check for differences among means and variances across quarters. No significant differences were found.

- The following tests indicate that the time series is non-stationary in seasonality for a Quarterly period: OCSB. On the other hand, other tests (Wang-Smith-Hyndman) fail to reject the stationary null hypothesis.
- Overall, there is a strong evidence that the time series is not stationary around a constant level. But, the data is constant around a level in each Quarter.
- **\*\*Forecasting experiments\*\***: There is evidence for a quarterly seasonal pattern from statistical tests. Besides, including information about this period in the forecasting model improved its performance.

## Variance

Variance measures how data points spread around the average value in your time series. This section examines whether the variability remains stable (homoskedastic) or changes (heteroskedastic) over time. Understanding variance patterns is crucial for selecting appropriate modeling techniques, which can have a significant impact on forecasting accuracy.

## Heteroskedasticity Testing




Figure 11: Time series residuals analysis. Difference in the distribution of the residuals (left) and the series (right) in the first and last thirds of the series, following a Goldfeld-Quandt partition.

- In the analysis of seasonality we did not find significant differences in the dispersion among periodic groups of observations.
- The following tests suggest that the time series is heteroskedastic: White and Breusch-Pagan. But, other tests (Goldfeld-Quandt) fail to reject the hypothesis that the time series has a constant variance.
- **\*\*Forecasting experiments\*\***: Transforming the series with either the logarithm or the Box-Cox method did not improve forecasting performance.



- In the original scale, the Logarithm distribution is a reasonable fit to the data. But, after taking the Gamma transformation, a Chi-squared distribution was found to be a better fit.

## Change Detection

Change points denote significant shifts in the underlying distribution of time series. These structural changes can manifest as sudden shifts in level, trend, variance, or seasonal patterns. Detecting and understanding these points is crucial as they often indicate important events or regime changes that affect modeling decisions. This section identifies potential change points and assesses their impact on the overall analysis strategy.

### Change Points




Figure 12: Time series plot with marked change points according to the PELT method.

- A single change point was found in the time series.
- The change point was found at February 1992 where the time series shows an decreasing tendency.

## Changes in Distribution




Figure 13: Time series analysis before and after the first detected change point occurs. Paired distributions before and after change (left), and overlapped density plot (right).

- Before the change point, the data follows a Logistic distribution. But, after the first change point, a Gamma distribution is a better fit.