A Dialectica Model of the Lambek Calculus (DRAFT)

Valeria de Paiva

This note discusses the Lambek Calculus ([Lam58], [vBen88]) and some variants from the point-of-view of Linear Logic ([Gir87]). The Lambek Calculus was first introduced, with the name of Syntactic Calculus, by J. Lambek in 1958 ([Lam58]) as an explanation of the mathematics of sentence structure. After a long period of ostracism, around 1980, the Syntactic Calculus, now called the Lambek Calculus, was taken up by logicians interested in Computational Linguistics, especially in the area of Categorial Grammar. The calculus was considered "like a logical system" but much too weak to be taken seriously as a logic. To quote from [vBen88],

> ...One fundamental system of this kind is the so-called 'Lambek Calculus', whose type-change rules show a close analogy with the inference rules of constructive propositional logic.

This situation could change drastically with the advent of Girard's Linear Logic [Gir'87]. Girard showed that there is a full embedding, preserving proofs, of Intuitionistic Logic into Linear Logic with modality '!'. That means, in loose terms, that Linear Logic is a logic[1] as expressive as Intuitionistic Logic and the claim is that it is more precise, i.e it makes finer distinctions. As the Lambek Calculus is the (multiplicative) fragment of non-commutative Intuitionistic Linear Logic, several interesting questions can be asked. One of them, posed by Morrill et al in [Mor] is whether we can extend the Lambek Calculus with a modality that does for the structural rule of (*exchange*) what the modality '!' does for the rules of (*weakening*) and (*contraction*). A very preliminary proposal, which answers this question affirmatively, is set forward in this paper. But it must be said from the start that the 'answer' is only provided in semantical terms. The Proof Theory of the systems considered should be investigated in future work. Another warning is that the perspective of this note is basically from Category Theory as a branch of Mathematics, so words like categories and functors are always meant in their mathematical, rather than linguistical or philosophical sense.

We first recall Linear Logic and provide the transformations to show that the Lambek Calculus **L** really is the multiplicative fragment of (non-commutative) Intuitionistic Linear Logic. In the second section we describe the usual String Semantics for the Lambek Calculus **L** and generalise it, using a categorical perspective. In the third section we describe our Dialectica model for the Lambek Calculus. In the last section we discuss modalities and some 'untidiness' of the Curry-Howard correspondence for the fragments of Linear Logic in question.

I would like to thank Jan van Eijck for inviting me to give the talk that became this note, thereby gently 'forcing' me to think about the subject, as well as, for his generous hospitality. I also would like to thank Martin Hyland, Harold Schellinx, Dirk Roorda, Mark Hepple, Glyn Morrill and Michael Moortgat for several useful discussions. Many of the ideas in this paper have been shaped by these discussions, but of course the mistakes are all mine. Finally I want to thank Jim Lambek for 'putting me right' about how completeness has nothing to do with the existence of two disjunctions, in the most friendly possible way.

1 From Linear Logic to the Lambek Calculus

Intuitionistic Linear Logic, henceforth **ILL**, was described in Girard and Lafont's TAPSOFT [G/L] paper. One of the best ways of thinking about the system **ILL**

Intuitionistic Linear Logic (ILL)

Axioms:

$$A \vdash A \quad (identity)$$
$$\vdash I$$
$$\Gamma \vdash 1 \qquad \Gamma, 0 \vdash A$$

Structural Rules:

$$\frac{\Gamma, A, B, \Gamma' \vdash C}{\Gamma, B, A, \Gamma' \vdash C} \quad (permutation) \qquad \frac{\Gamma \vdash A \qquad A, \Gamma' \vdash B}{\Gamma, \Gamma' \vdash B} \quad (cut)$$

Logical Rules:
Multiplicatives:

$$(unit)\frac{\Gamma \vdash A}{\Gamma, I \vdash A}$$

$$(\otimes_l)\frac{\Gamma, A, B \vdash C}{\Gamma, A \otimes B \vdash C} \qquad (\otimes_r)\frac{\Gamma \vdash A \qquad \Gamma' \vdash B}{\Gamma, \Gamma' \vdash A \otimes B}$$

$$(-\circ_l)\frac{\Gamma \vdash A \qquad \Gamma', B \vdash C}{\Gamma, \Gamma', A -\circ B \vdash C} \qquad (-\circ_r)\frac{\Gamma, A \vdash B}{\Gamma \vdash A -\circ B}$$

Additives:

$$(\&_r)\frac{\Gamma \vdash A \qquad \Gamma \vdash B}{\Gamma \vdash A \& B} \qquad (\&_l)\frac{\Gamma, A \vdash C}{\Gamma, A \& B \vdash C} \qquad \frac{\Gamma, B \vdash C}{\Gamma, A \& B \vdash C}$$

$$(\oplus_l)\frac{\Gamma, A \vdash C \qquad \Gamma, B \vdash C}{\Gamma, A \oplus B \vdash C} \qquad (\oplus_r)\frac{\Gamma \vdash A}{\Gamma \vdash A \oplus B} \qquad \frac{\Gamma \vdash B}{\Gamma \vdash A \oplus B}$$

A difference between sequent-calculus for traditional intuitionistic logic and the one above is that one thinks of Γ as a set in the traditional case, while here Γ means a multiset (or a bag) of assumptions.

Note that in the multiplicative fragment, we can put different contexts Γ and Γ' together - rules $(\otimes_r)$ and $(-\circ_l)$ - while in the additive fragment, we must have exactly the same context Γ - rules $(\&_r)$ and $(\oplus_r)$ - to apply a rule. Observe that, as the system **ILL** satisfies Cut-elimination, there is only one structural rule left, namely (*permutation*). Also in **ILL** we have only one (additive) disjunction '$\oplus$', which contrasts with Classical Linear Logic [Gir] or Full Intuitionistic Linear Logic [HdP].

If we delete the rules for the additives (and their axioms) from the system above, we have the system **LP** studied by van Benthem and Moorgat ([vBen], [Moo]), that is the Lambek Calculus with Permutation. Note that this system could also be called $\mathbf{ILL}_m$, that is the multiplicative fragment of **ILL**. For the sake of completeness we list the axioms and rules of **LP**.

Lambek Calculus with Permutation (LP)

Axioms:

$$A \vdash A \quad (identity)$$
$$\vdash I$$

Structural Rules:

$$\frac{\Gamma, A, B, \Gamma' \vdash C}{\Gamma, B, A, \Gamma' \vdash C} \quad (permutation) \qquad \frac{\Gamma \vdash A \qquad A, \Gamma' \vdash B}{\Gamma, \Gamma' \vdash B} \quad (cut)$$

Logical Rules:

$$(unit)\frac{\Gamma \vdash A}{\Gamma, I \vdash A}$$

$$(\otimes_l)\frac{\Gamma, A, B \vdash C}{\Gamma, A \otimes B \vdash C} \qquad (\otimes_r)\frac{\Gamma \vdash A \qquad \Gamma' \vdash B}{\Gamma, \Gamma' \vdash A \otimes B}$$

$$(-\circ_l)\frac{\Gamma \vdash A \qquad \Gamma', B \vdash C}{\Gamma, \Gamma', A -\circ B \vdash C} \qquad (-\circ_r)\frac{\Gamma, A \vdash B}{\Gamma \vdash A -\circ B}$$

Now if we remove the last structural rule of (*permutation*), which means "the commutativity of $\otimes$", the rules for '$-\circ$' can be given in two (now different) guises, eg

$$(\rightharpoonup_r)\frac{\Gamma, A \vdash B}{\Gamma \vdash A \rightharpoonup B} \qquad (\leftharpoondown_r)\frac{A, \Gamma \vdash B}{\Gamma \vdash A \leftharpoondown B}$$

Thus we get two directional implications and the system is called the Lambek Calculus **L**. The rules and axioms of **L** are as follows:

Lambek Calculus (L)

Axioms:

$$A \vdash A \quad (identity)$$
$$\vdash I$$

Structural Rule:

$$\frac{\Gamma \vdash A \qquad \Gamma', A, \Delta \vdash B}{\Gamma', \Gamma, \Delta \vdash B} \quad (cut)$$

Logical Rules:

$$(unit)\frac{\Gamma \vdash A}{\Gamma', I, \Gamma'' \vdash A} \quad {\scriptstyle (\Gamma', \Gamma'' = \Gamma)}$$

$$(\otimes_l)\frac{\Gamma, A, B, \Gamma' \vdash C}{\Gamma, A \otimes B, \Gamma' \vdash C} \qquad (\otimes_r)\frac{\Gamma \vdash A \qquad \Gamma' \vdash B}{\Gamma, \Gamma' \vdash A \otimes B}$$

$$(\rightharpoonup_l)\frac{\Gamma \vdash A \qquad \Gamma', B, \Delta \vdash C}{\Gamma', A \rightharpoonup B, \Gamma, \Delta \vdash C} \qquad (\rightharpoonup_r)\frac{\Gamma, A \vdash B}{\Gamma \vdash A \rightharpoonup B}$$

$$(\leftharpoondown_l)\frac{\Gamma \vdash A \qquad \Gamma', B, \Delta \vdash C}{\Gamma', \Gamma, A \leftharpoondown B, \Delta \vdash C} \qquad (\leftharpoondown_r)\frac{A, \Gamma \vdash B}{\Gamma \vdash A \leftharpoondown B}$$

Note that, due to the absence of permutation, Γ is not a multiset, but a sequence of assumptions. The lack of rule (*permutation*) splits the connective '$\multimap$' into two implications '$\rightharpoonup$' and '$\leftharpoondown$', which one can think of as *over* and *under* or predictive and retroactive implication. For that reason, we call '$\rightharpoonup$' *forward* and '$\leftharpoondown$' *back*. Lack of (*permutation*) also accounts for the small modifications in the rules (*cut*), (*unit*) and $(\otimes_l)$.

The system above is called the *syntactic calculus* in Lambek[88] p 305 but in Lambek[87] we have a version without the constant I. Lambek showed in 1958 cut-elimination for this calculus. Moorgat in [Moo90] presents a version without (explicit) $\otimes$ and I, which he calls the Lambek-Gentzen Calculus **L**, but we shall call the tensor-free system $\mathbf{L}_-$. Moorgat also gives a Hilbert-style presentation of this logic, which he calls **Z** for Zielonka and says that the equivalence between **Z** and $\mathbf{L}_-$ was shown by Buszkowski (1989).

In Lambek[87] we also found the definition,

Definition 1 *A* categorial grammar *of a language may be viewed as consisting of the syntactic calculus* **L** *freely generated from a finite set* $\{S, N, \ldots\}$ *of basic types together with a dictionary which assigns to each word of the language a finite set of types composed from the basic types and* I *by the three binary operations,* $\otimes$, $\rightharpoonup$ *and* $\leftharpoondown$.

but we will only mention the definition of a categorial grammar and not try anything closer to real Linguistics in this note.

Both Moorgat and Lambek consider briefly the system where the operation $\otimes$ - usually denoted by a dot '$\cdot$' - is *not* associative, but we will not do this.

More importantly Moorgat and van Benthem argue that for certain linguistics applications (non-peripheral extraction is one of the examples) one would like to have some amount of permutation in the system. We will try and say something about that in the last section. Before going on to semantics we present a last sequent-calculus system, very natural from the Linear Logic viewpoint, namely the Lambek calculus with additives, which we call **LA**.

Lambek Calculus with Additives (LA)

Axioms:

$$A \vdash A \quad (identity)$$

$$\vdash I$$

$$\Gamma \vdash 1 \qquad \Gamma, 0, \Gamma' \vdash A$$

Structural Rule:

$$\frac{\Gamma \vdash A \qquad \Gamma', A, \Delta \vdash B}{\Gamma', \Gamma, \Delta \vdash B} \quad (cut)$$

Logical Rules:

Multiplicatives:

$$(unit)\frac{\Gamma, \Gamma' \vdash A}{\Gamma, I, \Gamma' \vdash A}$$

$$(\otimes_l)\frac{\Gamma, A, B, \Delta \vdash C}{\Gamma, A \otimes B, \Delta \vdash C} \qquad (\otimes_r)\frac{\Gamma \vdash A \qquad \Gamma' \vdash B}{\Gamma, \Gamma' \vdash A \otimes B}$$

$$(\rightharpoonup_l)\frac{\Gamma \vdash A \qquad \Gamma', B, \Delta \vdash C}{\Gamma', A \rightharpoonup B, \Gamma, \Delta \vdash C} \qquad (\rightharpoonup_r)\frac{\Gamma, A \vdash B}{\Gamma \vdash A \rightharpoonup B}$$


$$(\leftharpoondown_l)\frac{\Gamma \vdash A \qquad \Gamma', B, \Delta \vdash C}{\Gamma', \Gamma, A \leftharpoondown B, \Delta \vdash C} \qquad (\leftharpoondown_r)\frac{A, \Gamma \vdash B}{\Gamma \vdash A \leftharpoondown B}$$

Additives:

$$(\&_r)\frac{\Gamma \vdash A \qquad \Gamma \vdash B}{\Gamma \vdash A \& B} \qquad (\&_l)\frac{\Gamma, A, \Delta \vdash C}{\Gamma, A \& B, \Delta \vdash C} \qquad \frac{\Gamma, B, \Delta \vdash C}{\Gamma, A \& B, \Delta \vdash C}$$

$$(\oplus_l)\frac{\Gamma, A, \Delta \vdash C \qquad \Gamma, B, \Delta \vdash C}{\Gamma, A \oplus B, \Delta \vdash C} \qquad (\oplus_r)\frac{\Gamma \vdash A}{\Gamma \vdash A \oplus B} \qquad \frac{\Gamma \vdash B}{\Gamma \vdash A \oplus B}$$

Note that the system **LA** could also be called non-commutative intuitionistic linear logic. That is **ILL=LP +A**. As a summary of the systems discussed we draw a diagram



But remember that to obtain **LP** from **L** we add an inference rule, namely (*permutation*), whereas to obtain **LA** from **L** we add the additive connectives.

2 Semantics

The usual [Lam58] model for the logical system **L** is given by sets of expressions from a (generally taken as finite) non-empty universe V^+. This model is called in [Emms'90] the **String Semantics** for the Lambek Calculus **L**. To describe it, given subsets of V^+, $A \subseteq V^+, B \subseteq V^+$ define:

$$A \cdot B = \{xy \in V^+ |\ x \in A \text{ and } y \in B\}$$

$$C/B = \{x \in V^+ |\ \text{ for all } y \in B, xy \in C\}$$

$$A\backslash C = \{y \in V^+ |\ \text{ for all } x \in A, xy \in C\}$$

If you call $\mathcal{N}$ the set of all subsets of V^+, $\mathcal{N}$ has a natural order induced by *set containment*. If $A, B \subseteq V^+$ we say $A \leq B$ iff $A \subseteq B$. Concatenation, represented by the dot $\cdot$ above, gives $\mathcal{N}$ a (non-commutative) monoidal structure. Concatenation is associative, has identity given by the empty sequence, and is, in general, non-commutative. Given $A, B \in \mathcal{N}$

- $A \cdot (B \cdot C) = (A \cdot B) \cdot C$ (associativity of concatenation);
- If we denote by I the empty sequence $A \cdot I = I \cdot A = A$ (identity);
- $A \cdot B \neq B \cdot A$, in general.

Now, we want to generalise the structure above, taking our clues from Category Theory [Mac]. We first recall the definition of an ordered (non-commutative) monoid. Ordered (commutative) monoids are used in Concurrency Theory and were called *and-posets* in [HdP'90].

Definition 2 *An* ordered (non-commutative) monoid *is a poset $(M, \leq)$ with a given compatible monoidal structure $(M, \circ, e)$. That is, a set M equipped with a binary relation '$\leq$' satisfying:*

- $a \leq a$ *for all* a *in* M $(reflexivity)$
- $a \leq b$ *and* $b \leq c \Rightarrow a \leq c$ $(transitivity)$
- $a \leq b$ *and* $b \leq a \Rightarrow a = b$ $(antisymmetry)$

together with a monoid structure $(\circ, e)$ consisting of a 'multiplication' $\circ\colon M \times M \to M$ and a distinguished object 'e' of M, such that the following hold:

- $(a \circ b) \circ c = a \circ (b \circ c)$ $(associativity)$
- $a \circ e = e \circ a = a$ $(identity)$

The structures are compatible in the sense that, if $a \leq b$, we have

- $a \circ c \leq b \circ c$, *for all* c *in* M *and*
- $c \circ a \leq c \circ b$, *for all* c *in* M.

We write an ordered (non-commutative) monoid as a quadruple $(M, \leq, \circ, e)$. Note that

$$(\mathcal{N}, \leq, \cdot, I)$$

in the string semantics above is an ordered (non-commutative) monoid as the order $\leq$ is compatible with the multiplication given by concatenation.

$$A \leq B \Rightarrow A \cdot C \leq B \cdot C$$

$$A \leq B \Rightarrow C \cdot A \leq C \cdot B$$

But ordered monoids are not the structure we are after, we need some more structure to model the "slashes" or $\leftharpoonup$ and $\rightharpoonup$. Following the example of traditional lattice theory we define *left(right)-pseudocomplements* and more importantly the notion of a *biclosed poset.*

Definition 3 *Suppose M is a ordered monoid and $a, b \in M$. If there exists a largest $x \in M$ such that $a \circ x \leq b$ then this element is denoted $a \rightharpoonup b$ and it is called the* left pseudocomplement of a wrt b. *Similarly, if there exists a largest $x \in M$ such that $x \circ a \leq b$ then this element is denoted $a \leftharpoonup b$ and it is called the* right pseudocomplement of a wrt b.

A biclosed poset *is an ordered monoid $(M, \leq, \circ, e)$ such that $a \rightharpoonup b$ and $a \leftharpoonup b$ exist for all a and b in M. We write a biclosed poset as 6-tuple $(M, \leq, \circ, e, \rightharpoonup, \leftharpoonup)$.*

It is clear that $(\mathcal{N}, \leq, \cdot, I)$ with A/B and $A\backslash B$ defined for the string semantics is a biclosed poset. We now state some propositions, which are well-known fact¡s:

FACT 1 *In a biclosed poset $(M, \leq, \circ, e, \rightharpoonup, \leftharpoonup)$ the binary operations $\leftharpoonup$ and $\rightharpoonup$ are antimonotonic in their first coordinate, monotonic in their second coordinate and such that the biclosed poset satisfies:*

1. *$a \circ b \leq c$ iff $a \leq b \rightharpoonup c$ in M;*
2. *$b \circ a \leq c$ iff $a \leq c \leftharpoonup b$ in M.*

The next fact is the important one:

FACT 2 *Any biclosed poset $(M, \leq, \circ, e, \rightharpoonup, \leftharpoonup)$ is a model for the Lambek Calculus* **L**.

By that we mean that if $\Gamma \vdash A$ is a derivable sequent in **L**, and $(\ )^0$ assigns to each basic type B an element of the biclosed poset M, B^0, then $\Gamma^0 \leq A^0$. Note that a biclosed poset is the natural simplification to the poset set up of the notion of a (non-symmetric) monoidal biclosed category, see [Eilenberg/Kelly].

Note also that if the monoidal operation '$\circ$' is commutative, a left pseudocomplement is also a right pseudocomplement, and the two implications '$\rightharpoonup$' and '$\leftharpoonup$' collapse back into a single one. The *commutative* structure corresponding to a biclosed poset is a *closed poset*, that is a closed poset is a biclosed poset where the monoidal structure is commutative; examples can be found in [Law], [Flagg] and [HdP].

FACT 3 *Any (symmetric) closed poset $(M, \leq, \circ, e, \multimap)$ is a model for the Lambek Calculus with permutation* **LP**.

Recall that **LA** denotes the Lambek calculus with additives. The system **LA** satisfies cut-elimination, as the proof for **ILL** - with the modification made by Lambek to account for two implications - also work for **LA**.

This calculus has been given string semantics by Morril [Mor], Moorgat [Moo] and others. One way of adding additives to the string model above is defining:

$$A \cap B = \{x \in V^+ | x \in A \text{ and } x \in B\}$$

$$A \cup B = \{x \in V^+ | x \in A \text{ or } x \in B\}$$

To generalise the string semantics with additives, we could say:

FACT 4 *Any biclosed poset with (finite) meets and joins $(M, \leq, \circ, e, \rightharpoonup, \leftharpoonup, \wedge, \vee)$ is a model for the Lambek Calculus with additives* **LA**.

But note that adding booleans is stronger than adding additives, as for instance additives do not necessarily satisfy distributivity $A\&(B \oplus C) \ncong A\&B0 \oplus (A\&C)$ whereas booleans do (see [Roo]). A biclosed poset with finite meets and joins is the natural simplification to the poset set up of the notion of a (non-symmetric) monoidal biclosed category with finite products and coproducts. For the sake of completeness we state yet another fact.

FACT 5 *A (symmetric) closed poset with (finite) meets and joins* $(M, \leq, \circ, e, \multimap, \wedge, \vee)$ *is a model for the Lambek Calculus with additives and full permutation* **ILL**.

Note that a symmetric closed poset with meets and joins is almost a quantale [Yet], i.e a quantale without completeness. Many people have considered quantales as models for Intuitionistic Linear Logic, but the completeness of a quantale is not necessary to model the propositional logic and it distracts from the essential structure. Another observation is that additives do not have to be distributive, whereas products do distribute over coproducts, so our categorical framework is dealing with a special class of additives.

Moorgat also uses the boolean connectives to define a commutative concatenation and a corresponding - non-directional - internal-hom as follows:.

$$A \cdot_* B =_{def} (A \cdot B) \cup (B \cdot A)$$

$$A|B =_{def} (B \backslash A) \cap (A / B)$$

These are used to account for a phenomenon called *local discontinuity* in Linguistics. But the non-directional internal-hom gives us no notion of implication, as $A \to B$ is clearly not the same as $B \to A$ and $A|B = B|A$. In the next short subsections we discuss particular cases of facts 2 and 3 above, that is algebraic models for **L** and **LP**.

Matrices as a Model for L

In this subsection we show that mathematical structures, which look superficially very different from the traditional model, can model **L** as well as strings. To do that we use an example where we calculate with matrices.

Consider the set of square matrices with elements from N the set of natural numbers. More precisely we call $\mathcal{N}$ the set of invertible matrices or

$$\mathcal{N} = \{\text{invertible square matrices } k \times k \text{ with elements in } N\}$$

We use Greek letters α, β for our matrices. Of course we have the usual operation of matrix multiplication, denoted

$$\otimes : \mathcal{N} \times \mathcal{N} \to \mathcal{N}$$

Recall that matrix multiplication is a non-commutative monoidal operation on $\mathcal{N}$. That is

- There is an identity for matrix multiplication I, simply the identity matrix of dimension k,
- For any matrix α, $\alpha \otimes I = I \otimes \alpha = \alpha$,
- Associativity of matrix multiplication $(\alpha \otimes \beta) \otimes \gamma = \alpha \otimes (\beta \otimes \gamma)$ holds and
- The matrix product is, in general, non-commutative $\alpha \otimes \beta \neq \beta \otimes \alpha$.

We can define a pre-order on $\mathcal{N}$, saying that

$$\alpha \leq \beta \qquad \text{iff} \qquad \alpha^{-1} \otimes \beta \leq I$$

This clearly determines a pre-order as:

- $\alpha \leq \alpha$ as $\alpha \otimes \alpha^{-1} = I$,
- If $\alpha \leq \beta$ and $\beta \leq \gamma$ then $\alpha \leq \gamma$.

Given any pre-order we can have an induced partial order by taking equivalence classes, the traditional construction. The order induced on $\mathcal{N}$ is compatible with the matrix multiplication. If $\alpha \leq \beta$ - meaning $\alpha^{-1} \otimes \beta = I$ then $(\alpha \otimes \gamma) \leq (\beta \otimes \gamma)$. To show that, we calculate $(\alpha \otimes \gamma)^{-1} \otimes (\beta \otimes \gamma) = \gamma^{-1} \otimes \alpha^{-1} \otimes \beta \otimes \gamma = \gamma^{-1} \otimes I \otimes \gamma = I$. Thus $\langle \mathcal{N}, \leq, \otimes, I \rangle$ is another non-commutative ordered monoid. Moreover, we can use the invertibility of our matrices to define two operations that make $\mathcal{N}$ a biclosed poset.

Definition 4 *We can define two binary operations on elements of $\mathcal{N}$, $[(-) \rightharpoonup (-)]$ and $[(-) \leftharpoonup (-)]$, by saying $\alpha \rightharpoonup \beta = \alpha^{-1} \otimes \beta$ and $\alpha \leftharpoonup \beta = \beta^{-1} \otimes \alpha$.*

Note that these are operations of the form $\mathcal{N}^{op} \times \mathcal{N} \to \mathcal{N}$. By that we mean that the operations are antimonotonic in the first coordinate and monotonic in the second, or:

- If $\alpha_1 \leq \alpha_2$ then $\alpha_2 \rightharpoonup \beta \leq \alpha_1 \rightharpoonup \beta$
- If $\beta_1 \leq \beta_2$ then $\alpha \rightharpoonup \beta_1 \leq \alpha \rightharpoonup \beta_2$
- If $\alpha_1 \leq \alpha_2$ and $\beta_1 \leq \beta_2$ then $\alpha_2 \rightharpoonup \beta_1 \leq \alpha_1 \rightharpoonup \beta_2$.

Similarly for the operation '$\leftharpoonup$'. It is also clear that

- $\alpha \otimes (\alpha \rightharpoonup \beta) \leq \beta$ and
- $\beta \otimes (\alpha \leftharpoonup \beta) \leq \alpha$

Proposition 1 *The ordered monoid of invertible $k \times k$ matrices over N, $(\mathcal{N}, \leq, \otimes, I)$ with the operations of matrix multiplication '$\otimes$', and '$\leftharpoonup$' and '$\rightharpoonup$' defined above constitutes a biclosed poset.*

Thus $(\mathcal{N}, \leq, \otimes, I, \leftharpoonup, \rightharpoonup)$ is another model for the Lambek Calculus and there are lots of other models like modules of rings, etc. According to Lambek the idea for the Syntactic Calculus came from algebraic models, like the ones above. Actually it is easy to see that if one has a group, i.e the possibility of inverting elements of an (non-symmetric) ordered monoid, we will have a biclosed poset.

2.1 N as a Model for LP

Another observations is that if we wanted to produce algebraic models for **LP**, the calculus with permutation, instead of the Lambek Calculus, we would have many more 'familiar' structures, like for instance the set of natural numbers.

The set of the natural numbers N is clearly a (symmetric) *closed poset*, where conjunction '$\circ$' is given by addition of natural numbers, the identity for this conjunction is 0 and the internal-hom is given by truncated subtraction, cf. [Law] [dP91]. It is easy to prove the adjunction, see [dP91] if you are feeling lazy and another easy argument on Lindenbaum algebras (following Girard's proof) proves completeness of the models above.

In the next section we describe another model for the Lambek Calculus, but this time a categorical one, that is, one where we take in consideration different proofs of the same entailment. Note that we used a similar notation for matrices and strings, to emphasize that what matters is the *structure* of a biclosed (or closed) poset.

3 A Dialectica Construction

In this section we show that we can construct a Dialectica[2] category model of the Lambek Calculus in a straightforward way. The construction is meant both as evidence of the flexibility and powerfulness of the dialectica-like constructions and also as a way of, semantically, showing that we can define a modality that adds restricted permutation to the Lambek Calculus. The construction is given with all details, which might make it look too complicated and a bit boring at first reading. The reader should be warned that some familiarity with Category Theory is presupposed in this section.

Now suppose $\mathbf{C}$ is the category of sets and usual maps, so, in particular, $\mathbf{C}$ is a monoidal closed - actually cartesian closed - category with finite products and coproducts. As $\mathbf{C}$ has finite products, in particular it has a terminal object 1, any singleton set, and an initial object 0, the empty set. We write - as usual - $A \times B$ for the categorical product and $A + B$ for the coproduct in $\mathbf{C}$. Consider as well a biclosed poset $(\mathcal{N}, \leq, \otimes, e, \leftharpoonup, \rightharpoonup)$ in $\mathbf{C}$, you could think about it as either the string semantics (or the matrices model) above.

Note that the partial order on $\mathcal{N}$ induces an order on the hom-set $\mathbf{C}(X, \mathcal{N})$ given by

$$\text{for } f, g\colon X \to \mathcal{N} \quad f \leq g \qquad \text{iff} \qquad \forall x \in X \quad f(x) \leq g(x)$$

Using the data above we can define a (dialectica) category $D_{\mathcal{N}}\mathbf{C}$ as follows.

Definition 5 *Given a biclosed poset $(\mathcal{N}, \leq, \otimes, e, \leftharpoonup, \rightharpoonup)$ and $\mathbf{C}$ the category of* **Sets** *and usual functions, the category $D_{\mathcal{N}}\mathbf{C}$ consists of:*

- *Objects of $D_{\mathcal{N}}\mathbf{C}$ are triples (U, X, α) where α is a map in $\mathbf{C}$ of the form*

$$U \times X \xrightarrow{\quad\alpha\quad} \mathcal{N}$$

  We call the object above A, because it is determined by the function α; and write it as $A = (U \overset{\alpha}{\Leftarrow} X)$.

- *Morphisms in $D_{\mathcal{N}}\mathbf{C}$ from an object A to an object B of the form $V \times Y \overset{\beta}{\to} \mathcal{N}$ correspond to pairs of maps in $\mathbf{C}$, $f\colon U \to V$ and $F\colon Y \to X$ such that in the following diagram*

$$\begin{array}{ccc} U \times Y & \xrightarrow{\quad U \times F \quad} & U \times X \\ {\scriptstyle f \times Y}\downarrow & & \downarrow{\scriptstyle \alpha} \\ V \times Y & \xrightarrow[\quad\beta\quad]{} & \mathcal{N} \end{array}$$

  we have $\alpha \circ (U \times F) \leq \beta \circ (f \times Y)$ as morphisms in $\mathbf{C}(U \times Y, \mathcal{N})$.

We write morphisms in $D_{\mathcal{N}}\mathbf{C}$ as diagrams:

[2] Dialectica categories were introduced by the author in [dP87] as an internal categorical version of Gödel's Dialectica Interpretation. They turned out to be also an interesting model of Intuitionistic Linear Logic.

$$\begin{array}{ccc} U & \xleftarrow{\ \alpha\ } & X \\ {\scriptstyle f}\downarrow & \Downarrow & \uparrow{\scriptstyle F} \\ V & \xleftarrow[\ \beta\ ]{} & Y. \end{array} \qquad \forall u \in U, \forall y \in Y \quad \alpha(u, Fy) \leq \beta(f\,u\,y)$$

Note that checking that the data above really defines a category is not totally trivial. The category $D_{\mathcal{N}}\mathbf{C}$ above is a variant of the construction in the author's thesis [dP88], fully described in [dP'91], which is related to Chu's construction [Barr'79] and to $GAME_k$, [L/S'91].

Proposition 2 *The description above defines a category $D_{\mathcal{N}}\mathbf{C}$.*

Composition of morphisms (f, F) and (g, G) is given by the morphism (gf, FG). Identities are identities in each coordinate and associativity is given by the associativity of function composition in $\mathbf{C}$, that is **Sets**. But note that composition really corresponds to the following diagram:

$$\begin{array}{ccccc} U \times Z & \xrightarrow{U \times G} & U \times Y & \xrightarrow{U \times F} & U \times X \\ {\scriptstyle f \times Z}\downarrow & = & \downarrow{\scriptstyle f \times Y} & & \downarrow{\scriptstyle \alpha} \\ V \times Z & \xrightarrow[V \times G]{} & V \times Y & & | \\ {\scriptstyle g \times Z}\downarrow & & & \searrow^{\beta} & \downarrow \\ W \times Z & & \xrightarrow[\gamma]{} & & \mathcal{N} \end{array}$$

□

Having defined the category we want to use as a model for the Lambek Calculus, we now embark in a series of boring and relatively easy calculations to show that this category is monoidal biclosed. We start by defining a monoidal structure on $D_{\mathcal{N}}\mathbf{C}$.

Definition 6 *Given two objects $(U \overset{\alpha}{\leftarrow} X)$ and $(V \overset{\beta}{\leftarrow} Y)$ in $D_{\mathcal{N}}\mathbf{C}$ we define $A \otimes B$ their tensor product as follows:*

$$A \otimes B = (U \otimes V \xleftarrow{\ \alpha \otimes \beta\ } X^V \times Y^U)$$

The morphism "$\alpha \otimes \beta$" is given intuitively by $\alpha \otimes \beta(u, v, f, g) = \alpha(u, fv) \otimes \beta(v, gu)$, where the second $\otimes$ refers to the (non-commutative) monoidal operation in $\mathcal{N}$.

To give a formal definition of $\alpha \otimes \beta$ consider the map

$$U \times V \times X^V \times Y^U \xrightarrow{\text{"}\langle ev, ev\rangle\text{"}} U \times Y \times V \times X \xrightarrow{\alpha \times \beta} \mathcal{N} \times \mathcal{N} \xrightarrow{\otimes} \mathcal{N}$$

Note that we are using the cartesian closed - hence commutative - structure of **Sets** - to write X^V and Y^U- and the monoidal structure of $\mathcal{N}$ to define $\alpha \otimes \beta$.

This operation clearly defines a bifunctor, which is a tensor product. Associativity is straightforward and the object $I = (1 \stackrel{I}{\leftarrow} 1)$ - the map $1 \times 1 \stackrel{I}{\rightarrow} \mathcal{N}$, which only picks the identity I in $\mathcal{N}$ - is the identity for this tensor product.

Note that the 'carrier' of the tensor product is the same for $A \otimes B$ and $B \otimes A$, due to commutativity of products in $\mathbf{C}$, but as objects in $D_{\mathcal{N}}\mathbf{C}$ they are different, so the tensor product in $D_{\mathcal{N}}\mathbf{C}$ is *not* commutative.

One observation is that if one tries to get a (Dialectica) model for the Lambek calculus using a (non-symmetric) monoidal closed category $\mathbf{C}$ that leads into 'braiding' problems [J/S], which seems worth investigating.

Actually this odd-looking tensor product, analogous to the one in $\mathbf{GC}$ cf. [dP89], is the right one to prove monoidal-biclosedness of $D_{\mathcal{N}}\mathbf{C}$ with respect to the two internal-homs. To define the internal-homs:

Definition 7 *Given two objects $(U \stackrel{\alpha}{\leftarrow} X)$ and $(V \stackrel{\beta}{\leftarrow} Y)$ in $D_{\mathcal{N}}\mathbf{C}$ we define $[A,B]_l$ their left -respectively right - internal hom as follows:*

$$[A,B]_l = (V^U \times X^Y \stackrel{\alpha \rightharpoonup \beta}{\longleftarrow} U \times Y)$$

$$[A,B]_r = (V^U \times X^Y \stackrel{\alpha \leftharpoonup \beta}{\longleftarrow} U \times Y)$$

The morphisms "$\alpha \rightharpoonup \beta$" and "$\alpha \leftharpoonup \beta$" are given intuitively by $(\alpha \rightharpoonup \beta)(f,F,u,y) = \alpha(u,Fy) \rightharpoonup \beta(fu,y)$, respectively, $(\alpha \leftharpoonup \beta)(f,F,u,y) = \alpha(u,Fy) \leftharpoonup \beta(fu,y)$, where '$\rightharpoonup$' and '$\leftharpoonup$' are the corresponding 'internal-homs' or 'slashes' in $\mathcal{N}$.

The definition of the morphisms $(\alpha \rightharpoonup \beta)$ and $(\alpha \leftharpoonup \beta)$ is similar to the definition of $\otimes$ above. First consider maps $\overline{\alpha}$ and $\overline{\beta}$:

$$([U,V] \times [Y,X]) \otimes (U \otimes Y) \stackrel{\pi_1 \otimes U \otimes Y}{\longrightarrow} [U,V] \otimes U \otimes Y \stackrel{eval \otimes Y}{\longrightarrow} V \otimes Y \stackrel{\beta}{\longrightarrow} \mathcal{N}$$

$$([U,V] \times [Y,X]) \otimes (U \otimes Y) \stackrel{\pi_2 \otimes U \otimes Y}{\longrightarrow} [Y,X] \otimes U \otimes Y \stackrel{U \otimes eval}{\longrightarrow} U \otimes X \stackrel{\alpha}{\longrightarrow} \mathcal{N}$$

Then to obtain $\alpha \rightharpoonup \beta$ - resp. $\alpha \leftharpoonup \beta$ - we pair $\overline{\alpha}$ and $\overline{\beta}$ and compose the result with $\rightharpoonup$ and $\leftharpoonup$, considered as maps from $\mathcal{N} \times \mathcal{N}$ to $\mathcal{N}$:

$$([U,V] \times [Y,X]) \otimes (U \otimes Y) \stackrel{<\overline{\alpha},\overline{\beta}>}{\longrightarrow} \mathcal{N} \times \mathcal{N} \stackrel{\rightharpoonup}{\longrightarrow} \mathcal{N}$$

$$([U,V] \times [Y,X]) \otimes (U \otimes Y) \stackrel{<\overline{\alpha},\overline{\beta}>}{\longrightarrow} \mathcal{N} \times \mathcal{N} \stackrel{\leftharpoonup}{\longrightarrow} \mathcal{N}$$

Proposition 3 *The construction above defines bifunctors $[-,-]_l \colon D_{\mathcal{N}}\mathbf{C}^{op} \times D_{\mathcal{N}}\mathbf{C} \rightarrow D_{\mathcal{N}}\mathbf{C}$ and $[-,-]_r \colon D_{\mathcal{N}}\mathbf{C}^{op} \times D_{\mathcal{N}}\mathbf{C} \rightarrow D_{\mathcal{N}}\mathbf{C}$.*

Having defined internal homs and a tensor product we state and prove the desired theorem.


Theorem 1 *The category $D_{\mathcal{N}}\mathbf{C}$ is a (non-symmetric) monoidal biclosed category.*

The proof is very simple, one has to verify the natural isomorphisms


$$Hom_{D_{\mathcal{N}}\mathbf{C}}(A \otimes B, C) \cong Hom_{D_{\mathcal{N}}\mathbf{C}}(A, [B,C]_l)$$

$$Hom_{D_{\mathcal{N}}\mathbf{C}}(B \otimes A, C) \cong Hom_{D_{\mathcal{N}}\mathbf{C}}(A, [C,B]_r)$$

This can be done by looking at the diagrams



and



and calculating the sums.

If the morphism $(f, \langle F_1, F_2 \rangle)$ is in $Hom(A \otimes B, C)$, then we know $(\alpha \otimes \beta) \leq \gamma$, that is

$$(\alpha \otimes \beta)(u, v, F_1 z, F_2 z u) \leq \gamma(f(u, v), z)$$

This means that $\alpha(u, F_1 z v) \otimes \beta(v, F_2 z u) \leq \gamma(f(u, v), z)$ in $\mathcal{N}$. But $\mathcal{N}$ is a biclosed poset, which implies that $\alpha \leq \beta \rightharpoonup \gamma$ or $\alpha(u, F_1 z v) \leq \beta(v, F_2 z u) \rightharpoonup \gamma(f(u, v), z)$. To show that the corresponding morphism $(\langle f, \overline{F_2} \rangle, \overline{F_1})$ is in $Hom(A, [B, C]_l)$ we have to show $\alpha(u, F_1(v, z)) \leq (\beta \rightharpoonup \gamma)(\langle f, F_2 \rangle u, \langle v, z \rangle)$ that means $\alpha(u, F_1(v, z) \leq (\beta \rightharpoonup \gamma)(fu, F_2 u, v, z)$. As $(\beta \rightharpoonup \gamma)(fu, F_2 u, v, z) = \beta(v, F_2 uz) \rightharpoonup \gamma(fuv, z)$ we know what we need, if transposing is allowed in the category $\mathbf{C}$. Similarly for the other isomorphism. □

That is all we need to show that the category $D_{\mathcal{N}}\mathbf{C}$ with the structure described is a model of the Lambek Calculus.

FACT 6 *A monoidal biclosed category* $\mathbf{M}$ *is a model for the Lambek Calculus* $\mathbf{L}$.

The notion of a monoidal biclosed category reduces to the notion of a biclosed poset in the category of partial orders. Thus fact 6 is a generalisation of fact 2.

FACT 7 *The category* $D_{\mathcal{N}}\mathbf{C}$ *is a sound model for the Lambek Calculus* $\mathbf{L}$*, meaning that if a sequent* $\Gamma \vdash A$ *is proved in* $\mathbf{L}$*, there is a morphism* $f\colon |\Gamma| \to |A|$ *in the category.*

Atomic formulae are objects of $D_{\mathcal{N}}\mathbf{C}$, $\otimes$ is the non-commutative conjunction and '$\leftharpoonup$' and '$\rightharpoonup$' are the left and right internal-homs. The existence of the adjunction shows the validity of the inference rules for $\leftharpoonup$ and $\rightharpoonup$. □

The model $D_{\mathcal{N}}\mathbf{C}$ above is an easy model to calculate with as it is made out of sets, but it is not the syntactical traditional one. Recall that as we mentioned before there are lots of other algebraic models. But the reason to give the construction above - admittedly not the easiest algebraic model - is that we can define a non-trivial of course '!' operator in $D_{\mathcal{N}}\mathbf{C}$, more about that in the next section.

Moreover we can define additives $A \& B$ and $A \oplus B$ very easily and independently of the $\mathcal{N}$ chosen.

Proposition 4 *The category $D_{\mathcal{N}}\mathbf{C}$ has binary (hence finite) products and coproducts given by*

$$A\&B = (U \times V \xleftarrow{\alpha\&\beta} X + Y)$$

$$A \oplus B = (U + V \xleftarrow{\alpha \oplus \beta} X \times Y).$$

Note that $\alpha\&\beta$ and $\alpha \oplus \beta$ actually only use either 'α' or 'β', as the coproduct forces us to choose, say between x and y, in the product case.

FACT 8 *The category $D_{\mathcal{N}}\mathbf{C}$ is a model for the Lambek Calculus with additives* **LA**.

The attentive reader will have noticed that we could state three other facts, to the effect that symmetric monoidal closed categories are categorical models for **LP**, monoidal biclosed categories with (finite) products and coproducts are categorical models for **LA** and symmetric monoidal closed categories with products and coproducts are categorical models for **ILL**.

4 Modalities

We start our discussion of modalities by recalling the rules for the modality "!" in usual *commutative* Intuitionistic Linear Logic. These are:

$$\frac{\Gamma, A \vdash B}{\Gamma, !A \vdash B}\ (dereliction) \qquad \frac{\Gamma \vdash B}{\Gamma, !A \vdash B}\ (weakening)$$

$$\frac{\Gamma, !A, !A \vdash B}{\Gamma, !A \vdash B}\ (contraction) \qquad \frac{!\Gamma \vdash A}{!\Gamma \vdash !A}\ (!)$$

As observed by several people, the four rules above fall neatly into two pairs. One pair, namely (*weakening*) and (*contraction*), has to do with putting back into the logic, in a controlled way, contraction and weakening for specific formulae and the other pair - (*dereliction*) and (!) - make '!' look like the (necessitation) $\Box$ modal operator of **S4**.

In the categorical models one usually interprets '!' as a functor with a natural comonad structure, while the objects $!A$ have a natural comonoid structure wrt the tensor product of the category. The comonad structure of '!' is used to interpret the **S4**-like rules for the modality !, but this interpretation also seems to use the fact that there are additives in the logic. Thus if one knows that

$$!(A\&B) \cong !A \otimes !B \text{ and } !0 \cong 1$$

and that there exists a comonad '!', we can reason as follows. There is always a map $!A \xrightarrow{\delta} !!A$ and if $!G \xrightarrow{f} A$, then we can apply the functor '!' to f, to get $!!G \xrightarrow{!f} !A$ and precompose it with $!G \xrightarrow{\delta} !!G$ to get $!G \longrightarrow !A$, which shows the rule (!) is satisfied. For this we are assuming that $!\Gamma \vdash A$ corresponds to a morphism $!G \to A$, but to know it we use that

$$!\Gamma = !G_1 \otimes !G_2 \otimes \ldots \otimes !G_k \cong !(G_1\&G_2\& \ldots G_k).$$

Thus we are using the fact that 'commas' really are instances of the connective '$\otimes$' and that the sequence $!G_1, !G_2, \ldots, !G_n$ can be transformed into '!' applied to the object $G_1 \& G_2 \ldots \& G_n$. The reasoning above seems to hint at the usefulness of treating additives as well as multiplicatives, if one wants to re-introduce the structural rules into the system. It may not be necessary, but it certainly makes it semantically easier, as we shall see.

The little work that has been done in non-commutative Linear Logic, [Yetter], [Lincoln et al] has maintened a (weaker) notion of commutativity given by circular permutations. For the linguistic purposes envisaged for the Lambek calculus, the idea of allowing circular permutations is very odd. But for linguistical purposes no one is interested in having a single dualising element, admittedly at least in Yetter's paper, the main reason for the circular permutations rule.

It seems clear that formulas of the type $!A$, that we are allowed to contract and weaken over, should commute with any other formula, not only with some special ones. Also additives commute.

Hence what follows is a preliminary proposal, based mainly on [Morril et al] and [Yetter]'s ideas. We start with the calculus **LA** the Lambek calculus with additives, we add a modality like Yetter's 'κ' - *kappa* - with its rules and call the new calculus $\mathbf{LA}_\kappa$, for Lambek calculus with additives and (very restricted) permutation. Then we add Girard's [Gir'87] four rules for the modality '!' and call the resulting system $\mathbf{LA}_{\kappa!}$.

The reason for doing that is merely semantical, as we can interpret these modalities κ and ! in a variant of the Dialectica category $D_{\mathcal{N}}\mathbf{C}$ described in the previous section.

Lambek Calculus with Additives and κ ($\mathbf{LA}_\kappa$)

Axioms:

$$A \vdash A \quad (identity)$$

$$\vdash I$$

$$\Gamma \vdash 1 \qquad \Gamma, 0, \Gamma' \vdash A$$

Structural Rule:

$$\frac{\Gamma \vdash A \qquad \Gamma', A, \Delta \vdash B}{\Gamma', \Gamma, \Delta \vdash B} \quad (cut)$$

Logical Rules:
Multiplicatives:

$$(unit)\frac{\Gamma \vdash A}{\Gamma', I, \Gamma'' \vdash A}$$

$$(\otimes)\frac{\Gamma, A, B, \Delta \vdash C}{\Gamma, A \otimes B, \Delta \vdash C} \qquad (\otimes)\frac{\Gamma \vdash A \qquad \Gamma' \vdash B}{\Gamma, \Gamma' \vdash A \otimes B}$$

$$(\rightharpoonup_l)\frac{\Gamma \vdash A \qquad \Gamma', B, \Delta \vdash C}{\Gamma', A \rightharpoonup B, \Gamma, \Delta \vdash C} \qquad (\rightharpoonup_r)\frac{\Gamma, A \vdash B}{\Gamma \vdash A \rightharpoonup B}$$

$$(\leftharpoonup_l)\frac{\Gamma' \vdash A \qquad \Gamma, B, \Delta \vdash C}{\Gamma, \Gamma', A \leftharpoonup B, \Delta \vdash C} \qquad (\leftharpoonup_r)\frac{A, \Gamma \vdash B}{\Gamma \vdash A \leftharpoonup B}$$

Additives:

$$(\&_r)\frac{\Gamma \vdash A \qquad \Gamma \vdash B}{\Gamma \vdash A \& B} \qquad (\&_l)\frac{\Gamma, A, \Delta \vdash C}{\Gamma, A \& B, \Delta \vdash C} \qquad \frac{\Gamma, B, \Delta \vdash C}{\Gamma, A \& B, \Delta \vdash C}$$

$$(\oplus_l)\frac{\Gamma, A, \Delta \vdash C \qquad \Gamma, B, \Delta \vdash C}{\Gamma, A \oplus B, \Delta \vdash C} \qquad (\oplus_r)\frac{\Gamma \vdash A}{\Gamma \vdash A \oplus B} \qquad \frac{\Gamma \vdash B}{\Gamma \vdash A \oplus B}$$

Yetter's Rules for κ (kappa)

$$\frac{\Gamma, A, \kappa B, \Gamma' \vdash C}{\Gamma, \kappa B, A, \Gamma' \vdash C}\ (e1) \qquad \frac{\Gamma, \kappa A, B, \Gamma' \vdash C}{\Gamma, B, \kappa A, \Gamma' \vdash C}\ (e2)$$

$$\frac{\Gamma, A, \Delta \vdash C}{\Gamma, \kappa A, \Delta \vdash C}\ (der_k) \qquad \frac{\kappa\Gamma \vdash C}{\kappa\Gamma \vdash \kappa C}\ (\kappa)$$

Note that the difference between the systems $\mathbf{(LA}_\kappa\mathbf{)}$ and **ILL** is that permutation is restricted in $\mathbf{(LA}_\kappa\mathbf{)}$ to the formulas marked with a kappa. Also note that the rules for kappa are like **S4**-rules.

Recall that $D_{\mathcal{N}}\mathbf{C}$ models **LA** for any biclosed poset $\mathcal{N} = \langle N, \circ, \leq, e, \rightharpoonup, \leftharpoonup \rangle$. In particular, we can take as $\mathcal{N}$ a (non-commutative) quantale Q. That is not really necessary, but simplifies parts of the exposition, as we can quote from Yetter's paper, [Yet'90]. Recall that by proposition 1.9 in page 44 of [Yet], if Q is a quantale, there is a unique maximal central modality in Q, denoted $c_m : Q \to Q$. Use this modality c_m to define a modality κ in $D_{\mathcal{N}}\mathbf{C}$ as follows:

Definition 8 *Given a category $D_{\mathcal{N}}\mathbf{C}$, where $\mathcal{N}$ is a quantale Q we define*

$$\kappa : D_{\mathcal{N}}\mathbf{C} \to D_{\mathcal{N}}\mathbf{C}$$

which acts on carriers of objects of $D_{\mathcal{N}}\mathbf{C}$ as the identity endofunctor. Thus

$$\kappa(U \overset{\alpha}{\leftarrow} X) = (U \overset{c_m \alpha}{\leftarrow} X)$$

where $U \times X \overset{\alpha}{\to} Q$, a morphism in $\mathbf{C}$, is taken by κ to the composition


$$U \times X \xrightarrow{\alpha} Q \xrightarrow{c_m} Q$$

and c_m is the corresponding unique maximal central *modality in $\mathcal{Q}$.*

Moreover it is very easy to see that:

Proposition 5 *The endofunctor $\kappa : D_{\mathcal{N}}\mathbf{C} \to D_{\mathcal{N}}\mathbf{C}$ has a natural structure of comonad.*

The natural transformations are induced by the 'natural transformations' in Q:



All of this can be made much more concrete if we one takes a quantale of relations (cf. [BG'91]). The point here is that, the same way the completeness of a quantale is *not* necessary to model propositional Linear Logic, the completeness of a quantale is not necessary to model the existence of an of course "!" operator. But completeness is an added bonus (extra structure), which makes some calculations simpler.

Because the modality is central is makes the kappa rules sound, cf. Yetter. Thus we have a proposition:

Proposition 6 *The category $D_{\mathcal{N}}\mathbf{C}$ - where $\mathcal{N}$ is a quantale Q - together with the comonad κ defined above is a categorical model for the Lambek Calculus with additives and restricted permutation or system $\mathbf{LA}_{\kappa}$.*

Now we add to the system $\mathbf{LA}_{\kappa}$ the Girard's rules for the modality '!'.

Lambek Calculus with Additives, κ and '!' ($\mathbf{LA}_{\kappa!}$)

Axioms:

$$A \vdash A \quad (identity)$$
$$\vdash I$$
$$\Gamma \vdash 1 \qquad \Gamma, 0, \Gamma' \vdash A$$

Structural Rule:

$$\frac{\Gamma \vdash A \qquad \Gamma', A, \Delta \vdash B}{\Gamma', \Gamma, \Delta \vdash B} \quad (cut)$$

Logical Rules:
Multiplicatives:

$$(unit)\frac{\Gamma \vdash A}{\Gamma', I, \Gamma'' \vdash A}$$

$$(\otimes)\frac{\Gamma, A, B, \Delta \vdash C}{\Gamma, A \otimes B, \Delta \vdash C} \qquad (\otimes)\frac{\Gamma \vdash A \qquad \Gamma' \vdash B}{\Gamma, \Gamma' \vdash A \otimes B}$$

$$(\multimap_l)\frac{\Gamma \vdash A \qquad \Gamma', B, \Delta \vdash C}{\Gamma', A \multimap B, \Gamma, \Delta \vdash C} \qquad (\multimap_r)\frac{\Gamma, A \vdash B}{\Gamma \vdash A \multimap B}$$

$$(\multimap_l)\frac{\Gamma \vdash A \qquad \Gamma', B, \Delta \vdash C}{\Gamma', \Gamma, A \multimap B, \Delta \vdash C} \qquad (\multimap_r)\frac{A, \Gamma \vdash B}{\Gamma \vdash A \multimap B}$$

Additives:

$$(\&_r)\frac{\Gamma \vdash A \qquad \Gamma \vdash B}{\Gamma \vdash A\&B} \qquad (\&_l)\frac{\Gamma, A, \Delta \vdash C}{\Gamma, A\&B, \Delta \vdash C} \qquad \frac{\Gamma, B, \Delta \vdash C}{\Gamma, A\&B, \Delta \vdash C}$$

$$(\oplus_l)\frac{\Gamma, A, \Delta \vdash C \qquad \Gamma, B, \Delta \vdash C}{\Gamma, A \oplus B, \Delta \vdash C} \qquad (\oplus_r)\frac{\Gamma \vdash A}{\Gamma \vdash A \oplus B} \qquad \frac{\Gamma \vdash B}{\Gamma \vdash A \oplus B}$$

Yetter's Rules for κ (kappa)

$$\frac{\Gamma, A, \kappa B, \Gamma' \vdash C}{\Gamma, \kappa B, A, \Gamma' \vdash C}\ (e1) \qquad \frac{\Gamma, \kappa A, B, \Gamma' \vdash C}{\Gamma, B, \kappa A, \Gamma' \vdash C}\ (e2)$$

$$\frac{\Gamma, A, \Delta \vdash C}{\Gamma, \kappa A, \Delta \vdash C}\ (der_k) \qquad \frac{\kappa\Gamma \vdash C}{\kappa\Gamma \vdash \kappa C}\ (\kappa)$$

Girard's Rules for ! (of course!)

$$\frac{\Gamma, A \vdash B}{\Gamma, !A \vdash B}\ (dereliction) \qquad \frac{\Gamma \vdash B}{\Gamma, !A \vdash B}\ (weakening)$$

$$\frac{\Gamma, !A, !A \vdash B}{\Gamma, !A \vdash B}\ (contraction) \qquad \frac{!\Gamma \vdash A}{!\Gamma \vdash !A}\ (!)$$

Rules connecting κ and !\[3\]

$$\frac{\Gamma, !A, \Gamma' \vdash B}{\Gamma, \kappa A, \Gamma' \vdash}\ (!\kappa_l) \qquad \frac{\Gamma \vdash \kappa C}{\Gamma \vdash !C}\ (!\kappa_r)$$

The next step is to define a modality

$$!: D_{\mathcal{N}}\mathbf{C} \to D_{\mathcal{N}}\mathbf{C}$$

which given an object $A = (U \overset{\alpha}{\leftarrow} X)$ returns an object $(U \overset{!\alpha}{\leftarrow} X^{*U})$. This can be shown to be an endofunctor with a natural comonad structure inherited from the monoidal structure of X^{*U}, cf. [dP91]. There are lots of details to check, but as the category $\mathbf{C}$ is **Sets**, which is cartesian closed, the constructions above work and give you a comonad and comonoids in $D_{\mathcal{N}}\mathbf{C}$, cf [dP'91].

After checking all the details, we should have:

Proposition 7 *The category $D_{\mathcal{N}}\mathbf{C}$ is a categorical model for the Lambek Calculus with additives, restricted permutation κ and Girard modality '!' (of course!).*

As we mentioned before, this is all done semantically, it would be very interesting to check what happens with the Proof Theory of the systems ($\mathbf{LA}_\kappa$) and ($\mathbf{LA}_{\kappa!}$). On the one hand, the proof theory should not be too complicated, as we do not have the *par* operator interacting with linear implication, but on the other hand, people interested in proof-nets find weak forms of commutativity like (*cycling*) extremely natural. More work is needed here.

[3] The necessity of rule ($!\kappa_l$) to (eventually) obtain cut-elimination was explained to me by Dirk Roorda.

Loose Ends

If one takes the linguistics motivation seriously, the modality κ above is not exactly what we want to do. There might be some use in having κ around [Hepple, pers. comm], but it seems that it would be more useful to have a modality which did a *very* limited amount of permutation. In particular it would be interesting to produce rules for a modality that applied to two formulas A and B said something like *It is known that A and B commute*. The modality κ above says something like *κA commutes with everything*. There were some attempts at less contraining commutative modalities and the interested reader should consult [B/M], for several possibilities, amongst them we found the rules below:

Rules for Permutators


$$\frac{\Gamma, A, \triangleleft B \vdash C}{\begin{array}{l}\Gamma, \triangleleft B, A \vdash C \\ \Gamma, A\triangleright, B, \Delta \vdash C\end{array}} \qquad \frac{\Gamma, A, \Delta \vdash C}{\begin{array}{l}\Gamma, \triangleleft A, \Delta \vdash C \\ \Gamma, A, \Delta \vdash C\end{array}}$$

$$\frac{}{\Gamma, B, A\triangleright, \Delta \vdash C} \qquad \frac{}{\Gamma, A\triangleright, \Delta \vdash C}$$

We do not have a proposal of our own, as we have not had time to check in the algebraic - or categorical - literature for examples of such phenomena. In particular I would like to investigate the "quantor types" of Moortgat [Moo], as they look a bit like the comonad T in my previous work.

Curry-Howard Correspondences

It seems worth thinking about the Curry-Howard correspondence for some of the fragments of Linear Logic mentioned. We have four diagrams showing some untidiness on the Curry-Howard correspondence. Notice that the pointing of the arrows means nothing, it is just a typographical device to draw the squares.




The first diagram consists of logics:

- at the top Intuitionistic Logic, with three structural rules, (*weakening*), (*contraction*) and (*permutation*).

- on the left, what is called Direct Logic [KW], [Bel], which has (*weakening*), but not (*contraction*),
- on the right, a variant of Relevance Logic, which allows (*contraction*), but not (*weakening*), as weakening corresponds to vacuous implication,
- at the bottom of the square, without (*contraction*) and (*weakening*), we have the Lambek Calculus with permutation **LP**, or the multiplicative fragment of ILL,
- below that, when we remove the structural rule of (*permutation*) or exchange we have the Lambek calculus **L**
- finally, we have **J** the version of the calculus that would appear if we removed the associativity of the tensor product.

The second diagram consists of (untyped) λ-calculi corresponding to the logics. In particular, the fact that one does not allow (*weakening*) in Relevance Logic changes the lambda-calculus so that we get λI, the calculus originally proposed by Church, instead of the (now usual) λK [Bar]. Note that one is not allowed to abstract over a variable that is not present, but if a variable appears more than once the lambda-abstraction binds all the occurrences. The fact that we do not allow (*contraction*) in Direct Logic changes the lambda-calculus in a different way so one has to be careful when dealing with variables. That is lambda-abstraction has to bind a single occurrence of a variable in λK_s, when in λK it usually binds multiple (or no) ocurrences. But one is still allowed to abstract over a variable not present, i.e binding no variables. If we do the two restrictions we get λI_s, no vacuous abstraction and no multiple bindings either. I don't know what is the effect, if any, of not allowing commutativity or associativity of products, hence the question marks.



The third diagram gives the categorical correspondence, as far as I know it at the moment, but [Mere] is working on categorical models for Relevance Logic. Finally the fourth diagram shows the combinators version, but it is mostly guessed.

References

[Bar] H.P BARENDREGT **The Lambda Calculus - Its syntax and Semantics**, Studies in Logic and the Foundations of Mathematics vol 103, North-Holland, 1984.

[Barr] M. BARR $*$-**Autonomous Categories**, LNM 752, Springer-Verlag, 1979.

[Bel] G. BELLIN **Mechanizing Proof-Theory:Resource-Aware Logics and Proof-transformations to Extract Implicit Information** ,Ph D thesis, Stanford Computer Science Dept. TR STAN-CS-90-1319.

[vBen] J. van BENTHEM *The Lambek Calculus*, **Categorial Grammars and Natural Language Structures**, eds R. Oehrle, E. Bach and D. Wheeler, 1988.

[B/M] G. BARRY and G. MORRILL *Studies in Categorial Grammar*, Edinburgh Working Papers in Cognitive Science, vol 5, 1990.

[BG] C. BROWN and D. GURR *A Representation Theorem for Quantales*, manuscript, 1991.

[BGdP] C. BROWN, D. GURR and V.de PAIVA *General Petri Nets and Simulation Morphisms*, manuscript, Feb'91.

[Bus] W. BUSZKOWSKI *Completeness Results for Lambek Syntactic Calculus*, Zeitschr. f. Math, Logik und Grundlagen d. Math 32, S.13-28, (1986).

[E/K] S. EILENBERG and M. KELLY *Closed Categories*, in: Proc. Conf. on Categorical Algebra, La Jolla, 1965 (Springer-Verlag 1966) pp 421-562.

[Emms] M. EMMS *Polymorphic Quantifiers*, in *Studies in Categorial Grammar* eds Guy Barry and Glyn Morrill, Edinburgh Working Papers in Cognitive Science, vol 5, 1990.

[Fla] R. C. FLAGG *A Generalised Logic*, manuscript, Fall'90.

[G/L] J- Y. GIRARD and Y. LAFONT *Linear Logic and Lazy Computation*, TAPSOFT-87, 1987.

[Gir] J- Y. GIRARD *Linear Logic* , Theoretical Computer Science, 50, 1987 , 1-102.

[HdP] M. HYLAND and V.de PAIVA *Lineales*, manuscript Sept'90.

[J/S] A. JOYAL and R. STREET *Braided Monoidal Categories*, Macquarie Mathematics Report 860081 (1986).

[L/S] Y. LAFONT and T. STREICHER *Game Semantics for Linear Logic*, LICS'91.

[K/W] J. KETONEN and R. WEYHRAUCH *A Decidable Fragment of Predicate Calculus*, TCS 32, 297–307, (1984).

[Lam] J. LAMBEK *The Mathematics of Sentence Structure*, Amer. Math. Monthly 65, (1958) 154-69.

[Lam] J. LAMBEK *Categorial and Categorical Grammmas*, in **Categorial Grammars and Natural Language Structures**, eds R. Oehrle, E. Bach and D. Wheeler, 1988.

[Lam] J. LAMBEK *Multicategories Revisited*, in **Categories in Computer Science and Logic - 87** eds. J. Gray and A. Scedrov, 1989.

[Law] F.W. LAWVERE *Metric Spaces, Generalized Logic, and Closed Categories*, Rendiconti del Seminario Matematico e Fisico di Milano, 43 (1973).

[Linc] P. LINCOLN, J. MITCHELL, A. SCEDROV and N. SHANKAR *Decision Problems for Propositional Linear Logic*, in **Proc. of FOCS - 90**.

[Mac] S. MACLANE **Categories for the Working Mathematician**, Springer-Verlag, 1971.

[Mere] M.C. MERÉ *Categorical Models of Relevance Logic*, thesis, ITT.

[Moo] M. MOORGAT *The Logic of Discontinous Type Constructors*, manuscript 1990.

[Mor] G. MORRILL *Grammar and Logical Types*, in *Studies in Categorial Grammar*, Edinburgh Working Papers in Cognitive Science, vol 5, 1990.

[Mor] G. MORRILL, N. LESLIE, M. HEPPLE and G. BARRY *Categorial Deductions and Structural Operations*, in *Studies in Categorial Grammar*, Edinburgh Working Papers in Cognitive Science, vol 5, 1990.

[dP] V.C.V. de PAIVA *The Dialectica Categories*, In "Categories in Computer Science and Logic", Proc. Summer Research Conf. AMS 92, Boulder, CO, June 1987.

[dP] V.C.V. de PAIVA *A Dialectica-like Model of Linear Logic*, LNCS 389, Manchester, Sept. 1989.

[dP] V.C.V. de PAIVA *The Dialectica Categories*, thesis, Nov'88, issued as TR213 from Computer Lab, Jan'91.

[dP] V.C.V. de PAIVA *Petri Nets, Linear Logic and Categorical Relations*, TR 225, Computer Lab, 1991.

[Roo] D. ROORDA **Resource Logics:Proof-theoretical Investigations**,Ph. D Thesis, Univ. van Amsterdam, Sept. 1991 The Netherlands.

[Roo] D. ROORDA personal communication, Oct. 1991.

[Tak] G. TAKEUTI **Proof theory**, Studies in Logic and the Foundations of Mathematics vol 81, North-Holland, 1975.

[Yet] D. YETTER *Quantales and (non-commutative) Linear Logic*, JSL 55, March '90.