

# Efficient Gaussian Processes for data-driven decision making



**Vincent Dutordoir**

Department of Engineering  
University of Cambridge

Report submitted to be registered for the PhD Degree  
*First-Year-Report*



## Abstract

This is where you write your abstract ...



# Table of contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Contributions and Layout of this Report . . . . .	1
<b>2</b>	<b>Theoretical Framework</b>	<b>3</b>
2.1	Gaussian Processes . . . . .	3
2.1.1	The Beauty of Gaussian Process Regression: Exact Bayesian Inference .	4
2.2	Approximate Inference with Sparse Gaussian Processes . . . . .	6
2.3	Interdomain Inducing Variables . . . . .	6
2.3.1	Example: heavyside inducing variable . . . . .	6
2.4	Deep Gaussian Processes . . . . .	7
2.5	Covariance Functions . . . . .	7
<b>3</b>	<b>Spherical Harmonic Variational Gaussian Processes</b>	<b>9</b>
3.1	Spherical Harmonics . . . . .	9
3.1.1	How do we construct a fundamental system of points? . . . . .	12
<b>4</b>	<b>Deep Neural Networks as Point Estimates for Deep Gaussian Processes</b>	<b>13</b>
<b>5</b>	<b>Future Research</b>	<b>15</b>
5.1	Geometric Gaussian Decision Systems . . . . .	15
5.2	Projects in Progress . . . . .	17



# Chapter 1

## Introduction

1. Neural networks and Gaussian processes: complementary strengths and weaknesses
2. Ideally we have a single model that can handle low and high dimensional inputs, make robust uncertainty-aware predictions and can be used in big and small data regimes.
3. Gaussian processes and Bayesian Neural networks: connections [Neal, 1992; 1995; Williams and Rasmussen, 1996]
4. Problem with BNN is that (approximate) Bayesian inference is challenging
5. Deep Gaussian processes [Damianou and Lawrence, 2013]
6. Require accurate and scalable approximate Bayesian inference procedures

### 1.1 Contributions and Layout of this Report

This report represents my learning and the research that I conducted during the first year of my PhD degree. Most notably, we developed a novel sparse approximation for (deep) Gaussian processes based on the decomposition of the kernel in Spherical harmonics. In chapter 2 we cover the necessary theoretical background.

**Chapter 3** In this chapter we introduce a new class of inter-domain variational GPs where data is mapped onto the unit hypersphere in order to use spherical harmonic representations. The inference scheme is comparable to Variational Fourier Features, but it does not suffer from the curse of dimensionality, and leads to diagonal covariance matrices between inducing variables. This enables a speed-up in inference, because it bypasses the need to invert large covariance matrices. The experiments show that our model is able to fit a regression model for a dataset with 6 million entries two orders of magnitude faster compared to standard sparse GPs, while retaining state of the art accuracy.

The content of this chapter is largely based on:

Vincent Dutordoir, Nicolas Durrande, and James Hensman [2020]. “Sparse Gaussian Processes with Spherical Harmonic Features”. In: *Proceedings of the 37th International Conference on Machine Learning (ICML)*,

with the exception of the algorithm for computing the spherical harmonics in high dimensions.

**Chapter 3** Following up on the previous chapter, we use the decomposition of zonal kernels to design an interdomain inducing variable that mimics the behaviour of activation functions in neural network layers.

The content of this chapter is largely based on:

Vincent Dutordoir, James Hensman, Mark van der Wilk, Carl Henrik Ek, Zoubin Ghahramani, and Nicolas Durrande [2021a]. “Deep Neural Networks as Point Estimate for Deep Gaussian Processes”. In: *submission to NeurIPS*.

**Chapter 4** In the last chapter of the report we will shed a light on what the future work will focus on: “Gaussian Decision Systems with Geometric Gaussian processes”.



## Chapter 2

# Theoretical Framework

This chapter discusses Gaussian processes (GP) and deep Gaussian processes (DGPs), the composite model obtained by stacking multiple GP models on top of each other. We also review how to perform approximate Bayesian inference in these models, with a particular attention to Variational Inference. We also cover the theory of positive definite kernels and the Reproducing Kernel Hilbert Spaces (RKHS) they characterise.

### 2.1 Gaussian Processes

Gaussian processes (GPs) [Rasmussen and Williams, 2006] are non-parametric distributions over functions similar to Bayesian Neural Networks (BNNs). The core difference is that neural networks represent distributions over functions through distributions on weights, while a Gaussian process specifies a distribution on function values at a collection of input locations. This representation allows us to use an infinite number of basis functions, while still enables Bayesian inference [Neal, 1995].

Following from the Kolmogorov extension theorem, we can construct a real-valued stochastic process (i.e. function) on a non-empty set  $\mathcal{X}$ ,  $f : \mathcal{X} \rightarrow \mathbb{R}$ , if there exists on all finite subsets  $\{x_1, \dots, x_N\} \subset \mathcal{X}$ , a *consistent* collection of finite-dimensional marginal distributions over  $f(\{x_1, \dots, x_n\})$ . For a Gaussian process, in particular, the marginal distribution over every finite-dimensional subset is given by a multivariate normal distribution. This implies that, in order to fully specify a Gaussian process, it suffices to only define the mean and covariance (kernel) function because they are the sufficient statistics for every finite-dimensional marginal distribution. We can therefore denote the GP as

$$f \sim \mathcal{GP}(\mu, k), \tag{2.1}$$

where  $\mu : \mathcal{X} \rightarrow \mathbb{R}$  is the mean function, which encodes the expected value of  $f$  at every  $x$ ,  $\mu(x) = \mathbb{E}_f[f(x)]$ , and  $k : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  is the covariance (kernel) function that describes the

covariance between function values,  $k(x, x') = \text{Cov}(f(x), f(x'))$ . The covariance function has to be a symmetric, positive-definite function. The Gaussianity, and the fact that we can manipulate function values at some finite points of interest without taking the behaviour at any other points into account (the marginalisation property) make GPs particularly convenient to manipulate and use as priors over functions in Bayesian models – as we will show next.

Throughout this report, we will consider  $f$  to be the complete function, and intuitively manipulate it as an infinitely long vector. Moreover,  $f(\mathbf{x}) \in \mathbb{R}^N$  denotes the function evaluated at a finite set of points, whereas  $f^{\setminus \mathbf{x}}$  denotes another infinitely long vector similar to  $f$  but excluding  $f(\mathbf{x})$ . From the marginalisation property it follows that integrating out over the infinitely many points that are not included in  $\mathbf{x}$ , we obtain a valid finite-dimensional density for  $f(\mathbf{x})$

$$p(f(\mathbf{x})) = \int p(f) \mathrm{d}f^{\setminus \mathbf{x}}. \quad (2.2)$$

In the case of GPs, this finite-dimensional marginal is given by a multivariate Gaussian distribution, fully characterised by the mean  $\mu$  and the covariance function  $k$

$$p(f(\mathbf{x})) = \mathcal{N}(\boldsymbol{\mu}_{\mathbf{f}}, \mathbf{K}_{\mathbf{ff}}), \quad \text{where} \quad [\boldsymbol{\mu}_{\mathbf{f}}]_i = \mu(x_i) \text{ and } [\mathbf{K}_{\mathbf{ff}}]_{i,j} = k(x_i, x_j). \quad (2.3)$$

Conditioning the GP at this finite set of points leads to a conditional distribution for  $f^{\setminus \mathbf{x}}$ , which is given by another Gaussian process

$$p(f^{\setminus \mathbf{x}} | f(\mathbf{x}) = \mathbf{f}) = \mathcal{GP}(\mathbf{k}_{\mathbf{f}}^{\top} \mathbf{K}_{\mathbf{ff}}^{-1} (\mathbf{f} - \boldsymbol{\mu}_{\mathbf{f}}), \quad k(\cdot, \cdot) - \mathbf{k}_{\mathbf{f}}^{\top} \mathbf{K}_{\mathbf{ff}}^{-1} \mathbf{k}_{\mathbf{f}}), \quad (2.4)$$

where  $[\mathbf{k}_{\mathbf{f}}]_i = k(x_i, \cdot)$ . The conditional distribution over the whole function  $p(f | f(\mathbf{x}) = \mathbf{f})$  has the exact same form as in eq. (2.4). This is mathematically slightly confusing because the random variable  $f(\mathbf{x})$  is included both on the left and right-hand-side of the conditioning, but the equation is technically correct [Matthews et al., 2016].

### 2.1.1 The Beauty of Gaussian Process Regression: Exact Bayesian Inference

One of the key advantages of Gaussian processes for regression is that we can perform exact Bayesian inference. Assume a supervised learning setting where  $x \in \mathcal{X}$  (typically,  $\mathcal{X} = \mathbb{R}^d$ ) and  $y \in \mathbb{R}$ , and we are given a dataset  $\mathcal{D} = \{(x_i, y_i)\}_{i=1}^N$  of input and corresponding output pairs. For convenience, we sometimes group the inputs in  $\mathbf{x} = \{x_i\}_{i=1}^N$  into a single design matrix and outputs  $\mathbf{y} = \{y_i\}_{i=1}^N$  into a vector. We further assume that the data is generated by an unknown function  $f : \mathcal{X} \rightarrow \mathbb{R}$ , such that the outputs are perturbed versions of functions evaluations at the corresponding inputs:  $y_i = f(x_i) + \epsilon_i$ . In the case of regression we assume a Gaussian noise model  $\epsilon_i \sim \mathcal{N}(0, \sigma^2)$ . We are interested in learning the function  $f$  that generated the data.

[General introduction to Bayesian modelling] The key idea in Bayesian modelling is to specify a prior distribution over the quantity of interest. The prior encodes what we know at that point in time about the quantity. In general term, this can be a lot or a little. We encode

this information in the form of a distribution. Then, as more data becomes available, we use the rules of probability, in particular Bayes' rule, to update our prior beliefs and compute a posterior distribution (see **bisschop**; MacKay [2003] for a thorough introduction).

Following the Bayesian approach, we specify a *prior* over the parameters of interests, which in the case of GPs is the function itself. The prior is important because it characterises the search space over possible solutions for  $f$ . Through the prior, we can encode strong assumptions, such as linearity, differentiability, periodicity, etc. or any combination thereof, which makes it possible to generalise well from very limited data. Compared to (Bayesian) parametric models, it is much more convenient and intuitive to specify priors directly in *function-space*, rather than on the weights of a parametric model [Rasmussen and Williams, 2006].

Following eq. (2.1) the prior over function evaluations at the datapoints is defined by the covariance function  $k$ . As we assume a  $\hat{a}$ -priori zero mean function  $\mu$  (without loss of generality) this can be written as:

$$p(f(\mathbf{x})) = \mathcal{N}(\mathbf{0}, \mathbf{K}_{\mathbf{ff}}), \quad \text{where} \quad [\mathbf{K}_{\mathbf{ff}}]_{i,j} = k(x_i, x_j). \quad (2.5)$$

Given the function  $f$  the likelihood factorises over datapoints and is given by a Gaussian:

$$p(\mathbf{y} | f) = \prod_{i=1}^N p(y_i | f) = \prod_{i=1}^N \mathcal{N}(y_i | f(x_i), \sigma^2) \quad (2.6)$$

We can obtain the posterior over the function using Bayes' rule and the marginalisation property

$$p(f | \mathbf{y}) = \frac{p(f) p(\mathbf{y} | f)}{p(\mathbf{y})} \quad (2.7)$$

$$= p(f \setminus \mathbf{x} | f(\mathbf{x})) \frac{p(f(\mathbf{x})) \prod_{i=1}^N \mathcal{N}(y_i | f(x_i), \sigma^2)}{p(\mathbf{y})} \quad (2.8)$$

$$= \mathcal{GP}(\mathbf{k}_{\mathbf{f}}^\top \mathbf{K}_{\mathbf{ff}}^{-1} \mathbf{f}, \quad k(\cdot, \cdot) - \mathbf{k}_{\mathbf{f}}^\top \mathbf{K}_{\mathbf{ff}}^{-1} \mathbf{k}_{\mathbf{f}}), \quad (2.9)$$

The marginal likelihood (model evidence)

$$p(\mathbf{y}) = \mathcal{N}(\mathbf{y} | \mathbf{0}, \mathbf{K}_{\mathbf{ff}} + \sigma^2 \mathbf{I}) \quad (2.10)$$

1. Plot: Prior, Data, Posterior
2. occam's razor

**Problems** A common criticism for GPs is that any modification to this approach breaks the Gaussian assumption.

1. Non-Gaussian likelihoods
2. Large datasets

3. Transformations: log or square transform

### Solutions

1. Laplace
2. Expectation Propagation
3. Sparse Variational Inference

## 2.2 Approximate Inference with Sparse Gaussian Processes

1. General introduction to Variational inference [Blei et al., 2017] variational inference (VI), where the problem of Bayesian inference is cast as an optimization problem—namely, to maximize a lower bound of the logarithm of the marginal likelihood.
2. Sparse approximations [Snelson and Ghahramani, 2005; Quiñonero-Candela and Rasmussen, 2005]

## 2.3 Interdomain Inducing Variables

### 2.3.1 Example: heavyside inducing variable

$f \sim \mathcal{GP}$  defined on  $\mathbb{S}^1$  (the unit circle),  $f : [-\pi, \pi] \rightarrow \mathbb{R}, \theta \mapsto f(\theta)$ .

kernel (Arc Cosine order 0):

$$k(\theta, \theta') = \kappa(\rho) = \pi - |\theta - \theta'| \quad (2.11)$$

$$\frac{d}{d\theta} \Big|_{\theta=\theta_m} k(\theta, \theta') = \quad (2.12)$$

$$u_m = \mathcal{L}_m(f) \quad (2.13)$$

$$\mathcal{L}_m = \frac{d}{d\theta} \Big|_{\theta=\theta_m} + \int d\theta \quad (2.14)$$

$$\text{Cov}(u_m, f(\theta')) = \mathbb{E}_f[\mathcal{L}_m(f) f(\theta')] \quad (2.15)$$

$$= \mathcal{L}_m k(\theta', \cdot) \quad (2.16)$$

$$= \frac{d}{d\theta} \Big|_{\theta=\theta_m} k(\theta', \theta) + \int_{-\pi}^{\pi} k(\theta', \theta) d\theta \quad (2.17)$$

## 2.4 Deep Gaussian Processes

Vincent Dutoit, Hugh Salimbeni, Eric Hambro, John McLeod, Felix Leibfried, Artem Artemev, Mark van der Wilk, James Hensman, Marc P Deisenroth, and ST John [2021b]. “GPflux: A Library for Deep Gaussian Processes”. In: *arXiv preprint arXiv:2003.01115*

## 2.5 Covariance Functions

1. Positive Definite and Symmetry
2. RKHS
3. Bochner’s theorem
4. Mercer Decomposition
5. Examples of RKHS
6. RKHS through Spectral Decomposition
7. Representer Theorem
8. Show how sparse approximation links anchor points



## Chapter 3

# Spherical Harmonic Variational Gaussian Processes

1. Related work: Variational Fourier Features
2. Zonal kernels
  - (a) Examples
  - (b) Decomposition in spherical harmonics
  - (c) Laplace Beltrami operator and Zonal kernel operator commute (this is why they share the same eigenfeatures)
3. How to compute the Spherical Harmonics: greedy algorithm
4. Experiments

### 3.1 Spherical Harmonics

Spherical harmonics are a special set of functions defined on the hypersphere and play a central role in harmonic analysis and approximation theory [Wendland, 2005]. They originate from solving Laplace’s equation, and form a complete set of orthogonal functions. Any sufficiently regular function defined on the sphere can be written as a sum of these spherical harmonics, similar to the Fourier series with sines and cosines. Spherical harmonics are defined in arbitrary dimensions [Efthimiou and Frye, 2014; Dai and Xu, 2013], but lack practical implementations in dimensions larger than three. This is mainly due to the complexity of solving Laplace’s equation in higher dimensions.

In this section, we propose a novel algorithm to construct spherical harmonics in  $d$  dimensions. The algorithm is based on the existence of a fundamental system of points which we select in a greedy fashion. The result are spherical harmonics that are a linear combination of zonal

functions and form an orth-normal basis on  $\mathbb{S}^{d-1}$ . The algorithm lends itself well for a practical implementation in Python and TensorFlow, which we provide at: <https://github.com/vdutor/SphericalHarmonics>. The code is accompanied by a series of tests that show the properties of the functions, as detailed below. Before outlining the algorithm in XXX we briefly define spherical harmonics in  $\mathbb{R}^d$ . We refer the interested reader to Dai and Xu [2013] and Efthimiou and Frye [2014] for a comprehensive review.

We adopt the usual  $L_2$  inner product for functions  $f : \mathbb{S}^{d-1} \rightarrow \mathbb{R}$  and  $g : \mathbb{S}^{d-1} \rightarrow \mathbb{R}$  restricted to the sphere

$$\langle f, g \rangle_{L_2(\mathbb{S}^{d-1})} = \frac{1}{\Omega_{d-1}} \int_{\mathbb{S}^{d-1}} f(x) g(x) d\omega, \quad (3.1)$$

where  $d\omega(x)$  is the surface area measure such that  $\Omega_{d-1}$  denotes the surface area of  $\mathbb{S}^{d-1}$

$$\Omega_{d-1} = \int_{\mathbb{S}^{d-1}} d\omega(x) = \frac{2\pi^{d/2}}{\Gamma(d/2)}. \quad (3.2)$$

For  $x = (x_1, \dots, x_d) \in \mathbb{R}^d$  and  $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{N}^d$ , a monomial  $x^\alpha$  is a product  $x^\alpha = x_1^{\alpha_1} \dots x_d^{\alpha_d}$ , which has degree  $|\alpha| = \alpha_1 + \dots + \alpha_d$ . A real homogeneous polynomial  $P(x)$  of degree  $n$  is a linear combination of monomials of degree  $n$  with real coefficients, that is  $P(x) = \sum_{|\alpha|=n} c_\alpha x^\alpha$ , with  $c_\alpha \in \mathbb{R}$ . We denote  $\mathcal{P}_n^d$  as the space of real homogeneous polynomials of degree  $n$ , and can show that, counting the cardinality of the set  $\{\alpha \in \mathbb{N}^d : |\alpha| = n\}$ , that  $\dim(\mathcal{P}_n^d) = \binom{n+d-1}{n}$ . A function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is said to be *harmonic* if  $\Delta f = 0$ , where  $\Delta = \partial_{x_1}^2 + \dots + \partial_{x_d}^2$  and  $\partial_{x_i}$  the partial derivate w.r.t. the  $i$ -th variable.

**Definition 1.** The spherical harmonics of degree  $n$  of  $d$  variables, denoted by  $\mathcal{H}_n^d$ , is the linear space of harmonic and homogeneous in degree  $n$  polynomials on  $\mathbb{S}^{d-1}$ , that is

$$\mathcal{H}_n^d = \{p \in \mathcal{P}_n^d : \Delta p = 0 \text{ and } p : \mathbb{S}^{d-1} \rightarrow \mathbb{R}\}. \quad (3.3)$$

The dimensionality of  $\mathcal{H}_n^d$  is given by

$$\dim(\mathcal{H}_n^d) = \frac{2n+d-2}{n} \binom{n+d-3}{d-1} := N_n^d. \quad (3.4)$$

The space  $\mathcal{H}_n^d$  has an orthonormal basis consisting of  $N_n^d$  functions, denoted by  $\{\phi_{n,j}\}_{j=1}^{N_n^d}$ . This implies that

$$\mathcal{H}_n^d = \text{span}(\phi_{n,1}, \dots, \phi_{n,N_n^d}), \quad \text{and} \quad \langle \phi_{n,j}, \phi_{n',j'} \rangle_{L_2(\mathbb{S}^{d-1})} = \delta_{nn'} \delta_{jj'}. \quad (3.5)$$

From the completeness and orthonormality of the spherical harmonic basis  $\{\phi_{n,j}\}_{n=0,j=1}^{\infty,N_n^d}$ , it can be shown that they also form a basis of square-integrable functions [Efthimiou and Frye,



2014]. This means that we can decompose a function  $f : \mathbb{S}^{d-1} \rightarrow \mathbb{R}$  as

$$f = \sum_{n=0}^{\infty} \sum_{j=1}^{N_n^d} \hat{f}_{n,j} \phi_{n,j}, \quad \text{with} \quad \hat{f}_{n,j} = \langle f, \phi_{n,j} \rangle_{L_2(\mathbb{S}^{d-1})}, \quad (3.6)$$

which can be seen as the spherical analogue of the Fourier decomposition of periodic functions onto a basis of sines and cosines. TODO: discuss uniqueness: any rotations is also a basis.

Subsequently, we will coin the set  $\{\phi_{n,j}\}$  as the spherical harmonics. They are indexed by  $n$  and  $j$ , where  $n = 0, 1, 2, \dots$  denotes the degree (or level) and  $j = 1, \dots, N_n^d$  denotes the orientation of the spherical harmonic. We are interested in finding  $\{\phi_{n,j}\}$  in arbitrary dimension. For  $d = 2$ , is solving Laplace's equation ( $\Delta p = 0$ ) directly relatively straightforward. Doing so reveals that  $N_0^2 = 1$  with  $\phi_{0,1} = 1$  and  $N_n^2 = 2$  for all  $n > 0$  with  $\phi_{n,1}(\theta) = \sqrt{2} \cos(n\theta)$  and  $\phi_{n,2}(\theta) = \sqrt{2} \sin(n\theta)$ . This shows that on the unit circle  $\mathbb{S}^1$ , the spherical harmonics correspond to the Fourier basis. For  $d = 3$ , we can also directly solve Laplace's differential equation to find  $N_n^d = 2n + 1$  and a closed form solution for  $\{\phi_{n,j}\}$ . However, for  $d > 3$ , explicit formulations for the spherical harmonics become very rare. To the best of our knowledge, the only explicit formulation we could find is in Dai and Xu [2013, Theorem 5.1], which consists of a product over polynomials. This makes the implementation and its derivate numerically unstable, and only practically useful up to 10 dimensions [Dutordoir et al., 2020]. To present our algorithm we first need to cover a few more properties of spherical harmonics.

**Theorem 1** (Addition). *Let  $\{\phi_{n,j}\}_{j=1}^{N_n^d}$  be an orthonormal basis for the spherical harmonics of degree  $n$  and  $x, x' \in \mathbb{S}^{d-1}$ . Then the Gegenbauer polynomial  $C_n^{(\alpha)} : [-1, 1] \rightarrow \mathbb{R}$  of degree  $n$  can be written as*

$$\sum_{j=1}^{N_n^d} \phi_{n,j}(x) \phi_{n,j}(x') = \frac{n + \alpha}{\alpha} C_n^{(\alpha)}(x^\top x') \quad \text{with} \quad \alpha = \frac{d-2}{2}. \quad (3.7)$$

As a result of the relation between the Gegenbauer polynomial and the spherical harmonics, are the Gegenbauer polynomial sometimes referred to as ultraspherical polynomials. For  $d = 2$ , Theorem 1 recovers the addition formula of the cosine function, as indeed  $\cos(\theta) \cos(\theta') + \sin(\theta) \sin(\theta') = \cos(\theta - \theta')$  and  $C_n^{(0)}(t) = \cos(n \arccos(t))$ . The Gegenbauer polynomials with  $\alpha = 0$  are better known as the Chebyshev polynomials. Another connection between spherical harmonics and Gegenbauer polynomials is given by the Funk-Hecke theorem and applies to zonal functions. A zonal function on  $\mathbb{S}^{d-1}$  is a function that is rotationally invariant w.r.t. to a vector  $\eta \in \mathbb{S}^{d-1}$ . This means that the function only depends on the inner product  $\eta^\top x$ .

**Theorem 2** (Funk-Hecke). *Let  $f$  be an integrable function such that  $\int_{-1}^1 \|f(t)\| (1-t^2)^{(d-3)/2} dt$  is finite and  $d \geq 2$ . Then for every  $\phi_{n,j}$*

$$\frac{1}{\Omega_{d-1}} \int_{\mathbb{S}^{d-1}} f(x^\top x') \phi_{n,j}(x') d\omega(x') = \lambda_n \phi_{n,j}(x), \quad (3.8)$$

where  $\lambda_n$  is a constant defined by

$$\lambda_n = \frac{\omega_d}{C_n^{(\alpha)}(1)} \int_{-1}^1 f(t) C_n^{(\alpha)}(t) (1-t^2)^{\frac{d-3}{2}} dt, \quad (3.9)$$

with  $\alpha = \frac{d-2}{2}$  and  $\omega_d = \frac{\Omega_{d-2}}{\Omega_{d-1}}$ .

Zonal function?

In light of the close connection between spherical harmonics and Gegenbauer polynomials

Construct a basis for spherical harmonics  $\mathcal{H}_n^d$  that consists of a linear combination of zonal functions

**Definition 2** (Fundamental System). *A collection of points  $\{\eta_1, \dots, \eta_N\} \in \mathbb{S}^{d-1}$  is called a fundamental system of degree  $n$  on the sphere if*

$$\det\left([C_n^{(\alpha)}(\eta_i^\top \eta_j)]_{i,j=1}^N\right) > 0, \quad [C_n^{(\alpha)}(\eta_i^\top \eta_j)]_{i,j=1}^N = \begin{bmatrix} C_n^{(\alpha)}(1) & \dots & C_n^{(\alpha)}(\eta_1^\top \eta_N) \\ \vdots & & \vdots \\ C_n^{(\alpha)}(\eta_N^\top \eta_1) & \dots & C_n^{(\alpha)}(1) \end{bmatrix} \quad (3.10)$$

**Lemma 3.** *There exists a fundamental system of degree  $n$  on the sphere.*

**Theorem 4.** *If  $\{\eta_1, \dots, \eta_N\} \in \mathbb{S}^{d-1}$  is a fundamental system of points on the sphere, then  $\{C_n^{(\alpha)}(\eta_i \cdot)\}_{i=1}^N$  is a basis for  $\mathcal{H}_n^d$ .*

Orth-normal basis using Gram-Schmidt.

### 3.1.1 How do we construct a fundamental system of points?

$$\mathbf{C}_n^{(\alpha)}(\boldsymbol{\eta}, \eta_*) = \left[ \begin{array}{c|c} C_n^{(\alpha)}(\boldsymbol{\eta} \boldsymbol{\eta}^\top) \in \mathbb{R}^{M \times M} & C_n^{(\alpha)}(\boldsymbol{\eta} \eta_*^\top) \in \mathbb{R}^{M \times 1} \\ \hline C_n^{(\alpha)}(\boldsymbol{\eta} \eta_*^\top)^\top \in \mathbb{R}^{1 \times M} & C_n^{(\alpha)}(1) \in \mathbb{R} \end{array} \right] \quad (3.11)$$

Determinant using Schur complement

$$\det(\mathbf{C}_n^{(\alpha)}(\boldsymbol{\eta}, \eta_*)) = \det(C_n^{(\alpha)}(\boldsymbol{\eta} \boldsymbol{\eta}^\top)) \left( C_n^{(\alpha)}(1) - C_n^{(\alpha)}(\eta_* \boldsymbol{\eta}^\top) [C_n^{(\alpha)}(\boldsymbol{\eta} \boldsymbol{\eta}^\top)]^{-1} C_n^{(\alpha)}(\boldsymbol{\eta} \eta_*^\top) \right) \quad (3.12)$$

$$\boldsymbol{\eta} = \operatorname{argmax}_{\eta_* \in \mathbb{S}^{d-1}} \det(\mathbf{C}_n^{(\alpha)}(\boldsymbol{\eta}, \eta_*)) \quad (3.13)$$

$$= \operatorname{argmin}_{\eta_* \in \mathbb{R}^d} C_n^{(\alpha)}\left(\frac{\eta_*}{\|\eta_*\|} \boldsymbol{\eta}^\top\right) [C_n^{(\alpha)}(\boldsymbol{\eta} \boldsymbol{\eta}^\top)]^{-1} C_n^{(\alpha)}\left(\boldsymbol{\eta} \frac{\eta_*^\top}{\|\eta_*\|}\right) \quad (3.14)$$

## Chapter 4

# Deep Neural Networks as Point Estimates for Deep Gaussian Processes

1. Introduction
2. Related work: connection between BNN and GPs
3. Activated Interdomain features
4. Experiments



# Chapter 5

## Future Research

### 5.1 Geometric Gaussian Decision Systems

#### Problem setting

1. Black Box Functions  $f : \mathcal{X} \rightarrow \mathcal{Y}$
2. We want to estimate a computable property  $\mathcal{O}_{\mathcal{A}}(f)$
3.  $\mathcal{A}$  is an algorithm  $\mathcal{O}_{\mathcal{A}}(f) = \mathcal{A}(f)$
4. Evaluating  $f$  is *very* expensive (we can only evaluate it a limited amount of times)

#### Examples

1. Bayesian Optimisation:  $\mathcal{A}(f) = \operatorname{argmax}_{x \in \mathcal{X}} f(x)$ , which implies  $\mathcal{O}_{\mathcal{A}}(f) = x^*$ .
2. Sensor Placement (Active Learning):  $\mathcal{O}_{\mathcal{A}}(f) = \operatorname{argmax}_{X \subset \mathcal{X}, |X|=T} \operatorname{MI}(f, f(X))$ .
3. Level sets:  $\mathcal{O}_{\mathcal{A}}(f) = \{X \subset \mathcal{X} : f(x) > C, \forall x \in X\}$ .
4. Shortest path:  $\mathcal{O}_{\mathcal{A}}(f) = \text{shortest path between two nodes in a graph.}$

**Model** We model  $f$  by a Gaussian process

$$f \sim \mathcal{GP} \tag{5.1}$$

1. Low dimensions
2. Prior knowledge
3. Limited and very expensive data

Basically settings where DNNs are never going to be competitive with GPs - low-dim, very data-efficient, high-cost - not even if someone figures out how to do DNN uncertainty right, due to GP regret guarantees (under reasonable assumptions) matching the best possible regret achievable by any model/decision system.




Fig. 5.1 Domains (top) and applications (bottom)

## Objectives

1. Theory and analysis
2. Getting these ideas into people's hands to get applications off the ground
  - (a) Graph GPs for combinatorial optimization use cases
  - (b) Manifold GPs for scientific use cases

## Collaborators

1. Alex Terenin (Imperial College London)
2. Willie Neiswanger (Stanford University)

## 5.2 Projects in Progress

1. “Pay Attention to Deep Gaussian Processes”

Transformer Layer Gaussian Processes using an explicit feature representation of the attention operation.

$$\exp(\mathbf{x}^\top \mathbf{y}) = \Phi^\top(\mathbf{x})\Phi(\mathbf{y})$$

2. A Unifying Theory for Interdomain Gaussian Processes.
3. VISH-PI: Probabilistic Integration with Variational Inducing Spherical Harmonics.





# References

- David M Blei, Alp Kucukelbir, and Jon D McAuliffe (2017). “Variational Inference: A Review for Statisticians”. In: *Journal of the American Statistical Association*.
- Feng Dai and Yuan Xu (2013). *Approximation Theory and Harmonic Analysis on Spheres and Balls*. Springer.
- Andreas Damianou and Neil D. Lawrence (2013). “Deep Gaussian Processes”. In: *Proceedings of the 16th International Conference on Artificial Intelligence and Statistics (AISTATS)*.
- Vincent Dutordoir, Nicolas Durrande, and James Hensman (2020). “Sparse Gaussian Processes with Spherical Harmonic Features”. In: *Proceedings of the 37th International Conference on Machine Learning (ICML)*.
- Vincent Dutordoir, James Hensman, Mark van der Wilk, Carl Henrik Ek, Zoubin Ghahramani, and Nicolas Durrande (2021a). “Deep Neural Networks as Point Estimate for Deep Gaussian Processes”. In: *submission to NeurIPS*.
- Vincent Dutordoir, Hugh Salimbeni, Eric Hambro, John McLeod, Felix Leibfried, Artem Artemev, Mark van der Wilk, James Hensman, Marc P Deisenroth, and ST John (2021b). “GPflux: A Library for Deep Gaussian Processes”. In: *arXiv preprint arXiv:2003.01115*.
- Costas Efthimiou and Christopher Frye (2014). *Spherical Harmonics in p Dimensions*. World Scientific Publishing.
- David J. C. MacKay (2003). *Information Theory, Inference and Learning Algorithms*. Cambridge University Press.
- Alexander G. de G. Matthews, James Hensman, Turner E. Richard, and Zoubin Ghahramani (2016). “On Sparse Variational Methods and the Kullback-Leibler Divergence between Stochastic Processes”. In: *Proceedings of the 19th International Conference on Artificial Intelligence and Statistics (AISTATS)*.
- Radford M. Neal (1992). “Bayesian Mixture Modeling”. In: *Maximum Entropy and Bayesian Methods*.
- Radford M. Neal (1995). *Bayesian Learning for Neural Networks*. Springer.
- Joaquin Quiñonero-Candela and Carl E. Rasmussen (2005). “A Unifying View of Sparse Approximate Gaussian Process Regression”. In: *Journal of Machine Learning Research*.
- Carl E. Rasmussen and Christopher K. I. Williams (2006). *Gaussian Processes for Machine Learning*. MIT Press.
- Edward Snelson and Zoubin Ghahramani (2005). “Sparse Gaussian Processes using Pseudo-inputs”. In: *Advances in Neural Information Processing Systems 4 (NIPS 2005)*.
- Holger Wendland (2005). *Scattered Data Approximation*. Cambridge University Press.
- Christopher K. I. Williams and Carl E. Rasmussen (1996). “Gaussian processes for regression”. In: