Fitting Models to Data in Ecology and Evolution

Samraat Pawar

Imperial College London

October 22, 2018

What does "modelling data" mean to you?


 Mechanistic models aim to explain the PROCESSES or MECHANISMS underlying PATTERNS or PHENOMENA in empirical data

- Mechanistic models aim to explain the PROCESSES or MECHANISMS underlying PATTERNS or PHENOMENA in empirical data
 - These models have a THEORETICAL BASIS


- Mechanistic models aim to explain the PROCESSES or MECHANISMS underlying PATTERNS or PHENOMENA in empirical data
 - These models have a THEORETICAL BASIS
- Empirical/Phenomenological models establish the existence of STATISTICALLY SIGNIFICANT, NON-RANDOM PATTERNS or PHENOMENA in empirical data

- Mechanistic models aim to explain the PROCESSES or MECHANISMS underlying PATTERNS or PHENOMENA in empirical data
 - These models have a THEORETICAL BASIS
- Empirical/Phenomenological models establish the existence of STATISTICALLY SIGNIFICANT, NON-RANDOM PATTERNS or PHENOMENA in empirical data
 - They make no assumptions about the processes or mechanisms that generate the patterns

- Mechanistic models aim to explain the PROCESSES or MECHANISMS underlying PATTERNS or PHENOMENA in empirical data
 - These models have a THEORETICAL BASIS
- Empirical/Phenomenological models establish the existence of STATISTICALLY SIGNIFICANT, NON-RANDOM PATTERNS or PHENOMENA in empirical data
 - They make no assumptions about the processes or mechanisms that generate the patterns
 - That is, these models lack a THEORETICAL BASIS




source: https://www.cds.caltech.edu/~murray/amwiki/images/8/8f/LHgraph.gif



source: https://www.cds.caltech.edu/~murray/amwiki/images/8/8f/LHgraph.gif

• Mechanistic model: The Lynx-Hare Cycle is driven by density-dependent population growth in hares



source: https://www.cds.caltech.edu/~murray/amwiki/images/8/8f/LHgraph.gif

- Mechanistic model: The Lynx-Hare Cycle is driven by density-dependent population growth in hares
- Phenomenological model: The Lynx and Hare Cycles have a significant asynchrony (period shift) of xx years

• It's not really one vs. the other, Both types of models play a role in science (and Biology)

- It's not really one vs. the other, Both types of models play a role in science (and Biology)
- Phenomenological model-fitting reveals patterns in data that generate HYPOTHESES


- It's not really one vs. the other, Both types of models play a role in science (and Biology)
- Phenomenological model-fitting reveals patterns in data that generate HYPOTHESES
 - These can be tested using further model fitting

- It's not really one vs. the other; Both types of models play a role in science (and Biology)
- Phenomenological model-fitting reveals patterns in data that generate HYPOTHESES
 - These can be tested using further model fitting
 - Example: Whether climatic temperature affects the Lynx-Hare cycle (using Generalized Linear Model-fitting)

- It's not really one vs. the other, Both types of models play a role in science (and Biology)
- Phenomenological model-fitting reveals patterns in data that generate HYPOTHESES
 - These can be tested using further model fitting
 - Example: Whether climatic temperature affects the Lynx-Hare cycle (using Generalized Linear Model-fitting)
- Mechanistic model-fitting tries to validate a mechanistic model that can explain the observed phenomenological pattern and generate MORE ACCURATE, MECHANISTIC HYPOTHESES

- It's not really one vs. the other; Both types of models play a role in science (and Biology)
- Phenomenological model-fitting reveals patterns in data that generate HYPOTHESES
 - These can be tested using further model fitting
 - Example: Whether climatic temperature affects the Lynx-Hare cycle (using Generalized Linear Model-fitting)
- Mechanistic model-fitting tries to validate a mechanistic model that can explain the observed phenomenological pattern and generate MORE ACCURATE, MECHANISTIC HYPOTHESES
 - Example: How climatic temperature drives the Lynx-Hare cycle

- It's not really one vs. the other; Both types of models play a role in science (and Biology)
- Phenomenological model-fitting reveals patterns in data that generate HYPOTHESES
 - These can be tested using further model fitting
 - Example: Whether climatic temperature affects the Lynx-Hare cycle (using Generalized Linear Model-fitting)
- Mechanistic model-fitting tries to validate a mechanistic model that can explain the observed phenomenological pattern and generate MORE ACCURATE, MECHANISTIC HYPOTHESES
 - Example: How climatic temperature drives the Lynx-Hare cycle
- Ultimately, successful, EMPIRICALLY-GROUNDED mechanistic models are the best path towards a THEORY in any scientific discipline (including ecology and evolution)



 Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological Why?

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological Why?
 - Partly because we are still establishing the existence of GENERAL PATTERNS/PHENOMENA,

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological Why?
 - Partly because we are still establishing the existence of GENERAL PATTERNS/PHENOMENA,
 - ... and partly because we are (or are forced to be) interested in FORECASTING rather than EXPLAINING.

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological Why?
 - Partly because we are still establishing the existence of GENERAL PATTERNS/PHENOMENA,
 - ... and partly because we are (or are forced to be) interested in FORECASTING rather than EXPLAINING.
- So the big question is, can we FORECAST WITHOUT EXPLAINING?

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological Why?
 - Partly because we are still establishing the existence of GENERAL PATTERNS/PHENOMENA,
 - ... and partly because we are (or are forced to be) interested in FORECASTING rather than EXPLAINING.
- So the big question is, can we FORECAST WITHOUT EXPLAINING?
 - For example, disease outbreaks: Do we really need to care about the underlying mechanisms if we can predict a future event using phenomenological modelling (e.g., Machine-learning of time series patterns)?

Somewhat subjective!

- Somewhat subjective!
- For example, the Ricker model can be thought of as mechanistic:

$$N_{t+1} = N_t e^{r\left(1 - \frac{N_t}{k}\right)}$$

- Somewhat subjective!
- For example, the Ricker model can be thought of as mechanistic:

$$N_{t+1} = N_t e^{r\left(1 - \frac{N_t}{k}\right)}$$

• What is the mechanism?

- Somewhat subjective!
- For example, the Ricker model can be thought of as mechanistic:

$$N_{t+1} = N_t e^{r\left(1 - \frac{N_t}{k}\right)}$$

• What is the mechanism?

- Somewhat subjective!
- For example, the Ricker model can be thought of as mechanistic:

$$N_{t+1} = N_t e^{r\left(1 - \frac{N_t}{k}\right)}$$

- What is the mechanism? Density dependence through scramble competition (Brannstrom & Sumpter 2005)
- If the Ricker model and another model with contest competition were compared with data — some would call it mechanistic modelling because one is trying to get at the underlying mechanism, scramble or contest competition

- Somewhat subjective!
- For example, the Ricker model can be thought of as mechanistic:


$$N_{t+1} = N_t e^{r\left(1 - \frac{N_t}{k}\right)}$$

- What is the mechanism? Density dependence through scramble competition (Brannstrom & Sumpter 2005)
- If the Ricker model and another model with contest competition were compared with data — some would call it mechanistic modelling because one is trying to get at the underlying mechanism, scramble or contest competition
- But is this REALLY mechanistic? What are r and k really?




 Proponents of Metabolic Theory of Ecology argue that we have not progressed far enough towards mechanistic modelling because metabolism has been ignored

 Proponents of Metabolic Theory of Ecology argue that we have not progressed far enough towards mechanistic modelling because metabolism has been ignored



 Proponents of Metabolic Theory of Ecology argue that we have not progressed far enough towards mechanistic modelling because metabolism has been ignored



$$B = B_0 e^{-\frac{E}{kT}} f(T, T_{pk}, E_D)$$

T = temperature (K)

 $k = \text{Boltzmann constant (eV K}^{-1})$


E = Activation energy (eV)

 T_{pk} = Temperature of peak performance

 E_D = Deactivation energy (eV)

(J H van't Hoff 1884, S Arrhenius 1889)

 Proponents of Metabolic Theory of Ecology argue that we have not progressed far enough towards mechanistic modelling because metabolism has been ignored



$$B=B_0 e^{-\frac{E}{kT}} f(T,T_{pk},E_D)$$

T = temperature (K)

 $k = \text{Boltzmann constant (eV K}^{-1})$

E = Activation energy (eV)

 T_{pk} = Temperature of peak performance

 $F_{D} = Deactivation energy (eV)$

(J H van't Hoff 1884, S Arrhenius 1889)

- Surely there is more to thermal responses?
- What about alternative models?



• If possible, use biological knowledge to construct models

- If possible, use biological knowledge to construct models
- See if the models "agree well" with data

- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms

- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms
- That's the one that's best for predictions (e.g. population cycles), estimating rates (e.g. population or individual growth rates), etc

- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms
- That's the one that's best for predictions (e.g. population cycles), estimating rates (e.g. population or individual growth rates), etc
- Don't use models you already know have the wrong mechanisms just because they are popular!

- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms
- That's the one that's best for predictions (e.g. population cycles), estimating rates (e.g. population or individual growth rates), etc
- Don't use models you already know have the wrong mechanisms just because they are popular!
- Phenomenological/statistical models often perform better than mechanistic ones.



- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms
- That's the one that's best for predictions (e.g. population cycles), estimating rates (e.g. population or individual growth rates), etc
- Don't use models you already know have the wrong mechanisms just because they are popular!
- Phenomenological/statistical models often perform better than mechanistic ones.



- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms
- That's the one that's best for predictions (e.g. population cycles), estimating rates (e.g. population or individual growth rates), etc
- Don't use models you already know have the wrong mechanisms just because they are popular!
- Phenomenological/statistical models often perform better than mechanistic ones. Why?



- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms
- That's the one that's best for predictions (e.g. population cycles), estimating rates (e.g. population or individual growth rates), etc
- Don't use models you already know have the wrong mechanisms just because they are popular!
- Phenomenological/statistical models often perform better than mechanistic ones. Why? — because they have lest restrictive assumptions

MODELS: HOW TO BUILD THEM?

- It's an art, takes practice
- Build models one mechanism at a time in biology, it means start at the right level of organization!
- Always consider an alternative that is more parsimonious, even if it is phenomenological (the thermal performance curves example: Sharpe-Schoolfield, Briere, or Polynomial?)!
- For example, the Boltzmann-Arrhenius model is a good first try describe and uncover mechanisms underlying individual level rates (e.g., vector fecundity or development rate)
- The next step would be to include species interactions with temperature dependence of individuals (or go in an evolutionary direction)

FITTING MODELS TO DATA

Multiple ways to do it:

- Least Squares methods
 - Linear
 - Non-linear
- Likelihood-based methods
 - Maximum Likelihood Estimation (MLE)
 - Bayesian
- Artificial intelligence and Machine learning
 - Focus in on maximizing ability to discover pattern and predict at the cost of mechanistic insights

METHODS YOU CAN USE IN THE MINIPROJECT

- Least squares: along with Linear Model fitting, Non-linear Least Squares (NLLS) fitting is a particularly versatile and powerful approach because many mechanisms in biology and inherently non-linear
- MLE/Bayesian methods: more robust if you are able to calculate the likelihood function — you will learn this in Term 2, so not recommended
- Al/machine Learning: most versatile for large amounts of noisy data — you will be introduced to these at the end of term 2, so definitely not recommended

 Ideally, several competing (meaningful, not just null) hypotheses (mathematical models) should be fitted to data and compared using statistical theory

- Ideally, several competing (meaningful, not just null) hypotheses (mathematical models) should be fitted to data and compared using statistical theory
- This is an advance over the traditional "null hypothesis" approach in Biology

- Ideally, several competing (meaningful, not just null) hypotheses (mathematical models) should be fitted to data and compared using statistical theory
- This is an advance over the traditional "null hypothesis" approach in Biology
- Necessary for developing the advancement of Biology from from an observational and axiomatic discipline to one with general theories.

- Ideally, several competing (meaningful, not just null) hypotheses (mathematical models) should be fitted to data and compared using statistical theory
- This is an advance over the traditional "null hypothesis" approach in Biology
- Necessary for developing the advancement of Biology from from an observational and axiomatic discipline to one with general theories.
- Necessary for understanding the mechanisms underlying biological patterns/phenomena

READINGS

- Levins, R. (1966) The strategy of model building in population biology. Am. Sci. 54, 421–431.
- Johnson, J. B. & Omland, K. S. (2004) Model selection in ecology and evolution. Trends Ecol. Evol. 19, 101–108.
- Bolker, B. M. et al. (2013) Strategies for fitting nonlinear ecological models in R, AD Model Builder, and BUGS. Methods Ecol. Evol. 4, 501–512.
- Some illustrative examples of (non-linear) model fitting to ecological/evolutionary data https://groups.nceas.ucsb. edu/non-linear-modeling/projects
- Additional readings in the git repository