

# Evaluating the performance and energy efficiency of COSMO-ART, a fully online coupled model system composed of a numerical weather forecast model and a chemical transport model.

J.Charles · W.Sawyer · M.Dolz · C.Malossi

Received: date / Accepted: date

**Abstract** In this paper we present COSMO-ART, an extension of the operational weather forecast model of the German Weather Service (DWD), developed for the evaluation of the interactions of reactive gases and aerosol particles with the state of atmosphere at the regional scale. It includes secondary aerosols, directly emitted components like soot, mineral dust, sea salt and biological material as pollen. Processes such as emissions, coagulation, condensation, dry deposition, wet removal, and sedimentation of aerosols are taken into account. The overall performance of this application on HPC systems is analysed by a profiling and tracing study to determine hotspots and identify critical paths. Moreover, we describe measurement devices and energy-aware techniques employed to evaluate the en-

ergy footprint of the considered application and to get detailed insights about power bottlenecks. Our motivation is to improve corresponding code sections to sustain high performance while minimizing energy-to-solution. This preliminary study sets the basis of subsequent considerations to tackle challenges related to energy efficient high performance computing in the framework of the Exa2Green project (<http://exa2green.eu/>).

**Keywords** High performance computing · Energy-aware computing · Green computing · Numerical weather prediction · Atmospheric chemistry · Aerosols modelling · Profiling methods · Benchmark analysis · COSMO-ART coupled model

---

Grants or other notes about the article that should go on the front page should be placed here.

---

Dr. Joseph Charles  
Swiss National Supercomputing Centre (CSCS)  
CH-6900 Lugano, Switzerland  
Tel: +41 (0) 91 610 8216  
E-mail: joseph.charles@cscs.ch

Dr. William Sawyer  
Swiss National Supercomputing Centre (CSCS)  
CH-6900 Lugano, Switzerland  
Tel: +41 (0) 91 610 8229  
E-mail: william.sawyer@cscs.ch

Dr. Manuel Dolz  
University of Hamburg  
20148 Hamburg, Germany  
Tel: +49 (0) 40 460094-404  
E-mail: manuel.dolz@informatik.uni-hamburg.de

Dr. Cristiano Malossi  
IBM Zurich Research Laboratory  
CH-8000 Zurich, Switzerland  
Tel: +41 (0) 44 724 8616  
E-mail: ACM@zurich.ibm.com

## 1 Introduction

Comprehensive scientific assessments of climate-warming trends over the past 50 years have shown that the state and dynamics of the Earth's climate system have undergone unprecedented major changes, as evidenced by increases in global average atmosphere and ocean temperatures, widespread melting of snow and ice, and rising sea levels. The last report of the Intergovernmental Panel on Climate Change, released in September 2013 (6), concluded that most of the observed temperature increases since the mid 20th Century have been caused by increasing concentrations of greenhouse gases resulting from anthropogenic activities such as fossil fuel burning and deforestation. Long-lived greenhouse gases, for example carbon dioxide, methane and nitrous oxide, are chemically stable and linger homogeneously well-mixed and over time scales of a decade to thousands of years in the atmosphere; whereas short-lived gases such as water, sulphur dioxide and carbon monoxide which respond physically or chemically to changes

in temperature, act primarily as a feedback mechanism. Some extremely powerful greenhouse gases, such as fluorinated gases can even have a warming effect on the atmosphere up to 23000 times greater than carbon dioxide (28). Fortunately the European Parliament, which has gained worldwide recognition as a leader in climate policy, has passed legislation in March 2014 to phase out fluorinated gas emissions by two-thirds by 2030, towards the internationally agreed goal of keeping global warming below 2 degrees Celsius compared to the temperature in pre-industrial times.

Current cumulative releases of human-induced gas emissions enhances the greenhouse effect to the Earth's atmosphere and thus creates a large imbalance between incoming solar absorbed radiation and outgoing long-wave radiation emitted back to space, through radiative trapping. While the Earth's temperature is dependent upon the greenhouse-like action of the atmosphere, the Earth's radiation balance is strongly influenced by several other factors such as the type of surface that sunlight first encounters. Forests, grasslands, ocean surfaces, ice caps, deserts, and cities all absorb, reflect, and radiate radiation differently. Sunlight falling on a white glacier surface strongly reflects back into space, resulting in minimal heating of the surface and lower atmosphere. Sunlight falling on a dark desert soil is strongly absorbed, on the other hand, and contributes to significant heating of the surface and lower atmosphere. Cloud cover also affects greenhouse warming by both reducing the amount of solar radiation reaching the earth's surface and by reducing the amount of radiation energy emitted into space.

Aerosols are suspensions of liquid, solid or mixed particles in the air (including sea salt, mineral dust, sulphate, nitrate, organic carbon and black carbon), with highly variable chemical composition and size distribution (18). Although they are not considered a heat-trapping greenhouse gas and have shorter atmospheric lifetimes, they significantly affect the atmospheric radiative fluxes and are acknowledged as one of the most significant and uncertain aspects in anthropogenic forcing over the last 150 years (6, 11). Enhanced aerosols concentrations can impact the climate system by reflecting (e.g. pure sulfates and nitrates) or absorbing (e.g. black carbon) solar radiation and thereby exert a cooling or warming effect on the Earth-atmosphere system, causing a so-called direct radiative forcing (3–5, 14, 16, 19). Depending on their size and chemical composition, they can also act as cloud condensation and ice nuclei, and thereby influence the cloud microphysical processes and optical properties (cloud albedo

effect, 25) through the contribution to cloud formation. This results in changes in droplet concentrations (1) and precipitation (7, 8, 13, 17, 20, 22, 24), producing a so-called negative indirect radiative forcing (5, 15, 21, 26). Wet scavenging of aerosol particles represents a major removal mechanism for air pollutants from the atmosphere, however the effects of anthropogenic aerosols on clouds and the hydrological cycle as well as the cloud lifetime effect are especially hard to assess and quantify (6), and remain one of the largest uncertainties in climate modeling and in climate change prediction due to the lack of knowledge in cloud feedbacks (12, 21, 23).

Hence to improve our understanding of aerosol-cloud interactions and reduce uncertainties of aerosol effects in climate, the research community is making a concerted international effort to represent the underlying chemical processes through models. These models, such as ART (Aerosols and Reactive Trace gases) extension, developed at the Karlsruhe Institute of Technology (KIT), offer a key opportunity to reduce the climate uncertainty, particularly on the regional scale (2, 9, 10). These models can be coupled with climate models, such as the regional weather forecast model COSMO (Consortium for Small-scale Modelling), jointly developed by a consortium of European weather centers including the German weather service DWD and MeteoSwiss, and used in the climate version COSMO-CLM by a wide research community. The extended COSMO-ART model provides a detailed description of air pollution chemistry and aerosol processes, and is mainly designed to study air quality and aerosol meteorology feedbacks on short, episodic to annual time scales. It is capable of simulating aerosol distributions as well as their interactions over Europe as well as other regional domains.

COSMO-ART is computationally much more demanding than the COSMO core since a large number of additional tracers and processes have to be considered. Thus this model is currently severely-limited in terms of applicability and expensive in terms of energy consumption. COSMO has recently been ported to GPUs within the framework of the High Performance and High Productivity Computing (HP2C) Initiative to optimize it for computational and energy efficiency. Although these developments will facilitate the application of COSMO for numerical weather prediction and climate simulations, they do little to address the coupling with the ART model extension, for which significant investments are still required to take it to a similar level. The efficiency of ART is being addressed in the EU Exa2Green project. The ultimate goal of the project is to deliver

a prototype code, which provides an energy efficiency of at least five times of the baseline value. Such an implementation would allow the community to investigate critical questions at higher resolution and over longer periods, at reduced cost to the environment. In Sec. 2, we introduce the online coupled mesoscale atmosphere and chemistry model COSMO-ART (27) that accounts for feedbacks between chemistry, aerosols, radiation, and clouds. Sec. 3 and Sec. 4 successively describe the hardware and measurement devices used in evaluation as well as energy-aware techniques employed to determine the energy footprint of the considered application. Sec. 5 shows benchmark results from the Exa2Green project and highlight the critical components of the application. Finally, a summary and directions for model optimization are given in Sec. 6.

## 2 The operational COSMO-ART

### 2.1 Model description

### 2.2 Model setup

## 3 Hardware description

## 4 Power measurement systems

## 5 Benchmark results

## 6 Conclusion

**Acknowledgements** This research was supported by the Exa2Green research project co-funded under the EU 7th Framework Program Future and Emerging Technologies (FET) Proactive Initiative: Minimising Energy Consumption of Computing to the Limit (MINECC). The authors would like to thank the High Performance and High Productivity Computing Initiative ([www.hp2c.ch](http://www.hp2c.ch)) for results that will be leveraged in subsequent code refactoring.

## References

1. B. A. Albrecht. Aerosols, Cloud Microphysics, and Fractional Cloudiness. *Science*, 15(4923):1227–1230, 1989. doi: 10.1126/science.245.4923.1227.
2. M. Bangert, C. Kottmeier, B. Vogel, and H. Vogel. Regional scale effects of the aerosol cloud interaction simulated with an online coupled comprehensive chemistry model. *Atmospheric Chemistry and Physics*, 11:4411–4423, 2011. doi: 10.5194/acp-11-4411-2011.
3. M. Bangert, A. Nenes, B. Vogel, H. Vogel, D. Barahona, V. A. Karydis, P. Kumar, C. Kottmeier, and U. Blahak. Saharan dust event impacts on cloud formation and radiation over Western Europe. *Atmospheric Chemistry and Physics Discussions*, 11(12):31937–31982, 2011. doi: 10.5194/acpd-11-31937-2011.
4. R. J. Charlson et al. Perturbation of the Northern Hemisphere radiative balance by backscattering from anthropogenic sulfate aerosols. *Tellus Series A Dynamic Meteorology and Oceanography*, 43(4):152–163, 1991. doi: 10.1034/j.1600-0870.1991.00013.x.
5. J. M. Haywood and O. Boucher. Estimates of the direct and indirect radiative forcing due to tropospheric aerosols: A review. *Reviews of Geophysics*, 38(4):513–543, 2000. doi: 10.1029/1999RG000078.
6. IPCC - Intergovernmental Panel on Climate Change. *Climate Change 2013: The Physical Science Basis*. Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
7. A. Khain, N. Benmoshe, and A. Pokrovsky. Factors determining the impact of aerosols on surface precipitation from clouds: an attempt at classification. *Journal of the Atmospheric Sciences*, 65(6):1721–1748, 2008. doi: 10.1175/2007JAS2515.1.
8. A. P. Khain. Notes on state-of-the-art investigations of aerosol effects on precipitation: a critical review. *Environmental Research Letters*, 4(015004), 2009. doi: 10.1088/1748-9326/4/1/015004.
9. C. Knöte and D. Brunner. An advanced scheme for wet scavenging and liquid-phase chemistry in a regional online-coupled chemistry transport model. *Atmospheric Chemistry and Physics*, 13:1177–1192, 2013. doi: 10.5194/acp-13-1177-2013.
10. C. Knöte, D. Brunner, H. Vogel, J. Allan, A. Asmi, M. Äijälä, S. Carbone, H. D. van der Gon, J. L. Jimenez, A. Kiendler-Scharr, C. Mohr, L. Poulain, A. S. H. Prévôt, E. Swietlicki, and B. Vogel. Towards an online-coupled chemistry-climate model: evaluation of trace gases and aerosols in COSMO-ART. *Geoscientific Model Development*, 4(4):1077–1102, 2011. doi: 10.5194/gmd-4-1077-2011.
11. D. Koch et al. Distinguishing Aerosol Impacts on Climate over the Past Century. *Journal of Climate*, 22(10):2659–2677, 2009. doi: 10.1175/2008JCLI2573.1.
12. L. A. Lee, K. J. Pringle, C. L. Reddington, G. W. Mann, P. Stier, D. V. Spracklen, J. R. Pierce, and K. S. Carslaw. The magnitude and causes of uncertainty in global model simulations of cloud condensation nuclei. *Atmospheric Chem-*

- istry and Physics*, 13:8879–8914, 2013. doi: 10.5194/acp-13-8879-2013.
13. S. S. Lee, G. Feingold, and P. Y. Chuang. Effect of aerosol on cloud-environment interactions in trade cumulus. *Journal of the Atmospheric Sciences*, 69: 3607–3632, 2012. doi: 10.1175/JAS-D-12-026.1.
14. H. Liao and J. H. Seinfeld. Global impacts of gas-phase chemistry-aerosol interactions on direct radiative forcing by anthropogenic aerosols and ozone. *Journal of Geophysical Research-Atmospheres*, 110(D18), 2005. doi: 10.1029/2005JD005907.
15. U. Lohmann and J. Feichter. Global indirect aerosol effects: a review. *Atmospheric Chemistry and Physics*, 5:715–737, 2005. doi: 10.5194/acp-5-715-2005.
16. K. Lundgren, B. Vogel, H. Vogel, and C. Kottmeier. Direct radiative effects of sea salt for the mediterranean region at conditions of low to moderate wind speeds. *Journal of Geophysical Research*, 118(4): 1906–1923, 2013. doi: 10.1029/2012JD018629.
17. H. R. Pruppacher and J. D. Klett. *Microphysics of Clouds and Precipitation*, volume 18 of *Atmospheric and Oceanographic Sciences Library*. Kluwer Academics, 2010.
18. J.-P. Putaud et al. A European aerosol phenomenology 3: Physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe. *Atmospheric Environment*, 44(10):1308–1320, 2010. doi: 10.1016/j.atmosenv.2009.12.011.
19. V. Ramanathan et al. Indian Ocean Experiment: An integrated analysis of the climate forcing and effects of the great Indo-Asian haze. *Journal of Geophysical Research-Atmospheres*, 106(D22): 28371–28398, 2001.
20. D. Rosenfeld. Suppression of Rain and Snow by Urban and Industrial Air Pollution. *Science*, 287(5459):1793–1796, 2000. doi: 10.1126/science.287.5459.1793.
21. D. Rosenfeld, R. Wood, L. J. Donner, and S. C. Sherwood. Aerosol Cloud-Mediated Radiative Forcing: Highly Uncertain and Opposite Effects from Shallow and Deep Clouds. *Climate Science for Serving Society: Research, Modelling and Prediction Priorities*, pages 105–149, 2013. doi: 10.1007/978-94-007-6692-1.5.
22. A. Seifert, C. Köhler, and K. D. Beheng. Aerosol-cloud-precipitation effects over Germany as simulated by a convective-scale numerical weather prediction model. *Atmospheric Chemistry and Physics*, 12(2):709–725, 2012. doi: 10.5194/acp-12-709-2012.
23. S. C. Sherwood, M. J. Alexander, A. R. Brown, N. A. McFarlane, E. P. Gerber, G. Feingold, A. A. Scaife, and W. W. Grabowski. Climate Processes: Clouds, Aerosols and Dynamics. *Climate Science for Serving Society: Research, Modelling and Prediction Priorities*, pages 73–103, 2013. doi: 10.1007/978-94-007-6692-1.4.
24. W.-K. Tao, J. P. Chen, Z. Li, C. Wang, and C. Zhang. Impact of aerosols on convective clouds and precipitation. *Reviews of Geophysics*, 50(2), 2012. doi: 10.1029/2011RG000369.
25. S. Twomey. The influence of pollution on the short-wave albedo of clouds. *Journal of the Atmospheric Sciences*, 34(7):1149–1152, 1977. doi: 10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2.
26. S. C. Van den Heever, G. L. Stephens, and N. B. Wood. Aerosol indirect effects on tropical convection characteristics under conditions of radiative-convective equilibrium. *Journal of the Atmospheric Sciences*, 68(4):699–718, 2011. doi: 10.1175/2010JAS3603.1.
27. B. Vogel, H. Vogel, D. Bäumer, M. Bangert, K. Lundgren, R. Rinke, and T. Stanelle. The comprehensive model system COSMO-ART radiative impact of aerosol on the state of the atmosphere on the regional scale. *Atmospheric Chemistry and Physics Discussions*, 9(4):14483–14528, 2009. doi: 10.5194/acpd-9-14483-2009.
28. O. Zehner. *Green Illusions: The Dirty Secrets of Clean Energy and the Future of Environmentalism*. Technology and Engineering. University of Nebraska Press, 2012.