

ASSUMPTIONS WITH EQUILIBRIUM

ASSUMPTIONS WITH EQUILIBRIUM


Example #1

At 2000°C, K_{eq} is 6.40×10^{-7} for the decomposition of CO_2 into CO and O_2 . Calculate all equilibrium concentrations if 0.250 mol of CO_2 is placed in a 1.00 L container at the given temperature.

ASSUMPTIONS WITH EQUILIBRIUM

Example #1

At 2000°C, K_{eq} is 6.40×10^{-7} for the decomposition of CO_2 into CO and O_2 .
Calculate all equilibrium concentrations if 0.250 mol of CO_2 is placed in a 1.00 L container at the given temperature.




How do we solve this cubic equation?!

ASSUMPTIONS WITH EQUILIBRIUM

Example #1

At 2000°C, K_{eq} is 6.40×10^{-7} for the decomposition of CO_2 into CO and O_2 . Calculate all equilibrium concentrations if 0.250 mol of CO_2 is placed in a 1.00 L container at the given temperature.



Look at the K_{eq} (which is 0.000000640): VERY little of the CO_2 actually decomposes.

So x must be a VERY VERY small number.

So we assume that $0.250-2x = 0.250$

It's **as if** we're assuming that x is 0.

ASSUMPTIONS WITH EQUILIBRIUM

Example #1



I	0.250	0	0
C	-2x	+2x	+x
E	0.250-2x	2x	x

$$K_{\text{eq}} = \frac{[\text{CO}]^2[\text{O}_2]}{[\text{CO}_2]^2}$$

$$6.40 \times 10^{-7} = \frac{[2x]^2[x]}{[0.250-2x]^2}$$

Assumption: $0.250-2x = 0.250$

$$6.40 \times 10^{-7} = \frac{[2x]^2[x]}{[\mathbf{0.250}]^2}$$

$$6.40 \times 10^{-7} = \frac{4x^3}{0.0625}$$

$$4.00 \times 10^{-8} = 4x^3$$

$$1.00 \times 10^{-8} = x^3$$

$$\sqrt[3]{(1.00 \times 10^{-8})} = \sqrt[3]{(x^3)}$$

$$2.15 \times 10^{-3} = x$$

ASSUMPTIONS WITH EQUILIBRIUM

Example #1

At 2000°C, K_{eq} is 6.40×10^{-7} for the decomposition of CO_2 into CO and O_2 .

Calculate all equilibrium concentrations

if 0.250 mol of CO_2 is placed in a 1.00 L container at the given temperature.

	$2\text{CO}_{2(g)}$	\rightleftharpoons	$2\text{CO}_{(g)}$	$+$	$\text{O}_{2(g)}$
I	0.250		0		0
C	-2x		+2x		+x
E	0.250-2x		2x		x

$$x = 2.15 \times 10^{-3}$$

$$\begin{aligned} &= 0.250 - 2(2.15 \times 10^{-3}) \\ &= 0.247\text{M} \end{aligned}$$

$$\begin{aligned} &= 2(2.15 \times 10^{-3}) \\ &= 0.0043\text{M} \end{aligned}$$

$$\therefore [\text{CO}_{2(g)}] = 2.47 \times 10^{-1}\text{M}, [\text{CO}_{(g)}] = 4.30 \times 10^{-3}\text{M}, \text{ \& } [\text{O}_{2(g)}] = 2.15 \times 10^{-3}\text{M}$$

ASSUMPTIONS WITH EQUILIBRIUM

Example #1

	$2\text{CO}_{2(g)} \rightleftharpoons 2\text{CO}_{(g)} + \text{O}_{2(g)}$		
I	0.250	0	0
C	-2x	+2x	+x
E	0.250-2x	2x	x

0.247 and 0.250 are very close

The difference is 1.2%. As long as the difference is less than 5%, you can use the assumption.

You can also divide the initial concentration by k_{eq} . If the answer is AT LEAST 100, you can use the assumption.

$0.250/6.40 \times 10^{-7} = 3.91 \times 10^5$, which is **MUCH** bigger than 100

ASSUMPTIONS WITH EQUILIBRIUM

Simplification of equilibrium equations may usually be used if K_{eq} values are:

1.greater than 10^3

2.less than 10^{-3}

Questions involving a lot of polynomial expansion is a good indication that an assumption should be used.

ASSUMPTIONS WITH EQUILIBRIUM

Example #2


What is molar solubility of $\text{PbCl}_{2(s)}$ in a 0.2 mol / L $\text{NaCl}_{(aq)}$ solution? $K_{sp} = 1.7 \times 10^{-5}$

K_{sp} = solubility product constant


ASSUMPTIONS WITH EQUILIBRIUM

Example #2

What is molar solubility of $\text{PbCl}_{2(s)}$ in a 0.2 mol / L $\text{NaCl}_{(aq)}$ solution? $K_{sp} = 1.7 \times 10^{-5}$



The Cl^{-} ions *ALREADY* in solution will limit the amount of PbCl_2 that can be dissolved



ASSUMPTIONS WITH EQUILIBRIUM

Example #2

What is molar solubility of $\text{PbCl}_{2(s)}$ in a $0.2 \text{ mol / L NaCl}_{(aq)}$ solution? $K_{sp} = 1.7 \times 10^{-5}$

	$\text{PbCl}_{2(s)}$	\rightleftharpoons	$\text{Pb}^{2+}_{(aq)}$	$+$	$2\text{Cl}^{-}_{(aq)}$
I	-		0		0.2
C	-		+x		+2x
E	-		x		0.2+2x

$$K_{sp} = [\text{Pb}^{2+}][\text{Cl}^{-}]^2$$

$$1.7 \times 10^{-5} = [x][0.2+2x]^2$$

Because K_{sp} is SO small, very little PbCl_2 actually dissolves

Assumption: $0.2+2x = 0.2$

$$1.7 \times 10^{-5} = [x][\mathbf{0.2}]^2$$

$$\frac{1.7 \times 10^{-5}}{0.04} = x$$

$$\frac{0.04}{4.2 \times 10^{-4}} = x$$

\therefore the molar solubility of PbCl_2 in $0.2\text{M NaCl}_{(aq)}$ is $4.2 \times 10^{-4} \text{ mol/L}$


ASSUMPTIONS WITH EQUILIBRIUM

Net Ionic Equations

Do you remember what this is?


How does it affect equilibrium?

Chemical equation:




Ionic equation:

→ Anything that is aqueous must be separated into its component ions



Net Ionic equation:

→ **Spectator ions** are eliminated



ASSUMPTIONS WITH EQUILIBRIUM

Common Ion Effect

For aqueous solutions, the addition of ions may or may not affect equilibrium.

The added ion will only affect equilibrium when it is one that is **common** with those involved in the net ionic equation.

Spectator ions will not affect equilibrium.