

Name – Vedanti Rahul Bhanuse

Batch – DSC 46 July 2022

## Assignment-base Subjective Questions

1. From your analysis of the categorical variables from the dataset, what could you infer about their effect on the dependent variable?

Answer:

- Here are some of the inferences I made from my analysis of categorical variables from the dataset on the dependent variable (Count)
- Fall has the highest median, which is expected as weather conditions are most optimal to ride bike followed by summer.
- Median bike rents are increasing year on as year 2019 has a higher median than 2018, it might be due the fact that bike rentals are getting popular and people are becoming more aware about environment.
- Overall spread in the month plot is reflection of season plot as fall months have higher median.
- People rent more on non - holidays compared to holidays, so reason might be they prefer to spend time with family and use personal vehicle instead of bike rentals.  
Overall median across all days is same but spread for Saturday and Wednesday is bigger may be evident that those who have plans for Saturday might not rent bikes as it a non-working day.
- Working and non-working days have almost the same median although spread is bigger for non-working days as people might have plans and do not want to rent bikes because of that.
- Clear weather is most optimal for bike renting, as temperate is optimal, humidity is less, and temperature is less.

2. Why is it important to use drop\_first=True during dummy variable creation?

Answer: A variable with n levels can be represented by n-1 dummy variables. So, if we remove the first column then also, we can represent the data. If the value of variable from 2 to n is 0, it means that the value of 1st variable is 1.

3. Looking at the pair-plot among the numerical variables, which one has the highest correlation with the target variable?

Answer: 'temp' had the highest correlation coefficient 0.48.

4. How did you validate the assumptions of Linear Regression after building the model on the training set?

Answer: By plotting the residuals distribution. It came out to be a normal distribution with a mean value of 0.

5. Based on the final model, which are the top 3 features contributing significantly towards explaining the demand of the shared bikes?

Answer: The Following are the top 3 features contributing significantly the demands of the shared bikes:

- temp (0.45)
- yr\_2019 (0.23)
- mnth\_Sep (0.05)

## General Subjective Questions

1. Explain the linear regression algorithm in detail.

Answer: A linear regression algorithm tries to explain the relationship between independent and dependent variable using a straight line. It is applicable to numerical variables only.

Following steps are performed while doing linear regression:

- The dataset is divided into test and training data
- Train data is divided into features(independent) and target (dependent) datasets
- A linear model is fitted using the training dataset. Internally the api's from python uses gradient descent algorithm to find the coefficients of the best fit line. The gradient descent algorithm works by minimising the cost function. A typical example

of cost function is residual sum of squares.

- In case of multiple features, the predicted variable is a hyperplane instead of line.

The predicted variable takes the following form:

$$Y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3 + \dots + \beta_n x_n$$

- The predicted variable is then compared with test data and assumptions are checked.

## 2. Explain the Anscombe's quartet in detail.

Answer: Anscombe's quartet comprises of four data sets that have nearly identical simple descriptive statistics but have quite different distribution when visualized graphically. The simple statistics consist of mean, sample variance of  $x$  and  $y$ , correlation coefficient, linear regression line and R-Square value. Anscombe's Quartet shows that multiple data sets with many similar statistical properties can still be vastly different from one another when graphed.

The graphs are shown below:




Image source - [https://en.wikipedia.org/wiki/Anscombe%27s\\_quartet](https://en.wikipedia.org/wiki/Anscombe%27s_quartet)

3. First plot (top left) appears to be simple linear relationship

4. The second plot (top right) is not distributed normally and correlation coefficient is

irrelevant as it shows a nonlinear relationship

5. The third plot (bottom left) is linear but has different regression line. This is happening because of the outliers present in the data

6. The fourth plot (bottom right) does not show linear relationship however due to

outliers the statistics got adjusted.

In a nutshell, it is a better practice to visualize data and remove outliers before analysing it.

3. What is Pearson's R?

Answer: Pearson's R measures the strength of association of two variables. It is the covariance of two variables divided by the product of their standard deviation. It has a value from +1 to -1.

- A value of 1 means a total positive linear correlation. It means that if one variable increase then other will also increase
- A value of 0 means no correlation
- A value of -1 means a total negative correlation. It means that if one variable increase then other will decrease.

4. What is scaling? Why is scaling performed? What is the difference between normalized scaling and standardized scaling?

Answer: Scaling of a variable is performed to keep a variable in certain range. Scaling is a pre-processing step in linear regression analysis. The reason we scale variable is to make the computation of gradient descent faster. The step size of gradient descent is generally low for accuracy, if the data has some small variables (values in the range of 0-1) and some big variables (values in the range of 0 -1000) then the time taken by the gradient descent algorithm will be huge.

Normalised Scaling	Standardized scaling
Called min max scaling, scales the variable such that the range is 0-1	Values are centred around mean with a unit standard deviation
Good for non- gaussian distribution	Good for gaussian distribution
Value is bounded between 0 and 1	Value is not bounded
Outliers are also scaled	Does not affect outliers

5. You might have observed that sometimes the value of VIF is infinite. Why does this happen?

Answer: If there is perfect correlation, then  $VIF = \infty$ . This shows a perfect correlation between two independent variables. In the case of perfect correlation, we get  $R^2 = 1$ , which leads to  $1/(1-R^2)$  infinity. To solve this problem, we need to drop one of the variables from the dataset which is causing this perfect multicollinearity. An infinite VIF value indicates that the corresponding variable may be expressed exactly by a linear combination of other variables (which show an infinite VIF as well).

6. What is a Q-Q plot? Explain the use and importance of a Q-Q plot in linear regression.

Answer: Q-Q plots are also known as Quantile-Quantile plots. As the name suggests, they plot the quantiles of a sample distribution against quantiles of a theoretical distribution. Doing this helps us determine if a dataset follows any particular type of probability distribution like normal, uniform, exponential. Before we dive into the Q-Q plot, let's discuss some of the probability distributions.