

FOURIER SERIES

Find the Fourier series for the following functions.

FOURIER EXPANSION OF $f(x)$ IN THE INTERVAL $(0, 2\pi)$

1. $f(x) = x^2$ in $(0, 2\pi)$ Hence deduce that $\frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$

$$[\text{Ans: } f(x) = \frac{4\pi^2}{3} + 4 \sum_{n=1}^{\infty} \frac{1}{n^2} \cos nx - 4\pi \sum_{n=1}^{\infty} \frac{1}{n} \sin nx]$$

2. $f(x) = e^{-x}$, $0 < x < 2\pi$ & $f(x+2\pi) = f(x)$ Hence deduce the value of $\sum_{n=2}^{\infty} \frac{(-1)^n}{n^2 + 1}$

$$[\text{Ans: } f(x) = \frac{1 - e^{-2\pi}}{\pi} \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2}]$$

3. $f(x) = x \sin x$ in the interval $0 \leq x \leq 2\pi$. Hence deduce that $\sum_{n=1}^{\infty} \frac{1}{n^2 - 1} = \frac{3}{4}$

$$[\text{Ans: } f(x) = -1 - \frac{1}{2} \cos x + \sum_{n=2}^{\infty} \frac{2}{n^2 - 1} \cos nx + \pi \sin x]$$

4. $f(x) = \sqrt{1 - \cos x}$ in $(0, 2\pi)$ Hence deduce that $\sum_{n=1}^{\infty} \frac{1}{4n^2 - 1} = \frac{1}{2}$

5. $f(x) = x$, $0 < x \leq \pi$
 $= 2\pi - x$, $\pi \leq x < 2\pi$ Hence deduce that $\frac{\pi^2}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$

$$[\text{Ans: } f(x) = \frac{\pi}{2} - \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^2} \cos nx]$$

6. $f(x) = x$ in $(0, 2\pi)$

$$[\text{Ans: } f(x) = \pi - 2 \sum_{n=1}^{\infty} \frac{\sin nx}{n}]$$

7. $f(x) = \frac{3x^2 - 6x\pi + 2\pi^2}{12}$ in $(0, 2\pi)$ Hence deduce that $\frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$

$$[\text{Ans: } f(x) = \sum_{n=1}^{\infty} \frac{\cos nx}{n^2}]$$

8. $f(x) = \left(\frac{\pi - x}{2}\right)$ in the interval $0 \leq x \leq 2\pi$ Also deduce that $\frac{\pi}{4} = \frac{1}{1} - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots$

9. $f(x) = 1$, $0 < x \leq \pi$
 $= 2 - \frac{x}{\pi}$, $\pi \leq x < 2\pi$

$$[\text{Ans: } f(x) = \frac{3}{4} - \frac{2}{\pi^2} \left[\frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \dots \right] + \frac{1}{\pi} \left[\frac{\sin x}{1} + \frac{\sin 2x}{2} + \dots \right]]$$

10. $f(x) = 2x$ in $(0, 2\pi)$ Also find a_4 & b_{10} .

[Ans: $f(x) = 2\pi - 4 \sum_{n=1}^{\infty} \frac{\sin nx}{n}$, $0, -0.4$]

11. $f(x) = \cos px$, in $(0, 2\pi)$ where p is not an integer.

12. $f(x) = kx$, $0 \leq x \leq 2\pi$.Also find a_4 & b_{10} .

13. $f(x) = e^{2x}$ in $(0, 2\pi)$

14. $f(x) = e^{-2x}$ in $(0, 2\pi)$

FOURIER EXPANSION OF $f(x)$ IN THE INTERVAL $(-\pi, \pi)$

15. state the value of $f(x)$ at $x = 0$ if $f(x) = \begin{cases} -\pi, & -\pi < x < 0 \\ x, & 0 < x < \pi \end{cases}$ and hence deduce that $\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}$

[Ans: $f(x) = -\frac{\pi}{4} + \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{[(-1)^n - 1]}{n^2} \cos nx + \sum_{n=1}^{\infty} \frac{[1 - 2(-1)^n]}{n} \sin nx$]

16. $f(x) = 1/2$, $-\pi < x < 0$
 $= x/\pi$, $0 < x < \pi$ Hence deduce that $\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}$

[Ans: $f(x) = \frac{1}{2} - \frac{2}{\pi^2} \sum_{n=1}^{\infty} \frac{\cos(2n-1)x}{(2n-1)^2} - \frac{1}{2} \sum_{n=1}^{\infty} \frac{\sin 2nx}{n}$]

17. $f(x) = -x - \pi$, $-\pi \leq x \leq 0$
 $= x + \pi$, $0 \leq x \leq \pi$

[Ans: $f(x) = \frac{\pi}{2} - \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{\cos(2n-1)x}{(2n-1)^2} + 4 \sum_{n=1}^{\infty} \frac{\sin(2n-1)x}{(2n-1)}$]

18. $f(x) = 0$, $-\pi \leq x \leq 0$
 $= x$, $0 \leq x \leq \pi$

Hence, deduce that i) $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots$ ii) $\frac{\pi}{4} = \frac{1}{1} - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots$

19. Obtain Fourier Series for $f(x) = e^{-|x|}$, $-\pi \leq x \leq \pi$

20. $f(x) = 0$, $-\pi \leq x \leq 0$
 $= \sin x$, $0 \leq x \leq \pi$, Hence, deduce that i) $\frac{1}{2} = \frac{1}{1 \cdot 3} + \frac{1}{3 \cdot 5} + \frac{1}{5 \cdot 7} + \dots$

ii) $\frac{1}{4}(\pi - 2) = \frac{1}{1 \cdot 3} - \frac{1}{3 \cdot 5} + \frac{1}{5 \cdot 7} - \dots$ [Ans: $f(x) = \frac{1}{\pi} + \frac{\sin x}{2} - \frac{2}{\pi} \left[\frac{\cos 2x}{4 \cdot 1^2 - 1} + \frac{\cos 4x}{4 \cdot 2^2 - 1} + \dots \right]$]

21. It is given that for $-\pi < x < \pi$, $x^2 = \frac{\pi^2}{3} + 4 \sum_{n=1}^{\infty} (-1)^n \frac{\cos nx}{n^2}$

Using Parseval's identity prove that $\sum_{n=1}^{\infty} \frac{1}{n^4} = \frac{\pi^4}{90}$.

$$f(x) = 1 + \frac{2x}{\pi}, \quad -\pi \leq x \leq 0$$

$$= 1 - \frac{2x}{\pi}, \quad 0 \leq x \leq \pi$$

Deduce that $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} \dots$

22. [Ans: $f(x) = \sum_{n=1}^{\infty} \frac{4}{\pi^2 n^2} [1 - (-1)^n] \cos nx$]

$$f(x) = x + \frac{\pi}{2}, \quad -\pi < x < 0$$

$$= \frac{\pi}{2} - x, \quad 0 < x < \pi$$

Hence, Deduce that

i) $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} \dots$

ii) $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} \dots$

23. [Ans: $f(x) = \sum_{n=1}^{\infty} \frac{2}{\pi n^2} [1 - (-1)^n] \cos nx$]

24. Prove that $\sinh ax = \frac{2}{\pi} \sinh a\pi \left[\sum_{n=1}^{\infty} \frac{(-1)^{n+1} \cdot n}{n^2 + a^2} \sin nx \right]$

25. $f(x) = x \cos x$, in $(-\pi, \pi)$ [Ans: $f(x) = -\frac{1}{2} \sin x + 2 \sum_{n=2}^{\infty} (-1)^n \cdot \frac{n}{n^2 - 1} \sin nx$]

26. $f(x) = x + x^2$, in $(-\pi, \pi)$. Hence deduce that i) $\frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} \dots$ ii) $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} \dots$

27. $f(x) = \cos px$, in $(-\pi, \pi)$. Where p is not an integer. Hence, prove that

$$\cot p\pi = \frac{2p}{\pi} \left[\frac{1}{2p^2} - \frac{1}{p^2 - 1^2} + \frac{1}{p^2 - 2^2} - \frac{1}{p^2 - 3^2} + \dots \right]$$

And deduce that $\cos \theta = \frac{1}{\theta} - \sum_{n=1}^{\infty} \frac{2\theta}{n^2 \pi^2 - \theta^2}$

Also deduce that $\frac{1}{2} - \frac{\pi\sqrt{3}}{18} = \frac{1}{9 \cdot 1^2 - 1} + \frac{1}{9 \cdot 2^2 - 1} + \frac{1}{9 \cdot 3^2 - 1} + \dots$

28. $f(x) = |\sin x|$, in $(-\pi, \pi)$ [Ans: $f(x) = \frac{2}{\pi} - \frac{4}{\pi} \left[\frac{\cos 2x}{3} + \frac{\cos 4x}{15} + \frac{\cos 6x}{35} + \dots \right]$]

29. $f(x) = |x|$, in $(-\pi, \pi)$ Hence deduce that $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} \dots$

[Ans: $f(x) = \frac{\pi}{2} - \frac{\pi}{2} \left[\frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} + \dots \right]$]

30. $f(x) = x \sin x$, in $(-\pi, \pi)$. Hence deduce that $\frac{1}{4}(\pi - 2) = \frac{1}{1 \cdot 3} - \frac{1}{3 \cdot 5} + \frac{1}{5 \cdot 7} - \dots$

[Ans: $f(x) = 1 - \frac{1}{2} \cos x - 2 \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2 - 1} \cos nx$]

31. $f(x) = \frac{e^{ax} - e^{-ax}}{e^{a\pi} - e^{-a\pi}}$, in $(-\pi, \pi)$

[Ans: $f(x) = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^{n+1} \cdot n}{n^2 + a^2} \sin nx$]

32. $f(x) = \frac{x(\pi^2 - x^2)}{12}$, in $(-\pi, \pi)$

[Ans: $f(x) = \frac{\sin x}{1^3} - \frac{\sin 2x}{2^3} + \frac{\sin 3x}{3^3} - \dots$]

$$f(x) = 0, \quad -\pi \leq x \leq 0$$

$$= x^2, \quad 0 \leq x \leq \pi$$

33. $x^2 = \frac{\pi^2}{3} + 4 \sum (-1)^n \cdot \frac{\cos nx}{n^2}$ for $-\pi < x < \pi$, prove that $\sum \frac{1}{n^4} = \frac{\pi^4}{90}$

34. $f(x) = \sin x$, in $(-\pi, \pi)$

35. $f(x) = \frac{\pi^2}{12} - \frac{x^2}{4}$, in $(-\pi, \pi)$ [Ans: $f(x) = \frac{\cos x}{1^2} - \frac{\cos 2x}{2^2} + \frac{\cos 3x}{3^2} - \dots$]

36. $f(x) = x$, $-\pi < x < 0$
 $= 0$, $0 < x < \pi/2$
 $= x - \pi/2$, $\pi/2 < x < \pi$

37. $f(x) = x^2$, in $(-\pi, \pi)$

38. $f(x) = \begin{cases} 0, & -\pi < x < 0 \\ x^2, & 0 < x < \pi \end{cases}$

39. $f(x) = x \cos x$ in $(-\pi, \pi)$

40. $f(x) = \cosh px$ in $(-\pi, \pi)$, p is not an integer

41. $f(x) = \frac{x(\pi-x)(\pi+x)}{12}$ in $(-\pi, \pi)$

42. $f(x) = x|x|$, $-\pi \leq x \leq \pi$

43. $f(x) = e^{-|x|}$, $-\pi \leq x \leq \pi$

FOURIER EXPANSION OF $f(x)$ IN THE INTERVAL $(0, 2l)$

44. $f(x) = x^2$, in $(0, a)$ Hence deduce that $\frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots$

[Ans: $f(x) = \frac{a^2}{3} + \sum_{n=1}^{\infty} \frac{a^2}{n^2 \pi^2} \cos\left(\frac{n\pi x}{a}\right) - \sum_{n=1}^{\infty} \frac{a^2}{n \pi} \sin\left(\frac{n\pi x}{a}\right)$]

45. $f(x) = 2x - x^2$, $0 \leq x \leq 3$

[Ans: $f(x) = \sum_{n=1}^{\infty} \frac{-9}{n^2 \pi^2} \cos\left(\frac{2n\pi x}{3}\right) + \sum_{n=1}^{\infty} \frac{3}{n \pi} \sin\left(\frac{2n\pi x}{3}\right)$]

46. $f(x) = \pi x$, $0 < x < 1$
 $= 0$, $1 < x < 2$

[Ans: $f(x) = \frac{\pi}{4} - \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^2} \cos n\pi x - \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \sin n\pi x$]

48. $f(x) = \pi x, \quad 0 \leq x \leq 1$
 $= \pi(2-x), \quad 1 \leq x \leq 2$, with period 2, show that $f(x) = \frac{\pi}{2} - \frac{4}{\pi} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} \cos(2n+1)\pi x$

49. $f(x) = \pi x, \quad 0 < x < 1$
 $= 0, \quad x \equiv 1$, Hence show that $\frac{\pi}{4} = \frac{1}{1} - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots$ [Ans: $f(x) = \frac{\pi}{4} + 2 \sum_{n=1}^{\infty} (-1)^{n+1} \sin n\pi x$]
 $= \pi(2-x), \quad 1 < x < 2$

50. $f(x) = 3kx/l, \quad 0 < x < (l/3)$
 $= 3k(l-2x)/l, \quad (l/3) < x < (2l/3),$
 $= \pi(2-x), \quad (2l/3) < x < l$ [Ans: $\frac{9k}{\pi^2} \sum \frac{1}{n^2} \sin \frac{2n\pi}{3} \cdot \sin \frac{2n\pi x}{l}$]

51. If $x^2 = \frac{4l^2}{3} + \frac{4l^2}{\pi^2} \sum \frac{1}{n^2} \cos\left(\frac{n\pi x}{l}\right) - \frac{4l^2}{\pi} \sum \frac{1}{n} \sin\left(\frac{n\pi x}{l}\right)$ in $0 < x < 2l$, find the sum of the series

$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots$ [Ans: $\frac{\pi^2}{6}$]

52. $f(x) = kx$ in the interval $0 \leq x \leq 2$. Hence deduce that $\sum_{n=1}^{\infty} \frac{1}{n^2} \equiv \frac{\pi^2}{6}$

53. Find Fourier series to represent $f(x) = 2x - x^2$ in $(0,3)$ and prove that $\frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$

54. $f(x) = 2 - \frac{x^2}{2}$ in $0 \leq x \leq 2$

FOURIER EXPANSION OF $f(x)$ IN THE INTERVAL $(-l, l)$

55. $f(x) = 0, \quad -c < x < 0$
 $= a, \quad 0 < x < c$

[Ans: $f(x) = \frac{a}{2} + \frac{2a}{\pi} \left[\frac{1}{1} \sin \frac{\pi x}{c} + \frac{1}{3} \sin \frac{3\pi x}{c} + \dots \right]$

56. $f(x) = -x, \quad -1 < x < 0$
 $= x, \quad 0 < x < 1$

[Ans: $f(x) = 1 - \frac{4}{\pi^2} \sum \frac{1}{(2n-1)^2} \cos n\pi x$]

57. $f(x) = x, \quad -1 < x < 0$
 $= x+2, \quad 0 < x < 1$

[Ans: $f(x) = 1 + \frac{2}{\pi} \sum_{n=1}^{\infty} [1 - 2(-1)^n] \sin n\pi x$]

58. $f(x) = |x|, \quad -2 < x < 2$, Hence deduce that $\sum_{n=1}^{\infty} \frac{1}{(2n-1)^4} = \frac{\pi^4}{96}$

[Ans: $f(x) = 1 - \frac{8}{\pi^2} \sum \frac{1}{(2n-1)^2} \cos \left[\frac{(2n-1)\pi x}{2} \right]$]

59. $f(x) = 1 - x^2, \quad -1 < x < 1,$

[Ans: $f(x) = \frac{2}{3} - \frac{4}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \cos n\pi x$]

60. $f(x) = \sin ax, \quad -l < x < l,$

[Ans: $f(x) = 2\pi \sin al \sum \frac{(-n)(-1)^n}{n^2 \pi^2 - a^2 l^2} \sin \frac{n\pi x}{l}$]

$$61. f(x) = x - x^2, \quad -1 < x < 1,$$

$$[\text{Ans: } f(x) = -\frac{1}{3} - \frac{4}{\pi^2} \sum \frac{(-1)^n}{n^2} \cos n\pi x - \frac{2}{\pi} \sum \frac{(-1)^n}{n} \sin n\pi x]$$

$$62. f(x) = a^2 - x^2, \quad -a < x < a,$$

$$[\text{Ans: } f(x) = \frac{2a^2}{3} + \frac{4a^2}{\pi^2} \left[\frac{1}{1^2} \cos \frac{\pi x}{a} - \frac{1}{2^2} \cos \frac{2\pi x}{a} + \frac{1}{3^2} \cos \frac{3\pi x}{a} - \dots \right]]$$

$$63. f(x) = x^2, \quad -1 < x < 1,$$

$$[\text{Ans: } f(x) = \frac{1}{3} - \frac{4}{\pi^2} \left[\frac{1}{1^2} \cos \pi x - \frac{1}{2^2} \cos 2\pi x + \frac{1}{3^2} \cos 3\pi x - \dots \right]]$$

$$64. f(x) = 9 - x^2, \quad -3 < x < 3,$$

$$[\text{Ans: } f(x) = 6 + \frac{36}{\pi^2} \left[\frac{1}{1^2} \cos \frac{\pi x}{3} - \frac{1}{2^2} \cos \frac{2\pi x}{3} + \frac{1}{3^2} \cos \frac{3\pi x}{3} - \dots \right]]$$

$$65. f(x) = x - x^3 \text{ in } (-1, 1)$$

$$[\text{Ans: } f(x) = -\frac{12}{\pi^3} \sum_{n=1}^{\infty} \frac{(-1)^n \sin n\pi x}{n^3}]$$

$$66. f(x) = x - x^2 \text{ in } (-1, 1)$$

$$[\text{Ans: } f(x) = -\frac{1}{3} - \frac{4}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^n \cos n\pi x}{n^2} - \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n \sin n\pi x}{n}]$$

$$67. f(x) = x^2 - 2, \quad -2 \leq x \leq 2$$

$$[\text{Ans: } f(x) = -\frac{2}{3} - \frac{16}{\pi^2} \left[\cos \frac{\pi x}{2} - \frac{1}{4} \cos \pi x + \frac{1}{9} \cos \frac{3\pi x}{2} - \dots \right]]$$

$$68. f(x) = \begin{cases} 0, & -2 < x < -1 \\ 1+x, & -1 < x < 0 \\ 1-x, & 0 < x < 1 \\ 0, & 1 < x < 2 \end{cases}$$

$$[\text{Ans: } f(x) = \frac{1}{4} + \sum \frac{4}{n^2 \pi^2} \left(1 - \cos \frac{n\pi}{2} \right) \cdot \cos \left(\frac{n\pi x}{2} \right)]$$

$$69. f(x) = e^{-x}, \quad (-a, a)$$

$$[\text{Ans: } f(x) = \frac{\sinh a}{a} + 2a \sinh a \sum \frac{(-1)^n}{a^2 + n^2 \pi^2} \cos \frac{n\pi x}{a} + 2\pi \sinh a \sum \frac{(-1)^{n+1} \cdot n}{a^2 + n^2 \pi^2} \sin \frac{n\pi x}{a}]$$

$$70. f(x) = |x|, \quad -1 < x < 1$$

$$71. f(x) = \begin{cases} 0, & -2 < x < -1 \\ k, & -1 < x < 1 \\ 0, & 1 < x < 2 \end{cases}$$

$$72. f(x) = \begin{cases} 0, & -5 < x < 0 \\ 7, & 0 < x < 5 \end{cases} \quad \text{period of the function is 10.}$$

$$73. f(x) = \begin{cases} 0, & -2 < x < 0 \\ x+5, & 0 < x < 2 \end{cases}$$

$$74. f(x) = 1 - x^2 \text{ in } (-1, 1) \text{ hence find } \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$$

$$75. f(x) = \begin{cases} -\sin \frac{\pi x}{k}, & -k < x < 0 \\ \sin \frac{\pi x}{k}, & 0 < x < k \end{cases}$$

$$76. f(x) = \begin{cases} 2(x-4), & -4 < x < 0 \\ 2(x+4), & 0 < x < 4 \end{cases}$$

77. $f(x) = x^2 - 2$ on $(-2, 2)$

HALF RANGE SERIES

78. Obtain half range sine series for $f(x) = x, \quad 0 < x < \pi/2$
 $= \pi - x, \quad \pi/2 < x < \pi$ Hence find $\sum_{n=1}^{\infty} \frac{1}{(2n-1)^4}$.

[Ans: $f(x) = \sum \frac{4}{\pi} \frac{\sin(n\pi/2)}{n^2} \cdot \sin nx$]

79. Find half range cosine series for $f(x) = x, \quad (0, 2)$. Using Parseval's identity, deduce that

i) $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$ ii) $\sum \frac{1}{n^4} = \frac{\pi^4}{90}$.

80. Obtain the expression of $f(x) = x(\pi - x), \quad 0 < x < \pi$ as a half-range cosine series. Hence, show that i) $\frac{\pi^2}{6} = \sum_{n=1}^{\infty} \frac{1}{n^2}$

ii) $\frac{\pi^2}{12} = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^2}$ iii) $\sum \frac{1}{n^4} = \frac{\pi^4}{90}$. [Ans: $f(x) = \frac{\pi^2}{6} - \left[\frac{1}{1^2} \cos 2x + \frac{1}{2^2} \cos 4x + \frac{1}{3^2} \cos 6x + \dots \right]$

81. Show that if $0 < x < \pi$, $\cos x = \frac{8}{\pi} \sum_{m=1}^{\infty} \frac{m}{4m^2 - 1} \sin 2mx$

82. Expand $f(x) = lx - x^2, \quad 0 < x < l$ in a half range i) cosine series, ii) sine series.

Hence from sine series deduce that $\frac{\pi^3}{32} = 1 - \frac{1}{3^3} + \frac{1}{5^3} - \frac{1}{7^3} + \dots$

[Ans: i) $f(x) = \frac{l^2}{6} - \frac{4l^2}{\pi^2} \left[\frac{1}{2^2} \cos \frac{2\pi x}{l} + \frac{1}{4^2} \cos \frac{4\pi x}{l} + \frac{1}{6^2} \cos \frac{6\pi x}{l} + \dots \right]$
 ii) $f(x) = \frac{8l^2}{\pi^3} \left[\frac{1}{1^3} \sin \frac{\pi x}{l} + \frac{1}{3^3} \sin \frac{3\pi x}{l} + \frac{1}{5^3} \sin \frac{5\pi x}{l} + \dots \right]$

83. Find half range cosine series for $f(x) = \begin{cases} x, & 0 < x < (\pi/2) \\ \pi - x, & (\pi/2) < x < \pi \end{cases}$

[Ans: $f(x) = \frac{\pi}{4} - \frac{8}{\pi} \left[\frac{1}{2^2} \cos 2x + \frac{1}{6^2} \cos 6x + \frac{1}{10^2} \cos 10x + \dots \right]$

84. Prove that in the interval $0 < x < \pi$, $\frac{e^{ax} - e^{-ax}}{e^{a\pi} - e^{-a\pi}} = \frac{2}{\pi} \left[\frac{\sin x}{a^2 + 1} - \frac{2 \sin 2x}{a^2 + 4} + \frac{3 \sin 3x}{a^2 + 9} - \dots \right]$

85. Obtain half-range sine series for $f(x) = x(2 - x)$ in $0 < x < 2$ and hence find $\sum \frac{1}{n^6} = \frac{\pi^6}{945}$

86. Obtain half range sine series for $f(x) = \begin{cases} (1/4) - x, & 0 < x < (1/2) \\ x - (3/4), & (1/2) < x < 1 \end{cases}$

$$[\text{Ans: } f(x) = \left(\frac{1}{\pi} - \frac{4}{\pi^2}\right) \sin \pi x + \left(\frac{1}{3\pi} - \frac{4}{3^2 \pi^2}\right) \sin 3\pi x + \left(\frac{1}{5\pi} - \frac{4}{5^2 \pi^2}\right) \sin 5\pi x + \dots]$$

87. Obtain half-range cosine series for $f(x) = x$ in $0 < x < l$. Hence deduce that $\frac{1}{2^4} + \frac{1}{4^4} + \frac{1}{6^4} + \dots = \frac{\pi^4}{1440}$

88. Obtain half-range cosine series for $f(x) = \begin{cases} kx, & 0 < x < (l/2) \\ l-x, & (l/2) < x < l \end{cases}$

Hence, deduce that i) $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$ ii) $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$

$$[\text{Ans: } f(x) = \frac{kl}{4} - \frac{8kl}{\pi^2} \left[\frac{1}{2^2} \cos \frac{2\pi x}{l} + \frac{1}{6^2} \cos \frac{6\pi x}{l} + \frac{1}{10^2} \cos \frac{10\pi x}{l} + \dots \right]]$$

89. Find half range sine series of period $2l$ for $f(x) = \begin{cases} \frac{2x}{l}, & 0 < x < (l/2) \\ \frac{2}{l}(l-x), & (l/2) < x < l \end{cases}$

$$[\text{Ans: } f(x) = \frac{8}{\pi^2} \sum \frac{1}{n^2} \sin \frac{n\pi}{2} \cdot \sin \frac{n\pi x}{l}]$$

90. Obtain sine series for $f(x) = \begin{cases} mx, & 0 < x \leq (\pi/2) \\ m(\pi-x), & (\pi/2) \leq x < \pi \end{cases}$ [Ans: $f(x) = \frac{4m}{\pi} \left[\frac{\sin x}{1^2} - \frac{\sin 3x}{3^2} + \frac{\sin 5x}{5^2} - \dots \right]$]

91. Obtain half range cosine series for $f(x) = \sin\left(\frac{\pi x}{l}\right)$ in $0 < x < l$.

$$[\text{Ans: } f(x) = \frac{2}{\pi} - \frac{4}{\pi} \left[\frac{1}{1 \cdot 3} \cos \frac{2\pi x}{l} + \frac{1}{3 \cdot 5} \cos \frac{4\pi x}{l} + \dots \right]]$$

92. Obtain half-range cosine series for $f(x) = (x-1)^2$ in $0 < x < 1$. Hence, find $\sum_{n=1}^{\infty} \frac{1}{n^2}$ & $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^2}$

$$[\text{Ans: } f(x) = \frac{1}{3} + \frac{4}{\pi^2} \sum_{n=1}^{\infty} \frac{\cos n\pi x}{n^2}]$$

93. Find HRSS for $f(x) = \begin{cases} \frac{2x}{3}, & 0 \leq x \leq \frac{\pi}{3} \\ \frac{\pi-x}{3}, & \frac{\pi}{3} \leq x \leq \pi \end{cases}$

$$[\text{Ans: } f(x) = \frac{\sqrt{3}}{\pi} \left[\frac{1}{1^2} \sin x + \frac{1}{2^2} \sin 2x - \frac{1}{4^2} \sin 4x - \frac{1}{5^2} \sin 5x + \dots \right]]$$

94. Obtain the half range sine series for $f(x) = x(\pi-x)$, $0 < x < \pi$ Hence, find $\sum_{n=1}^{\infty} \frac{(-1)^3}{(2n-1)^3}$

95. Show that in the interval $0 < x < \pi$, $\sin x = \frac{2}{\pi} - \frac{4}{\pi} \left[\frac{\cos 2x}{2^2-1} + \frac{\cos 4x}{4^2-1} + \dots \right]$

96. Obtain half-range sine series for $f(x) = x^2$ in $0 < x < 3$.

97. Obtain HRCS for $f(x) = x(2-x)$ in $0 < x < 2$ [Ans: $f(x) = \frac{2}{3} - \frac{8}{\pi^2} \sum_{n=1}^{\infty} \frac{[1+(-1)^n]}{n^2} \cos\left(\frac{n\pi x}{2}\right)$]

98. Find half range cosine series for $f(x) = \begin{cases} kx & 0 \leq x \leq l/2 \\ 0 & l/2 < x \leq l \end{cases}$ Hence deduce that $\frac{\pi^2}{90} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2} + \dots$

99. Find half range cosine series for $f(x) = x$ on $(0, 2)$ hence deduce that $\frac{\pi^4}{90} = \frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \dots$

COMPLEX FORM OF FOURIER SERIES

Obtain complex form of Fourier series for the following functions:

100. $f(x) = e^{ax}$ in $(-\pi, \pi)$ where a is not an integer

[Ans: $f(x) = \sum_{n=-\infty}^{\infty} \frac{(-1)^n \sinh a\pi \cdot (a + in)}{\pi(a^2 + n^2)} e^{inx}$]

101. $f(x) = e^{ax}$ in $(-l, l)$, where a is not an integer.

[Ans: $f(x) = \sum_{n=-\infty}^{\infty} \frac{(-1)^n \sinh al \cdot (al + in\pi)}{(a^2 l^2 + n^2 \pi^2)} e^{in\pi x/l}$]

102. $f(x) = \cosh ax + \sinh ax$ in $(-l, l)$

[Ans: $f(x) = \sum_{n=-\infty}^{\infty} \frac{(-1)^n \sinh al \cdot (al + in\pi)}{(a^2 l^2 + n^2 \pi^2)} e^{in\pi x/l}$]

103. $f(x) = \cosh ax$ in $(-l, l)$

[Ans: $f(x) = \sum_{n=-\infty}^{\infty} \frac{(-1)^n al \sinh al}{(a^2 l^2 + n^2 \pi^2)} e^{in\pi x/l}$]

104. $f(x) = \sin ax$ in $(-\pi, \pi)$, where a is not an integer.

[Ans: $f(x) = \frac{\sin a\pi}{\pi} \sum_{n=-\infty}^{\infty} (-1)^n \cdot \frac{n}{(a^2 - n^2)} \cdot e^{inx}$]

105. $f(x) = e^{ax}$ in $(0, a)$

[Ans: $f(x) = (e^{a^2} - 1) \sum_{n=-\infty}^{\infty} \frac{e^{2in\pi/a}}{(a^2 - 2in\pi)}$]

106. $f(x) = e^{ax}$ in $(-1, 1)$

[Ans: $f(x) = \sum_{n=-\infty}^{\infty} \frac{(-1)^n \sinh a \cdot (a + in\pi)}{(a^2 + n^2 \pi^2)} e^{in\pi x}$]

107. $f(x) = \cosh 3x + \sinh 3x$ in $(-\pi, \pi)$

[Ans: $f(x) = \sum_{n=-\infty}^{\infty} \frac{(-1)^n \sinh 3\pi \cdot (3 + in)}{(9 + n^2)\pi} e^{inx}$]

108. $f(x) = 0, \quad 0 < x < l$
 $= a, \quad l < x < 2l$

[Ans: $f(x) = \frac{a}{2} + \frac{ai}{\pi} \left[(e^u - e^{-u}) + \frac{1}{3}(e^{3u} - e^{-3u}) + \dots \right]$ where $u = \frac{i\pi x}{l}$]

109. $f(x) = e^{-x}$ in $(-\pi, \pi)$ and $(-1, 1)$.

[Ans: $f(x) = \sum_{n=-\infty}^{\infty} \frac{(-1)^n (1 - in\pi) \sinh 1}{1 + n^2 \pi^2} e^{in\pi x}$]

110. $f(x) = \cos ax$ in $(-\pi, \pi)$, where a is not an integer.

[Ans: $f(x) = \frac{a \sin a\pi}{\pi} \sum_{n=-\infty}^{\infty} \frac{(-1)^n}{(a^2 - n^2)} e^{inx}$]

111. $f(x) = e^{ax}$ in $(0, 2\pi)$ where a is not an integer.

112. $f(x) = 2x$ in $(0, 2\pi)$ [Ans: $f(x) = \sum_{n=-\infty}^{\infty} \frac{2i}{n} e^{inx}$]

113. $f(x) = \cosh 2x + \sinh 2x$ in $(-5, 5)$

[Ans: $f(x) = \sum_{n=-\infty}^{\infty} \frac{(-1)^n \sinh 10 \cdot (10 + in\pi)}{(100 + n^2 \pi^2)} e^{in\pi x/5}$]

114. $f(x) = \cosh 2x + \sinh 2x$ in $(-2, 2)$

[Ans: $f(x) = \sum_{n=-\infty}^{\infty} \frac{(-1)^n \sinh 4 \cdot (4 + in\pi)}{(16 + n^2 \pi^2)} e^{in\pi x/2}$]

115. $f(x) = 1, \quad 0 < x < 1$

$= 0, \quad 1 < x < 2$

116. $f(x) = \cosh ax$ in $(-\pi, \pi)$

[Ans: $f(x) = \frac{a \sinh a\pi}{\pi} \sum_{n=-\infty}^{\infty} \frac{(-1)^n}{(a^2 + n^2)} e^{inx}$]

117. $f(x) = x^2 + x$ in $(-\pi, \pi)$

118. $f(x) = e^{-ax}$ in $(-2, 2)$ where a is not an integer.

119. $f(x) = \cos hx$ in $(-1, 1)$

120. If $f(x) = \begin{cases} 1, & 0 < x < 1 \\ 0, & 1 < x < 2 \end{cases}$ and $f(x) = f(x+2)$ for all x

And hence prove that $f(x) = \frac{1}{2} + \frac{2}{\pi} \left\{ \frac{\sin(\pi x)}{1} + \frac{\sin(3\pi x)}{3} + \frac{\sin(5\pi x)}{5} + \dots \right\}$

FOURIER INTEGRAL

121. Express the function $f(x) = \begin{cases} -e^{kx}, & \text{for } x < 0 \\ e^{-kx}, & \text{for } x > 0 \end{cases}$ as Fourier Integral and prove that

$$\int_0^{\infty} \frac{\omega \sin \omega x}{\omega^2 + k^2} d\omega = \frac{\pi}{2} e^{-kx} \quad \text{if } x > 0, k > 0$$

122. Using Fourier Cosine integral prove that $e^{-x} \cos x = \frac{2}{\pi} \int_0^{\infty} \left(\frac{\omega^2 + 2}{\omega^4 + 4} \right) \cdot \cos \omega x d\omega$

123. Find Fourier Integral for $f(x) = \begin{cases} 1-x^2 & \text{for } |x| \leq 1 \\ 0 & \text{for } |x| > 1 \end{cases}$ [Ans: $f(x) = \frac{4}{\pi} \int_0^{\infty} \frac{\sin \omega + \omega \cos \omega}{\omega^3} \cdot \cos \omega x d\omega$]

124. Express the function $f(x) = \begin{cases} \sin x, & 0 < x \leq \pi \\ 0, & x < 0, x > \pi \end{cases}$ as Fourier Integral. and prove that

$$f(x) = \frac{1}{\pi} \int_0^{\infty} \frac{\sin \omega x + \cos[\omega(\pi - x)]}{1 - \omega^2} d\omega \quad \text{Hence deduce that} \quad \int_0^{\infty} \frac{\cos(\omega\pi/2)}{1 - \omega^2} d\omega = \frac{\pi}{2}$$

125. Find Fourier Integral representation for $f(x) = \begin{cases} e^{ax} & x \leq 0, a > 0 \\ e^{-ax} & x \geq 0, a > 0 \end{cases}$

Hence show that $\int_0^{\infty} \frac{\cos \omega x}{\omega^2 + a^2} d\omega = \frac{\pi}{2a} e^{-ax}, \quad x > 0, a > 0$

126. Find Fourier Sine integral representation for $f(x) = \frac{e^{-ax}}{x}$ [Ans: $f(x) = \frac{2}{\pi} \int_0^{\infty} \sin \omega x \cdot \tan^{-1}(\omega/a) d\omega$]

127. Find Fourier Cosine integral for $f(x) = \begin{cases} 1-x^2 & 0 \leq x \leq 1 \\ 0 & x > 1 \end{cases}$ Hence evaluate $\int_0^{\infty} \frac{\sin \omega - \omega \cos \omega}{\omega^3} \cos \frac{\omega}{2} d\omega$

[Ans: $\frac{3\pi}{16}$] Hint: put $x = 1/2$

D1. Express the function $f(x) = \begin{cases} \sin x, & 0 < x \leq \pi \\ 0, & x < 0, x > \pi \end{cases}$ as Fourier sine Integral and evaluate $\int_0^{\infty} \frac{\sin \omega x \cdot \sin \pi \omega}{1 - \omega^2} d\omega$

D2. Find Fourier sine integral of $f(x)$ where $f(x) = \begin{cases} x, & 0 < x < 1 \\ 2-x, & 1 < x < 2 \\ 0, & x > 2 \end{cases}$

D3. Express $f(x) = e^{-kx}$ ($k > 0$) as Fourier Sine and Cosine Integral and show respectively that

$$\text{i) } \int_0^{\infty} \frac{\omega \sin \omega x}{k^2 + \omega^2} d\omega = \frac{\pi}{2} e^{-kx} \quad \text{ii) } \int_0^{\infty} \frac{\omega \cos \omega x}{k^2 + \omega^2} d\omega = \frac{\pi}{2k} e^{-kx}$$

D4. Find Fourier Integral representation for $f(x) = \begin{cases} x, & 0 < x < a \\ 0, & x > a \end{cases}$ and $f(-x) = f(x)$

FOURIER TRANSFORM

D5. Find Fourier Transform of $f(x) = \begin{cases} x, & |x| \leq a \\ 0, & |x| > a \end{cases}$

D6. Find Fourier Transform of $f(x) = \begin{cases} 1 - |x|, & |x| \leq 1 \\ 0, & |x| > 1 \end{cases}$, hence show that $\int_0^{\infty} \frac{\sin^2 x}{x^2} dx = \frac{\pi}{2}$

D7. Find Inverse Fourier Transform of $\Phi(s)$ defined as $\Phi(s) = \begin{cases} 1 + s^2, & |s| \leq 1 \\ 0, & |s| > 1 \end{cases}$

D8. Find Fourier Cosine Transform of e^{-ax} , $a > 0$

D9. Find Fourier Sine Transform of e^{-ax} , $a > 0$, hence find $F_S(xe^{-ax})$ and $F_S \frac{e^{-ax}}{x}$. Also deduce the value of $\int_0^{\infty} \frac{\sin sx}{x} dx$.

D10. Find Fourier Cosine Transform of $f(x) = \begin{cases} \cos x, & 0 < x < a \\ 0, & x > a \end{cases}$

D11. Find $f(x)$ if $\int_0^{\infty} f(x) \sin sx dx = e^{-as}$

D12. Find Inverse Fourier Cosine Transform of $\frac{1}{1+s^2}$

D13. Find Fourier Sine Transform and Fourier Cosine Transform of $f(x)$ if $f(x) = \begin{cases} 1, & 0 < x < 1 \\ 0, & x > 1 \end{cases}$

D14. Find $f(x)$ if $\int_0^{\infty} f(x) \cos sx dx = \frac{\sin s}{s}$