



4

Vedhavyas Pavankalyan G L 2023-IT-A ▾V2**Started on** Sunday, 19 October 2025, 5:56 PM**State** Finished**Completed on** Sunday, 19 October 2025, 5:56 PM**Time taken** 41 secs**Marks** 1.00/1.00**Grade** 4.00 out of 4.00 (100%)

Question 1 | Correct Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that $A[j] - A[i] = k$, $i \neq j$.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as $5 - 1 = 4$

So Return 1.

For example:

Input	Result
3	1
1 3 5	
4	

Answer: (penalty regime: 0 %)

Ace editor not ready. Perhaps reload page?

Falling back to raw text area.

```
#include <stdio.h>
```

```
int main() {
    int n;
    scanf("%d", &n);
    int a[n];
    for (int i = 0; i < n; i++)
        scanf("%d", &a[i]);
    int k;
    scanf("%d", &k);
    int i = 0, j = 1;
    while (i < n && j < n) {
        int diff = a[j] - a[i];
        if (i != j && diff == k) {
            printf("1");
            return 0;
        } else if (diff < k)
            j++;
    }
}
```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

[Back to Course](#)