F.No 3-6/2022-23/IPM/Advisory
भारत सरकार/ Government of India
कृषि एवं किसान कल्याण मंत्रालय / Ministry of Agriculture & Farmers Welfare
कृषि एवं किसान कल्याण विभाग / Department of Agriculture & Farmers Welfare
वनस्पति संरक्षण, संगरोध एवं संग्रह निदेशालय / Directorate of Plant Protection, Quarantine & Storage
एनएच4, फरीदाबाद (हरियाणा) – 121 001/ NH-IV, Faridabad (Haryana) – 121 001

Date: 19.06.2025

To

The Director of Agriculture
Government of Madhya Pradesh & Maharashtra

Sub: Preparatory advisory for timely management of vector, white fly and Yellow Mosaic Virus (YMV) in Soybean -reg.

Sir,

You are aware that the soybean yellow mosaic is a serious and widespread disease of soybean and its incidence was observed in Madhya Pradesh & Maharashtra states during last two years. The disease is caused by Soybean Yellow Mosaic Virus (SYMV), the virus is sap transmitted and spread by vector white fly (*Bemisia tabaci*). The disease symptoms appear as yellow spots which are either scattered or produced in indefinite bands along the major veins of soybean leaves. Rusty necrotic spots appear in the yellow areas as the leaves mature. Some time severe mottling and crinkling of leaves are also seen. Leaves of severely infected plants become yellow when they are young. Affected plants bear less flower and pods. The infection results in decrease in oil and increase in protein content.

It is observed that the disease incidence is progressing in soybean growing areas during past years and it may increase on other soybean growing areas. As the Soybean Yellow Mosaic Virus is transmitted by insect vector and its incidence is to be observed at early crop stage so there is requirement for adoption of integrated approach from sowing of the crop.

Keeping in view the ensuing Kharif season, the soybean crop should be monitored regularly for disease incidence and adopt Integrated Pest Management techniques beginning from sowing. Important immediate and long term IPM practices for management of Soybean Yellow Mosaic Virus are listed at **Annexure-I**

Further, it is informed that the Government of India, Ministry of Agriculture & Farmers Welfare has launched the AI-driven National Pest Surveillance System (NPSS) application which is designed to enhance pest surveillance and management across the nation, providing significant benefits to the farmers in the form of to gain real time advisories for pest management. NPSS also covers soybean crop under pest identification and pest surveillance module which provides complete solution for pest management at farm level.

In view of the above, it is advised to disseminate the above advisory among all extension functionaries of your state in soybean growing areas with advice to monitor soybean crop on regular basis for incidence of vector white fly and its timely and effective management so as to arrest the spread of Yellow Mosaic Virus (YMV). The detailed information on incidence of vector white fly and YMV may also be sent to this Directorate through e-mail: appa.ipm-ppqs@gov.in for further needful action.

Yours sincerely

(Dr. J. P. Singh)
Plant Protection Adviser

Copy for information:

1. PPS to Joint Secretary(PP), DA&FW, Krishi Bhawan, New Delhi- 110001
2. PPS to PPA for information.
3. Officer In-charges, all CIPMCs of soybean growing states with a direction to conduct regular survey in their areas to ascertain the vector/disease incidence and necessary follow-up with the state authorities

Annexure-I

Important immediate and long term IPM practices for management of Soybean Yellow Mosaic Virus

I. Immediate Measures

1. Regularly monitor population of whitefly and take effective IPM measures to manage its population.
2. Use of yellow sticky traps (20-25 traps/ha) in early crop stage to catch the flying adults of white fly thereby helps to minimizing the further transmission of YMV.
3. Use of CIBRC recommended chemicals at right time: For white fly management in standing crop, as soon as the symptoms appear, spray the crop with Acetamiprid 25% + Bifenthrin 25 % WG @ 250gm in 500 litre of water/ha. If the pest crossed Economic Threshold level (ETL) i.e., 21 Nymphs or Adults per leaf. **The list of CIB & RC approved pesticides for management of vector of disease (White fly) in soybean is available on** https://ppqs.gov.in/divisions/cib-rc/major-uses-of-pesticides
4. Identification, removal and destruction of infested plants from the field: Plants showing symptoms of Yellow Mosaic Virus (YMV) should be removed from the field to prevent further spread of the disease.
5. Avoid water stagnation in the field during crop stand.

II. Long term measure:

1. Advised to follow recommended sowing time.
2. Use of recommended tolerant/resistant varieties.
3. Use preventive methods (using disease-free seeds, adopting crop rotation, and inter cropping).
4. Seed treatment: Before sowing, treat the seed with Thiamethoxam 30 % WS or Thiamethoxam 30 % FS @ 6 ml/kg seed
5. Use of recommended seed rate, spacing and use of fertilizers.
6. The weeds in nearby area should be removed/ destroyed as virus survives in them in off season.
7. Removal and destruction of infested/ infected plants from the field after harvest of the crop.