Data handling & Visualization | Activity 9 | 24-10-2025 | Yaveeshtaa Reddy | 192825102 | S1CM-C

① Time Series & trend :-

```
n <- length(sales)
fit <- lm(sales~t)
plot(x)
coef(fit)[2]
```

Answers:- n=44, cycle length = 4 quarters, full cycles = 11, trend is rising, slope = 3.977 lakh/qtr

---

② Seasonal pattern :-

```
d <- decompose(x, type="multiply")
d$figure
which.max(d$figure)
which.min(d$figure)
diff(range(d$figure))
```

Answer:- Peak = Q4, trough = Q2 - Theoretical Seasonal factors are : 1.0, 0.7, 1.0, 1.3; theoretical swing = 0.60.

---

③ Consistency :-

```
cy <- matrix(sales, nrow=4)
sum(apply(cy, 2, which.max)==4)
matplot(1:4, cy, type="l", lty=1)
```

Answer:- Seasonality repeats every 4 quarters. The code counts how many of 11 cycles peak in Q4

(4). * Autocorrelation :-

```
e <- resid(lm(Sales ~t))
a<- acf(e, lag.max = 4, plot = FALSE)$acf
a[c(3,5)]
2/ sqrt(44)
```

Answer :- Significance bound = 0.302,
Compare absolute Acf values at lags 2 & 4 with 0.302.

---

(5). * Seasonality type :-

```
cy <- matrix(Sales, nrow=4)
r1 <- differ(range(cy[,1]))
r2 <- diff(range(cy[,11]))
c(r1, r2, r2/r1)
adj <- Sales / rep(cd$figure, 11)
Coef(lm(adj ~t))[2]
```

Answer :- Multiplicative seasonality, Sales are expected to peak in Q4 & reach a trough in Q2. plan inventory & staffing accordingly.