

Based on the likelihood ratio test, there **is** evidence of *episodic diversifying selection* in this dataset (p=0.01901).

BUSTED analysis (v4.0) was performed on the alignment from
/Users/sergei/Dropbox/Talks/VEME-current/data/spike.fas using HyPhy v2.5.40.
This analysis **did not include** site-to-site synonymous rate variation.

Suggested citation: *Gene-wide identification of episodic selection*, Mol Biol Evol.
32(5):1365–71, *Synonymous Site-to-Site Substitution Rate Variation Dramatically Inflates False Positive Rates of Selection Analyses: Ignore at Your Own Peril*, Mol Biol Evol.
37(8):2430–2439

Evidence ratio threshold

10

Update

118

sequences in the alignment

⋮

1273

codon sites in the alignment

⋯

1

partitions

^

162

median branches/partition used for testing

🔗

3 classes

non-synonymous rate variation

📦

None

synonymous rate variation

📁

0.019

p-value for episodic diversifying selection

⊕

11

Sites with ER≥10 for positive selection

⚡

N/A:N/A

Multiple hit rates (2H:3H)

🎯

Alignment-wide results

Model	Log (L)	AIC-c	Params.	Rate distribution	Rate plot
Unconstrained model	−9287.35	18937.1	181	Tested ω 0.000 (0.45262%) 0.2561 (97.532%) 14.88 (2.0156%) Mean = 0.5497 , CoV = 3.740	
Constrained model	−9290.62	18941.7	180	Tested ω 0.000 (46.919%) 1.000 (2.2974%) 1.000 (50.784%) Mean = 0.5308 , CoV = 0.9402	

Gene-wide selection analysis using a branch-site method (BUSTED), SARS-CoV-2 spike

```
hyphy busted --srv No --alignment data/spike.fas --tree data/spike.tree --starting-points 5
```

Produces *HIV-sets.fas.BUSTED.json* file
View in <http://vision.hyphy.org/BUSTED> or <https://observablehq.com/@spond/busted>

BUSTED analysis

- **West Nile Virus NS3 protein**

- No statistical support for selection; ML point estimate allocates a small proportion of sites ($\sim 1\%$) to the selected group ($dN/dS \sim 2$)
- The rest of the gene is very strongly conserved ($dN/dS = 0.004$)

- **HIV-1 transmission pair**

- Very strong evidence of strong episodic diversification ($dN/dS \sim 100$) on a small proportion of sites (2%)

- The rest of the gene evolves with weak purifying selection ($dN/dS = 0.6-0.7$)

- **SARS-CoV-2 spike**

- Evidence of episodic diversification ($dN/dS \sim 15$) on a small proportion of sites ($\sim 2\%$)
- Most of the rest of the gene evolves with purifying selection ($dN/dS = 0.2$)