

Composing the error budget of an optical clock: one rate scalar per atom

Michael Rudolph 

*Velar Scientific**

(Dated: August 30, 2026)

An optical clock’s systematic error budget is built today as a table: one row per physical effect (blackbody radiation, lattice or probe Stark shifts, the electric-quadrupole shift, motional time dilation, gravitational redshift), each row computed in its own formalism, the rows summed linearly, their uncertainties combined in quadrature, and fringe contrast treated as a separate measurement outside the budget. We describe an alternative composition already implemented and validated in the open-source CliffordClock pipeline: every systematic enters as a multiplicative factor on a single scalar rate at which an atom’s own proper time advances along its worldline, the accumulated phase is the time integral of that rate, the clock’s reported fractional shift is the mean of the resulting ensemble phase distribution, and the Ramsey fringe visibility is the modulus of the mean phasor, fixed by that same distribution’s phase variance under the standard Gaussian-phase closure. The additive table is the first-order term of this product’s own expansion, and we give the pipeline’s own bounds on the terms it drops: a DC-Stark $\times$ blackbody product correction of order 10^{-33} and a gravitational $\times$ field-driven cross term of order 10^{-31} , both at least twelve orders of magnitude below the 10^{-18} - 10^{-19} systematic floor of the clocks the budget describes. The value of the product form is structural. It changes no published total: the terms it adds beyond first order sit at least twelve orders of magnitude below current systematic floors, a bound this paper states directly from the pipeline’s own composition rules. We show three places the structure pays with numbers already validated in the companion paper: a multi-surface thermal-environment term whose higher-order moments must be evaluated per surface before they are combined; a trapped-ion secular-motion reconciliation where composing two systematics as correlated factors on one rate closes a discrepancy of 14σ down to 0.08σ against a published budget; and a Ramsey visibility read from the same worldline ensemble that sets the mean shift, computed with no separate fit. Relativistic timing already composes rate scalars for its own gravitational and kinematic terms; this paper’s own claim is one of scope: the same composition extended to the atomic-environment terms of a clock’s full budget, evaluated per quantum state, with fringe contrast recovered from the phase variance of the same object, in an open implementation checked against published budgets. Code, tests, and the ten-line composition rule itself are public under the AGPLv3.

I. INTRODUCTION

A modern optical clock’s systematic-uncertainty budget is built one row at a time. A group evaluates the blackbody-radiation shift from a Planck-integral model of the clock’s radiative environment, the lattice or probe AC-Stark shift from the trap-light polarizability, the DC-Stark shift and its spatial gradient from a field survey, the electric-quadrupole shift (for an ion with $J \geq 1$) from the local field-gradient tensor and an atomic theory coefficient, the motional time-dilation shift from sideband-thermometry-measured mode populations, and the gravitational redshift from a height survey against a local reference [1]. Each row is a physically independent calculation, in its own formalism, against its own inputs. The rows are then summed linearly to a total fractional shift, and their uncertainties are combined in quadrature to a total systematic uncertainty. The clock’s fringe contrast, the visibility that sets how well the interrogation resolves the transition, is measured separately, as an experimental observation outside the budget’s own rows.

Two recent publications make this practice concrete with real numbers. Aepli et al. report an ^{87}Sr optical lattice clock’s blackbody shift as $-4.841720 \times 10^{-15}$, with a stated uncertainty of $\pm 7.3 \times 10^{-19}$, from a Planck-integral model of the clock’s measured in-vacuum temperature combined with fitted dynamic polarizability coefficients [2]. Their total systematic uncertainty, after every such row is combined, is 8×10^{-19} [2]. Bothwell et al., on the same platform, publish a DC-Stark gradient row of $+0.3(0.2) \times 10^{-20}$ per millimetre from a field survey and, in the same table, a gravitational-redshift slope from two independent analysis methods that give $-9.8(2.3) \times 10^{-20}$ and $-12.8(2.7) \times 10^{-20}$ per millimetre respectively, the two methods differing from each other by 0.85σ even though both describe the same physical effect [3]. On a different platform and a different species, Brewer et al. report an $^{27}\text{Al}^+ / ^{25}\text{Mg}^+$ trapped-ion clock’s own motional time-dilation row as $-17.3(2.9) \times 10^{-19}$, one line among the rows that sum to their clock’s total

* michael.rudolph@velarscientific.com

systematic uncertainty, published as below 10^{-18} [4]. A later evaluation of the same species pair on a different trap, Marshall et al., reports the same row at $-114.6(3.8) \times 10^{-19}$ and a total systematic uncertainty of 5.5×10^{-19} [5]. Four papers, two atomic species, two very different orders of magnitude for nominally the same physical row: the practice that produces these numbers is precise and well established. It is a practice, and this paper asks what one composed object, replacing the table of separately computed rows, would look like, and where it would matter.

CliffordClock is an open-source pipeline, described and validated against published clock budgets in a companion paper,¹ that already implements the alternative this paper describes. Every systematic in that pipeline, DC-Stark, blackbody radiation, the electric-quadrupole shift, motional time dilation, gravitational redshift, enters as a multiplicative factor on one scalar rate at which an atom's own proper time advances, and the clock's reported shift and its fringe visibility both come out of the same accumulated-phase object. Section II states that composition rule. Section III derives the additive table as its first-order expansion and gives the pipeline's own bound on what that expansion drops. Section IV shows three places, already validated against published numbers, where the fuller composition changes what a lab would conclude. Section V places this against relativistic-timing practice, which already composes rate factors for gravitational and kinematic terms. Section VI gives the composition rule in ten lines of code.

II. THE COMPOSITION LAW

Write $\nu(t)$ for the clock transition's instantaneous frequency along one atom's worldline, its own path through space, time, and internal quantum state. In the standard treatment, each systematic k contributes an independently computed fractional perturbation p_k , and the rows are added: $\nu(t)/\nu_0 - 1 \approx \sum_k p_k$. The composition rule this paper describes instead multiplies:

$$\nu(t) = \nu_0 \prod_k (1 + p_k(x(t))), \quad (1)$$

where $x(t)$ is the atom's full state along its worldline at time t , position, velocity, and any internal or motional quantum number each term's own physics depends on. Equation (1) is the local proper-time rate: general relativity already has names for a scalar factor multiplying an ideal clock's rate this way, the lapse function in a 3+1 decomposition of spacetime, or, for a static observer, the redshift factor $\sqrt{-g_{00}}$. Each term p_k is still computed by its own physics: the DC-Stark shift from the transition's differential polarizability and the local field, the blackbody shift from a Planck-integral model of the radiative environment, the quadrupole shift from the local field-gradient tensor, the motional shift from the atom's own velocity or vibrational occupation, the gravitational term from height in a local potential. Nothing about any individual term changes. What changes is how the terms are put back together. Equation (1) multiplies them as factors on one shared rate. A budget table instead adds them, one row per systematic, an addition Sec. III shows is this product's own first-order truncation.

An atom's accumulated phase over an interrogation of proper duration T is the time integral of that rate,

$$\Delta\Phi = 2\pi \int_0^T \nu(t) dt, \quad (2)$$

in radians. An ensemble of M atoms, each with its own worldline (its own trajectory, motional state, and local field environment), gives M accumulated phases $\Delta\Phi_1, \dots, \Delta\Phi_M$. The clock's reported fractional frequency shift is the ensemble mean of these phases, normalized by the unperturbed phase $2\pi\nu_0 T$:

$$\frac{\Delta\nu}{\nu_0} = \frac{\langle \Delta\Phi \rangle}{2\pi\nu_0 T} - 1, \quad (3)$$

the first moment of the ensemble's phase distribution, converted to a fractional frequency. The Ramsey fringe visibility, the same contrast a lab measures at the end of an interrogation, is the modulus of the mean phasor over that same ensemble,

$$V = |\langle e^{i\Delta\Phi} \rangle| \leq 1. \quad (4)$$

For phases that are Gaussian-distributed across the ensemble, the characteristic-function identity $\langle e^{i\Delta\Phi} \rangle = e^{i\langle \Delta\Phi \rangle - \sigma_\Phi^2/2}$ gives the closed form $V = \exp(-\sigma_\Phi^2/2)$ exactly: the visibility is fixed by the phase distribution's variance, the

¹ The companion paper is `paper/main.tex` in the project repository, <https://github.com/velar-mbr/CliffordClock>. Every case-study number in this paper is copied from that paper, from `docs/CONVENTIONS.md`, or from `notebooks/13.trapped_ion_quantum_motion.ipynb`; this paper's own text does not recompute any of them.

same distribution whose mean fixed the shift in Eq. (3). One worldline ensemble, run once, yields both numbers. Section IV C validates this identity against a live pipeline run.

Two constructions of the ensemble average must be told apart, because one of them silently destroys the visibility signal. Averaging the per-worldline phases first and only then taking the phasor, $|\exp(i\langle\Delta\Phi\rangle)|$, gives modulus 1 for any spread in the underlying phases, because the spread is discarded before the modulus is ever taken. The correct order, Eq. (4), sums the phasors themselves and takes the modulus of that sum, which is what lets a real phase spread show up as $V < 1$. Both orderings are encoded as opposing kill tests in the pipeline’s own test suite (`tests/test_coherent_visibility.py`), so a future change that collapses the two back together is caught mechanically.

III. EQUIVALENCE AND SCOPE

Expanding the product in Eq. (1) to first order in each p_k recovers the additive budget exactly:

$$\prod_k (1 + p_k) = 1 + \sum_k p_k + \sum_{j < k} p_j p_k + \dots \quad (5)$$

The standard table, one row per systematic, rows summed linearly, is the truncation of Eq. (5) at first order. Every cross term the truncation drops is a product of two or more systematics that are each already small, and at the field’s current performance those products are far smaller still. The pipeline’s own composition rules state each bound directly, as part of the codebase’s own documentation. The DC-Stark and blackbody-radiation terms are independent scalar perturbations of the atom’s internal energy. Their second-order cross term averages to exactly zero for an isotropic thermal field, an exact cancellation, and their multiplicative product correction, the Stark×BBR cross term of Eq. (5), is bounded at order 10^{-33} .² The gravitational-redshift term depends only on an atom’s position in the local potential, and never on its motional or internal state, so its cross term with any field-driven shift is bounded at order 10^{-31} .³ Both bounds sit at least twelve orders of magnitude below the 10^{-18} -to- 10^{-19} systematic floor the clocks of Sec. I report.

This comparison settles what the product form is for. It states one composition rule that explains why the additive table works as well as it does, and it gives that rule’s own quantitative boundary, the order at which it would stop working, in the same units the rows themselves are reported in. It does not correct any of the published totals in Sec. I: the additive budgets those papers report are correct to a precision no experiment now approaches, and nothing in this paper revises a single published number. The three cases of Sec. IV instead show where the product form’s other structural consequences change a conclusion: evaluating a nonlinear function of a distributed quantity before averaging it, composing two systematics as correlated factors on one shared rate, and reading a fringe visibility off the same object that sets the shift. None of the cross-term corrections above ever enters that arithmetic.

IV. WHERE THE STRUCTURE PAYS

A. Per-state evaluation before averaging: a multi-surface thermal environment

A blackbody-radiation budget row is ordinarily a single number: a lab characterizes its enclosure by one effective temperature and evaluates a Planck-integral shift formula at that one temperature. A clock deployed outside a temperature-stabilized laboratory chamber, in a vehicle or behind a viewport a diurnal cycle warms, sees an enclosure whose visible surfaces can differ by tens of kelvin, and the blackbody shift’s dynamic term scales as T^6 through T^{10} , higher powers than the T^4 power the static shift and the naturally chosen effective temperature both track. A single temperature chosen to match the static (T^4) moment of a non-uniform environment does not, in general, reproduce the environment’s higher dynamic moments, and the companion paper quantifies the resulting mismatch: it crosses 10^{-18} , comparable to a modern lattice clock’s total systematic budget, at roughly an 11 K spread across two comparably sized surfaces, and at roughly 19 K for a small warm viewport against a larger chamber body.⁴

The pipeline’s multi-surface extension evaluates the blackbody shift’s moments per surface, before they are combined into a single number. Writing w_i for each surface’s solid-angle fraction and T_i for its temperature, the environment’s n -th weighted moment is $M_n = \sum_i w_i (T_i/T_0)^n$, and the static and dynamic terms of the shift formula are each

² docs/CONVENTIONS.md, section 13, “(E33) Scalar pivot composition.”

³ Companion paper, Sec. II.E, “The gravitational-redshift pivot term and extended samples.”

⁴ Companion paper, Sec. II.D, “The multi-surface thermal environment.”

TABLE I. Six successive treatments of the same published Al^+ secular-motion total, computed by the CliffordClock pipeline against Marshall et al.’s own published six-mode inputs [5], copied from the companion paper’s own progression table and Fig. 8 there. σ is the distance from Marshall et al.’s published band, combining the published and predicted uncertainties in quadrature. Participation and micromotion enter as two multiplicative factors on the same per-mode rate (Eq. (1)).

Treatment	σ	Verdict
Single species mass, every mode	0.10	MET
Participation, mass-ratio closed form	14.11	NOT MET
Participation, measured-spectrum fit	14.01	NOT MET
Participation $\times$ leading-order micromotion	1.62	NOT MET
Participation $\times$ exact-Floquet micromotion	1.62	NOT MET
Coupled two-ion Floquet, constrained fit	0.08	MET

evaluated against their own moment M_n , in place of a single power $(T/T_0)^n$ of one effective temperature.⁵ A uniform, single-surface environment reduces this construction to the ordinary single- T formula term for term: the two paths share the same coefficient-evaluation code and agree at floating-point precision. The construction is validated against a real transportable clock’s own position-resolved measurement: replaying Nosske et al.’s published aperture geometry, interior emissivity, and shield temperatures through the pipeline’s enclosure-and-aperture model predicts a differential shift, as the atoms move from the shield’s center out through an aperture, of -3.325×10^{-15} , against their own closed-form model’s $-3.32(7) \times 10^{-15}$ and their own measurement’s $-3.33(3) \times 10^{-15}$, landing within 5.011×10^{-18} of their model and 4.989×10^{-18} of their measurement, well inside either one’s own stated uncertainty.⁶

The point generalizes past blackbody radiation. A single effective value stands in for a physically distributed quantity throughout an optical-clock budget, an effective lattice depth over a trapped ensemble’s spatial extent, an effective field over a cloud’s own spread, and each substitution risks the same failure mode a single effective temperature shows here: a nonlinear function of the distributed quantity, evaluated at one collapsed value, does not equal that same nonlinear function’s own moments evaluated across the real distribution and then combined. The composition rule of Sec. II states the same discipline once, for every systematic and every ensemble the pipeline evaluates: the distribution comes first, and the moments (the mean that sets the shift, the variance that sets the visibility) come from integrating over the full distribution, before any collapse to one representative value.

B. A cross-row correlation a table cannot see: reconciling a trapped-ion secular-motion budget

A trapped two-ion clock’s motional time-dilation shift depends on two physically distinct pieces of the same underlying motion: how much of each shared vibrational mode the clock ion itself carries (its participation), and how much extra oscillation, at the trap’s own radio-frequency drive, that secular motion itself induces (intrinsic micromotion). A budget table built one row at a time can compute either piece on its own, and Marshall et al.’s published $^{27}\text{Al}^+ / ^{25}\text{Mg}^+$ two-ion crystal is a real case where doing so goes badly wrong before it goes right. Table I reproduces the companion paper’s own six-step progression through this reconciliation, each row built from Marshall et al.’s own published six-mode frequencies and occupations, reported against Marshall et al.’s own published secular-motion band of $-114.6(3.8) \times 10^{-19}$.⁷

The first row carries one species mass through every one of the six modes, with no participation split and no micromotion correction, and lands inside Marshall et al.’s own published band at 0.10σ . That agreement is a near-cancellation of two large omissions moving in opposite directions. The true clock-ion participation in a shared mode is close to one-half, and the true intrinsic-micromotion enhancement on the radial modes is close to a factor of two; the single-mass row’s implicit choice of one for both factors happens to land close to their true product, an accident of the approximation the next two rows expose. Supplying only the participation correction, rows two and three, removes that accidental cancellation without replacing it: the resulting total under-predicts the published band by a factor of close to two, a 14σ discrepancy against Marshall et al.’s own stated uncertainty. Composing the intrinsic-micromotion enhancement back in as a second multiplicative factor on the same per-mode rate, rows four and five, closes most of that distance, to 1.62σ , still outside the published band. The remaining gap traces to a specific mechanism the companion paper identifies: the two-mode pairs sharing an axis run at Coulomb-shifted collective frequencies a single-ion enhancement factor cannot see. Solving the two ions’ coupled equations of motion, then

⁵ docs/CONVENTIONS.md, section 13, “(E37) Multi-surface thermal environment.”

⁶ Companion paper, Sec. IV.J, “The PTB position-resolved thermal-environment validation,” citing Nosske et al. [6].

⁷ Companion paper, Secs. IV.H-I, “The Al -ion secular-motion arithmetic reproduction” and “Per-mode participation and the intrinsic-micromotion enhancement”; docs/CONVENTIONS.md section 16.

fitting the resulting two-parameter trap model to all four measured radial frequencies at once, closes the budget to 0.08σ , row six, comfortably inside Marshall et al.’s own published band.

A budget table computes participation and micromotion as separate rows, or omits the smaller of the two once the first row already looks close to the published value. Composing them as two correlated factors on the same per-mode rate exposes the near-cancellation the single-mass row hides. It also lets the coupled two-ion physics that resolves it enter as one more correction to that shared rate, the same object every other correction in this section already multiplies.

C. Shift and coherence from one object: Ramsey visibility

Section II states that the same worldline ensemble that sets the mean fractional shift, Eq. (3), also sets the Ramsey fringe visibility, Eq. (4). The visibility calculation uses the same accumulated-phase draw as the shift calculation, with no free parameter of its own. The companion paper validates the Gaussian closure $V = \exp(-\sigma_\Phi^2/2)$ this identity predicts against a live pipeline run: sweeping a classical worldline ensemble’s squeezing parameter over several values at 4000 worldlines each, the largest relative deviation between the pipeline’s own reported visibility and the same run’s independently computed Gaussian-closure check lands at roughly 0.65%, at the largest squeezing swept, consistent with the $O(\sigma^4/N)$ Monte Carlo scaling the pipeline’s own test-tolerance derivation documents for this comparison.⁸ That number comes from the same accumulated-phase distribution Eq. (3) already builds to report the mean shift, with the same worldline draw and the same weights supplying both outputs.

V. RELATION TO PRIOR PRACTICE

Relativistic timing already composes a rate scalar for a clock’s own systematics, for the terms general relativity itself supplies. Ashby’s treatment of relativity in the Global Positioning System expresses each GPS satellite clock’s rate, relative to a ground clock, as a product of the gravitational potential term and the special-relativistic velocity term, composed multiplicatively along the satellite’s worldline, the same proper-time-rate object this paper composes for an optical clock’s atomic-environment terms [7]. Chronometric geodesy extends the same idea in the other direction: an optical clock’s own measured rate, relative to a reference, is read as a measurement of the gravitational potential difference between the clock and the reference, a practice a recent review of atomic clocks for geodesy surveys across a range of clock platforms and baselines, each one treating the clock’s fractional frequency as a rate factor tied to the local metric [8].

Both of these established practices compose rate scalars for the gravitational and kinematic terms general relativity itself supplies. This paper’s own claim is about scope. What CliffordClock adds is the same composition, extended to a clock’s full budget, including the atomic-environment terms, the DC-Stark shift, blackbody radiation, and the electric-quadrupole shift, that have no counterpart in a GPS satellite’s or a geodetic baseline’s own rate factor. Those terms are evaluated per quantum state where the physics calls for it, the per-surface thermal moments of Sec. IV A and the per-mode motional factors of Sec. IV B among them, and fringe contrast comes out of that same object, Sec. IV C, as a byproduct of the shift calculation itself. The implementation is open, and every claim above is checked against a published budget in the repository’s own test suite. No claim is made that this is the only way to compose a clock’s systematics, or the first time a rate scalar has been composed for any of its individual terms.

VI. THE COMPOSITION, IN TEN LINES

Every equation of Sec. II, the multiplicative rate, the phase integral, the shift as a mean, and the visibility as a modulus of the mean phasor, is this short:

```
def clock_phase(worldline, terms, nu_0):
    rate = lambda t: nu_0 * prod(1.0 + p(worldline.state(t)) for p in terms)
    return 2.0 * pi * integrate(rate, worldline.proper_time)

phases = [clock_phase(w, terms, NU_0) for w in ensemble]
shift = mean(phases) / (2.0 * pi * NU_0 * T) - 1.0
visibility = abs(mean(exp(1j * array(phases))))
```

⁸ notebooks/13_trapped_ion_quantum_motion.ipynb, Sec. 5, “Ramsey visibility under squeezing”; companion paper, Sec. II.H, “Coherent Ramsey visibility,” Eq. (E39).

Each entry of `terms` is one systematic’s own physics, computed from its own formalism exactly as any published budget row is; the composition itself is the four lines that use them. The source code, test suite, and the companion validation paper underlying every number in this paper are available at <https://github.com/velar-mbr/CliffordClock> under the GNU Affero General Public License v3.0 or later (AGPLv3).

-
- [1] A. D. Ludlow, M. M. Boyd, J. Ye, E. Peik, and P. O. Schmidt, Optical atomic clocks, *Rev. Mod. Phys.* **87**, 637 (2015).
 - [2] A. Aepli, K. Kim, W. Warfield, M. S. Safronova, and J. Ye, A clock with 8×10^{-19} systematic uncertainty, *Phys. Rev. Lett.* **133**, 023401 (2024), arXiv:2403.10664.
 - [3] T. Bothwell, C. J. Kennedy, A. Aepli, D. Kedar, J. M. Robinson, E. Oelker, A. Staron, and J. Ye, Resolving the gravitational redshift across a millimetre-scale atomic sample, *Nature* **602**, 420 (2022), arXiv:2109.12238.
 - [4] S. M. Brewer, J.-S. Chen, A. M. Hankin, E. R. Clements, C. W. Chou, D. J. Wineland, D. B. Hume, and D. R. Leibbrandt, An $^{27}\text{Al}^+$ quantum-logic clock with systematic uncertainty below 10^{-18} , *Phys. Rev. Lett.* **123**, 033201 (2019), arXiv:1902.07694.
 - [5] M. C. Marshall, D. A. Rodriguez Castillo, W. J. Arthur-Dworschack, A. Aepli, K. Kim, D. Lee, W. Warfield, J. Hinrichs, N. V. Nardelli, T. M. Fortier, J. Ye, D. R. Leibbrandt, and D. B. Hume, High-stability single-ion clock with 5.5×10^{-19} systematic uncertainty, *Phys. Rev. Lett.* **135**, 033201 (2025), arXiv:2504.13071.
 - [6] I. Nosske, C. Vishwakarma, T. Lücke, J. Rahm, N. Poudel, S. Weyers, E. Benkler, S. Dörscher, and C. Lisdat, Transportable strontium lattice clock with 4×10^{-19} blackbody radiation shift uncertainty, *arXiv preprint* (2025), arXiv:2507.14030.
 - [7] N. Ashby, Relativity in the global positioning system, *Living Rev. Relativity* **6**, 1 (2003).
 - [8] T. E. Mehlstäubler, G. Grosche, C. Lisdat, P. O. Schmidt, and H. Denker, Atomic clocks for geodesy, *Rep. Prog. Phys.* **81**, 064401 (2018).