

PMATH 352 Complex Analysis
velo.x

Contents

1	Differentiation	2
1.1	Introduction to Complex Number - May 2	2
1.2	Limit, Continuity, Differentiability - May 4	3
1.3	Derivative Continued - May 6	5
1.4	Holomorphic and Cauchy-Riemann Equation - May 9	7
1.5	Holomorphic and Cauchy-Riemann Equation (Continued) - May 11	9
1.6	Branch - May 13	11
1.7	Trigonometric Functions and Power - May 16	13
1.8	Rotalion Approximation - May 18	14
2	Integration	16
2.1	Integrability - May 20	16
2.2	May 25	18
2.3	May 27	19
2.4	May 30	21
2.5	June 1	23
2.6	Cauchy Integral Formula - June 3	24
2.7	June 6	26
2.8	June 8	26
2.9	June 10	28
2.10	June 13	29
2.11	Midterm Review - June 15	30
3	Applications of The Cauchy Theory	32
3.1	Taylor Series - June 20	32
3.2	June 22	34
3.3	Singularities - June 24	36
3.4	Singularities and Residue Theory - June 27	38
3.5	June 29	41
3.6	Extended Complex Plane - July 4	43
3.7	Cauchy Principal Value - July 6	45
3.8	July 11	48
3.9	July 11	48
3.10	July 13	50
3.11	July 15	51
3.12	July 18	53
3.13	July 20	54
3.14	July 22	56
3.15	July 25	58
3.16	Final Review	59

1 Differentiation

1.1 Introduction to Complex Number - May 2

DEFINITION 1.1.1 (Complex Number).

Define i to be which $i^2 = -1$, define the set of complex numbers to be

$$\mathbb{C} = \{a + bi : a, b \in \mathbb{R}\}$$

Note that there is no a priori distinction between i and $-i$. Thus all behaviour in \mathbb{C} should be invariant under a map $i \leftrightarrow -i$.

Operations:

- Addition:

$$(a + bi) + (c + di) = (a + c) + (b + d)i$$

- Multiplication

$$(a + bi)(c + di) = ac + (bc + ad)i + bdi^2 = (ac - bd) + (bc + ad)i$$

- Division: assume $c \neq 0$ or $d \neq 0$,

$$\frac{a + bi}{c + di} = \frac{a + bi}{c + di} \cdot \frac{c - di}{c - di} = \frac{ac + bd + (bc - ad)i}{c^2 + d^2}$$

- Conjugation:

$$\overline{a + bi} = a - bi$$

Note

$$(a + bi)(\overline{a + bi}) = (a + bi)(a - bi) = a^2 + b^2$$

Remark 1.1.2. There is a canonical bijection between \mathbb{R}^2 and \mathbb{C} . $a + bi \leftrightarrow (a, b)$. Therefore, every complex number $a + bi$ can be mapped as a point on the 2-dimensional axis.

Lemma 1.1.1. Every coordinate point (x, y) can be translated to polar coordinate (r, θ) where

$$r = \sqrt{x^2 + y^2}, \quad \theta = \arctan \frac{y}{x};$$

every polar coordinate point (r, θ) can be translated to (x, y) where

$$x = r \cos \theta, \quad y = r \sin \theta.$$

DEFINITION 1.1.3 (Norm).

$$|a + bi| = \sqrt{a^2 + b^2}$$

Lemma 1.1.2. Powers of i loop in a circle of 4.

$$i^1 = i \Rightarrow i^2 = -1 \Rightarrow i^3 = -i \Rightarrow i^4 = 1 \Rightarrow i^5 = i \Rightarrow \dots$$

Lemma 1.1.3. Let $x \in \mathbb{R}$ and $z = a + bi, w = c + di \in \mathbb{C}$. We have $\frac{d}{dz}e^z = e^z$ and $e^{w+z} = e^w e^z$ and $\frac{d}{d(iy)}e^{iy} = e^{iy}$.

Proof. UNFINISHED □

1.2 Limit, Continuity, Differentiability - May 4

DEFINITION 1.2.1 (Distance).

The distance between two points $w, z \in \mathbb{C}$ is $|w - z| = |z - w|$.

Remark 1.2.2. \mathbb{C} and \mathbb{R}^2 are isomorphic as metric spaces.

DEFINITION 1.2.3 (Open Set and Closed Set).

An **open set** $S \subseteq \mathbb{C}$ is a set such that for every $z \in S$, there is ε such that $|z - w| < \varepsilon \Rightarrow w \in S$.

A **closed set** is if $\mathbb{C} \setminus S$ is open.

DEFINITION 1.2.4 (Limit).

Let $f : \mathbb{C} \rightarrow \mathbb{C}$. We say $\lim_{z \rightarrow w} f(z) = L$ if for all $\varepsilon > 0 \in \mathbb{C}$, $\exists \delta > 0$ s.t. for all $z \in \mathbb{C}$, if $|z - w| < \delta$ then $|f(z) - L| < \varepsilon$.

Example 1.2.1. Consider $\lim_{z \rightarrow 0} \frac{\bar{z}}{z}$.

Try approaching in different directions:

- Let $z = x, x \in \mathbb{R}$, then,

$$\lim_{z \rightarrow 0} \frac{\bar{z}}{z} = \lim_{x \rightarrow 0} \frac{\bar{x}}{x} = \lim_{x \rightarrow 0} \frac{x}{x} = 1 .$$

- Try $z = iy, y \in \mathbb{R}$,

$$\lim_{z \rightarrow 0} \frac{\bar{z}}{z} = \lim_{y \rightarrow 0} \frac{\overline{iy}}{iy} = \lim_{y \rightarrow 0} \frac{-iy}{iy} = -1 .$$

Therefore, the limit does not exist.

DEFINITION 1.2.5 (Continuity at a Point).

A function f is **continuous at the point** z_0 if $\lim_{z \rightarrow z_0} f(z) = f(z_0)$.

DEFINITION 1.2.6 (Continuity on a Set).

A function is **continuous on a set** S if it is continuous at all point $z \in S$.

Example 1.2.2. Consider $f(z) = z^2$.

Let $\Delta z = z - z_0$. Then,

$$\begin{aligned}\lim_{z \rightarrow z_0} z^2 &= \lim_{\Delta z \rightarrow 0} (z_0 + \Delta z)^2 \\ &= \lim_{\Delta z \rightarrow 0} z_0^2 + 2\Delta z z_0 + \Delta z^2 = z_0^2\end{aligned}$$

So f is continuous everywhere.

DEFINITION 1.2.7 (Connectedness).

A set S is **connected** if S cannot be written as $S = S_1 \cup S_2$, where S_1, S_2 are open and $S_1 \cap S_2 = \emptyset$.

DEFINITION 1.2.8 (Path).

A **path** is the image of $[0, 1]$ under a continuous function.

DEFINITION 1.2.9 (Path-Connectedness).

A set S is **path-connected** if $\forall z_1, z_2 \in S$, there exists a path from z_1 to z_2 lying in S .

DEFINITION 1.2.10 (Domain).

A **domain** is a path-connected open set.

Remark 1.2.11. path-connected \Rightarrow connected; connected $\not\Rightarrow$ path-connected

Example 1.2.3. Consider the set $S \subseteq \mathbb{R}^2$, $S = \{(x, \sin \frac{1}{x}), x > 0\} \cup \{(0, y) : y \in \mathbb{R}\}$, this set S is connected but not path connected.

DEFINITION 1.2.12 (Derivative).

Let $f : \mathbb{C} \rightarrow \mathbb{C}$, if

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

exists, then f is **differentiable** at z_0 and that the limit is its derivative $f'(z_0)$.

Example 1.2.4. Consider $f(z) = z^2$.

$$\begin{aligned}f'(z_0) &= \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{\Delta z_0^2 + 2z_0\Delta z + \Delta z^2 - \Delta z_0^2}{\Delta z} \\ &= \lim_{\Delta z \rightarrow 0} 2z_0 + \Delta z = 2z_0 = f'(z_0) .\end{aligned}$$

Example 1.2.5. Consider $f(z) = |z|$.

$$\lim_{\Delta z_0} \frac{f(z_0) + \Delta z - f(z_0)}{\Delta z} = \lim_{\Delta z_0} \frac{|z_0 + \Delta z| - |z_0|}{\Delta z} .$$

Try $\Delta z = x \in \mathbb{R}$, let $z_0 = a + bi$, $a, b \in \mathbb{R}$, then, because let $g(x, y) = \sqrt{x^2 + y^2}$, then, $\frac{\partial g}{\partial x} = \frac{x}{\sqrt{x^2 + y^2}}$. Therefore

$$\begin{aligned} \lim_{\Delta z_0 \rightarrow 0} \frac{|z_0 + \Delta z| - |z_0|}{\Delta z} &= \lim_{x \rightarrow 0} \frac{|a + x + bi| - |a + bi|}{x} \\ &= \lim_{x \rightarrow 0} \frac{\sqrt{(a+x)^2 + b^2} - \sqrt{a^2 + b^2}}{x} = \frac{a}{a^2 + b^2} . \end{aligned}$$

Similarly, try $\Delta z = yi$, $y \in \mathbb{R}$, then,

$$\begin{aligned} \lim_{\Delta z \rightarrow 0} \frac{|z_0 + \Delta z| - |z_0|}{\Delta z} &= \lim_{yi \rightarrow 0} \frac{\sqrt{a^2 + (b+y)^2} - \sqrt{a^2 + b^2}}{yi} \\ &= \frac{1}{i} \cdot \lim_{y \rightarrow 0} \frac{\partial g}{\partial y} = \frac{1}{i} \cdot \frac{b}{\sqrt{a^2 + b^2}} . \end{aligned}$$

Thus, $f(z) = |z|$ **is differentiable nowhere**.

1.3 Derivative Continued - May 6

Proposition 1.3.1 (Derivative Operations).

- $(f + g)'(z) = f'(z) + g'(z)$
- $(cf)'(z) = cf'(z)$, where $c \in \mathbb{C}$
- $(fg)'(z) = f(z)g'(z) + f'(z)g(z)$, product rule
- $(f \circ g)'(z) = f'(g(z))g'(z)$, chain rule
- $\left(\frac{f}{g}\right)'(z) = \frac{g(z)f'(z) - f(z)g'(z)}{g^2(z)}$

DEFINITION 1.3.1 (Real and Imaginary Parts).

Let $z \in \mathbb{C}$, $z = a + bi$, $a, b \in \mathbb{R}$. Then a and b are called **the real and imaginary parts** of z respectively, denoted $\text{Re}(z)$ and $\text{Im}(z)$.

Example 1.3.1. $f(z) = \text{Re}(z)$,

$$\lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h}$$

Let $h = h_x \in \mathbb{R}$, then,

$$\lim_{h_x \rightarrow 0} \frac{\operatorname{Re}(a + h_x + bi) - \operatorname{Re}(a + bi)}{h_x} = \lim_{h_x \rightarrow 0} \frac{a + h_x - a}{h_x} = 1 .$$

Now let $h = ih_y$, $h_y \in \mathbb{R}$, then,

$$\lim_{h_y \rightarrow 0} \frac{\operatorname{Re}(a + bi + ih_y) - \operatorname{Re}(a + bi)}{ih_y} = \lim_{h_y \rightarrow 0} \frac{a - i}{ih_y} = 0 .$$

Therefore, $\operatorname{Re}(z)$ is differentiable nowhere.

And $\operatorname{Im}(z)$ is differentiable nowhere because $\operatorname{Im}(z) = \operatorname{Re}(-iz)$. Then, $\bar{z} = \operatorname{Re}(z) - i\operatorname{Im}(z)$ is differentiable nowhere.

Intuition: Differentiable functions are these that act on z and are blind to $\operatorname{Re}(z)$, $\operatorname{Im}(z)$.

Example 1.3.2. $f(z) = z^2$, let $z = a + bi$, $f(z) = a^2 + 2ab - b^2$.

Recall $z = a + bi = re^{i\theta}$, where $r = \sqrt{a^2 + b^2}$, $\theta = \arctan \frac{b}{a}$, therefore, $f(z) = r^2 e^{2i\theta}$.

DEFINITION 1.3.2 (Modulus).

The **modulus** (or magnitude, sbolute value) of z is $r = |z|$.

DEFINITION 1.3.3 (Argument).

The **argument** of z is $\theta = \arg(z)$. Note the argument of z is NOT unique to a difference of multiple of 2π .

Example 1.3.3. what is $i^{1/2}$?

$i = e^{i\frac{\pi}{2}}$, so

$$i^{1/2} = (e^{i\frac{\pi}{2}})^{1/2} = e^{i\frac{\pi}{4}} = \frac{1}{\sqrt{2}} + \frac{i}{\sqrt{2}} .$$

$i = e^{i\frac{5\pi}{2}}$, so

$$i^{1/2} = (e^{i\frac{5\pi}{2}})^{1/2} = e^{i\frac{5\pi}{4}} = -\frac{1}{\sqrt{2}} - \frac{i}{\sqrt{2}} = -e^{i\frac{\pi}{4}} .$$

Therefore, $i^{1/2} = \pm e^{i\frac{\pi}{4}}$.

Proposition 1.3.2 (Number of Roots). If $n > 0 \in \mathbb{Z}$, $z = re^{i\theta}$, then

$$\begin{aligned} z^{1/n} &= (r \exp(i\theta))^{1/n} \\ &= \left\{ r^{1/n} e^{i\frac{\theta}{n}}, r^{1/n} e^{i(\frac{\theta+2\pi}{n})} \right\} \end{aligned}$$

So any nonzero $z \in \mathbb{C}$ has exactly n distinct n -th roots, which are $r^{1/n} e^{i\frac{\theta+2\pi k}{n}}$, $0 \leq k < n$.

Example 1.3.4. $(-1)^{\frac{1}{4}} = \exp(i\frac{\pi}{4}), \exp(i\frac{3\pi}{4}), \exp(i\frac{5\pi}{4}), \exp(i\frac{7\pi}{4})$.

1.4 Holomorphic and Cauchy-Riemann Equation - May 9

DEFINITION 1.4.1 (Holomorphic).

If $f : \mathbb{C} \rightarrow \mathbb{C}$ is differentiable on a domain D , we say f is **holomorphic** on D . Also called **(complex) analytic**, regular.

Remark 1.4.2. Sloppy terminology warning: A function is said to be **holomorphic at a point** z_0 if it is holomorphic on some open set containing z_0 .

Proposition 1.4.1 (Cauchy-Riemann Equation). Let $f = u + iv : \mathbb{C} \rightarrow \mathbb{C}$ be holomorphic on a domain D , then f satisfy the following **Cauchy-Riemann(CR) Equations** on D ,

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}.$$

Proof. Let $z_0 \in D$, since f is holomorphic on D , $\lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h}$ exists.

Consider $h = h_x \in \mathbb{R}$, then

$$f'(z) = \lim_{h_x \rightarrow 0} \frac{f(z + h_x) - f(z)}{h_x}.$$

Let $z = x + iy$, then

$$f'(z) = \lim_{h_x \rightarrow 0} \frac{f(x + h_x + iy) - f(x + iy)}{h_x},$$

let $f(z) = f(x + iy) = u(x, y) + iv(x, y)$ where $u, v : \mathbb{R}^2 \rightarrow \mathbb{R}$, then

$$\begin{aligned} f'(x + iy) &= \lim_{h_x \rightarrow 0} \frac{u(x + h_x, y) + iv(x + h_x, y) - u(x, y) - iv(x, y)}{h_x} \\ &= \lim_{h_x \rightarrow 0} \frac{u(x + h_x, y) - u(x, y)}{h_x} + i \lim_{h_x \rightarrow 0} \frac{v(x + h_x, y) - v(x, y)}{h_x} \\ &= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}. \end{aligned}$$

Now consider $h = ih_y$, $h_y \in \mathbb{R}$, then

$$\begin{aligned} f'(x + iy) &= \lim_{h_y \rightarrow 0} \frac{u(x, y + h_y) + iv(x, y + h_y) - u(x, y) - iv(x, y)}{ih_y} \\ &= \frac{1}{i} \left(\lim_{h_y \rightarrow 0} \frac{u(x, y + h_y) - u(x, y)}{h_y} + \lim_{h_y \rightarrow 0} \frac{iv(x, y + h_y) - iv(x, y)}{h_y} \right) \\ &= \frac{1}{i} \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y} = \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial y}. \end{aligned}$$

Therefore,

$$\begin{aligned} f'(x + iy) &= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial y} \\ \Rightarrow \frac{\partial u}{\partial x} &= \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}. \end{aligned}$$

□

Example 1.4.1. Let f be holomorphic on a domain D and let $v(x, y) = \text{Im}(f) = xy$ on D . Find $u(x, y)$. Let $f = u + iv$, then,

$$\begin{aligned}\frac{\partial u}{\partial x} &= x = \frac{\partial v}{\partial y} \\ \frac{\partial v}{\partial x} &= y = -\frac{\partial u}{\partial y}\end{aligned}$$

Therefore, $\frac{\partial u}{\partial x} = x$, $\frac{\partial u}{\partial y} = -y$. Hence,

$$\begin{aligned}u &= \int \frac{\partial u}{\partial x} dx = \frac{1}{2}x^2 + C_1(y) \\ u &= \int \frac{\partial u}{\partial y} dy = \frac{1}{2}y^2 + C_2(x) \\ \Rightarrow \quad u(x, y) &= \frac{1}{2}x^2 - \frac{1}{2}y^2 + C.\end{aligned}$$

Therefore,

$$f(x + iy) = \left(\frac{1}{2}x^2 - \frac{1}{2}y^2 + C\right) + xyi = \frac{z^2}{2} + C.$$

Example 1.4.2. Let f be holomorphic on a domain D and let $\text{Re}(f) = x^2y$, prove such function D.N.E..

Let $z = x + iy$, $f(x + iy) = u(x, y) + iv(x, y)$, $u(x, y) = x^2y$.

$$\begin{aligned}\frac{\partial u}{\partial x} &= 2xy = \frac{\partial v}{\partial y} \quad \Rightarrow \quad v = \int 2xy dy = xy^2 + C_1(x) \\ \frac{\partial v}{\partial x} &= -x^2 = -\frac{\partial u}{\partial y} \quad \Rightarrow \quad v = \int x^2 dx = \frac{1}{3}x^3 + C_2(y).\end{aligned}$$

No such v exists because $C_2(y)$ cannot contain x .

Remark 1.4.3.

$$f'(z) = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} - i\frac{\partial u}{\partial y} = \frac{\partial v}{\partial y} + i\frac{\partial v}{\partial x} = \frac{\partial u}{\partial x} - i\frac{\partial u}{\partial y}$$

Let $f = u + iv$ be holomorphic and let $v(x, y) = xy$. Then

1.5 Holomorphic and Cauchy-Riemann Equation (Continued) - May 11

Example 1.5.1. $f(z) = e^z$,

$$f(x + iy) = e^x e^{iy} = e^x (\cos y + i \sin y) = e^x \cos y + i e^x \sin y$$

Then,

$$\begin{aligned} \frac{\partial u}{\partial x} &= e^x \cos y = \frac{\partial v}{\partial y} \\ \frac{\partial u}{\partial y} &= -e^x \sin y = -\frac{\partial v}{\partial x} \end{aligned}$$

THEOREM 1.5.1 (Cauchy-Riemann Equation and Holomorphicity).

Let $u, v : \mathbb{R}^2 \rightarrow \mathbb{R}$, have continuous partial derivatives at (x_0, y_0) which satisfy

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x},$$

then $f(x + iy) = u(x, y) + iv(x, y)$ is holomorphic at $z_0 = x_0 + iy_0$.

Proof. Let $D \subseteq \mathbb{C}$ be an open set with $z_0 \in D$, let $z = x + iy \in D$. Then,

$$u(x, y) = u(x_0, y_0) + (x - x_0) \left(\frac{\partial u}{\partial x} + \varepsilon_1(x, y) \right) + (y - y_0) \left(\frac{\partial u}{\partial y} + \varepsilon_2(x, y) \right),$$

where $\varepsilon_1, \varepsilon_2$ are continuous at (x_0, y_0) and $\varepsilon_1(x_0, y_0) = \varepsilon_2(x_0, y_0) = 0$. And,

$$v(x, y) = v(x_0, y_0) + (x - x_0) \left(\frac{\partial v}{\partial x} + \varepsilon_3(x, y) \right) + (y - y_0) \left(\frac{\partial v}{\partial y} + \varepsilon_4(x, y) \right),$$

So

$$f(x + iy) = u(x, y) + iv(x, y) = f(z_0) + (z - z_0) \left(\frac{\partial u}{\partial x}(x_0, y_0) + i \frac{\partial v}{\partial x}(x_0, y_0) + \varepsilon(z) \right),$$

where

$$\varepsilon(z) = \frac{x - x_0}{z - z_0} (\varepsilon_1 + i\varepsilon_3) + \frac{y - y_0}{z - z_0} (\varepsilon_2 + i\varepsilon_4).$$

satisfying ε continuous at z_0 , $\varepsilon(z_0) \rightarrow 0$. So

$$\begin{aligned} \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} &= \lim_{z \rightarrow z_0} \frac{\partial u}{\partial x}(x_0, y_0) + i \frac{\partial v}{\partial x}(x_0, y_0) + \varepsilon(z) \\ &= \frac{\partial u}{\partial x}(x_0, y_0) + i \frac{\partial v}{\partial x}(x_0, y_0) \\ &= f'(z_0) \end{aligned}$$

exists and thus. □

Remark 1.5.2. We now have a good test for holomorphicity.

Example 1.5.2. Consider $f(z) = \bar{z}$.

$f(x + iy) = x - iy$, $u = x$, $v = -y$. then

$$\frac{\partial u}{\partial x} = 1 \neq \frac{\partial v}{\partial y} = -1 \quad \frac{\partial u}{\partial y} = 0 \neq -\frac{\partial v}{\partial x} = 1.$$

Therefore, $f(z)$ is nowhere holomorphic at complex plane.

Example 1.5.3. Consider $f(z) = \frac{1}{z}$.

Method 1:

$$f(x + iy) = \frac{1}{x + iy} = \frac{x - iy}{x^2 + y^2} = \frac{x}{x^2 + y^2} + i \frac{-y}{x^2 + y^2}$$

Therefore, $u = \frac{x}{x^2 + y^2}$, $v = \frac{-y}{x^2 + y^2}$.

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{(x^2 + y^2) - x(2x)}{(x^2 + y^2)^2} = \frac{y^2 - x^2}{(x^2 + y^2)^2} \\ \frac{\partial v}{\partial y} &= \frac{y^2 - x^2}{(x^2 + y^2)^2} . \end{aligned}$$

And

$$\frac{\partial u}{\partial y} = \frac{-2xy}{(x^2 + y^2)^2} = -\frac{\partial v}{\partial x} .$$

So f is holomorphic on $\mathbb{C} \setminus \{0\}$.

Method 2: $f(z) = \frac{1}{re^{i\theta}} = \frac{1}{r}e^{-r\theta}$ is well-behaved.

Lemma 1.5.1. If f, g are holomorphic at z_0 and $g(z_0) \neq 0$, then $\frac{f}{g}$ is holomorphic at z_0 .

However, the converse is false, that is if $\frac{f}{g}$ is holomorphic at z_0 , it does not imply that f, g are holomorphic at z_0 .

Proposition 1.5.1 (Polar Form of Cauchy-Riemann Equation). Let $z = re^{i\theta}$, $z_0 = r_0e^{i\theta_0}$, let $f(z)$ be holomorphic, and let $f(z) = u(r, \theta) + iv(r, \theta)$, we have

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} .$$

First, fix $\theta = \theta_0$ and let $r \rightarrow r_0$, then

$$\begin{aligned} f'(r_0e^{i\theta_0}) &= \lim_{r \rightarrow r_0} \frac{u(r, \theta_0) + iv(r, \theta_0) - (u(r_0, \theta_0) + iv(r_0, \theta_0))}{re^{i\theta_0} - r_0e^{i\theta_0}} \\ &= e^{-r\theta_0} \left(\frac{\partial u}{\partial r}(r_0, \theta_0) + i \frac{\partial v}{\partial r}(r_0, \theta_0) \right) \end{aligned}$$

Next let $r = r_0$, $\theta \rightarrow \theta_0$, then,

$$\begin{aligned}
f'(r, e^{i\theta_0}) &= \lim_{\theta \rightarrow \theta_0} \frac{u(r_0, \theta) + iv(r_0, \theta) - (u(r_0, \theta_0) + iv(r_0, \theta_0))}{r_0 e^{i\theta} - r_0 e^{i\theta_0}} \\
&= \frac{1}{r_0} \lim_{\theta \rightarrow \theta_0} \left[\frac{u(r_0, \theta) - u(r_0, \theta_0)}{\theta - \theta_0} + i \frac{v(r_0, \theta) - v(r_0, \theta_0)}{\theta - \theta_0} \right] \cdot \frac{\theta - \theta_0}{e^{i\theta} - e^{i\theta_0}} \\
&= \frac{1}{r} \frac{1}{ie^\theta} \left(\frac{\partial u}{\partial \theta} + i \frac{\partial v}{\partial \theta} \right) \\
&= e^{-i\theta} \left(\frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r} \right)
\end{aligned}$$

Therefore,

$$\begin{aligned}
\frac{\partial u}{\partial r} &= \frac{1}{r} \frac{\partial v}{\partial \theta} \\
\frac{\partial v}{\partial r} &= -\frac{1}{r} \frac{\partial u}{\partial \theta}
\end{aligned}$$

1.6 Branch - May 13

Remark 1.6.1. Assuming all partial derivatives are continuous and $z = re^{i\theta}$,

$$\begin{aligned}
f &= u + iv \text{ is holomorphic on a domain } D \\
\iff \frac{\partial u}{\partial x} &= \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \text{ on } D \\
\iff \frac{\partial u}{\partial r} &= \frac{1}{r} \frac{\partial v}{\partial \theta} \text{ and } \frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta} \text{ on } D
\end{aligned}$$

Remark 1.6.2. Recall $z^{1/n} = (re^{i\theta})^{1/n} = r^{1/n} e^{i\frac{\theta+2\pi m}{n}}$, $0 \leq m < n$. Consider $z^{1/n} = r^{1/n} e^{i\frac{\theta}{n}} = r^{1/n} \cos \frac{\theta}{n} + ir^{1/n} \sin \frac{\theta}{n}$. Then let

$$u(r, \theta) = r^{1/n} \cos \frac{\theta}{n} \quad v(r, \theta) = r^{1/n} \sin \frac{\theta}{n},$$

$$\begin{aligned}
\frac{\partial u}{\partial r} &= \frac{1}{n} r^{\frac{1}{n}-1} \cos \frac{\theta}{n} = \frac{1}{r} \frac{\partial v}{\partial \theta} \\
\frac{\partial v}{\partial r} &= \frac{1}{n} r^{\frac{1}{n}-1} \sin \frac{\theta}{n} = -\frac{1}{r} \frac{\partial u}{\partial \theta}.
\end{aligned}$$

This function is not continuous along $\text{Arg}(z) = \pi$. This function is holomorphic along $\mathbb{C} \setminus \mathbb{R}_{\leq 0}$ (Note 0 is included). This is called the **branch cut**.

We can put the branch cut somewhere else, i.e.

$$f(re^{i\theta}) = \theta, 0 \leq \theta < 2\pi$$

then, f is not continuous at $\text{Arg}(z) = 2\pi$.

DEFINITION 1.6.3.

Define $\arg(re^{i\theta}) = \theta + 2\pi m$, $m \in \mathbb{Z}$, and $\text{Arg}(re^{i\theta}) = \theta$, $-\pi < \theta \leq \pi$.

This is a **branch** of the multivalued function \arg . In particular Arg is called the **principle branch**.

$\text{Arg}(z)$ is not continuous at $\text{Arg}(z) = \pi$.

Remark 1.6.4. Other branches:

- $f(re^{i\theta}) = \theta$, $\pi < \theta \leq 3\pi = \text{Arg}(re^{i\theta}) + 2\pi$
- $\text{Arg}(z) + 2\pi n$, $n \in \mathbb{Z}$ is a branch of $\arg(z)$.

DEFINITION 1.6.5 (Logarithm).

For $z \neq 0 \in \mathbb{C}$ define

$$\begin{aligned}\log(z) &= \log(re^{i\theta}) = \log(r) + \log(e^{i\theta}) \\ &= \log r + i\theta \\ &= \ln r + i\theta \\ &= \ln |z| + i\theta + 2\pi im, m \in \mathbb{Z}.\end{aligned}$$

Note \log is a multivalued function.

Notation 1.6.6. Use \log for the multivalued function and Log for the principal branch:

$$\text{Log}(z) = \ln |z| + i\text{Arg}(z) .$$

$$u = \ln r, v = \theta,$$

$$\frac{\partial u}{\partial r} = \frac{1}{r} = \frac{1}{r} \frac{\partial v}{\partial \theta} \quad \frac{\partial v}{\partial r} = 0 = -\frac{1}{r} \frac{\partial u}{\partial \theta} .$$

Example 1.6.1. $-\pi < \theta \leq \pi$

$$\begin{aligned}\text{Log}(z^{1/2}) &= \text{Log}((re^{i\theta})^{1/2}) \\ &= \text{Log}(r^{1/2} e^{i\frac{\theta}{2}}) \\ &= \ln |r^{1/2}| + i\frac{\theta}{2} \\ &= \frac{1}{2}(\ln |r| + i\theta) = \frac{1}{2}\text{Log}(z) .\end{aligned}$$

Remark 1.6.7. Note that $\text{Log}(wz) \neq \text{Log}(z) + \text{Log}(w)$ in general.

1.7 Trigonometric Functions and Power - May 16

DEFINITION 1.7.1 (Trigonometric Functions on Complex Plane).

Let $z \in \mathbb{C}$,

$$\cos z = \frac{e^{iz} + e^{-iz}}{2} \quad \text{and} \quad \sin z = \frac{e^{iz} - e^{-iz}}{2i}.$$

\sin and \cos are immediately holomorphic in the entire \mathbb{C} , since the exponential functions are.

Remark 1.7.2. If two holomorphic functions are equal on "enough" of a set, they must agree on their domains.

Lemma 1.7.1 (Derivatives of Trig Functions).

$$\begin{aligned} \frac{d}{dz} \sin(z) &= \frac{d}{dz} \left(\frac{e^{iz} - e^{-iz}}{2i} \right) \\ &= \frac{ie^{iz} + ie^{-iz}}{2i} = \frac{e^{iz} + e^{-iz}}{2} = \cos z \\ \frac{d}{dz} \cos(z) &= \frac{d}{dz} \left(\frac{e^{iz} + e^{-iz}}{2i} \right) \\ &= \frac{ie^{iz} - ie^{-iz}}{2} = \frac{-e^{iz} + e^{-iz}}{2i} = -\sin z. \end{aligned}$$

Remark 1.7.3 (Trig Functions are unbounded in \mathbb{C}). All the trig identities in \mathbb{R} carry over in the obvious way to \mathbb{C} . However, \cos , \sin have properties that are NOT true in \mathbb{R} .

Consider $\cos(iy)$,

$$|\cos(iy)| = \left| \frac{e^{i(iy)} + e^{-i(iy)}}{2} \right| = \left| \frac{e^{-y} + e^y}{2} \right|,$$

then, $|\cos(iy)| \rightarrow \infty$ as $y \rightarrow \infty$ or $y \rightarrow -\infty$. Hence, $\sin z$, $\cos z$ are NOT bounded functions on \mathbb{C} .

DEFINITION 1.7.4 (Hyperbolic Functions).

Let $z \in \mathbb{C}$,

$$\cosh z = \frac{e^z + e^{-z}}{2} \quad \text{and} \quad \sinh z = \frac{e^z - e^{-z}}{2}.$$

Then,

$$\sinh(iz) = i \sin z \quad \text{and} \quad \cosh(iz) = \cos z$$

Example 1.7.1. Consider i^i .

$$i^i = (e^{\log i})^i = e^{i \log i} = e^{i(\frac{\pi i}{2} + 2\pi i k)} = e^{-\frac{\pi}{2} - 2\pi k}, k \in \mathbb{Z}.$$

Example 1.7.2.

$$\log i = \log e^{i\frac{\pi}{2}} = -\frac{i\pi}{2} + 2\pi ik, k \in \mathbb{Z}$$

DEFINITION 1.7.5 (Derivative of Power).

Define $z^w = e^{w \log z}$, then its derivative is

$$\frac{d}{dz}(z^w) = \frac{d}{dz}(e^{w \log z}) = e^{w \log z} \frac{d}{dz}(w \log z) = w \frac{1}{z} e^{w \log z} = \frac{w}{z} z^w = w z^{w-1}.$$

Remark 1.7.6. How many values does z^w have?

$$z^w = e^{w \log z} = e^{w(\log z + 2\pi ik)} = e^{w \log z} e^{2\pi i k w}$$

When is $e^{2\pi i k w} = e^{2\pi i n w}$, $n, w \in \mathbb{Z}, n \neq k$?

If $e^{2\pi i k w} = e^{2\pi i n w}$, these are equal when

$$w\pi i k w = 2\pi i n w + 2\pi i m, m \in \mathbb{Z}.$$

$k w = n w + m$, $m \in \mathbb{Z}, w = \frac{m}{k-n}$, $m, n, k \in \mathbb{Z}$. Thus the powers z^w repeat if and only if $w \in \mathbb{Q}$. Then if $w = \frac{p}{q}$, $z^w = z^{p/q} = (z^p)^{1/q}$ has q distinct values.

Proposition 1.7.1. If $z \neq 0$,

$$z^w \text{ take on } \begin{cases} 1, & \text{if } w \in \mathbb{Z} \\ q, & \text{if } w = \frac{p}{q} \in \mathbb{Q} \\ \infty, & \text{otherwise} \end{cases}.$$

1.8 Rotalion Approximation - May 18

Algorithm 1.8.1 (Rotalion Approximation). Let f be holomorphic at z_0 , we have

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}.$$

The modulus and argument must converge individually:

•

$$|f'(z_0)| = \lim_{z \rightarrow z_0} \left| \frac{f(z) - f(z_0)}{z - z_0} \right|$$

So near z_0 ,

$$|f(z) - f(z_0)| \approx |f'(z_0)| \cdot |z - z_0|.$$

- and

$$\arg(f'(z_0)) = \lim_{z \rightarrow z_0} \arg\left(\frac{f(z) - f(z_0)}{z - z_0}\right).$$

for some branch of \arg holomorphic near $z_0, f(z_0), f'(z_0)$.

$$\begin{aligned} \arg(f'(z)) &= \arg(f(z) - f(z_0)) - \arg(z - z_0) \\ \Rightarrow \arg(f(z) - f(z_0)) &\approx \arg(f'(z_0)) + \arg(z - z_0). \end{aligned}$$

So near z_0 we have

$$f(z) \approx f(z_0) + e^{i \arg(f'(z_0))} |f'(z_0)| (z - z_0).$$

This is a rotation of $z - z_0$ by $\arg(f'(z_0))$ and a scaling by $|f'(z_0)|$.

Example 1.8.1. Consider $f(z) = z^2, f(re^{i\theta}) = r^2 e^{i2\theta}$.

Let $z_0 = 1 + i = \sqrt{2}e^{i\frac{\pi}{4}}, f(z_0) = z_0^2 = 2i, f'(z) = 2z, f'(z_0) = 2(1 + i) = 2\sqrt{2}e^{i\frac{\pi}{4}}$.

So for small $h = z - z_0$,

$$f(z_0 + h) \approx f(z_0) + e^{i \arg f'(z_0)}$$

THEOREM 1.8.1.

If f is holomorphic on a domain D and $f'(z) = 0$ for all $z \in D$, then f is constant on D .

Proof.

$$f'(z) = 0 = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial y}$$

Then, all partial derivatives are 0:

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial y} = \frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} = 0.$$

Therefore, u, v are constant on any horizontal or vertical line segment in D . But D is domain, so D is open and path-connected. Then any two points in D can be connected by a path of horizontal and vertical segments. So u and v are constant on D so f . Therefore, $f = u + iv$ is constant on D . \square

Example 1.8.2. Find a branch of $(z^2 - 1)^{1/2}$ holomorphic on $|z| > 1$.

Note that the principal branch of $z^{1/2}$ does not work: $e^{1/2} \log(z^2 - 1)$.

Its branch cut is where $z^2 - 1 \in \mathbb{R}, z^2 - 1 \leq 0$. But let $z = 2i, z^2 - 1 = -4 - 1 = -5 \leq 0$.

Find $f(z)$ holomorphic on $|z| > 1$ such that $f(z)^2 = z^2 - 1$.

Consider the principal branch of $f(z) = z(1 - \frac{1}{z^2})^{1/2}$.

Its branch cut lies wherever $1 - \frac{1}{z^2} \leq 0$ in \mathbb{R} , which is $\frac{1}{z^2} \geq 1$ in $\mathbb{R} \Rightarrow z^2 \leq 1$ in \mathbb{R} , so $|z| < 1$.

2 Integration

2.1 Integrability - May 20

DEFINITION 2.1.1 (Smooth Curve).

A **smooth curve** in \mathbb{C} is the image of a function $\gamma : [a, b] \rightarrow \mathbb{C}$ satisfying

- γ is continuously differentiable on $[a, b]$
- $\gamma' \neq 0$ on $[a, b]$
- γ is one to one

This definition rules out gaps, sharp corners, pausing(temporarily stopping at a point), retracing, self-intersection.

DEFINITION 2.1.2 (Directed Smooth Curve).

A **directed smooth curve** is a smooth curve with a fixed direction, *i.e.* the points on the curve are ordered and any γ must trace them in order.

DEFINITION 2.1.3 (Contour).

A **contour** is a directed piecewise smooth curve. *i.e.* $\Gamma = C_1 \cup C_2 \cup \dots \cup C_n$, where the C_j are directed small curves and the terminal point of C_j is the initial point of C_{j+1} .

DEFINITION 2.1.4 (Simple Contour).

A contour is **simple** if it has no self-intersections.

DEFINITION 2.1.5 (Closed Contour).

A contour is **closed** if its initial point coincides with its terminal point.

DEFINITION 2.1.6 (Simple Closed Contour).

A **simple closed contour** is a contour both simple and closed.

Example 2.1.1. $\Gamma : r_1(t) = z_0 t + z_1(1 - t), t \in [0, 1], z_0, z_1 \in \mathbb{C}$.

Note parametrizations are NOT unique, e.g. $r_2(t) = z_0(2t) + z_1(1 - 2t), t \in [0, \frac{1}{2}]$, $r_3(t) = z_0 t^2 + z_1(1 - t^2), t \in [0, 1]$.

DEFINITION 2.1.7 (Circular Contour).

$C_r(z_0)$ is the circular contour with radius r and center z_0 , traversed counterclockwise.

Example 2.1.2. $\Gamma = C_1 \cup C_2 \cup C_3$

$$C_1 : \gamma_1(t) = t, t \in [0, 1]$$

$$C_2 : \gamma_2(t) = ti + (1 - t), t \in [0, 1]$$

$$C_3 : \gamma_3(t) = (1 - t)i, t \in [0, 1]$$

$$\Rightarrow \gamma(t) = \begin{cases} t, & t \in [0, 1] \end{cases}$$

THEOREM 2.1.8 (Jordan Curve Theorem).

A simple closed contour divides \mathbb{C} into two disjoint regions, a bounded interior and an unbounded exterior.

DEFINITION 2.1.9 (Orientation).

A simple closed contour is **positively oriented** if its interior is to the left when traverse, **negatively oriented** otherwise.

DEFINITION 2.1.10 (Partition).

Let Γ be a directed smooth curve with initial point w_0 and terminal point w_1 , a **partition** of Γ is a set of points $Z_0 = w_0, z_1, z_2, \dots, z_n = w_1$ such that $\forall j, 0 \leq j < n, z_{j+1}$ is farther along Γ than z_j .

DEFINITION 2.1.11 (Mesh).

The **mesh** of a partition is the largest distance between two consecutive points z_j, z_{j+1} along Γ .

DEFINITION 2.1.12 (Riemann Sum).

Let Γ lie in a domain D and let $f : D \rightarrow \mathbb{C}$. The **Riemann sum** of f with respect to P_n is

$$S_f(P_n) = f(z_1)(z_1 - z_0) + f(z_2)(z_2 - z_0) + \dots + f(z_n)(z_n - z_{n-1}) .$$

DEFINITION 2.1.13 (Integrable).

f is integrable along Γ if

$$\lim_{\text{mesh}(P_n) \rightarrow 0} S(P_n) \text{ exists .}$$

DEFINITION 2.1.14.

If f is integrable along Γ , the integrable of f along Γ with partition $P_n = z_0, z_1, \dots, z_n$ is

$$\int_{\Gamma} f = \lim_{\text{mesh}(P_n) \rightarrow 0} S(P_n) = \lim_{\text{mesh}(P_n) \rightarrow 0} \sum_{j=0}^{n-1} f(z_j)(z_{j+1} - z_j)$$

Remark 2.1.15. Note in the above formula, we do not reference a parametrization of Γ and therefore, the integral of f is independent of any parametrization.

2.2 May 25

DEFINITION 2.2.1 (Integral).

Let Γ be a directed smooth curve, let $P_n = z_0, z_1, \dots, z_n$ be a partition of Γ and let

$$\int_{\Gamma} f(z) dz = \lim_{\text{mesh}(P_n) \rightarrow 0} \sum_{j=0}^{n-1} f(z_j) (z_{j+1} - z_j) .$$

Now let Γ be parametrized by $\gamma : [a, b] \rightarrow \Gamma$ and let t_0, t_1, \dots, t_n be a partition of $[a, b]$ such that $\gamma(t_j) = z_j, 0 \leq j \leq n$, then

$$\begin{aligned} \lim_{\text{mesh}(P_n) \rightarrow 0} \sum_{j=0}^{n-1} f(z_j) \Delta z_j &= \lim_{\Delta t_j \rightarrow 0, 0 \leq j < n} \sum_{j=0}^{n-1} f(\gamma(t_j)) \Delta z_j , \\ &\quad \text{where } \Delta t_j = t_{j+1} - t_j, \Delta z_j = z_{j+1} - z_j \\ &= \lim_{\Delta t_j \rightarrow 0} \sum_{j=0}^{n-1} f(\gamma(t_j)) \gamma'(t_j) \Delta t_j \\ &= \int_a^b f(\gamma(t)) \gamma'(t) dt \\ \Rightarrow \int_{\Gamma} f(z) dz &= \int_a^b f(\gamma(t)) \gamma'(t) dt . \end{aligned}$$

★ Integral of parametrization

DEFINITION 2.2.2.

Define the integral over a contour $\Gamma = \Gamma_1 + \Gamma_2 + \dots + \Gamma_n$, where Γ_j are smooth directed curves, to be

$$\int_{\Gamma} f = \int_{\Gamma_1} f + \dots + \int_{\Gamma_n} f .$$

We immediately have

- $$\int_{\Gamma} f + g = \int_{\Gamma} f + \int_{\Gamma} g .$$

- $$\int_{\Gamma_1 + \Gamma_2} f = \int_{\Gamma_1} f + \int_{\Gamma_2} f$$

- For $c \in \mathbb{C}$,

$$\int_{\Gamma} cf = c \int_{\Gamma} f$$

Example 2.2.1. $\int_{\Gamma} z^2 dz, \Gamma = \gamma(t) = e^{it}, 0 \leq t \leq \pi$.

$$\int_{\Gamma} z^2 dz = \int_0^{\pi} (e^{it})^2 (ie^{it}) dt = i \int_0^{\pi} e^{3it} dt = \frac{1}{3} (e^{3i\pi} - e^{3i(0)}) = -\frac{2}{3} .$$

Example 2.2.2. Let $C_1(0) = r(t) = e^{it}$, $0 \leq t \leq 2\pi$, then

$$\int_{C_1(0)} z dz = \int_0^{2\pi} e^{it} (ie^{it}) dt = i \int_0^{2\pi} e^{2it} dt = \frac{e^{2it}}{2} \Big|_0^{2\pi} = \frac{e^{4\pi i} - 1}{2} = 0 .$$

and

$$\int_{C_1(0)} \frac{1}{z} dz = \int_0^{2\pi} \frac{ie^{it}}{e^{it}} dt = 2\pi i .$$

DEFINITION 2.2.3 (Length).

The length of a contour Γ is parametrized by $\gamma : [a, b] \rightarrow \Gamma$, $\int_a^b |\gamma'(t)| dt$.

2.3 May 27

THEOREM 2.3.1 (ML bound).

Let f be integrable on Γ and let $|f(z)| \leq M$, $\forall z \in \Gamma$,

$$\begin{aligned} \left| \int_{\Gamma} f(z) dz \right| &= \left| \int_a^b f(\gamma(t)) \gamma'(t) dt \right| \\ &\leq \int_a^b |f(\gamma(t)) \gamma'(t)| dt \\ &= \int_a^b |f(\gamma(t))| |\gamma'(t)| dt \\ &\leq \int_a^b M |\gamma'(t)| dt \\ &= M \cdot \text{length}(\Gamma) . \end{aligned}$$

This is often called the ML bound.

DEFINITION 2.3.2 (Primitive).

A function F is a **primitive** (or **antiderivative**) for a function f on a domain D if F is holomorphic on D and $\forall z \in D$, $F'(z) = f(z)$.

DEFINITION 2.3.3.

Let f have a primitive F on D and let Γ lie on D . Consider

$$\int_{\Gamma} f(z) dz = \int_{\Gamma} F'(z) dz .$$

Let $\gamma : [a, b] \rightarrow \Gamma$ parametrize Γ . Then

$$\int_{\Gamma} F'(z) dz = \int_a^b F'(\gamma(t)) \gamma'(t) dt = \int_a^b \frac{dF}{d\gamma}(\gamma(t)) \frac{d\gamma}{dt} dt = \int_{\gamma(a)}^{\gamma(b)} \frac{dF}{d\gamma} d\gamma = F(\gamma(b)) - F(\gamma(a)) .$$

By the Fundamental Theorem of Calculus on \mathbb{R} .

THEOREM 2.3.4 (Fundamental Theorem of Calculus in \mathbb{C}).

If f has a primitive F on a domain in D and Γ lies in D with initial point z_0 and terminal point z_1 ,

$$\int_{\Gamma} f(z)dz = F(z_1) - F(z_0) .$$

Example 2.3.1. $f(z) = z$ has a primitive $F(z) = \frac{1}{2}z^2$ on all of \mathbb{C} . So for Γ running from z_0 to z_1 ,

$$\int_{\Gamma} z dz = \frac{1}{2}z^2 \Big|_{z_0}^{z_1} .$$

Example 2.3.2. $f(z) = \frac{1}{z}$ has primitive $\log z$. i.e. any branch of $\log z$ is a primitive of $\frac{1}{z}$ and its domain.

Corollary 2.3.1. If f has a primitive on a domain D and Γ is a closed contour lying in D ,

$$\int_{\Gamma} f(z)dz = 0 .$$

Proof.

$$\int_{\Gamma} f = F(z_1) - F(z_0) = F(z_0) - F(z_0) = 0 .$$

□

Lemma 2.3.1. Let f be continuous on a domain D and let $\oint_{\Gamma} f = 0$ for any closed Γ lying in D . Then given Γ_1, Γ_2 in D with the same initial and terminal points,

$$\int_{\Gamma_1} f = \int_{\Gamma_2} f .$$

Proof. Note that $\Gamma_1 + (-\Gamma_2)$ is closed, so

$$\int_{\Gamma_1 + (-\Gamma_2)} f = 0 = \int_{\Gamma_1} f - \int_{\Gamma_2} f = 0 .$$

□

Lemma 2.3.2. Let f be continuous on a domain D such that for Γ_1, Γ_2 in D sharing initial and terminal points,

$$\int_{\Gamma_1} f = \int_{\Gamma_2} f .$$

Then f has a primitive on D .

Remark 2.3.5. ★ important

THEOREM 2.3.6.

Let f be continuous on a domain D TFAE

1. f has a primitive on D
2. for all closed contours Γ lying in D , $\int_{\Gamma} f = 0$.
3. for any two contours Γ_1, Γ_2 in D sharing initial and terminal points

$$\int_{\Gamma_1} f = \int_{\Gamma_2} f .$$

2.4 May 30**DEFINITION 2.4.1.**

A **Cauchy sequence** is a sequence $\{z_n\}_{n=1}^{\infty}$ s.t. $\forall \varepsilon > 0, \exists N > 0 \forall n_1, n_2 > N, |z_{n_1} - z_{n_2}| < \varepsilon$.

Lemma 2.4.1. Closed bounded subsets of \mathbb{R}^n are compact.

Lemma 2.4.2 (Cauchy sequence convergence). A Cauchy sequence in a compact set $S \subseteq \mathbb{R}$ converges to a point in S .

Lemma 2.4.3. Let f be holomorphic at z_0 , then

$$f(z) = f(z_0) + f'(z_0)(z - z_0) + \varepsilon(z)(z - z_0) ,$$

for some $\varepsilon(z)$ satisfying $\lim_{z \rightarrow z_0} \varepsilon(z) = 0$.

Proof. Let $\varepsilon(z) = \frac{f(z) - f(z_0)}{z - z_0} - f'(z_0)$. Take $\lim_{z \rightarrow z_0}$. □

THEOREM 2.4.2 (Goursat's Theorem).

Let f be holomorphic on a domain D and let T be a triangle lying in D with interior in D . Then

$$\int_T f(z) dz = 0 .$$

Proof. Divide T into four triangles by connecting the midpoints of its side, now

$$\int_T f = \int_{T_{(1)}} f + \int_{T_{(2)}} f + \int_{T_{(3)}} f + \int_{T_{(4)}} f .$$

There exists a T_1 such that $|\int_T f| \leq 4 |\int_{T_1} f|$. Call it T_1 . Note that $\text{length}(T_1) \leq \frac{1}{2} \text{length}(T)$, hence

$$\text{diam}(T_1) \leq \frac{1}{2} \text{diam}(T) .$$

Repeat this process, yielding $T = T_0, T_1, T_2, T_3, T_4, \dots$,

$$\left| \int_T f \right| \leq 4^n \left| \int_{T_n} f \right| ,$$

then, $\text{length}(T_n) \leq \frac{1}{2^n} \text{length}(T)$ and $\text{diam}(T_n) \leq \frac{1}{2^n} \text{diam}(T)$.

Let z_n be a point in the interior of T_n for each n . Then, $\{z_n\}$ is a Cauchy sequence and $\lim_{n \rightarrow \infty} z_n = w$, where w is in the interior of the triangle.

f is holomorphic at w , so

$$f(z) = f(w) + f'(w)(z - w) + \varepsilon(z)(z - w)$$

, where $\lim_{z \rightarrow w} \varepsilon(z) = 0$. Now consider

$$\int_{T_n} f(z) dz = \int_{T_n} f(w) + f'(w)(z - w) + \varepsilon(z)(z - w) dz .$$

$f(w)$ has primitive $zf(w)$, $f'(w)(z - w)$ has primitive $\frac{1}{2}f'(w)(z - w)^2$ so

$$\int_{T_n} f(w) + f'(w)(z - w) dz = 0 .$$

therefore,

$$\int_{T_n} f(z) dz = \int_{T_n} f(w) + f'(w)(z - w) + \varepsilon(z)(z - w) dz = \int_{T_n} \varepsilon(z)(z - w) dz .$$

Let $\varepsilon_n = \sup_{z \in T_n} |\varepsilon(z)|$, $|z - w| \leq \text{diam}(T_n) \leq \frac{1}{2^n} \text{diam}(T)$. $\text{length}(T_n) \leq \frac{1}{2^n} \text{length}(T)$. So

$$\left| \int_{T_n} f(z) dz \right| = \left| \int_{T_n} \varepsilon(z)(z - w) dz \right| \leq \varepsilon_n \text{diam}(T_n) \text{length}(T_n) \leq \varepsilon_n \frac{1}{4^n} \text{diam}(T) \text{length}(T) ,$$

thus

$$\left| \int_T f(z) dz \right| \leq 4^n \left| \int_{T_n} f(z) dz \right| \leq 4^n \varepsilon_n \frac{1}{4^n} \text{diam}(T) \text{length}(T) = \varepsilon_n \text{diam}(T) \text{length}(T) .$$

Let $n \rightarrow \infty$ and $\varepsilon_n \rightarrow 0$,

$$\left| \int_{T_n} f(z) dz \right| = 0 \Rightarrow \int_{T_n} f(z) dz = 0 .$$

□

Corollary 2.4.1. If f is holomorphic on an open disk, f has primitive on that disk.

Proof. Choose $z_0 \in D$, define $F(z) = \int_{\Gamma} f(z) dz$. Then

$$F(z + h) - F(z) = \int_{\Gamma_n} f(z) dz + \int_{\Delta} f + \int_{\square} f = \int_{\Gamma_n} f(z) dz$$

so that

$$\frac{d}{dz} \int_{\Gamma_n} f(z) dz = f(z)$$

as in last lecture.

□

2.5 June 1

THEOREM 2.5.1 (Goursat's Theorem).

If f is holomorphic on and inside a polygon triangle T , then $\oint_T f = 0$.

THEOREM 2.5.2 (Cauchy's Theorem).

On an open disk f holomorphic on an open disk D , let $\Gamma \subseteq D$ be a closed contour, then

$$\oint_{\Gamma} f = 0 .$$

Example 2.5.1. $f(z) = \frac{1}{z}$ is holomorphic on $0 < |z|$ but has no primitive and

$$\int_{C_1(0)} \frac{1}{z} dz = 2\pi i .$$

DEFINITION 2.5.3 (Homotopic).

Let Γ_1, Γ_2 be two contours in a domain in D with the same initial and terminal point. Γ_0 is **homotopic** (or **continuously deformable**) to Γ_1 if there exists $\gamma : [0, 1]^2 \rightarrow D$ satisfying

- γ is continuous on $[0, 1]^2$
- for a fixed s , $\gamma(s, t)$ is a parametrization of a contour in D with initial and terminal point shared with Γ_1, Γ_2 .
- $\gamma(0, t)$ parametrizes Γ_0
- $\gamma(1, t)$ parametrizes Γ_1

DEFINITION 2.5.4 (Simply Connected).

A domain D is **simply connected** if any two contours in D sharing initial and terminal points are homotopic to each other.

THEOREM 2.5.5 (Cauchy's Theorem).

Let f be holomorphic on a simply connected domain D , and let Γ be a closed contour in D . Then $\int_{\Gamma} f = 0$.

Proof. Γ is homotopic on a triangle. □

Example 2.5.2. $\int_{\Gamma} z^2 dz$, then z^2 is entire, so by Cauchy's Theorem,

$$\int_{\Gamma} z^2 dz = \int_{z^2} dz = \int_1^{-1} x^2 dx = -\frac{2}{3} .$$

Example 2.5.3.

$$\int_{C_2(0)} \frac{1}{x^2 - 1} dz$$

$\frac{1}{z^2 - 1}$ is holomorphic on $\mathbb{C} \setminus \{1, -1\}$. Then

$$\int_{C_2(0)} \frac{1}{z^2 - 1} dz = \int_{C_\varepsilon(-1)} \frac{1}{z^2 - 1} dz + \int_{C_\varepsilon(1)} \frac{1}{z^2 - 1} dz, \text{ for } 0 < \varepsilon < 2.$$

2.6 Cauchy Integral Formula - June 3**THEOREM 2.6.1.**

$f(z)$ holomorphic on a shaded region plus a bit, then

$$\int_{\Gamma_1} F(z) dz = \int_{\Gamma_2} F(z) dz$$

where Γ_1, Γ_2 have standard anticlockwise orientations.

THEOREM 2.6.2 (Cauchy Integral Formula).

Let f be a function holomorphic on a domain $D \subseteq \mathbb{C}$, and Γ a Jordan curve contained in D and whose interior is contained in D . Let $z_0 \in \text{interior of } \Gamma$. Then

$$f(z_0) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(z)}{z - z_0} dz.$$

Proof. We can replace Γ with $C(r) = \{|z - z_0| = r\}$ for small enough r . Then

$$\begin{aligned} \frac{1}{2\pi i} \int_{\Gamma} \frac{f(z)}{z - z_0} dz &= \frac{1}{2\pi i} \int_{C(r)} \frac{f(z)}{z - z_0} dz \\ &= \frac{1}{2\pi i} \left(\int_{C(r)} \frac{f(z_0)}{z - z_0} dz + \int_{C(r)} \frac{f(z) - f(z_0)}{z - z_0} dz \right) \\ &= f(z_0) + \int_{C(r)} \frac{f(z) - f(z_0)}{z - z_0} dz \end{aligned}$$

Then, we claim that

$$\int_{C(r)} \frac{f(z) - f(z_0)}{z - z_0} dz = 0.$$

Since $f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$ exists,

$$\left| \frac{f(z) - f(z_0)}{z - z_0} \right|$$

is bounded on $C(r)$ and its interior. So

$$\int_{C(r)} \frac{f(z) - f(z_0)}{z - z_0} dz \rightarrow 0 \text{ as } z \rightarrow 0 ,$$

so since this integral is independent of r , it must equal 0. □

Example 2.6.1. Let $\Gamma = \{|z| = 1\}$. Compute the following:

- Consider $\int_{\Gamma} \frac{\cos z}{z} dz$,

$$\int_{\Gamma} \frac{\cos z}{z} dz = \int_{\Gamma} \frac{\cos z}{z - 0} dz = 2\pi i \cos(0) = 2\pi i .$$

- Consider $\int_{\Gamma} \frac{e^z}{z-2} dz$, because $2 \notin \{|z| \leq 1\}$, by Cauchy integral theorem, this equals 0.

$$\int_{\Gamma} \frac{e^z}{z-2} dz = 2\pi i e(2)$$

- consider $\int_{\Gamma} \frac{\cos(2\pi z)}{2z-1} dz$

Example 2.6.2. Say g is holomorphic on $0 < |z| < R$, which of the following implies that

$$\int_{C(r)} g(z) dz = 0?$$

1. g is holomorphic at 0
2. g identically 0 on $0 < |z| < R$
3. $|g|$ is bounded on $0 < |z| < R$
4. $g(z) = 2\pi i$ identically
5. g is defined and continuous at 0
6. $\lim_{z \rightarrow 0} g(z) = \infty$

2.7 June 6

Proposition 2.7.1 (Cauchy Integral Formula for Derivatives). Note that $(\frac{1}{1-z})'$ then by CIF

$$\begin{aligned} f(w) &= \frac{1}{2\pi i} \int_{\Gamma} \int_{\Gamma} \frac{f(z)}{z-w} dz \\ \Rightarrow \quad \frac{d}{dw} f(w) &= \frac{1}{2\pi i} \frac{d}{dw} \int_{\Gamma} \frac{f(z)}{z-w} dz \\ &= \frac{1}{2\pi i} \int_{\Gamma} \frac{d}{dw} \frac{f(z)}{z-w} dz \\ &= \frac{1}{2\pi i} \int_{\Gamma} \frac{f(z)}{(z-w)^2} dz \\ &= \frac{2}{\pi i} \int_{\Gamma} \frac{f(z)}{z-2} dz . \end{aligned}$$

Then, taking derivative n times we have

$$f^{(n)}(w) = \frac{n!}{2\pi i} \int_{\Gamma} \frac{f(z)}{(z-w)^{n+1}} dz .$$

So f is infinitely differentiable.

2.8 June 8

THEOREM 2.8.1 (Maximum Modulus Principle).

Let f be holomorphic on a connected open set Ω . If f achieves its maximum on Ω , then f is constant. That is if there is some $z_0 \in \Omega$ s.t.

$$|f(z_0)| \geq |f(z)| \text{ for all } z \in \Omega$$

then f is constant.

Proof. Let z_0 be a local maximum of $|f|$ on Ω . Let

$$D = \{|z - z_0| \leq r\} \subset \Omega$$

be a disc around z_0 . Then the Cauchy Integral Formula says let $C(r) = \partial D = \{|z - z_0| = r\}$, then

$$\begin{aligned}
 f(z) &= \frac{1}{2\pi i} \int_{C(r)} \frac{f(z)}{z - z_0} dz \\
 &= \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z_0 + re^{i\theta})}{re^{i\theta}} d(z_0 + re^{i\theta}) \\
 &= \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z_0 + re^{i\theta})}{re^{i\theta}} re^{i\theta} d\theta \\
 &= \frac{1}{2\pi i} \int_0^{2\pi} f(z) d\theta \\
 \Rightarrow |f(z_0)| &\leq \frac{1}{2\pi} \int_0^{2\pi} |f(z)| d\theta \\
 &\leq \max_{z \in C(r)} \{|f(z)|\}.
 \end{aligned}$$

with equality if and only if $|f|$ is constant on $C(r)$, with $|f(z_0)| = \max_{z \in C(r)} |f(z)|$. Then because r was arbitrary, then around z_0 , $|f|$ is constant on D .

Now to show that f is constant on D , write $f = u + iv$, then $u^2 + v^2$ is constant on D . Then

$$\begin{aligned}
 2uu_x + 2vv_x &= 0 \\
 2uu_y + 2vv_y &= 0 \\
 -2uv_x + 2vu_x &= 0 \\
 \Rightarrow \begin{pmatrix} u & v \\ v & -u \end{pmatrix} \begin{pmatrix} u_x \\ v_x \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix}
 \end{aligned}$$

So either $u^2 + v^2 = 0$ or $u_x = v_x = 0$, $u^2 - v^2 = 0 \Rightarrow f = 0$ constant, or $u_x = v_x = -u_y = v_y = 0$ so f is constant on D .

WLOG, assume $\bar{\Omega}$ is compact. We can cover Ω with a set of open discs. Then, therefore f is constant on Ω . □

THEOREM 2.8.2 (Morera).

Say f is continuous on a domain Ω , with

$$\int_{\Gamma} f(z) dz = 0$$

for all simple closed curves $\Gamma \subset \Omega$ whose interiors are contained in Ω . Then f is holomorphic on Ω .

Proof. We will find a holomorphic F with $F' = f$. This will prove that f is holomorphic. Since holomorphicity is local, we can assume that $\Omega = D$ is a disc. Choose $z \in D$, define □

2.9 June 10

THEOREM 2.9.1 (Cauchy Integral Formula, general form).

$$f^{(n)}(z) = \frac{n!}{2\pi i} \int_C \frac{f(w)}{(w-z)^{n+1}} dw .$$

THEOREM 2.9.2.

Let f be holomorphic on and inside a circle C . Then for any z inside C , all derivatives $f^{(n)}(z)$ exist.

Note: this also means all partial derivatives exists and continuous.

Remark 2.9.3.

$$|f^{(n)}(z_0)| \leq \frac{n! \max_{z \in C_R(z_0)} |f(z)|}{R^n}$$
$$|f(z_0)| \leq \max_{z \in C_R(z_0)} |f(z)| .$$

THEOREM 2.9.4 (Liouville's Theorem).

If f is entire and bounded, f is constant.

THEOREM 2.9.5 (Fundamental Theorem of Algebra).

Every nonconstant polynomial over \mathbb{C} has a zero in \mathbb{C} .

Corollary 2.9.1. A degree- n polynomial over \mathbb{C} has exactly n zeros in \mathbb{C} , counted with multiplicity.

THEOREM 2.9.6 (Morera's Theorem).

Let f be continuous on a simply connected domain D . If $\oint_{\Gamma} f = 0$ for every closed contour Γ in D , then f is holomorphic on D .

THEOREM 2.9.7.

Let f be continuous on a simply connected domain. Then TFAE

- f has a primitive on D
- $\oint_{\Gamma} f = 0$ for all closed Γ in D
- $\int_{\Gamma_1} f = \int_{\Gamma_2} f$ for any Γ_1, Γ_2 in D sharing same initial and terminal point
- f is holomorphic on D

2.10 June 13

Lemma 2.10.1 (Symmetry Principle). Let D be a domain symmetric across \mathbb{R} . Let D^+, D^-, I be as indicated. Let f^+ be holomorphic on D^+ , f^- be holomorphic on D , both extended continuously to I and $f^+(z) = f^-(z)$ for $z \in I$.

Then

$$f(z) = \begin{cases} f^+(z), & z \in D^+ \\ f^+(z) = f^-(z), & z \in I \\ f^-(z), & z \in D^- \end{cases}$$

is holomorphic on D .

Proof. Note that f is continuous on D .

$$\left| \int_T f(z) dz - \int_{T_\varepsilon} f(z) dz \right| \leq \varepsilon \left(\max_{z \in T} |f'(z)| \right) (\text{length}(T)) \rightarrow 0 \text{ as } \varepsilon \rightarrow 0.$$

□

Lemma 2.10.2 (Schwartz's Lemma). Let $D = \{z \in \mathbb{C} : |z| < 1\}$. Let f be holomorphic on D , $f(0) = 0$, and $|f(z)| < 1 \forall z \in D$. Then $|f(z)| \leq |z|, \forall z \in D$ and $|f'(0)| \leq 1$. Furthermore, if $|f(z)| = |z|$ for any $z \in D$, then f is a rotation $f(z) = \lambda z, |\lambda| \leq 1$ constant.

Proof. Let

$$g(z) = \begin{cases} \frac{f(z)}{z}, & z \neq 0 \\ f'(0), & z = 0 \end{cases}$$

Note that g is holomorphic on D , since $\lim_{z \rightarrow 0} \frac{f(z)}{z} = \lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} = f'(0)$.

Consider g on $|z| < r < 1$. Then

$$g(z) \leq \max_{|w|=r} |g(w)| \leq \max_{|w|=r} \frac{|f(w)|}{|w|} \leq \frac{1}{r} \rightarrow 1 \text{ as } r \rightarrow 1$$

So $|g(z)| \leq 1$ on D .

For $z \neq 0$,

□

2.11 Midterm Review - June 15

COMPLEX NUMBER:

- $\bar{z} = x - iy$,
- $|z| = \sqrt{z\bar{z}} = \sqrt{x^2 + y^2} = r$,
- $\operatorname{Re}(z) = x, \operatorname{Im}(z) = y$.
- $\arg z = \theta = \arctan \frac{y}{x}, \operatorname{Arg} z = \theta \in (-\pi, \pi]$.

STANDRAD FUNCTIONS

- for $\theta \in \mathbb{R}$, $e^{\theta} = \cos \theta + i \sin \theta$
- $e^z = e^x(\cos y + i \sin y)$
- $\log z = \ln r + i\theta$

$$\sin z = \frac{e^{iz} - e^{-iz}}{zi} \quad \cos z = \frac{e^{iz} + e^{-iz}}{z}.$$

$$z^w = e^{w \log z}$$

$$z^{1/n} = r^{1/n} e^{i\frac{\theta+2\pi k}{n}}, k = 0, 1, 2, \dots, n$$

DERIVATIVE:

$$f'(z) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}.$$

f is holomorphic on a domain if it is differentiable at every point in the domain.

CR eqautsion: f is holomorphic on D if and only if $f = u + iv$ satisfies the CR equation:

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

and these partials are continuous.

Also

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta}, \frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta}.$$

Also,

$$f'(z) = \frac{\partial u}{\partial x} - i \frac{\partial v}{\partial y} = e^{i\theta} \left(\frac{\partial u}{\partial r} - i \frac{\partial v}{\partial \theta} \right) = etc...$$

Harmonic: A function u is harmonic on a domain D if

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \text{ on } D \iff \forall z \in D, \exists f \text{ holomorphic at } z, \operatorname{Re}(f(x + iy)) = u(x, y).$$

A contour is a piecewise smooth directed curve.

- **simple:** no self-intersections
- **closed:** end point = initial point
- **smooth:** if a parametrization γ is continuously differentiable
- **positively oriented:** interior is on left side as you traverse

INTEGRATION:

For $\Gamma, \gamma : [a, b] \rightarrow \mathbb{C}$,

$$\int_{\Gamma} f(z) dz = \int_a^b f(\gamma(t)) \gamma'(t) dt$$

For f continuous on a simply connected domain D , TFAE:

- f is holomorphic on D
- f has a primitive on D
- $\oint_{\Gamma} f = 0$ for all closed $\Gamma \subseteq D$
- $\int_{\Gamma_1} f = \int_{\Gamma_2} f$ for all $\Gamma_1, \Gamma_2 \subseteq D$ sharing start and end points.

Cauchy's Theorem: If f is holomorphic on a simply connected domain D , and let Γ be a closed contour in D . Then $\int_{\Gamma} f = 0$.

Cauchy's Integral Formula: If f is holomorphic on and inside a simple, closed, positively oriented contour, then $\forall z_0$ inside the contour

$$f^{(n)}(z) = \frac{n!}{2\pi i} \int_{\Gamma} \frac{f(z)(z - z_0)^{n+1}}{d} dz ,$$

in particular for $n = 0$ case

$$f(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(z)}{(z - z_0)} dz .$$

f holomorphic if and only if f is infinitely differentiable.

Liouville's Theorem: every bounded entire(holomorphic on all of \mathbb{C}) function is constant.

Fundamental Theorem of Algebra: every nonconstant polynomial has a zero in \mathbb{C} .

Maximum Modulus Principle: A nonconstant holomorphic f on a domain D cannot attain a maximum modulus in D .

Schwarz Reflection Principle: $f(z)$

3 Applications of The Cauchy Theory

3.1 Taylor Series - June 20

Remark 3.1.1. Consider a function f harmonic on a punctured disk $0 < |z - z_0| < r$. Let Γ be a simple closed, positively oriented contour wrapping about z_0 once. Consider

$$\int_{\Gamma} f(z) dz.$$

This will be a value independent of the choice of Γ . If f is holomorphic on a domain D , f is infinitely differentiable on D . In \mathbb{R} an infinitely differentiable function has a Taylor series representation.

DEFINITION 3.1.2 (Convergent Series).

A series $\sum_{n=1}^{\infty} z_n \in \mathbb{C}$ is **convergent** if

$$\lim_{k \rightarrow \infty} \sum_{n=1}^k z_n$$

converges.

DEFINITION 3.1.3 (Cauchy Series).

A series is **Cauchy** if

$$\lim_{n \rightarrow \infty} \sum_{n=k}^{\infty} z_n = 0 .$$

DEFINITION 3.1.4.

A sequence $\{f_n\}$ is **uniformly convergent** on a set S if for all $\varepsilon > 0$, exists some $N > 0 \in \mathbb{Z}$, such that $\forall z \in S, \exists L, \forall n > N$,

$$|f_n(z) - L| < \varepsilon .$$

Lemma 3.1.1. If $f_n \rightarrow f$ uniformly on S , then

$$\int_S f_n \rightarrow \int_S f .$$

Proof. $\forall \varepsilon > 0, \exists N, \forall n > N$,

$$|f_n(z) - f(z)| < \varepsilon \text{ on } S .$$

thus

$$\left| \int_S f - \int_S f_n \right| \leq \left| \int_S f - f + n \right| \leq \int_S |f - f_n| \leq \varepsilon \text{length}(s) \rightarrow 0 \text{ as } \varepsilon \rightarrow 0 .$$

□

DEFINITION 3.1.5.

A series is uniformly convergent if its sequence of partial sums is.

DEFINITION 3.1.6.

A series $\sum_{n=0}^{\infty} z_n$ is **absolutely convergent** if the series $\sum_{n=0}^{\infty} |z_n|$ converges.

DEFINITION 3.1.7.

Let $D_r(z_0) = \{z \in \mathbb{C} : |z - z_0| < r\}$ be the open disk of radius r centered at z_0 . Let $\overline{D_r(z_0)} = D_r(z_0) \cup C_r(z_0)$ be its closure.

DEFINITION 3.1.8.

Let $\{x_n\} \subseteq \mathbb{R}$,

$$\limsup x_n = \lim_{n \rightarrow \infty} \sup_{k > n} x_k .$$

Lemma 3.1.2 (Ratio Test). If $\limsup \left| \frac{z_{n+1}}{z_n} \right| < 1$, then $\sum_{n=0}^{\infty} z_n$ converges absolutely. Otherwise if $\limsup \left| \frac{z_{n+1}}{z_n} \right| > 1$, then $\sum_{n=0}^{\infty} z_n$ diverges.

Lemma 3.1.3 (Root Test). If $\limsup |z_n|^{1/n} < 1$, $\sum_{n=0}^{\infty} z_n$ converges absolutely. If $\limsup |z_n|^{1/n} > 1$, $\sum_{n=1}^{\infty} z_n$ diverges.

Lemma 3.1.4 (Comparison Test). If $\sum_{n=0}^{\infty} x_n \in \mathbb{R}$ converges and $|z_n| \leq x_n$ for all n . Then $\sum_{n=0}^{\infty} z_n$ converges absolutely.

Lemma 3.1.5. Any Cauchy series is convergent, any uniformly Cauchy series is uniformly convergent.

Lemma 3.1.6 (Weierstrass M-test). Let $\{f_n\}_{n=1}^{\infty}$ satisfy $|f_n(z)| \leq M_n$ for all $z \in S$. If $\sum_{n=1}^{\infty} M_n$ converges, then $\sum_{n=1}^{\infty} f_n(z)$ converges uniformly on S .

Proof. Let $g_n(z) = \sum_{k=1}^n f_k(z)$, $\{g_n\}$ is uniformly Cauchy on S . □

DEFINITION 3.1.9 (Power Series).

A **power series** about a point z_0 is a series of the form

$$\sum_{n=0}^{\infty} a_n (z - z_0)^n, a_n \in \mathbb{C} .$$

THEOREM 3.1.10.

If a power series $\sum_{n=0}^{\infty} a_n (z - z_0)^n$ converges at a point z with $|z - z_0| = R$, then it converges absolutely on $D_R(z_0)$ and converges uniformly on any closed subdisk of $D_R(z_0)$.

Proof. Let $w \in D_R(z_0)$ and let $|w - z_0| < r < R$. Then

$$|a_n(w - z_0)^n| = \underbrace{|a_n(z - z_0)^n|}_{\rightarrow 0, \text{ so is bounded } \leq M} \underbrace{\left| \frac{(w - z_0)^n}{(z - z_0)^n} \right|}_{\leq \frac{r}{R}} \leq M \left(\frac{r}{R} \right)^n,$$

where $\frac{r}{R} < 1$. Since $M \left(\frac{r}{R} \right)^n$ is a convergent geometric series. So

$$\sum_{n=0}^{\infty} a_n(w - z_0)^n$$

converges absolutely by comparison test.

Apply the Weierstrass M-test to the above to get uniform convergence on $\overline{D_r(z_0)}$. □

3.2 June 22

THEOREM 3.2.1 (Taylor's Theorem).

Let f be holomorphic on $D_R(z_0)$. Then $\forall z \in D_R(z_0)$,

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n.$$

Proof. Choose $z \in D_R(z_0)$ and let $|z - z_0| < r < R$. By Cauchy's Integral Formula,

$$f(z) = \frac{1}{2\pi i} \int_{C_r(z_0)} \frac{f(w)}{w - z} dw.$$

$\forall w \in C_r(z_0)$,

$$\begin{aligned} \frac{f(w)}{w - z} &= \frac{f(w)}{(w - z_0) - (z - z_0)} = \frac{f(w)}{w - z_0} \frac{1}{1 - \frac{z - z_0}{w - z_0}} \\ &= \frac{f(w)}{w - z_0} \sum_{n=0}^{\infty} \left(\frac{z - z_0}{w - z_0} \right)^n = \sum_{n=0}^{\infty} f(w) \frac{(z - z_0)^n}{(w - z_0)^{n+1}}. \end{aligned}$$

Now

$$\left| f(w) \frac{(z - z_0)^n}{(w - z_0)^{n+1}} \right| \leq \max_{w \in C_r(z_0)} |f(w)| \cdot \frac{|z - z_0|^n}{r^{n+1}}.$$

so by the Weierstrass M-test, this series converges uniformly on C . Thus integrate term-by-term,

$$\begin{aligned} f(z) &= \frac{1}{2\pi i} \int_{C_r(z_0)} \sum_{n=0}^{\infty} f(w) \frac{(z - z_0)^n}{(w - z_0)^{n+1}} dw \\ &= \frac{1}{2\pi i} \sum_{n=0}^{\infty} \int_{C_r(z_0)} \frac{f(w)}{(w - z_0)^{n+1}} dw \cdot (z - z_0)^n = \sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n. \end{aligned}$$

□

Example 3.2.1.

- $R = \infty$

$$e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!} = 1 + z + \frac{z^2}{2!} + \cdots$$

- $R = \infty$,

$$\sin z = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)!} = z - \frac{z^3}{3!} + \frac{z^5}{5!} + \cdots$$

- $R = \infty$,

$$\cos z = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n}}{(2n)!}$$

- $R = 1$,

$$\frac{1}{1-z} = \sum_{n=0}^{\infty} z^n = 1 + z + z^2 + \cdots$$

Example 3.2.2. Expand $\frac{1}{\frac{1}{2}z^2+1}$ about 0.

Let $w = -\frac{1}{2}z^2$. Then

$$\begin{aligned} \frac{1}{\frac{1}{2}z^2+1} &= \frac{1}{1-w} = \sum_{n=0}^{\infty} w^n, |w| < 1 \\ &= \sum_{n=0}^{\infty} \left(-\frac{1}{2}z^2\right)^n, \left|-\frac{1}{2}z^2\right| < 1 \iff |z^2| < 2 \iff |z| < \sqrt{2}. \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n}{2^n} z^{2n}, |z| < \sqrt{2}. \end{aligned}$$

Example 3.2.3. Taylor series for $\cos z + i \sin z$ about 0.

$$\begin{aligned} \cos z + i \sin z &= 1 - \frac{z^2}{2} + \frac{z^4}{4!} + \cdots \\ &\quad + i\left(z - \frac{z^3}{3!} + \frac{z^5}{5!} - \cdots\right) \\ &= 1 + iz - \frac{z^2}{2!} - i\frac{z^3}{3!} + \frac{z^4}{4!} + \cdots \\ &= \sum_{n=0}^{\infty} \frac{i^n z^n}{n!} = e^{iz}. \end{aligned}$$

3.3 Singularities - June 24

Remark 3.3.1. Recall a function f holomorphic on a disk $D_R(z_0)$ has a Taylor series

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n ,$$

that converges on $D_R(z_0)$ and converges uniformly on any closed disk of $D_R(z_0)$.

THEOREM 3.3.2.

Let f be analytic on an open set D containing the annulus $\{z : r_1 \leq |z - z_0| \leq r_2\}$, $0 < r_1 < r_2 < \infty$, and let γ_1, γ_2 denote the positively oriented inner and outer boundaries of the annulus. Then for $r_1 < |z - z_0| < r_2$, we have

$$f(z) = \frac{1}{2\pi i} \int_{\gamma_2} \frac{f(w)}{w - z} dw - \frac{1}{2\pi i} \int_{\gamma_1} \frac{f(w)}{w - z} dw .$$

DEFINITION 3.3.3.

Define

$$\sum_{n=-\infty}^{\infty} a_n (z - z_0)^n = \sum_{n=0}^{\infty} a_n (z - z_0)^n + \sum_{n=1}^{\infty} a_n (z - z_0)^{-n} .$$

Note this means that e.g.

$$\cdots - \frac{1}{3} - \frac{1}{2} - 1 + 1 + \frac{1}{2} + \frac{1}{3} + \cdots$$

does not converge

THEOREM 3.3.4 (Laurent Series).

Let f be holomorphic on an annulus $\{z \in \mathbb{C} : r_1 < |z - z_0| < r_2\}$, then on that annulus, f has a unique **Laurent series** (generalization of Cauchy series),

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z - z_0)^n ,$$

which converges on the annulus and converges uniformly on closed subannuli, and the coefficients a_n are given by

$$a_n = \frac{1}{2\pi i} \int_{C(z_0, r)} \frac{f(w)}{(w - z_0)^{n+1}} dw, n = 0, \pm 1, \pm 2, \dots$$

Proof. HW

□

DEFINITION 3.3.5 (Isolated singularity).

f has an **isolated singularity** at z_0 if f is not analytic at z_0 but is analytic on the punctured disk $D(z_0, r) \setminus \{z_0\}$ for some $r > 0$.

Lemma 3.3.1. If $f(z)$ is analytic at z_0 and $f(z_0) = 0$, and f is not identically zero in any $D_r(z_0)$, then $\frac{1}{f(z)}$ has a singularity at z_0 .

DEFINITION 3.3.6.

An analytic function f has a **zero of order m at z_0** if $\frac{f(z)}{(z - z_0)^m}$ is analytic at z_0 but $\frac{f(z)}{(z - z_0)^{m+1}}$ is not. Equivalently if $f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n$, the order is the smallest n such that $a_n \neq 0$.

Example 3.3.1. Let $f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n$ about z_0 . Then

$$f(z) = 2(z - z_0)^4 - (z - z_0)^5 + \cdots$$

has a zero of order 4.

DEFINITION 3.3.7.

A **singularity** of f is a point where f is not analytic but is a limit point of points where f is analytic.

DEFINITION 3.3.8.

Let z_0 be an isolated singularity of f . Let $\sum_{n=-\infty}^{\infty} a_n(z - z_0)^n$ be the Laurent series of f at z_0 .

- If $a_{-m} \neq 0$ but $a_n = 0$ for all $n > m$, we call z_0 a **pole of order m** $\iff (z - z_0)^m f(z)$ is analytic at z_0 and $(z - z_0)^{m-1} f(z)$ is not.
- If $a_{-n} = 0$ for all $n > 0$ we call this a **removable singularity**. In this case,

$$f(z) = \begin{cases} f(z), & z \neq z_0 \\ \lim_{z \rightarrow z_0} f(z), & z = z_0 \end{cases}$$

is analytic at z_0 .

- If $a_{-n} \neq 0$ for infinitely many $n > 0$, we call this an **essential singularity**.

Example 3.3.2 (Removable Singularity).

$$f(z) = \frac{\sin z}{z} = \frac{1}{z} \cdot \left(z - \frac{z^3}{3!} + \frac{z^5}{5!} - \cdots \right) = 1 - \frac{z^2}{3!} + \frac{z^4}{5!} - \cdots.$$

Lemma 3.3.2. Let $f(z) = \sum_{n=-m}^{\infty} a_n(z - z_0)^n$, $a_m \neq 0$ on a punctured disk $0 < |z - z_0| < r$. Let Γ be a simple, closed, positively oriented contour in the annulus with z_0 inside the loop, then

$$\int_{\Gamma} f(z) dz = \int_{\Gamma} \sum_{n=-m}^{\infty} a_n(z - z_0)^n dz = \sum_{n=-m}^{\infty} a_n \int_{\Gamma} (z - z_0)^n dz = 2\pi i a_{-1}.$$

Proof.

$$a_{-1} = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(w)}{(w - z_0)^{-1+1}} dw$$

$$\Rightarrow 2\pi i a_{-1} = \int_{\Gamma} f(w) dw = \int_{\Gamma} \sum_{n=-m}^{\infty} a_n (z - z_0)^n dz$$

□

DEFINITION 3.3.9 (Residue).

Given f, z_0, Γ as before, we define the **residue** of f at z_0 to be

$$\frac{1}{2\pi i} \int_{\Gamma} f(z) dz = a_{-1} = \text{res}_{z_0}(f) .$$

DEFINITION 3.3.10 (Meromorphic).

A function f is called **meromorphic** on a domain D if it is holomorphic on all of D except for a set of isolated poles.

THEOREM 3.3.11 (The Residue Theorem).

Let f be meromorphic on a simply connected domain D and let Γ be a simple, closed, positively oriented contour lying in D . Let z_1, \dots, z_k be the poles of f inside Γ . Then

$$\int_{\Gamma} f(z) dz = 2\pi i \sum_{j=1}^k \text{res}_{z_j}(f) .$$

3.4 Singularities and Residue Theory - June 27

Recall let f be meromorphic with a pole at z_0 and let

$$\sum_{n=-m}^{\infty} a_n (z - z_0)^n, a_{-m} \neq 0$$

be the Laurent series for f valid in some punctured disk $0 < |z - z_0| < R, R > 0$. Then the order of the pole at z_0 is m and the residue is a_{-1} .

Example 3.4.1.

$$f(z) = \frac{3}{z^4} - \frac{5}{z^2} + \frac{7}{z} + 2 + z + \dots$$

f has an order 4 pole at 0 with residue 7.

Example 3.4.2. Consider $f(z) = \frac{1}{z^2+z} = \frac{1}{z(z+1)}$,

- $0 < |z| < 1$. Then

$$\begin{aligned}\frac{1}{z(z+1)} &= \frac{1}{z} \left(\frac{1}{1 - (-z)} \right) \\ &= \frac{1}{z} \sum_{n=0}^{\infty} (-z)^n, \quad | -z | < 1 \\ &= \frac{1}{z} - 1 + z - z^2 + z^3 - \dots\end{aligned}$$

order 1, residue 1

- $0 < |z+1| < 1$, then

$$\begin{aligned}\frac{z(z+1)}{z(z+1)} \frac{1}{z+1} &= \frac{1}{z+1} \left(-\frac{1}{1 - (z+1)} \right) \\ &= -\frac{1}{z+1} \sum_{n=0}^{\infty} (z+1)^n \\ &= -\frac{1}{z+1} - 1 - (z+1) - (z+1)^2 - \dots\end{aligned}$$

Simple pole, residue -1 .

- $|z| > 1$,

$$\begin{aligned}\frac{1}{z(z+1)} &= \frac{1}{z} \frac{1}{z+1} \frac{1/z}{1/z} \\ &= \frac{1}{z^2} \frac{1}{1 + \frac{1}{z}} \\ &= \frac{1}{z^2} \sum_{n=0}^{\infty} \left(-\frac{1}{z} \right)^n, \quad \left| -\frac{1}{z} \right| < 1 \\ &= \dots + \frac{1}{z^5} + \frac{1}{z^4} + \frac{1}{z^3} + \frac{1}{z^2}\end{aligned}$$

DEFINITION 3.4.1 (Simple Pole).

A pole of order 1 is called a **simple pole**.

Remark 3.4.2. Recall

f has a pole of order m at z_0

$\iff (z - z_0)^m f(z)$ has a removable singularity at z_0 , $(z - z_0)^{m-1} f(z)$ has a pole at z_0 .

Therefore,

f has a zero of order m at z_0

$\iff \frac{f(z)}{(z - z_0)^m}$ has a removable singularity at z_0 , $\frac{f(z)}{(z - z_0)^{m+1}}$ has a pole at z_0 .

$$\frac{1}{(z - z_0)^m} (a_m(z - z_0)^m + a_{m+1}(z - z_0)^{m+1} + \dots) = a_m + a_{m+1}(z - z_0) + \dots$$

Proposition 3.4.1. Let f and g be analytic at z_0 , let f have a zero of order m at z_0 and let g have a zero of order n at z_0 . Then

$$\frac{f(z)}{g(z)} = \frac{\frac{f(z)}{(z-z_0)^m} (z - z_0)^m}{\frac{g(z)}{(z-z_0)^n} (z - z_0)^n} = (z - z_0)^{m-n} h(z)$$

where $h(z_0) \neq 0$, h analytic at z_0 .

$$\frac{f(z)}{g(z)} \text{ has } \begin{cases} \text{a zero of order } m - n \text{ at } z_0, & \text{if } m > n \\ \text{a pole of order } n - m \text{ at } z_0, & \text{if } m < n \\ \text{a removable singularity,} & \text{otherwise} \end{cases}$$

Example 3.4.3. $\frac{1}{z(z+1)}$ has simple poles at $z = 0, z = -1$. Then

$$\frac{1}{z^3(z+1)^2(z-2)} \text{ has } \begin{cases} \text{order 3 pole at } 0 \\ \text{order 2 pole at } -1 \\ \text{simple pole at } 2 \end{cases}$$

Example 3.4.4.

$$\frac{\cos z - 1}{z^2} = \frac{1}{z^2} \left(1 - \frac{z^2}{2!} + \frac{z^4}{4!} - \dots - 1 \right) = -\frac{1}{2!} + \frac{z^2}{4!} - \dots$$

So $\frac{\cos z - 1}{z^2}$ has a removable singularity at $z = 0$.

Remark 3.4.3. Let f have a simple pole at z_0 and a Laurent series

$$f(z) = \frac{a_{-1}}{z - z_0} + a_0 + a_1(z - z_0) + \dots$$

in some punctured disk about z_0 . Then

$$\begin{aligned} (z - z_0)f(z) &= a_{-1} + a_0(z - z_0) + a_1(z - z_0)^2 + \dots \\ \Rightarrow \lim_{z \rightarrow z_0} ((z - z_0)f(z)) &= a_{-1} = \text{res}_{z_0} f. \end{aligned}$$

If f has a simple pole at z_0 ,

$$\text{res}_{z_0} f = \lim_{z \rightarrow z_0} (z - z_0)f(z).$$

Example 3.4.5.

$$f(z) = \frac{1}{z(z+1)}. \quad \text{res}_0 f = \lim_{z \rightarrow 0} z f(z) = \lim_{z \rightarrow 0} \frac{1}{z+1} = 1.$$

$$\text{res}_{-1} f = \lim_{z \rightarrow -1} (z+1) \frac{1}{z(z+1)} = -1.$$

Remark 3.4.4. Now let f have a pole of order m at z_0 ,

$$\begin{aligned} f(z) &= \frac{a_{-m}}{(z-z_0)^{-m}} + \frac{a_{-m+1}}{(z-z_0)^{-m+1}} + \cdots + \frac{a_{-1}}{(z-z_0)} + \cdots \\ \Rightarrow (z-z_0)^m f(z) &= a_{-m} + a_{-m+1}(z-z_0) + \cdots + a_{-1}(z-z_0)^{m-1} + \cdots \\ \Rightarrow \frac{d}{dz}((z-z_0)^m f(z)) &= a_{-m+1} + \cdots + (m-1)a_{-1}(z-z_0)^{m-2} + \cdots \\ \Rightarrow \frac{d^{m-1}}{dz^{m-1}}((z-z_0)^m f(z)) &= a_{-1}(m-1)! + \cdots \\ \lim_{z \rightarrow z_0} \frac{d^{m-1}}{dz^{m-1}}((z-z_0)^m f(z)) &= a_{-1}(m-1)! \\ a_{-1} = \text{res}_{z_0} f &= \frac{1}{(m-1)!} \lim_{z \rightarrow z_0} \frac{d^{m-1}}{dz^{m-1}}((z-z_0)^m f(z)) \end{aligned}$$

Example 3.4.6. $f(z) = \frac{e^z+1}{z^3}$. $e^0 + 1 = 2 \neq 0$, so f has a order 3 pole at 0.

$$\text{res}_0 f = \frac{1}{(3-1)!} \lim_{z \rightarrow 0} \frac{d^2}{dz^2} \left(z^3 \frac{e^z+1}{z^3} \right) = \frac{1}{2!} \lim_{z \rightarrow 0} e^z = \frac{1}{2}.$$

3.5 June 29

Proposition 3.5.1. Let $f(z) = \sum_{n=-\infty}^{\infty} a_n(z-z_0)^n$ be the Laurent series for f in some annulus.

$$a_{-1} = \frac{1}{2\pi i} \oint_{\Gamma} f(z) dz,$$

where Γ is a simple, closed, positively oriented contour looping around the inner circle of the annulus.

Now

$$(z-z_0)^{-m-1} f(z) = \cdots + \frac{a_m}{z-z_0} + \cdots \quad \Rightarrow \quad a_m = \frac{1}{2\pi i} \oint_{\Gamma} \frac{f(z)}{(z-z_0)^{m+1}} dz.$$

Note for a Taylor series, this is equivalent to $\frac{f^{(n)}(z)}{n!}$ by Cauchy's Integral Theorem.

Proposition 3.5.2. Let $f(z) = \frac{g(z)}{h(z)}$, where g, h are analytic at z_0 . Let $g(z_0) \neq 0, h(z) = 0, h'(z_0) \neq 0$, i.e., f has a simple pole at z_0 . Then,

$$\begin{aligned}\operatorname{res}_{z_0}(f) &= \lim_{z \rightarrow z_0} (z - z_0)f(z) \\ &= \lim_{z \rightarrow z_0} \frac{g(z)}{h(z)} \\ &= g(z_0) \lim_{z \rightarrow z_0} \frac{z - z_0}{h(z)} \\ &= g(z_0) \lim_{z \rightarrow z_0} \frac{z - z_0}{h(z) - h(z_0)} \\ &= \frac{g(z_0)}{h'(z_0)}.\end{aligned}$$

Example 3.5.1. Find residues of all poles of $f(z) = \frac{1}{z^3 - 1}$. Then

$$z^3 - 1 = 0 \iff z^3 = 1 \iff z \in \{1, e^{2\pi i/3}, e^{4\pi i/3}\}$$

Thus f has 3 simple poles. Residue at a simple pole z is $\frac{1}{3z^2}$

$$\begin{aligned}\operatorname{res}_1 f &= \frac{1}{3(1^2)} = \frac{1}{3} \\ \operatorname{res}_{e^{2\pi i/3}} f &= \frac{1}{3(e^{2\pi i/3})^2} = \frac{1}{3}e^{2\pi i/3} \\ \operatorname{res}_{e^{4\pi i/3}} f &= \frac{1}{3(e^{4\pi i/3})^2} = \frac{1}{3}e^{4\pi i/3}.\end{aligned}$$

Example 3.5.2. $\int_0^\infty \frac{1}{x^4 + 1} dz$

Let $I = \int_0^\infty \frac{1}{x^4 + 1} dx$, note that $2I = \int_{-\infty}^\infty \frac{1}{x^4 + 1} dz$.

Let Γ_R be the line segment running from $-R$ to $R \in \mathbb{R}$. Then $2I = \lim_{R \rightarrow \infty} \int_{\Gamma_R} \frac{1}{x^4 + 1} dz$. Let C_R be the upper semi circle running from R to $-R$. Not $\Gamma_R + C_R$ is a simple, closed, positively oriented contour, so we use the Residue Theorem. Consider

$$\left| \int_{C_R} \frac{1}{z^4 + 1} dz \right| \leq \left| \int_{C_R} \frac{1}{R^4} dz \right| \leq |\pi i R| \frac{1}{R^4} \leq \frac{\pi}{R^3} \rightarrow 0 \text{ as } R \rightarrow \infty.$$

Next we locate the poles of $\frac{1}{z^4 + 1}$ and find their residues. $z^4 + 1 = 0 \iff z^4 = -1 \iff z \in \{e^{i\pi/4}, e^{3\pi/4}, e^{5\pi/4}, e^{7\pi/4}\}$

3.6 Extended Complex Plane - July 4

DEFINITION 3.6.1.

Extended complex plane $\mathbb{C} \cup \{\infty\} = \hat{\mathbb{C}}$.

DEFINITION 3.6.2.

Define the behavior of $f(z)$ at ∞ to behavior of $f(\frac{1}{z})$ at 0.

Example 3.6.1. Let $f(z) = z^2 + 1$, so that $f(\frac{1}{z}) = \frac{1}{z^2} + 1$, so that it has order 2 and residue 0 and

$$\lim_{R \rightarrow \infty} \int_{C_R(0)} z^2 + 1 dz = 0$$

also

$$\int_{-C_\infty(0)} f(z) dz = -2\pi i \operatorname{res}_\infty(f).$$

Example 3.6.2. $f(z) = \frac{z+1}{z-i}$

$$f\left(\frac{1}{z}\right) = \frac{\frac{1}{z} + 1}{\frac{1}{z} - i} = \frac{1 + z}{1 - iz}$$

At $z = 0$, $f(\frac{1}{z}) = 1$, so f is analytic at ∞ .

Example 3.6.3. $f(z) = \sin z$, $f(\frac{1}{z}) = \sin \frac{1}{z}$ does not converge as $z \rightarrow 0$. So $\sin z$ has an essential singularity at ∞ .

$$\lim_{R \rightarrow \infty} \int_{C_R(0)} f(z) dz = 2\pi i \sum_{z_j \in \mathbb{C}, z \text{ pole of } f} \operatorname{res}_{z_j} f = -2\pi i \cdot \operatorname{res}_\infty f$$

DEFINITION 3.6.3.

At an isolated singularity z_0 ,

- If $\lim_{z \rightarrow z_0} f(z) = c \in \mathbb{C}$, then f is analytic at z_0 (removable singularity)
- If $\lim_{z \rightarrow z_0} |f(z)| = \infty$, then f has a pole at z_0 .
- If $\lim_{z \rightarrow z_0} f(z)$ does not exist in $\hat{\mathbb{C}}$, then f has essential singularity at z_0 .

Example 3.6.4.

$$\int_0^\infty \frac{1}{x^3 + 1} dx$$

Let $f(z) = \frac{1}{z^3 + 1}$, f has poles at $z = -1, e^{i\pi/3}, e^{5i\pi/3}$.

$$\int_{\Gamma_2} \frac{1}{z^3 + 1} dz = \int_0^R \frac{1}{(te^{2\pi i/3})^3 + 1} e^{2\pi i/3} dt = \int_0^R \frac{e^{2\pi i/3}}{t^3 + 1} dt = e^{2\pi i/3} \int_0^R \frac{1}{t^3 + 1} dt = e^{2\pi i/3} \int_{\Gamma_1} f ,$$

parametrize Γ_1 by $\Gamma_1 : \gamma_1(t) = t, t \in [0, R]$. Then

$$\int_{\Gamma_1} f(z) dz = \int_0^R \frac{1}{t^3 + 1} dt .$$

3.7 Cauchy Principal Value - July 6

DEFINITION 3.7.1.

$$\int_{-\infty}^{\infty} f(z)dz = \lim_{a \rightarrow -\infty} \int_a^0 f(x)dz + \lim_{b \rightarrow \infty} \int_0^b f(x)dz$$

if both limits converge.

Example 3.7.1. Note $\lim_{R \rightarrow \infty} \int_{-R}^R f(x)dx$ and $\lim_0^{\infty} f(z)dz$ are not necessarily the same thing. e.g. $\int_{-\infty}^{\infty} xdx$ diverges but $\lim_{\infty} \int_R^R xdx = 0$.

DEFINITION 3.7.2 (Cauchy Principal Value).

Given a continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$, define the **Cauchy principal value** of $\int_{-\infty}^{\infty} f(x)dx$ as

$$p.v. \int_{-\infty}^{\infty} f(x)dx = \lim_{R \rightarrow \infty} \int_{-R}^R f(x)dz .$$

Remark 3.7.3. If $\int_{-\infty}^{\infty} f(x)dx$ exists, then

$$\int_{-\infty}^{\infty} f(x)dx = p.v. \int_{-\infty}^{\infty} f(x)dx$$

Example 3.7.2. $p.v. \int_{-\infty}^{\infty} \frac{\cos x}{1+x^2} dx$.

Let $f(z) = \frac{\cos z}{1+z^2}$,

$$\left| \int \frac{\cos z}{1+z^2} dz \right| = \left| \int \frac{\frac{1}{2}(e^{iz} + e^{-iz})}{1+z^2} dz \right|$$

- Consider

$$I_1 = p.v. \int_{-\infty}^{\infty} \frac{e^{iz}}{1+z^2} dz$$

We have

$$\left| \int_{C_r} \frac{e^{iz}}{1+z^2} dz \right| \sim \frac{1}{R^2} R \sim \frac{1}{R} \rightarrow 0 \text{ as } R \rightarrow \infty ,$$

then

$$\int_{C_R + \Gamma} f(z)dz = 2\pi i \operatorname{res}_i(f) = 2\pi i \left. \frac{e^{iz}}{2z} \right|_{z=i} = 2\pi i \frac{e^{-1}}{2i} = \frac{\pi}{e} .$$

Hence, $I = \frac{\pi}{e} - 0 = \frac{\pi}{e}$.

- Consider

$$I_2 = p.v. \int_{-\infty}^{\infty} \frac{e^{-iz}}{1+z^2} dz .$$

We have

$$\left| \int_{C_R} \frac{e^{-iz}}{1+z^2} dz \right| \sim \frac{1}{R^2} \cdot R \sim \frac{1}{R} \rightarrow 0 \text{ as } R \rightarrow \infty .$$

$$\int_{C_R+\Gamma} f = -2\pi i \operatorname{res}_{-i} f = -2\pi i \frac{e^{-iz}}{2z} \Big|_{z=-i} = 2\pi i \frac{e^{-1}}{-2i} = \frac{\pi}{e}.$$

Hence $I_2 = \frac{\pi}{e}$.

Therefore,

$$p.v. \int_{-\infty}^{\infty} \frac{\cos z}{1+z^2} dz = \frac{1}{2} \left(p.v. \int_{-\infty}^{\infty} \frac{e^{iz}}{1+z^2} dz + p.v. \int_{-\infty}^{\infty} \frac{e^{-iz}}{1+z^2} dz \right) = \frac{1}{2} \left(\frac{\pi}{e} + \frac{\pi}{e} \right) = \frac{\pi}{e}.$$

Example 3.7.3.

$$\int_0^{2\pi} \sin^2 \theta d\theta$$

Let $z = e^{i\theta} = \cos \theta + i \sin \theta$.

$$\sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i} = \frac{1}{2i} \left(z + \frac{1}{z} \right).$$

thus

$$\int_0^{2\pi} \sin^2 \theta d\theta = \int_{C_1(0)} \left(\frac{1}{2i} \left(z + \frac{1}{z} \right) \right)^2 \frac{d\theta}{dz} dz = \int_{C_1(0)} \left(\frac{z + \frac{1}{z}}{2i} \right)^2 \frac{1}{iz} dz.$$

$$\begin{aligned} \int_{C_1(0)} \frac{1}{-4i} \left(z^2 + 2 + \frac{1}{z^2} \right) \left(\frac{1}{z} \right) dz &= -\frac{1}{4i} \int_{C_1(0)} \left(z - \frac{2}{z} + \frac{1}{z^3} \right) dz \\ &= -\frac{1}{4i} 2\pi i \operatorname{res}_0 \left(z - \frac{2}{z} + \frac{1}{z^3} \right) \\ &= -\frac{1}{4i} (2\pi i)(-2i) = \pi. \end{aligned}$$

DEFINITION 3.7.4.

Let f be continuous on $[a, b]$ except at c , $a < c < b$, Then

$$p.v. \int_a^b f(z) dz = \lim_{\varepsilon \rightarrow 0} \left(\int_a^{c-\varepsilon} f(x) dx + \int_{c+\varepsilon}^b f(x) dx \right)$$

Combine those definitions.

Proposition 3.7.1. Let $p(z), q(z)$ be polynomial with $\deg(p) \leq \deg(q) - 2$, then for any arc C_R of $C_R(0)$.

$$\lim_{R \rightarrow \infty} \left| \int_{C_R} \frac{p(z)}{q(z)} dz \right| = 0,$$

this is because

$$\left| \int_{C_R} \frac{p(z)}{q(z)} dz \right| \sim R \cdot \frac{R^{\deg(p)}}{R^{\deg(q)}} = R \cdot R^{-2} = \frac{1}{R} \rightarrow 0 \text{ as } R \rightarrow \infty$$

Lemma 3.7.1 (Jordan's Lemma). Let $a > 0$ and $\deg(q) \geq 1 + \deg(p)$, let C_R be the upper half of $C_R(0)$, then

$$\lim_{R \rightarrow \infty} \int_R e^{iaz} \frac{p(z)}{q(z)} dz = 0 .$$

Proof. Parametrize C_R by Re^{it} , $t \in [0, \pi]$, then

$$\int_R e^{iaz} \frac{p(z)}{q(z)} dz = \int_R e^{iaRe^{it}} \frac{p(Re^{it})}{q(Re^{it})} iRe^{it} dt$$

Then

•

$$\left| e^{iaRe^{it}} \right| = |\exp(iaR(\cos t + i \sin t))| = \exp(-aR \sin t)$$

- For sufficiently large R , $\exists K \in \mathbb{R}$ such that

$$\left| \frac{p(Re^{it})}{q(Re^{it})} \right| \leq \frac{K}{R} .$$

Therefore,

$$\begin{aligned} \left| \int_R e^{iaRe^{it}} \frac{p(Re^{it})}{q(Re^{it})} iRe^{it} dt \right| &\leq \int_0^\pi e^{-aR \sin t} \frac{K}{R} \cdot R dt \\ &= K \int_0^\pi e^{-aR \sin t} dt \\ &= 2K \int_0^{\pi/2} e^{-aR \sin t} dt \\ &\leq 2K \int_0^{\pi/2} e^{-aR \frac{2t}{\pi}} dt. \quad (\sin t \geq \frac{2t}{\pi} \text{ on } [0, \frac{\pi}{2}]) \\ &= 2K \left(-\frac{\pi}{2aR} \right) (e^{-aR} - 1) \rightarrow 0 \text{ as } R \rightarrow \infty . \end{aligned}$$

□

Lemma 3.7.2. Let f be meromorphic with a simple pole at z_0 and let Γ_r be parametrized by $\gamma(t) = z_0 + re^{i\theta}$, $\theta_1 < \theta < \theta_2$. Then,

$$\lim_{r \rightarrow 0^+} \int_{\Gamma_r} f(z) dz = i(\theta_2 - \theta_1) \text{res}_{z_0} f .$$

Proof.

$$f(z) = \frac{a_{-1}}{z - z_0} + \sum_{n=0}^{\infty} a_n (z - z_0)^n .$$

unfinished

□

3.8 July 11

Lemma 3.8.1. Let $a > 0$, $\deg(q) \geq 1 + \deg(p)$, let C_R be the upper half of $C_R(0)$, then

$$\lim_{R \rightarrow \infty} \int_{C_R} e^{-az} \frac{p(z)}{q(z)} dz = 0 .$$

Proof. Parameterize C_R by Re^{it} with $t \in [0, \pi]$, now □

Remark 3.8.1.

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{p(z)}{q(z)} dz & \quad \text{Need } \deg(q) \geq 2 + \deg(p) \\ \int_{-\infty}^{\infty} \cos(z) \frac{p(z)}{q(z)} dz & \quad \text{Need } \deg(q) \geq 1 + \deg(p) \end{aligned}$$

Lemma 3.8.2. Let f be meromorphic with a simple pole at z_0 , and Γ_r be parameterized by $r(t) = z_0 + re^{i\theta}$ with $\theta_1 < \theta < \theta_2$ then

$$\int_{r \rightarrow 0^+} \int_{\Gamma_r} f(z) dz = i(\theta_2 - \theta_1) \text{res}_{z_0} f .$$

Proof.

$$f(z) = \frac{a_{-1}}{z - z_0} + \sum_{n=0}^{\infty} a_n (z - z_0)^n = \frac{a_{-1}}{z - z_0} + g(z) ,$$

where g is analytic so g is continuous then $\exists R$ such that for $0 < r \leq R$, $\exists M > 0$ s.t. $|g(z)| \leq M$, so that

$$\left| \int_{\Gamma_r} g(z) dz \right| \leq M \cdot \text{length}(\Gamma_r) = M \cdot (\theta_2 - \theta_1) r \rightarrow 0 \text{ as } r \rightarrow 0^+ ,$$

then

$$\int_{\Gamma_r} f(z) dz = \int_{\Gamma_r} \frac{a_{-1}}{z - z_0} dz + 0 = a_{-1} \int_{\theta_1}^{\theta_2} \frac{1}{re^{i\theta}} rie^{i\theta} d\theta = a_{-1} \int_{\theta_1}^{\theta_2} i d\theta = i(\theta_2 - \theta_1) \text{res}_{z_0}(f) .$$

□

3.9 July 11

Recall

Lemma 3.9.1. Let f be meromorphic with a simple pole at z_0 . Then if C_r is parametrized by $\gamma(t) = z_0 + re^{it}$, $t \in [\theta_1, \theta_2]$,

$$\lim_{r \rightarrow 0^+} \int_{C_r} f(z) dz = i(\theta_2 - \theta_1) \text{res}_{z_0}(f) .$$

Example 3.9.1.

$$\begin{aligned} p.v. \int_{-\infty}^{\infty} \frac{e^{ix}}{x} dx &= \lim_{R \rightarrow \infty, r \rightarrow 0^+} \left(\int_{-R}^{-r} \frac{e^{ix}}{x} dx + \int_r^R \frac{e^{ix}}{x} dx \right) \\ &\Rightarrow \int_{C_R} f(z) dz \rightarrow 0 \text{ by Jordan's Lemma .} \end{aligned}$$

$$\begin{aligned} \int_{C_r} \frac{e^{iz}}{z} dz &= i(0 - \pi) \text{res}_0 f = -\pi i(1) = i\pi i , \\ \oint_{C_R + C_r + \Gamma} f(z) dz &= 0 \\ \Rightarrow \lim_{R \rightarrow \infty, r \rightarrow 0^+} \int_{\Gamma} f(z) dz &= \oint_{C_R + C_r + \Gamma} f - \int_{C_R} f - \int_{C_r} f = 0 - 0 - (-\pi i) = \pi i . \end{aligned}$$

Example 3.9.2. $\int_0^{\infty} \frac{x^{1/3}}{1+x^2} dx$, let $f(z) = \frac{z^{1/3}}{1+z^2}$ with branch cut along the positive real axis.

$$\begin{aligned} \left| \int_{C_R} \frac{z^{1/3}}{z^2+1} dz \right| &\sim \frac{R^{1/3}}{R^2} R \sim R^{-2/3} \rightarrow 0 \text{ as } R \rightarrow \infty \\ \left| \int_{C_r} \frac{z^{1/3}}{z^2+1} dz \right| &\sim r \cdot \frac{r^{1/3}}{1} R \sim r^{4/3} \rightarrow 0 \text{ as } r \rightarrow 0^+ . \\ \lim_{R \rightarrow \infty, r \rightarrow 0^+} \int_{\Gamma_1} f &\rightarrow \int_0^{\infty} f(z) dz = I . \end{aligned}$$

$$\begin{aligned} \int_{\Gamma_2} f(z) dz &= \int_{\Gamma_2} \frac{z^{1/3}}{1+z^2} dz \\ &= \int_{\Gamma_1} \frac{(ze^{2\pi i})^{1/3}}{1+z^2} dz \\ &= \int_{\Gamma_1} \frac{z}{1+z^2} e^{2\pi i/3} dz = I e^{\frac{2\pi i}{3}} . \end{aligned}$$

$f(z)$ has simple poles at $z = \pm i$ with residues.

$$\begin{aligned} \text{res}_i f &= \frac{i^{1/3}}{2i} = \frac{(e^{i\pi/2})^{1/3}}{2i} = \frac{e^{i\pi/6}}{2i} = \frac{\frac{\sqrt{3}}{2} + i\frac{1}{2}}{2i} = \frac{1}{4} - i\frac{\sqrt{3}}{4} \\ \text{res}_{-i} f &= \frac{(-i)^{1/3}}{-2i} = \frac{(e^{i\pi/2})^{1/3}}{-2i} = \frac{e^{i\pi/6}}{-2i} = -\frac{1}{2} . \end{aligned}$$

Then

$$\begin{aligned} 2\pi i \left(-\frac{1}{4} - i\frac{\sqrt{3}}{4} \right) &= \pi i \left(-\frac{1}{2} - i\frac{\sqrt{3}}{2} \right) = \pi i e^{-2\pi i/3} = I(1 - e^{2\pi i/3}) \\ I &= \frac{\pi i e^{-2\pi i/3}}{1 - e^{2\pi i/3}} = \frac{\pi}{\sqrt{3}} . \end{aligned}$$

Example 3.9.3. $\int_0^\infty \frac{1}{1+x^3} dx$

Let $f(z) = \frac{\log z}{1+z^3}$, branch cut along the positive real axis.

$$\int_{\Gamma_1} \frac{\log z}{1+z^3} dz = \int_r^R \frac{\ln x}{1+x^3} dx ,$$

$$\int_{\Gamma_2} \frac{\log z}{1+z^3} dz = \int_r^R \frac{\log(xe^{2\pi i})}{1+x^3} dx = \int_r^R \frac{\ln x + 2\pi i}{1+x^3} dx .$$

Therefore,

$$\int_{\Gamma_1} f dz - \int_{\Gamma_2} f dz = \int_r^R \frac{\ln x}{1+x^3} dx - \int_r^R \frac{\ln x + 2\pi i}{1+x^3} dx = -2\pi i \int_r^R \frac{1}{1+x^3} dx = 2\pi i I .$$

3.10 July 13

Remark 3.10.1. Let $f \neq 0$ be meromorphic on D and let Γ be a simple, positively oriented closed contour with Γ and its interior he is in D , consider $\frac{f'(z)}{f(z)}$ is meromorphic and its poles can only lie at poles and zeros of f .

Let z_0 be an order $-m$ zero of f . Then

$$f(z) = (z - z_0)^m g(z), g(z_0) \neq 0, g \text{ analytic} .$$

Now

$$f'(z) = m(z - z_0)^{m-1} g(z) + (z - z_0)^m g'(z),$$

so

$$\frac{f'(z)}{f(z)} = \frac{m(z - z_0)^{m-1} g(z) + (z - z_0)^m g'(z)}{(z - z_0)^m g(z)} = \frac{m}{z - z_0} + \frac{g'(z)}{g(z)} .$$

Let z_0 be an order $-m$ pole of f , then

$$f(z) = \frac{h(z)}{(z - z_0)^n}, h(z_0) \neq 0, h \text{ analytic} ,$$

$$f' = \frac{(z - z_0)^m h'(z) - m(z - z_0)^{m-1} h(z)}{(z - z_0)^{2m}} .$$

so

$$\frac{f'}{f} = \frac{-m(z - z_0)^{m-1} h(z) + (z - z_0)^m h'(z)}{(z - z_0)^m h(z)} = -\frac{m}{(z - z_0)} + \frac{h'(z)}{h(z)} .$$

3.11 July 15

THEOREM 3.11.1 (The Argument Principle).

Let f be meromorphic on and inside a simple, closed, positively oriented contour Γ . Let $N_0(f)$ and $N_p(f)$ be the number of zeros and number of poles of f inside Γ . (both counted with multiplicity) Then

$$\frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z)} dz = N_0(f) - N_p(f) .$$

Note if f is analytic on and inside Γ this becomes

$$\frac{1}{2\pi i} \int_{\Gamma} \frac{f'}{f} = N_0(f) .$$

DEFINITION 3.11.2.

Let Γ be a closed contour and let $z_0 \notin \Gamma$. The **curling number** of Γ about z_0 , denoted $n(\Gamma, z_0)$ is the unique integer n such that Γ is homeomorphic to $\underbrace{C_1(z_0) + C_1(z_0) + \cdots + C_1(z_0)}_{n \text{ copies}}$ in $\mathbb{C} \setminus \{z_0\}$.

Lemma 3.11.1. For $z_0 \in \Gamma$,

$$\oint_{\Gamma} \frac{1}{z - z_0} dz = 2\pi i n(\Gamma, z_0) ,$$

Proposition 3.11.1. Let $f(\Gamma)$ be which $\Gamma : \gamma(t) : [a, b] \rightarrow \Gamma$,

$$\begin{aligned} \int_{\Gamma} \frac{1}{z - z_0} dz &= \int_a^b \frac{1}{\gamma(t) - z_0} \gamma'(t) dt \\ \Rightarrow \int_{\Gamma} \frac{1}{f(z) - z_0} dz &= \int_a^b \frac{1}{f(\gamma(t)) - z_0} f'(\gamma(t)) \gamma'(t) dt = \int_{f(\Gamma)} \frac{f'(z)}{f(z) - z_0} dz . \\ \Rightarrow \frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z) - z_0} dz &= n(f(\Gamma), z_0) . \end{aligned}$$

Note for $z_0 = 0$, we have

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = n(f(\Gamma), 0) = N_0(f) - N_p(f) .$$

Example 3.11.1.

$$\frac{d}{dz} \log(f(z)) = \frac{f'(z)}{f(z)} ,$$

then

$$\oint_{\Gamma_{i\theta}} \frac{f'(z)}{f(z)} dz = \log(f(z)) \Big|_{z_0}^{z_1} = 2\pi i n(f(\Gamma), 0) .$$

Let $f = re^{i\theta}$,

$$\log(f(z)) = \ln r + i\theta .$$

Lemma 3.11.2 (The Dog-walking Theorem). Let Γ_1, Γ_2 be parametrized by $\gamma_1, \gamma_2 : [a, b] \rightarrow \mathbb{C}$, and $\forall t \in [a, b]$, let $|\gamma_1(t) - \gamma_2(t)| < |\gamma_1(t)|$. Then $n(\Gamma_1, 0) = n(\Gamma_2, 0)$.

Proof. Note that $\gamma_1(t), \gamma_2(t) \neq 0$, let $\Gamma : \gamma(t) = \frac{\gamma_2(t)}{\gamma_1(t)}$. Then

$$\begin{aligned} |1 - \gamma(t)| &= \left| 1 - \frac{\gamma_2(t)}{\gamma_1(t)} \right| \\ &= \left| \frac{\gamma_1(t) - \gamma_2(t)}{\gamma_1(t)} \right| < 1. \end{aligned}$$

Thus Γ lies in $D_1(1)$ so $n(\Gamma, 0) = 0$. Let $\gamma_1 = r_1 e^{i\theta_1}, \gamma_2 = r_2 e^{i\theta_2}$, where $r_1, r_2, \theta_1, \theta_2$ are functions of t . Then

$$\gamma = \frac{\gamma_2}{\gamma_1} = \frac{r_2}{r_1 e^{i(\theta_2 - \theta_1)}}.$$

$$n(\Gamma_1, 0) = \theta_1(b) - \theta_1(a) \quad n(\Gamma_2, 0) = \theta_2(b) - \theta_2(a)$$

so

$$0 = n(\Gamma, 0) = \theta_2(b) - \theta_2(a) - (\theta_1(b) - \theta_1(a)) = n(\Gamma_2, 0) - n(\Gamma_1, 0).$$

So $n(\Gamma_1, 0) = n(\Gamma_2, 0)$. □

Lemma 3.11.3 (Generalized Dog-walking Theorem). Let Γ_1, Γ_2 be parametrized by $\gamma_1, \gamma_2 : [a, b] \rightarrow \mathbb{C}$ and $\forall t \in [a, b]$ let

$$|\gamma_1(t) - \gamma_2(t)| < |\gamma_1(t)| + |\gamma_2(t)|.$$

Then $n(\Gamma_1, 0) = n(\Gamma_2, 0)$.

Proof. Let $\gamma(t) = \frac{\gamma_1(t)}{\gamma_2(t)}$. Assume for contradiction that $\exists c > 0, \exists t \in [a, b], \gamma(t) = -c$. Then $\gamma_1(t) = -c\gamma_2(t)$.

$$|\gamma_1(t) - \gamma_2(t)| = |(-c - 1)\gamma_2(t)| = (c + 1)|\gamma_2(t)|.$$

But $|\gamma_2(t)| + |\gamma_1(t)| = |\gamma_2(t)| + |-c\gamma_2(t)| = (1 + c)|\gamma_2(t)|$. This contradicts the condition of the lemma, so no such c exists. So $\Gamma : \gamma(t)$ lies in the $\mathbb{C} \setminus \mathbb{R}_{\leq 0}$. So $n(\Gamma, 0) = 0$, so $n(\Gamma_1, 0) = n(\Gamma_2, 0)$. □

3.12 July 18

THEOREM 3.12.1 (Rouche's Theorem).

Let f, g be analytic on and inside a simple, positively oriented closed contour Γ . Let $|f(z)| > |g(z)|$ for all $z \in \Gamma$. Then f and $f + g$ have the same number of zeros inside Γ . (counted with multiplicity).

Proof. Let $h = f + g$. Then

$$|h(z) + (-f(z))| = |g(z)| < |-f(z)| \text{ on } \Gamma.$$

Then by argument principle and dog-walking theorem,

$$n(h(\Gamma), 0) = n(f(\Gamma), 0) \Rightarrow N_0(h) = N_0(f).$$

□

Example 3.12.1. All 5 zeros of $h(z) = z^5 + 3z + 1$ lie inside $|z| < 2$.

Example 3.12.2. How many zeros does $z + 3 + 2e^z$ have in the left half-plane $\operatorname{Re}(z) < 0$?

Let Γ_R be the closed left semi circle .

Let $f(z) = z + 3, g(z) = 2e^z$. Then

•

$$|g(z)| = |2e^z| = 2|e^z| = 2e^{\operatorname{Re}(z)}.$$

So $|g(z)| \leq 2$ on Γ_R for all R .

• $|f(z)| = |z + 3|$, so

$$|f(z)| = |z + 3| \geq \begin{cases} |3 + iy|, & z = iy \\ R - e, & |z| = R \end{cases} \geq \begin{cases} 3, & z = iy \\ R - 3, & |z| = R \end{cases}$$

So $\forall R > 5, |f(z)| > |g(z)|$ on Γ_R . Thus f has the same number of zeros inside Γ_R as $z + 3 + 2e^z$, $f(z) = z + 3$ has one zero inside Γ_R , namely 3. So $z + 3 + 2e^z$ has exactly one zero in the left half-plane.

DEFINITION 3.12.2.

A point z is a limit point of a set S if there exists a sequence $\{z_n\}_{n=1}^{\infty} \subseteq S, z_n \neq z, \lim_{n \rightarrow \infty} z_n = z$.

THEOREM 3.12.3.

Let f be holomorphic on a domain D . Let $Z \subseteq D$ be the set of zeros of f in D . if Z has a limit point in D , f is identically zero on D .

Proof. Let z_0 be the limit of $\{w_n\}_{n=1}^\infty \subseteq Z$, $z_0 \neq w_n$ for all n . Consider $D_\varepsilon(z_0)$ for some sufficiently small $\varepsilon > 0$, $f(z) = \sum_{n=0}^\infty a_n(z - z_0)^n$ on $D_\varepsilon(z_0)$.

If f is not identically zero on $D_\varepsilon(z_0)$, then there exists a minimal $m \geq 0$ such that $a_m \neq 0$. Write

$$f(z) = a_m(z - z_0)^m(1 + g(z - z_0)) ,$$

where $g(z - z_0) \rightarrow 0$ as $z \rightarrow z_0$.

Let k be sufficiently large that $w_k \in D_\varepsilon(z_0)$ for all $K \geq k$. Now $f(w_k) = 0$, but

$$0 = f(w_k) = a_m(w_k - z_0)^m(1 + g(w_k - z_0))$$

and $a_m \neq 0$, $(w_k - z_0)^m \neq 0$, and $g(w_k - z_0) \rightarrow 0$ as $w_k \rightarrow z_0$, $k \rightarrow \infty$. So for sufficiently large k ,

$$|g(w_k - z_0)| < 1,$$

so $1 + g(w_k - z_0) \neq 0$. This is a contradiction, so $f = 0$ on $D_\varepsilon(z_0)$.

Let U be the interior of Z . We just showed that U is nonempty. U is open by definition. Let $\{z_n\} \subseteq U$ converging $z_n \rightarrow z$. f is continuous, so $f(z) = 0$. By the earlier argument, $z \in U$. \square

Corollary 3.12.1. Let f, g be analytic on D and $f(z) = g(z)$ on $S \subseteq D$ where S has limit point in D , then $f(z) = g(z)$ on D .

3.13 July 20

THEOREM 3.13.1 (open mapping theorem).

If f is holomorphic on a domain D , then f is an open map on D . (i.e. it maps open set to open sets).

Proof. It suffices to show that $f(D)$ is open. Let $z_0 \in D$, $f(z_0) = w_0$. Let $w \in \mathbb{C}$ and

$$g(z) = f(z) - w = f(z) - w_0 + w_0 - w .$$

Choose $\delta > 0$ such that $D_\delta \subseteq D$ and such that $f(z) \neq w_0$ on the circle $|z - z_0| = \delta$, which exists by the previous corollary.

Choose $\varepsilon > 0$ such that $|f(z) - w_0| \geq \varepsilon$ on $|z - z_0| = \delta$. Now for all $|w - w_0| < \varepsilon$, we have

$$|f(z) - w_0| \geq \varepsilon > |w - w_0|$$

on the circle $|z - z_0| = \delta$.

So by Rouché's Theorem, g and $f(z) - w_0$ have the same number of zeros in $D_\delta(z_0)$, namely by one. Thus $\exists z \in D_\delta(z_0) \subseteq D$, $g(z) = 0 = f(z) - w \Rightarrow f(z) = w \Rightarrow w \in f(D)$. Thus $D_\varepsilon(w) \subseteq f(D)$. \square

Example 3.13.1. Let f be analytic on a domain D and $\operatorname{Re} f(z)$ is constant. Then f is constant. $\operatorname{Re}(f(z)) = K$ contains no open sets. So f must be constant by the contrapositive of the open mapping theorem.

DEFINITION 3.13.2.

The **gamma function** is defined for $s > 0$ in \mathbb{R} by

$$\Gamma(s) = \int_0^\infty e^{-t} t^{s-1} dt .$$

Lemma 3.13.1. Γ extends to an analytic function on $\operatorname{Re}(s) > 0$ and $\Gamma(s) = \int_0^\infty e^{-t} t^{s-1} dt$ still holds here.

Proof. It suffices to show the lemma on

$$S = \{z \in \mathbb{C} : \delta < \operatorname{Re}(s) < M\} \text{ for any } 0 < \delta < M < \infty .$$

Let $\operatorname{Re}(s) = \sigma$. Now

$$\int_0^\infty e^{-t} t^{s-1} dt = \lim_{\varepsilon \rightarrow 0} \int_\varepsilon^{1/\varepsilon} e^{-t} t^{s-1} dt .$$

Let $F_\varepsilon(s) = \int_\varepsilon^{1/\varepsilon} e^{-t} t^{s-1} dt$. Note that $F_\varepsilon(s)$ is analytic. Recall that the limit of a uniformly convergent sequence of analytic functions is analytic, consider

$$\begin{aligned} |\Gamma(s) - F_\varepsilon(w)| &= \left| \int_0^\infty e^{-t} t^{s-1} dt - \int_\varepsilon^{1/\varepsilon} e^{-t} t^{s-1} dt \right| \\ &= \left| \int_0^\varepsilon e^{-t} t^{s-1} dt + \int_{1/\varepsilon}^\infty e^{-t} t^{s-1} dt \right| \\ &\leq \int_0^\varepsilon e^{-t} t^{\sigma-1} dt + \int_{1/\varepsilon}^\infty e^{-t} t^{\sigma-1} dt . \end{aligned}$$

\square

3.14 July 22

Recall

$$\Gamma(s) = \int_0^\infty e^{-t} t^{s-1} dt ,$$

is an analytic function on the half-plane $\operatorname{Re}(s) > 0$.

Lemma 3.14.1. If $\operatorname{Re}(s) > 0$, $\Gamma(s+1) = s\Gamma(s)$.

Proof. Consider

$$\int_\varepsilon^{1/\varepsilon} e^{-t} t^{s-1} dt = \int_\varepsilon^{1/\varepsilon} e^{-t} \frac{1}{s} \frac{d}{dt} (t^s) dt = \frac{1}{s} (e^{-t} t^s \Big|_\varepsilon^{1/\varepsilon} - \int_\varepsilon^{1/\varepsilon} -e^{-t} t^s dt) .$$

Taking $\varepsilon \rightarrow 0$, this is

$$\Gamma(s) = \frac{1}{s} \int_0^\infty e^{-t} t^{(s+1)-1} dt = \frac{1}{s} \Gamma(s+1) .$$

□

NOTE 3.14.1.

Start with $\operatorname{Re}(s) > -1$, let $F_1(s) = \frac{\Gamma(s+1)}{s}$. $\Gamma(s+1)$ is analytic on $\operatorname{Re}(s) > -1$, so $F_1(s)$ is meromorphic on $\operatorname{Re}(s) > -1$ with possible pole at only 0.

Since $\Gamma(0+1) = \Gamma(1) = 1$, $F_1(s)$ has a simple pole of residue 1 at $s = 0$. For $\operatorname{Re}(s) > 0$, $F_1(s) = \frac{\Gamma(s+1)}{s} = \Gamma(s)$.

So $F_1(s)$ is the unique analytic continuation of $\Gamma(s)$ onto $\{\operatorname{Re}(s) > -1\} \setminus \{0\}$.

Repeat, define $F_m(s)$ on $\operatorname{Re}(s) > -m$. $m > 0$, $m \in \mathbb{Z}$ as

$$F_m(s) = \frac{\Gamma(s+m)}{(s+m-1)(s+m-2)\cdots(s)} .$$

F_m is meromorphic on $\operatorname{Re}(s) > -m$ with simple poles at $s = 0, -1, -2, \dots, -m+1$, $F_m(s) = \Gamma(s)$ on $\operatorname{Re}(s) > 0$.

So we have an analytic continuation of γ onto $\mathbb{C} \setminus \{0, -1, -2, \dots\}$. For $m > n$, Γ has residues

$$\begin{aligned} \operatorname{res}_{-n} F_m(s) &= \lim_{s \rightarrow -n} (s+m) \frac{\Gamma(-n+m)}{(-n+m-1)(-n+m-2)\cdots(-n+1)(-n)} \\ &= \frac{\Gamma(-n+m)}{(-n+m)} \end{aligned}$$

THEOREM 3.14.2.

$$\Gamma(s)\Gamma(1-s) = \frac{\pi}{\sin(\pi s)} .$$

Proof. It suffices to prove this on $0 < \operatorname{Re}(s) < 1$.

$$\Gamma(1-s) = \int_0^\infty e^{-u} u^{-s} du$$

□

Lemma 3.14.2. For $0 < \operatorname{Re}(a) < 1$,

$$\int_0^\infty \frac{v^{a-1}}{1+v} dv = \frac{\pi}{\sin(\pi a)}.$$

Proof. Let $v = e^x$.

$$\int_0^\infty \frac{v^{a-1}}{1+v} dv = \int_0^\infty \frac{e^{a-1}x}{1+e^x} e^x dx = \int_{-\infty}^\infty \frac{e^{ax}}{1+e^x} dx.$$

Let $f(z) = \frac{e^{az}}{1+e^z}$ and integrate over

$$\begin{aligned} \left| \int_{\Gamma_2} f(z) dz \right| &= \left| \int_0^{2\pi} \frac{e^a(R+it)}{1+e^{R+it}} dt \right| \\ &\leq C \frac{e^{aR}}{e^R} \sim C e^{(a-1)R} \rightarrow 0 \text{ as } R \rightarrow \infty. \end{aligned}$$

$$\left| \int_{\Gamma_4} f(z) dz \right| \leq C e^{-aR} \rightarrow 0 \text{ as } R \rightarrow \infty.$$

$$\begin{aligned} \int_{\Gamma_1} f(z) dz &= \int_{-R}^R \frac{e^{ax}}{1+e^x} dx. \\ \int_{\Gamma_3} f(z) dz &= \int_{-R}^R \frac{e^{a(x+2\pi i)}}{1+e^{x+2\pi i}} dx = -e^{2\pi ia} \int_{-R}^R \frac{e^{ax}}{1+e^x} dx. \end{aligned}$$

f has a pole at $z = \pi i$.

$$\begin{aligned} \lim_{z \rightarrow \pi i} (z - \pi i) f(z) &= \lim_{z \rightarrow \pi i} (z - \pi i) \frac{e^{az}}{1+e^z} = \lim_{z \rightarrow \pi i} e^{az} \left(\frac{z - \pi i}{e^z - e^{\pi i}} \right) \\ &= e^{a\pi i} \left(\frac{e^z - e^{\pi i}}{z - \pi i} \right)^{-1} \\ &= e^{a\pi i} \left(\frac{d}{dz}(e^z) \Big|_{z=\pi i} \right)^{-1} \\ &= e^{a\pi i} (e^{\pi i})^{-1} = -e^{a\pi i} = \operatorname{res}_{\pi i} f \end{aligned}$$

Thus

$$\int f(z) dz = 2\pi i (-e^{a\pi i}) =$$

□

3.15 July 25

DEFINITION 3.15.1.

Define the Riemann **zeta function** for real $s > 1$ as

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} .$$

ζ immediately has an analytic continuation to $\operatorname{Re}(s) > 1$ and the formula $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$ is still valid.

If $s = \sigma + it$, $\sigma, t \in \mathbb{R}$. If $\sigma > 1 + \delta > 1$, then

$$\left| \sum_{n=1}^{\infty} \frac{1}{n^s} \right| \leq \sum_{n=1}^{\infty} \left| \frac{1}{n^s} \right| = \sum_{n=1}^{\infty} \left| \frac{1}{e^{s \log n}} \right| = \sum_{n=1}^{\infty} \frac{1}{e^{\sigma \log n}} = \sum_{n=1}^{\infty} \frac{1}{n^{\sigma}} \leq \sum_{n=1}^{\infty} \frac{1}{n^{1+\delta}} \text{ uniformly .}$$

Thus $\zeta(s)$ is analytic on $\operatorname{Re}(s) > 1$.

Consider the Euler product

$$\prod_{p \text{ prime}} \frac{1}{p^{-s}} .$$

For $\operatorname{Re}(s) > 1$.

$$\frac{1}{1 - p^{-s}} = \sum_{n=1}^{\infty} \frac{1}{p^{ns}} .$$

Thus **unique prime factorization**

THEOREM 3.15.2.

$\zeta(s) - \frac{1}{s-1}$ has an analytic continuation to $\operatorname{Re}(s) > 0$. Thus $\zeta(s)$ is meromorphic on $\operatorname{Re}(s) > 0$ with a simple pole at $s = 1$ with residue 1.

Proof. Consider

$$\sum_{1 \leq n < N} \frac{1}{n^s} - \int_1^N \frac{1}{x^s} dx .$$

Let $\zeta_n(s) = \int_n^{n+1} \frac{1}{x^s} - \frac{1}{x^s} dx$. By the mean value theorem

$$\left| \frac{1}{n^s} - \frac{1}{x^s} \right| \leq \frac{|s|}{n^{\sigma+1}} , \text{ on } n \leq x \leq n+1 .$$

Thus we have uniform convergence on $\delta_n(s)$ on $1 + \sigma > 0 \iff \operatorname{Re}(s) > 0$. Thus $\sum_{n=1}^{\infty} \delta_n(s)$ is analytic on $\operatorname{Re}(s) > 0$. Now

$$\sum_{1 \leq n < N} \frac{1}{n^s} = \sum_{n=1}^N \delta_n(s) + \int_1^N \frac{1}{x^s} dx .$$

Now

$$\lim_{n \rightarrow \infty} \int_1^N \frac{1}{x^s} dx = \int_1^\infty \frac{1}{x^s} dx = \frac{1}{1-s} x^{1-s} \Big|_1^\infty.$$

On $\operatorname{Re}(s) > 1$, this $= \frac{1}{s-1}$. Thus $\sum_{n=1}^N \delta_n(s) + \int_1^N \frac{1}{x^s} dx$ converges uniformly and matches $\zeta(s)$ on $\operatorname{Re}(s) > 1$, so $\zeta(s) - \frac{1}{s-1} = \sum_{n=1}^\infty \delta_n(s)$ is analytic on $\operatorname{Re}(s) > 0$.

Thus $\zeta(s)$ is meromorphic on $\operatorname{Re}(s) > 0$ with a simple pole of residue 1 at $s = 1$. This argument can be extended to get ζ meromorphic on \mathbb{C} . \square

THEOREM 3.15.3.

$\zeta(s)$ has no zeros on the line $\operatorname{Re}(s) = 1$.

Proof. Let $x, y \in \mathbb{R}$, $y \neq 0$ and define $h(x) = \zeta(x)^3, \zeta(s+iy)^4 \zeta(x+2iy)$. Now

$$\zeta(s) = \prod_p \frac{1}{1-p^{-s}}$$

So

$$\ln |\zeta(s)| = \ln \prod_p \left| \frac{1}{1-p^{-s}} \right| = - \sum_p \ln |1-p^{-s}| = -\operatorname{Re} \sum_p \operatorname{Log}(1-p^{-s}).$$

Now

$$-\operatorname{Log}(1-w) = \sum_{n=1}^\infty \frac{w^n}{n} \text{ for } |w| < 1.$$

so $\ln |\zeta(s)| = \operatorname{Re} \sum_p \sum_n \frac{1}{n} p^{-sn}$. Thus

$$\begin{aligned} \ln |h(x)| &= 3 \ln |\zeta(x)| + 4 \ln |\zeta(x+iy)| + \ln |\zeta(x+2iy)| \\ &= 3 \operatorname{Re} \sum_p \sum_n \frac{1}{n} p^{-sn} + 4 \operatorname{Re} \sum_p \sum_n \frac{1}{n} p^{-s} \end{aligned}$$

\square

3.16 Final Review

Let $i^2 = -1$. Define

$$f'(z) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}.$$

If f is holomorphic on a domain D , f is infinitely differentiable on D .

THEOREM 3.16.1.

Let f be continuous on a simply connected domain D . Then TFAE:

- f is holomorphic on D

- $\oint_{\Gamma} f = 0$ for any closed $\Gamma \subseteq D$.
- f has a primitive on D
- $\int_{\Gamma_1} f = \int_{\Gamma_2} f$ for $\Gamma_1, \Gamma_2 \subseteq D$ sharing initial and terminal points.

Cauchy's Integral Formula:

$$f^{(n)}(z) = \frac{n!}{2\pi i} \oint \frac{f(w)}{(z-w)^{n+1}} dw .$$

Maximum Modulus Principle: a nonconstant analytic function can only at the boundary of a region.

Liouville's Theorem: A bounded entire function is constant.

Fundamental Theorem of Algebra: every polynomial in \mathbb{C} has a zero in \mathbb{C} .

A function holomorphic in an open disk has a Taylor series representation in that disk.

A function analytic in an annulus has a laurent series representaion in the annulus.

Classification of Singularities:

- Removable singularities: $\lim_{z \rightarrow z_0} |f(z)| < \infty$.
- Pole: $\lim_{z \rightarrow z_0} |f(z)| = \infty$.
- Essential singularities: $\lim_{z \rightarrow z_0} |f(z)|$ erratic.

Order-m pole: $(z - z_0)^m f(z)$ is analytic at z_0

Residue:

$$\text{res}_{z_0} f = \frac{1}{2\pi i} \oint_{\Gamma} f(z) dz .$$

Computing Residues: $\text{res}_{z_0} = a_{-1}$, where $\sum_{n=-m}^{\infty} a_n (z - z_0)^n$ is the Laurent series in a punctured disk.

$$\text{res}_{z_0} f = (m-1)! \lim_{z \rightarrow z_0} \frac{d^{m-1}}{dz^{m-1}} ((z - z_0)^m f(z)) .$$

If $m = 1$,

$$\lim_{z \rightarrow z_0} (z - z_0) f(z) = \text{res}_{z_0} f ,$$

if $m = 1$, if $f(z) = \frac{g(z)}{h(z)}$, g, h holomorphic, then $\text{res}_{z_0} f = \frac{g(z)}{h'(z_0)}$.

Lemma 3.16.1. If z_0 is a simple pole,

$$\int_{\Gamma} f = i(\theta_2 - \theta_1) \operatorname{res}_{z_0} f .$$

$$\lim_{R \rightarrow \infty} \int_{C_R} \frac{p(z)}{q(z)} dz = 0 \text{ if } \deg(q) \geq 2 + \deg(p) .$$

$$\lim_{R \rightarrow \infty} \int_{C_R} e^{iaz} \frac{p(z)}{q(z)} dz = 0 \text{ if } \deg(q) \geq 1 + \deg(p) .$$

THEOREM 3.16.2 (The Argument Principle).

$$\frac{1}{2\pi i} \int_{\gamma} f' = N_0(f) - N_p(f) .$$