

# Disease → Medicine RAG Knowledge Base

Curated disease-to-medicine reference • 157 disease/condition records • Generated 2026-10-02

**Purpose:** A RAG-oriented reference connecting diseases/conditions with commonly used medicines or therapeutic classes. This is an educational knowledge base, not a prescribing system.

**Important:** A disease does not always have one universal medicine. Treatment can depend on diagnosis, severity, organism/resistance, age, pregnancy, kidney/liver function, allergies, interactions, comorbidities and local guidelines. Antibiotics and other prescription medicines should not be inferred or prescribed solely from this document.

**Source framework:** WHO's 2025 Model List of Essential Medicines is a guide to priority medicines and is updated every two years; WHO notes that essential medicines represent only a small proportion of medicines available globally. NIH/NLM MedlinePlus provides consumer medication information including uses, precautions and side effects.

Recommended RAG fields: disease\_name, disease\_category, meaning, common\_medicines, therapeutic\_classes, treatment\_notes, warnings, source, source\_date, version.

## 1. Common cold / viral upper respiratory infection

**Meaning:** Viral infection of the upper airways.

**Common medicine / therapy examples:** Supportive care; paracetamol; ibuprofen when appropriate; saline nasal preparations.

**RAG safety note:** Antibiotics do not treat uncomplicated viral colds.

## 2. Influenza

**Meaning:** Influenza virus infection.

**Common medicine / therapy examples:** Antivirals such as oseltamivir in appropriate patients; supportive care.

**RAG safety note:** Early antiviral treatment may be important for high-risk or severe cases.

## 3. COVID-19

**Meaning:** Illness caused by SARS-CoV-2.

**Common medicine / therapy examples:** Supportive care; selected high-risk patients may receive antivirals such as nirmatrelvir/ritonavir or remdesivir under current guidance.

**RAG safety note:** Treatment depends on risk, severity, timing, interactions and local guidance.

## 4. Strep throat

**Meaning:** Bacterial pharyngitis, commonly caused by *Streptococcus pyogenes*.

**Common medicine / therapy examples:** Penicillin or amoxicillin when confirmed/clinically indicated.

**RAG safety note:** Antibiotics should not be used for viral sore throat.

## 5. Acute sinusitis

**Meaning:** Inflammation/infection of the paranasal sinuses.

**Common medicine / therapy examples:** Many cases are supportive-care conditions; selected bacterial cases may use amoxicillin/clavulanate.

**RAG safety note:** Distinguish viral from bacterial disease.

## 6. Pneumonia

**Meaning:** Infection/inflammation of lung tissue.

**Common medicine / therapy examples:** Antibiotic regimen depends on organism, severity and setting; examples include amoxicillin, ceftriaxone or azithromycin in selected contexts.

**RAG safety note:** Severe disease may require hospital treatment.

## 7. Tuberculosis

**Meaning:** Infection caused by *Mycobacterium tuberculosis*.

**Common medicine / therapy examples:** Combination regimens including rifampicin, isoniazid, pyrazinamide and ethambutol for drug-susceptible disease.

**RAG safety note:** Must follow a TB program and susceptibility/local guidance.

## 8. Malaria

**Meaning:** Parasitic infection transmitted by *Anopheles* mosquitoes.

**Common medicine / therapy examples:** Artemisinin-based combination therapies; severe malaria commonly requires injectable artesunate.

**RAG safety note:** Species, resistance, severity and pregnancy affect treatment.

## 9. Dengue

**Meaning:** Mosquito-borne viral infection.

**Common medicine / therapy examples:** Supportive care; paracetamol for fever/pain when appropriate.

**RAG safety note:** Avoid aspirin/NSAIDs when bleeding risk is relevant.

## 10. Cholera

**Meaning:** Acute diarrheal infection caused by *Vibrio cholerae*.

**Common medicine / therapy examples:** Oral rehydration solution; IV fluids when severe; selected antibiotics.

**RAG safety note:** Rapid rehydration is central.

## 11. Typhoid fever

**Meaning:** Systemic infection usually caused by *Salmonella Typhi*.

**Common medicine / therapy examples:** Antibiotic selection should follow resistance patterns and local guidance; ceftriaxone or azithromycin may be used in selected settings.

**RAG safety note:** Resistance is important.

## 12. Urinary tract infection

**Meaning:** Infection of the urinary tract.

**Common medicine / therapy examples:** Examples include nitrofurantoin, fosfomycin or other guideline-directed antibiotics depending on site and susceptibility.

**RAG safety note:** Pyelonephritis requires different assessment.

## 13. Bacterial meningitis

**Meaning:** Serious bacterial infection of meninges.

**Common medicine / therapy examples:** Urgent hospital therapy commonly includes ceftriaxone/cefotaxime plus other agents according to age and risk.

**RAG safety note:** Medical emergency.

## 14. HIV infection

**Meaning:** Chronic infection caused by human immunodeficiency virus.

**Common medicine / therapy examples:** Combination antiretroviral therapy, commonly using an integrase inhibitor-based regimen.

**RAG safety note:** Regimen selection requires current guidelines and interaction review.

## 15. Hepatitis B

**Meaning:** Liver infection caused by hepatitis B virus.

**Common medicine / therapy examples:** Selected chronic cases may use tenofovir or entecavir.

**RAG safety note:** Not every infection requires antiviral treatment.

## 16. Hepatitis C

**Meaning:** Liver infection caused by hepatitis C virus.

**Common medicine / therapy examples:** Direct-acting antiviral combinations such as sofosbuvir/velpatasvir in appropriate patients.

**RAG safety note:** Regimen depends on prior treatment, liver status and interactions.

## 17. Herpes simplex

**Meaning:** Viral infection caused by HSV.

**Common medicine / therapy examples:** Acyclovir, valacyclovir or famciclovir.

**RAG safety note:** Treatment differs for first episode, recurrence and severe disease.

## 18. Varicella / chickenpox

**Meaning:** Primary infection with varicella-zoster virus.

**Common medicine / therapy examples:** Supportive care; acyclovir in selected high-risk/severe cases.

**RAG safety note:** Avoid inappropriate aspirin use in children.

## 19. Shingles / herpes zoster

**Meaning:** Reactivation of varicella-zoster virus.

**Common medicine / therapy examples:** Acyclovir, valacyclovir or famciclovir; pain management.

**RAG safety note:** Early treatment can be important.

## 20. Fungal skin infection / tinea

**Meaning:** Dermatophyte fungal infection of skin, hair or nails.

**Common medicine / therapy examples:** Topical terbinafine or azole antifungals; systemic therapy for selected extensive infections.

**RAG safety note:** Nail/scalp disease often needs different therapy.

## 21. Candidiasis

**Meaning:** Infection/overgrowth caused by Candida species.

**Common medicine / therapy examples:** Topical azoles, nystatin or fluconazole depending on site and severity.

**RAG safety note:** Recurrent or invasive disease requires evaluation.

## 22. Scabies

**Meaning:** Infestation by Sarcoptes scabiei mites.

**Common medicine / therapy examples:** Permethrin 5% topical treatment; ivermectin in selected situations.

**RAG safety note:** Treat close contacts/environment according to guidance.

## 23. Head lice

**Meaning:** Infestation by Pediculus humanus capitis.

**Common medicine / therapy examples:** Permethrin or other approved pediculicides.

**RAG safety note:** Resistance and reinfestation can occur.

## 24. Acne vulgaris

**Meaning:** Inflammatory disorder of pilosebaceous units.

**Common medicine / therapy examples:** Benzoyl peroxide, topical retinoids, topical antibiotics, or systemic therapies for selected cases.

**RAG safety note:** Treatment depends on severity; avoid antibiotic monotherapy where inappropriate.

## 25. Eczema / atopic dermatitis

**Meaning:** Chronic inflammatory skin disorder.

**Common medicine / therapy examples:** Emollients; topical corticosteroids; topical calcineurin inhibitors; systemic/biologic therapy for selected severe disease.

**RAG safety note:** Skin care and trigger control are important.

## 26. Psoriasis

**Meaning:** Chronic immune-mediated inflammatory skin disease.

**Common medicine / therapy examples:** Topical corticosteroids, vitamin-D analogues, phototherapy, systemic or biologic medicines depending on severity.

**RAG safety note:** Requires individualized treatment.

## 27. Allergic rhinitis

**Meaning:** Allergic inflammation of nasal passages.

**Common medicine / therapy examples:** Intranasal corticosteroids; antihistamines such as cetirizine or loratadine.

**RAG safety note:** Avoid unnecessary sedating medicines when alertness matters.

## 28. Asthma

**Meaning:** Chronic inflammatory airway disease with variable obstruction.

**Common medicine / therapy examples:** Inhaled corticosteroid-containing therapy; bronchodilators such as salbutamol for symptom relief when appropriate.

**RAG safety note:** Controller therapy is central; acute severe asthma needs urgent care.

## 29. COPD

**Meaning:** Chronic obstructive pulmonary disease.

**Common medicine / therapy examples:** Inhaled bronchodilators such as tiotropium or LABA/LAMA combinations; inhaled corticosteroids for selected patients.

**RAG safety note:** Smoking cessation and vaccination are important.

## 30. Gastroesophageal reflux disease

**Meaning:** Backflow of stomach contents causing reflux symptoms.

**Common medicine / therapy examples:** Proton-pump inhibitors such as omeprazole; H2 blockers in selected cases.

**RAG safety note:** Persistent symptoms may need evaluation.

## 31. Peptic ulcer disease

**Meaning:** Ulceration of stomach or duodenal mucosa.

**Common medicine / therapy examples:** PPI therapy; H. pylori eradication uses combination antibiotic/PPI regimens.

**RAG safety note:** Treatment depends on cause and local resistance.

## 32. H. pylori infection

**Meaning:** Gastric infection caused by Helicobacter pylori.

**Common medicine / therapy examples:** Combination eradication therapy containing acid suppression plus antibiotics and sometimes bismuth.

**RAG safety note:** Regimens vary by resistance and prior exposure.

## 33. Nausea/vomiting

**Meaning:** Symptom with multiple possible causes.

**Common medicine / therapy examples:** Oral rehydration; ondansetron or other antiemetics in selected cases.

**RAG safety note:** Underlying cause must be considered.

### 34. Diarrhea, acute infectious

**Meaning:** Frequent loose stools, often infectious.

**Common medicine / therapy examples:** Oral rehydration solution; selected cases may use zinc in children; antibiotics only for selected infections.

**RAG safety note:** Avoid unnecessary antibiotics/antidiarrheals in some infections.

### 35. Constipation

**Meaning:** Infrequent or difficult bowel movements.

**Common medicine / therapy examples:** Fiber; polyethylene glycol; lactulose; stimulant laxatives when appropriate.

**RAG safety note:** Persistent/new severe constipation needs evaluation.

### 36. Irritable bowel syndrome

**Meaning:** Functional gastrointestinal disorder with abdominal pain and altered bowel habits.

**Common medicine / therapy examples:** Dietary measures; polyethylene glycol for constipation; antispasmodics or selected other agents.

**RAG safety note:** Treatment depends on IBS subtype.

### 37. Hypertension

**Meaning:** Persistently elevated blood pressure.

**Common medicine / therapy examples:** ACE inhibitors/ARBs, calcium-channel blockers, thiazide-type diuretics and other agents.

**RAG safety note:** Choice depends on comorbidities and patient factors.

### 38. Heart failure

**Meaning:** Clinical syndrome caused by impaired cardiac function.

**Common medicine / therapy examples:** Diuretics for congestion; ACEi/ARB/ARNI, beta-blockers, mineralocorticoid antagonists and SGLT2 inhibitors for appropriate patients.

**RAG safety note:** Therapy differs by ejection fraction and clinical status.

### 39. Angina / coronary artery disease

**Meaning:** Reduced myocardial oxygen supply causing ischemic symptoms.

**Common medicine / therapy examples:** Antiplatelet therapy, statins, beta-blockers, nitrates and other therapies as indicated.

**RAG safety note:** Chest pain can be an emergency.

### 40. Heart attack / myocardial infarction

**Meaning:** Acute coronary artery blockage causing myocardial injury.

**Common medicine / therapy examples:** Urgent reperfusion plus antiplatelets, anticoagulation and other hospital therapies.

**RAG safety note:** Emergency; treatment is time-sensitive.

### 41. Atrial fibrillation

**Meaning:** Irregular atrial heart rhythm.

**Common medicine / therapy examples:** Rate-control drugs such as beta-blockers; anticoagulation such as apixaban in appropriate patients.

**RAG safety note:** Stroke-risk assessment guides anticoagulation.

## 42. High cholesterol

**Meaning:** Elevated blood lipids, especially LDL cholesterol.

**Common medicine / therapy examples:** Statins such as atorvastatin or rosuvastatin; other lipid-lowering agents in selected patients.

**RAG safety note:** Risk level determines treatment intensity.

## 43. Type 2 diabetes

**Meaning:** Metabolic disease with insulin resistance and progressive beta-cell dysfunction.

**Common medicine / therapy examples:** Metformin, SGLT2 inhibitors, GLP-1 receptor agonists, DPP-4 inhibitors, sulfonylureas and/or insulin depending on circumstances.

**RAG safety note:** Treatment should be individualized.

## 44. Type 1 diabetes

**Meaning:** Autoimmune destruction of pancreatic beta cells causing insulin deficiency.

**Common medicine / therapy examples:** Insulin therapy is essential.

**RAG safety note:** Insulin must be individualized and monitored.

## 45. Hypothyroidism

**Meaning:** Insufficient thyroid hormone production.

**Common medicine / therapy examples:** Levothyroxine.

**RAG safety note:** Dose is individualized using clinical assessment and thyroid tests.

## 46. Hyperthyroidism

**Meaning:** Excess thyroid hormone production.

**Common medicine / therapy examples:** Antithyroid medicines such as methimazole/carbimazole; beta-blockers for symptoms; definitive therapy in selected cases.

**RAG safety note:** Cause and pregnancy status affect treatment.

## 47. Gout

**Meaning:** Inflammatory arthritis caused by monosodium urate crystals.

**Common medicine / therapy examples:** Acute: NSAIDs, colchicine or corticosteroids; prevention: urate-lowering therapy such as allopurinol.

**RAG safety note:** Choice depends on kidney function and comorbidities.

## 48. Rheumatoid arthritis

**Meaning:** Autoimmune inflammatory arthritis.

**Common medicine / therapy examples:** DMARDs such as methotrexate; biologic/targeted therapies for selected patients.

**RAG safety note:** Early specialist management is important.

## 49. Osteoarthritis

**Meaning:** Degenerative joint disease with pain and stiffness.

**Common medicine / therapy examples:** Exercise/weight management; topical NSAIDs; oral analgesics/NSAIDs when appropriate.

**RAG safety note:** Non-drug management is important.

## 50. Osteoporosis

**Meaning:** Reduced bone strength with increased fracture risk.

**Common medicine / therapy examples:** Bisphosphonates, denosumab, teriparatide and other therapies in selected patients; calcium/vitamin D when indicated.

**RAG safety note:** Fracture risk determines treatment.

## 51. Migraine

**Meaning:** Neurological disorder causing recurrent headache attacks.

**Common medicine / therapy examples:** Acute: NSAIDs, triptans or other agents; prevention: selected beta-blockers, topiramate, CGRP-targeted therapies and others.

**RAG safety note:** Treatment depends on migraine pattern and contraindications.

## 52. Epilepsy

**Meaning:** Disorder characterized by recurrent unprovoked seizures.

**Common medicine / therapy examples:** Antiseizure medicines such as levetiracetam, lamotrigine, valproate and others depending on seizure type.

**RAG safety note:** Drug choice is seizure-type and patient dependent.

## 53. Parkinson disease

**Meaning:** Progressive neurological disorder affecting movement.

**Common medicine / therapy examples:** Levodopa/carbidopa; dopamine agonists, MAO-B inhibitors and other agents in selected patients.

**RAG safety note:** Treatment is individualized.

## 54. Depression

**Meaning:** Mood disorder with persistent depressive symptoms.

**Common medicine / therapy examples:** SSRIs such as sertraline, escitalopram or fluoxetine; other antidepressants in selected cases.

**RAG safety note:** Psychotherapy and other non-drug care may also be important.

## 55. Generalized anxiety disorder

**Meaning:** Persistent excessive anxiety and worry.

**Common medicine / therapy examples:** SSRIs/SNRIs; buspirone in selected patients.

**RAG safety note:** Benzodiazepines are not appropriate for routine long-term use.

## 56. Schizophrenia

**Meaning:** Psychotic disorder with disturbances in thought, perception and behavior.

**Common medicine / therapy examples:** Antipsychotics such as risperidone, olanzapine, aripiprazole or others.

**RAG safety note:** Requires specialist monitoring.

## 57. Bipolar disorder

**Meaning:** Mood disorder involving episodes of mania/hypomania and depression.

**Common medicine / therapy examples:** Lithium, valproate, lamotrigine and selected atypical antipsychotics.

**RAG safety note:** Treatment depends on phase and patient factors.

## 58. ADHD

**Meaning:** Neurodevelopmental disorder with inattention and/or hyperactivity-impulsivity.

**Common medicine / therapy examples:** Stimulants such as methylphenidate; non-stimulants such as atomoxetine or guanfacine in selected patients.

**RAG safety note:** Diagnosis and treatment require appropriate assessment.

## 59. Iron-deficiency anemia

**Meaning:** Anemia caused by insufficient iron.

**Common medicine / therapy examples:** Oral iron preparations such as ferrous sulfate; IV iron for selected cases.

**RAG safety note:** Cause of iron deficiency should be investigated.

## 60. Vitamin B12 deficiency

**Meaning:** Deficiency causing hematologic and neurological problems.

**Common medicine / therapy examples:** Vitamin B12 replacement, oral or injectable depending on cause.

**RAG safety note:** Neurological symptoms need prompt assessment.

## 61. Folate deficiency

**Meaning:** Deficiency of folate causing megaloblastic anemia.

**Common medicine / therapy examples:** Folic acid.

**RAG safety note:** Vitamin B12 deficiency should be excluded before folate-only treatment.

## 62. Vitamin D deficiency

**Meaning:** Low vitamin D status that may affect bone health.

**Common medicine / therapy examples:** Vitamin D supplementation when clinically indicated.

**RAG safety note:** Dose depends on deficiency severity and clinical context.

## 63. Iron overload / selected thalassemia

**Meaning:** Excess body iron, often from transfusions or inherited disorders.

**Common medicine / therapy examples:** Iron chelators such as deferasirox or deferoxamine in appropriate patients.

**RAG safety note:** Specialist monitoring required.

## 64. Sickle cell disease

**Meaning:** Inherited hemoglobin disorder causing vaso-occlusive and hemolytic complications.

**Common medicine / therapy examples:** Hydroxyurea; analgesia and other disease-specific therapies; transfusion or newer agents in selected patients.

**RAG safety note:** Specialist care is important.

## 65. Benign prostatic hyperplasia

**Meaning:** Non-cancerous enlargement of the prostate causing urinary symptoms.

**Common medicine / therapy examples:** Alpha-blockers such as tamsulosin; 5-alpha-reductase inhibitors such as finasteride in selected patients.

**RAG safety note:** Symptoms may have other causes.

## 66. Erectile dysfunction

**Meaning:** Difficulty achieving or maintaining an erection.

**Common medicine / therapy examples:** PDE-5 inhibitors such as sildenafil or tadalafil when appropriate.

**RAG safety note:** Do not combine PDE-5 inhibitors with nitrates.

## 67. Vaginal candidiasis

**Meaning:** Candida infection of the vagina.

**Common medicine / therapy examples:** Topical azole antifungals or fluconazole in selected non-pregnant patients.

**RAG safety note:** Pregnancy changes treatment choices.

## 68. Bacterial vaginosis

**Meaning:** Alteration of vaginal flora causing discharge/odor.

**Common medicine / therapy examples:** Metronidazole or clindamycin regimens.

**RAG safety note:** Diagnosis should distinguish other causes of discharge.

## 69. Pelvic inflammatory disease

**Meaning:** Upper genital tract infection, often sexually transmitted.

**Common medicine / therapy examples:** Combination antibiotic therapy according to current STI guidelines.

**RAG safety note:** Untreated disease can cause infertility and complications.

## 70. Chlamydia

**Meaning:** Sexually transmitted infection caused by Chlamydia trachomatis.

**Common medicine / therapy examples:** Doxycycline commonly used; pregnancy requires different regimen.

**RAG safety note:** Partner management and testing are important.

## 71. Gonorrhea

**Meaning:** Sexually transmitted infection caused by Neisseria gonorrhoeae.

**Common medicine / therapy examples:** Ceftriaxone is a key treatment in current guidance.

**RAG safety note:** Resistance is an important concern.

## 72. Syphilis

**Meaning:** Sexually transmitted infection caused by Treponema pallidum.

**Common medicine / therapy examples:** Penicillin G is standard therapy for most stages.

**RAG safety note:** Stage, pregnancy and allergy history affect management.

## 73. Endometriosis

**Meaning:** Endometrial-like tissue outside the uterus causing pain and other symptoms.

**Common medicine / therapy examples:** NSAIDs for pain; hormonal therapies; other specialist treatments.

**RAG safety note:** Treatment depends on symptoms and fertility goals.

## 74. Menopause symptoms

**Meaning:** Symptoms associated with decline in ovarian hormone production.

**Common medicine / therapy examples:** Hormone therapy for selected patients; nonhormonal options such as SSRIs/SNRIs or other therapies.

**RAG safety note:** Risks and contraindications must be assessed.

## 75. Glaucoma

**Meaning:** Progressive optic neuropathy often associated with elevated intraocular pressure.

**Common medicine / therapy examples:** Prostaglandin analogues, beta-blocker eye drops and other IOP-lowering medicines.

**RAG safety note:** Requires ophthalmic monitoring.

## 76. Conjunctivitis

**Meaning:** Inflammation of conjunctiva, often viral, bacterial or allergic.

**Common medicine / therapy examples:** Lubricants; allergy medicines for allergic disease; selected bacterial cases may use topical antibiotics.

**RAG safety note:** Many cases are viral and do not need antibiotics.

## 77. Dry eye disease

**Meaning:** Insufficient lubrication or excessive tear evaporation.

**Common medicine / therapy examples:** Artificial tears; selected anti-inflammatory eye treatments.

**RAG safety note:** Persistent symptoms may need ophthalmology review.

## 78. Otitis externa

**Meaning:** Inflammation/infection of the external ear canal.

**Common medicine / therapy examples:** Topical antibiotic/anti-inflammatory ear drops when indicated.

**RAG safety note:** Keep ear canal dry; treatment depends on eardrum status.

## 79. Acute otitis media

**Meaning:** Middle-ear infection, commonly following a respiratory infection.

**Common medicine / therapy examples:** Pain relief; selected bacterial cases may use amoxicillin.

**RAG safety note:** Age, severity and diagnostic certainty affect antibiotic use.

## 80. Dental caries

**Meaning:** Tooth decay caused by acid-producing plaque bacteria.

**Common medicine / therapy examples:** Fluoride toothpaste/preventive fluoride; dental restoration; antibiotics only for spreading infection.

**RAG safety note:** Dental treatment addresses the underlying lesion.

## 81. Gingivitis

**Meaning:** Inflammation of gums due mainly to plaque.

**Common medicine / therapy examples:** Oral hygiene and professional cleaning; antiseptic mouthwash in selected situations.

**RAG safety note:** Dental assessment may be needed.

## 82. Obesity

**Meaning:** Excess body fat associated with health risk.

**Common medicine / therapy examples:** Lifestyle intervention; anti-obesity medicines such as GLP-1 receptor agonists in selected patients; bariatric surgery for selected severe cases.

**RAG safety note:** Treatment is individualized.

### 83. Alcohol use disorder

**Meaning:** Problematic alcohol use causing impairment or distress.

**Common medicine / therapy examples:** Naltrexone, acamprosate or other medicines in selected patients; withdrawal may require supervised treatment.

**RAG safety note:** Abrupt withdrawal can be dangerous.

### 84. Nicotine dependence

**Meaning:** Dependence on nicotine from tobacco or nicotine products.

**Common medicine / therapy examples:** Nicotine replacement therapy, varenicline or bupropion where appropriate.

**RAG safety note:** Behavioral support improves cessation outcomes.

### 85. Anaphylaxis

**Meaning:** Severe, potentially life-threatening allergic reaction.

**Common medicine / therapy examples:** Intramuscular epinephrine/adrenaline is first-line emergency treatment.

**RAG safety note:** Emergency medical care is required.

### 86. Poisoning / overdose

**Meaning:** Harmful exposure to a drug or toxic substance.

**Common medicine / therapy examples:** Antidote depends on toxin; examples include naloxone for opioid toxicity and N-acetylcysteine for acetaminophen poisoning.

**RAG safety note:** Emergency toxicology assessment is required.

### 87. Severe allergic reaction

**Meaning:** Systemic allergic reaction that may progress to anaphylaxis.

**Common medicine / therapy examples:** Epinephrine/adrenaline for anaphylaxis; antihistamines for selected symptoms.

**RAG safety note:** Airway and circulation support may be needed.

### 88. Diabetic ketoacidosis

**Meaning:** Life-threatening metabolic complication of diabetes.

**Common medicine / therapy examples:** IV fluids, insulin and electrolyte replacement under hospital protocols.

**RAG safety note:** Emergency condition.

### 89. Hypoglycemia

**Meaning:** Abnormally low blood glucose.

**Common medicine / therapy examples:** Fast-acting glucose; glucagon for severe episodes when appropriate.

**RAG safety note:** Cause should be identified.

### 90. Acid reflux / heartburn

**Meaning:** Burning discomfort caused by reflux of stomach contents.

**Common medicine / therapy examples:** Antacids, H2 blockers or PPIs depending on severity/frequency.

**RAG safety note:** Alarm symptoms need medical evaluation.

### 91. Hemorrhoids

**Meaning:** Swollen veins/tissues around anus causing pain or bleeding.

**Common medicine / therapy examples:** Fiber, fluids, topical symptom treatments; procedures for selected cases.

**RAG safety note:** Rectal bleeding should not automatically be attributed to hemorrhoids.

## 92. Inflammatory bowel disease

**Meaning:** Chronic inflammatory disease including Crohn disease and ulcerative colitis.

**Common medicine / therapy examples:** Aminosalicylates, corticosteroids, immunomodulators and biologics depending on disease and severity.

**RAG safety note:** Specialist management is usually required.

## 93. Ulcerative colitis

**Meaning:** Chronic inflammatory disease of colon.

**Common medicine / therapy examples:** Mesalamine/5-ASA, corticosteroids for flares, immunomodulators or biologics for selected patients.

**RAG safety note:** Therapy depends on extent and severity.

## 94. Crohn disease

**Meaning:** Chronic inflammatory bowel disease affecting any GI segment.

**Common medicine / therapy examples:** Corticosteroids for selected flares; immunomodulators and biologics for maintenance/severe disease.

**RAG safety note:** Requires specialist management.

## 95. Chronic kidney disease

**Meaning:** Progressive loss of kidney function.

**Common medicine / therapy examples:** ACEi/ARB for selected albuminuric disease; SGLT2 inhibitors in appropriate patients; treatment of anemia/mineral disorders as indicated.

**RAG safety note:** Drug doses often require renal adjustment.

## 96. Kidney stones

**Meaning:** Crystals/stones forming in urinary tract.

**Common medicine / therapy examples:** NSAIDs for renal colic; alpha-blockers in selected ureteral stones; procedures for obstructing/large stones.

**RAG safety note:** Hydration and stone analysis may guide prevention.

## 97. Acute kidney injury

**Meaning:** Sudden decline in kidney function.

**Common medicine / therapy examples:** Treat underlying cause; fluids when indicated; stop/adjust offending medicines under clinical supervision.

**RAG safety note:** Some medicines require urgent adjustment.

## 98. Liver cirrhosis

**Meaning:** Advanced scarring of liver from chronic disease.

**Common medicine / therapy examples:** Cause-specific therapy plus management of complications such as ascites or variceal bleeding.

**RAG safety note:** Many medicines require liver-specific caution.

## 99. Hepatic encephalopathy

**Meaning:** Brain dysfunction associated with severe liver disease.

**Common medicine / therapy examples:** Lactulose; rifaximin in selected recurrent cases.

**RAG safety note:** Underlying triggers require treatment.

## 100. Anemia of chronic disease

**Meaning:** Anemia associated with chronic inflammation or disease.

**Common medicine / therapy examples:** Treat underlying condition; iron or erythropoiesis-stimulating agents in selected cases.

**RAG safety note:** Cause must be established.

## 101. Deep vein thrombosis

**Meaning:** Blood clot in a deep vein.

**Common medicine / therapy examples:** Anticoagulants such as apixaban, rivaroxaban, heparin or warfarin depending on context.

**RAG safety note:** Pulmonary embolism symptoms require emergency care.

## 102. Pulmonary embolism

**Meaning:** Blood clot obstructing pulmonary arteries.

**Common medicine / therapy examples:** Anticoagulation; thrombolysis/procedures for selected severe cases.

**RAG safety note:** Emergency condition.

## 103. Stroke / ischemic stroke

**Meaning:** Brain injury from blocked cerebral blood flow.

**Common medicine / therapy examples:** Urgent reperfusion in eligible patients; antiplatelet therapy and secondary prevention afterward.

**RAG safety note:** Time-sensitive emergency.

## 104. Transient ischemic attack

**Meaning:** Temporary neurological dysfunction caused by transient cerebral ischemia.

**Common medicine / therapy examples:** Urgent evaluation; antiplatelet/statin and risk-factor treatment when appropriate.

**RAG safety note:** Warning sign for stroke.

## 105. Epileptic seizure emergency / status epilepticus

**Meaning:** Prolonged or repeated seizure activity.

**Common medicine / therapy examples:** Emergency benzodiazepine followed by antiseizure therapy under emergency protocols.

**RAG safety note:** Medical emergency.

## 106. Alzheimer disease

**Meaning:** Progressive neurodegenerative disorder causing cognitive decline.

**Common medicine / therapy examples:** Cholinesterase inhibitors such as donepezil; memantine in selected moderate/severe disease.

**RAG safety note:** Benefits and risks vary; specialist assessment is useful.

## 107. Osteomyelitis

**Meaning:** Bone infection.

**Common medicine / therapy examples:** Prolonged pathogen-directed antibiotics; surgery in selected cases.

**RAG safety note:** Culture and imaging often guide therapy.

## 108. Cellulitis

**Meaning:** Bacterial infection of skin and soft tissue.

**Common medicine / therapy examples:** Antibiotics such as cephalexin or other guideline-directed agents.

**RAG safety note:** Rapid progression/systemic symptoms need urgent assessment.

## 109. Abscess

**Meaning:** Localized collection of pus.

**Common medicine / therapy examples:** Incision and drainage is primary for many abscesses; antibiotics for selected cases.

**RAG safety note:** Antibiotics alone may not cure a drainable abscess.

## 110. Burn injury

**Meaning:** Tissue injury from heat, chemicals, electricity or other causes.

**Common medicine / therapy examples:** Pain control, wound care and infection prevention; specialized treatment for severe burns.

**RAG safety note:** Severe burns need emergency/burn-center care.

## 111. Acne rosacea / rosacea

**Meaning:** Chronic inflammatory facial skin disorder.

**Common medicine / therapy examples:** Topical metronidazole, azelaic acid or ivermectin; doxycycline in selected cases.

**RAG safety note:** Triggers and subtype affect treatment.

## 112. Urticaria / hives

**Meaning:** Transient itchy wheals from mast-cell mediator release.

**Common medicine / therapy examples:** Second-generation antihistamines such as cetirizine or loratadine.

**RAG safety note:** Anaphylaxis signs require emergency treatment.

## 113. Motion sickness

**Meaning:** Nausea caused by motion.

**Common medicine / therapy examples:** Meclizine, dimenhydrinate or other agents in selected patients.

**RAG safety note:** Sedation is common with some agents.

## 114. Insomnia

**Meaning:** Difficulty initiating or maintaining sleep.

**Common medicine / therapy examples:** Sleep-behavior interventions first; selected medicines may be used short-term.

**RAG safety note:** Long-term sedative use has risks.

## 115. Chronic pain

**Meaning:** Pain persisting beyond expected healing time.

**Common medicine / therapy examples:** Cause-specific treatment; acetaminophen/NSAIDs in selected cases; neuropathic pain agents such as gabapentin or duloxetine for selected conditions.

**RAG safety note:** Opioids require careful specialist oversight.

## 116. Neuropathic pain

**Meaning:** Pain caused by nerve injury/dysfunction.

**Common medicine / therapy examples:** Duloxetine, amitriptyline, gabapentin or pregabalin in selected patients.

**RAG safety note:** Choice depends on cause and comorbidities.

## 117. Restless legs syndrome

**Meaning:** Urge to move legs with uncomfortable sensations, often worse at rest.

**Common medicine / therapy examples:** Iron replacement if deficient; selected gabapentinoids or dopamine-related treatments.

**RAG safety note:** Iron status and contributing medicines should be assessed.

## 118. Glomerulonephritis

**Meaning:** Inflammatory disease of kidney glomeruli.

**Common medicine / therapy examples:** Cause-specific; may include corticosteroids or immunosuppressants in selected immune-mediated disease.

**RAG safety note:** Specialist evaluation is required.

## 119. Nephrotic syndrome

**Meaning:** Heavy protein loss in urine with edema and other findings.

**Common medicine / therapy examples:** Cause-specific treatment; ACEi/ARB for proteinuria in selected patients; immunosuppression for selected causes.

**RAG safety note:** Treatment depends on underlying disease.

## 120. Polycystic ovary syndrome

**Meaning:** Hormonal/metabolic disorder with ovulatory dysfunction and hyperandrogenism.

**Common medicine / therapy examples:** Lifestyle treatment; combined hormonal contraceptives, metformin or antiandrogens in selected patients.

**RAG safety note:** Treatment depends on goals such as fertility or cycle control.

## 121. Infertility

**Meaning:** Failure to achieve pregnancy after appropriate period of trying.

**Common medicine / therapy examples:** Cause-specific fertility treatment; ovulation induction medicines such as letrozole in selected patients.

**RAG safety note:** Requires evaluation of both partners when applicable.

## 122. Ectopic pregnancy

**Meaning:** Pregnancy implanted outside uterine cavity.

**Common medicine / therapy examples:** Methotrexate for selected stable cases; surgery for others.

**RAG safety note:** Medical emergency if rupture/bleeding occurs.

## 123. Iron-deficiency in pregnancy

**Meaning:** Iron deficiency during pregnancy.

**Common medicine / therapy examples:** Iron supplementation, usually oral; IV iron in selected cases.

**RAG safety note:** Pregnancy-specific evaluation and dosing required.

## 124. Preeclampsia

**Meaning:** Pregnancy-related hypertension with organ involvement.

**Common medicine / therapy examples:** Antihypertensive treatment such as labetalol/nifedipine in selected cases; magnesium sulfate for seizure prevention in severe disease.

**RAG safety note:** Obstetric emergency when severe.

## 125. Postpartum hemorrhage

**Meaning:** Severe bleeding after childbirth.

**Common medicine / therapy examples:** Uterotonics such as oxytocin; tranexamic acid and other emergency measures depending on cause.

**RAG safety note:** Medical emergency.

## 126. Preterm labor

**Meaning:** Labor occurring before term.

**Common medicine / therapy examples:** Antenatal corticosteroids and tocolysis in selected situations; magnesium sulfate for fetal neuroprotection in selected gestations.

**RAG safety note:** Obstetric specialist management.

## 127. Malnutrition / undernutrition

**Meaning:** Insufficient energy or nutrient intake causing impaired health.

**Common medicine / therapy examples:** Nutritional rehabilitation; micronutrient supplementation; therapeutic foods in selected severe cases.

**RAG safety note:** Underlying cause and severity matter.

## 128. Vitamin A deficiency

**Meaning:** Insufficient vitamin A causing visual/immune problems.

**Common medicine / therapy examples:** Vitamin A replacement under appropriate protocols.

**RAG safety note:** High-dose vitamin A can be toxic.

## 129. Scurvy

**Meaning:** Vitamin C deficiency.

**Common medicine / therapy examples:** Vitamin C replacement and nutritional rehabilitation.

**RAG safety note:** Underlying dietary/medical causes should be addressed.

## 130. Rickets / osteomalacia

**Meaning:** Defective bone mineralization, often due to vitamin D deficiency or other causes.

**Common medicine / therapy examples:** Vitamin D and calcium when deficient; cause-specific therapy.

**RAG safety note:** Diagnosis should distinguish different causes.

## 131. Rabies exposure

**Meaning:** Potentially fatal viral infection after exposure to infected animals.

**Common medicine / therapy examples:** Post-exposure prophylaxis: wound care, rabies vaccine and rabies immunoglobulin when indicated.

**RAG safety note:** Urgent public-health assessment is essential.

### 132. Tetanus

**Meaning:** Serious neurologic disease caused by *Clostridium tetani* toxin.

**Common medicine / therapy examples:** Tetanus immune globulin plus wound care and antibiotics in selected cases; vaccination for prevention.

**RAG safety note:** Emergency condition.

### 133. Leprosy

**Meaning:** Chronic infection caused by *Mycobacterium leprae*.

**Common medicine / therapy examples:** Multidrug therapy with rifampicin, dapsone and clofazimine according to disease classification.

**RAG safety note:** Requires prolonged supervised treatment.

### 134. Schistosomiasis

**Meaning:** Parasitic infection caused by *Schistosoma* species.

**Common medicine / therapy examples:** Praziquantel.

**RAG safety note:** Species and exposure history guide treatment.

### 135. Hookworm infection

**Meaning:** Intestinal infection by hookworms.

**Common medicine / therapy examples:** Albendazole or mebendazole.

**RAG safety note:** Treat anemia when present.

### 136. Pinworm infection

**Meaning:** Intestinal infection by *Enterobius vermicularis*.

**Common medicine / therapy examples:** Mebendazole, albendazole or pyrantel; repeat treatment is commonly needed.

**RAG safety note:** Household hygiene reduces reinfection.

### 137. Amebiasis

**Meaning:** Intestinal or extraintestinal infection caused by *Entamoeba histolytica*.

**Common medicine / therapy examples:** Metronidazole/tinidazole followed by a luminal agent such as paromomycin when indicated.

**RAG safety note:** Differentiate from nonpathogenic *Entamoeba* species.

### 138. Giardiasis

**Meaning:** Intestinal infection caused by *Giardia*.

**Common medicine / therapy examples:** Tinidazole, metronidazole or other approved agents.

**RAG safety note:** Hydration is important.

### 139. Toxoplasmosis

**Meaning:** Parasitic infection caused by *Toxoplasma gondii*.

**Common medicine / therapy examples:** Pyrimethamine plus sulfadiazine and folinic acid in selected severe/high-risk cases.

**RAG safety note:** Pregnancy and immune status substantially affect management.

### 140. Leishmaniasis

**Meaning:** Parasitic disease caused by Leishmania species.

**Common medicine / therapy examples:** Treatment varies by species and form; liposomal amphotericin B and other antileishmanial therapies are used in selected cases.

**RAG safety note:** Specialist/public-health guidance is important.

#### 141. Filariasis

**Meaning:** Parasitic infection caused by filarial worms.

**Common medicine / therapy examples:** Diethylcarbamazine or other regimens depending on species and setting.

**RAG safety note:** Treatment requires disease-specific guidance.

#### 142. Meningitis, viral

**Meaning:** Inflammation of meninges caused by viruses.

**Common medicine / therapy examples:** Supportive care; acyclovir for suspected herpesvirus causes in selected patients.

**RAG safety note:** Diagnosis is urgent because bacterial meningitis must be excluded.

#### 143. Encephalitis

**Meaning:** Inflammation of brain tissue, often infectious or autoimmune.

**Common medicine / therapy examples:** Empiric IV acyclovir is commonly used when herpes encephalitis is suspected; cause-specific therapy thereafter.

**RAG safety note:** Medical emergency.

#### 144. Multiple sclerosis

**Meaning:** Immune-mediated disease of central nervous system.

**Common medicine / therapy examples:** Disease-modifying therapies such as interferon beta, glatiramer acetate, or newer agents; corticosteroids for acute relapses.

**RAG safety note:** Specialist management required.

#### 145. Myasthenia gravis

**Meaning:** Autoimmune disorder causing fluctuating skeletal muscle weakness.

**Common medicine / therapy examples:** Pyridostigmine; immunosuppressants; IVIG/plasma exchange for selected crises.

**RAG safety note:** Respiratory weakness is an emergency.

#### 146. Systemic lupus erythematosus

**Meaning:** Systemic autoimmune disease affecting multiple organs.

**Common medicine / therapy examples:** Hydroxychloroquine; corticosteroids and immunosuppressants/biologics depending on organ involvement.

**RAG safety note:** Treatment is individualized.

#### 147. Ankylosing spondylitis

**Meaning:** Inflammatory arthritis affecting spine and sacroiliac joints.

**Common medicine / therapy examples:** NSAIDs; biologic therapies such as TNF or IL-17 inhibitors in selected patients.

**RAG safety note:** Exercise/physiotherapy is important.

#### 148. Sepsis

**Meaning:** Life-threatening organ dysfunction caused by dysregulated response to infection.

**Common medicine / therapy examples:** Immediate source control, IV antibiotics, fluids and organ support according to sepsis protocols.

**RAG safety note:** Emergency condition.

## 149. Shock

**Meaning:** Life-threatening failure of circulation.

**Common medicine / therapy examples:** Cause-specific fluids/blood products and vasopressors such as norepinephrine when indicated.

**RAG safety note:** Emergency resuscitation is required.

## 150. Asthma exacerbation

**Meaning:** Acute worsening of asthma symptoms and airflow limitation.

**Common medicine / therapy examples:** Inhaled/nebulized bronchodilator such as salbutamol plus systemic corticosteroid for moderate/severe attacks.

**RAG safety note:** Severe attacks require urgent care.

## 151. COPD exacerbation

**Meaning:** Acute worsening of COPD symptoms.

**Common medicine / therapy examples:** Short-acting bronchodilators; systemic corticosteroids; antibiotics for selected bacterial exacerbations.

**RAG safety note:** Severity determines need for hospital care.

## 152. Heartburn

**Meaning:** Common symptom of acid reflux.

**Common medicine / therapy examples:** Antacids, H2 blockers or PPIs depending on frequency.

**RAG safety note:** Frequent or alarm symptoms need evaluation.

## 153. Fever

**Meaning:** Elevated body temperature, usually from infection or inflammation.

**Common medicine / therapy examples:** Treat underlying cause; paracetamol or ibuprofen may reduce fever when appropriate.

**RAG safety note:** Fever itself is a symptom, not a diagnosis.

## 154. Headache

**Meaning:** Symptom with many possible causes.

**Common medicine / therapy examples:** Cause-specific treatment; paracetamol/NSAIDs for selected uncomplicated headaches.

**RAG safety note:** Sudden severe/new neurological symptoms require urgent evaluation.

## 155. Back pain

**Meaning:** Pain in back with many possible causes.

**Common medicine / therapy examples:** Activity, physiotherapy and analgesics/NSAIDs when appropriate.

**RAG safety note:** Neurological deficits or red flags need assessment.

## 156. Menstrual pain / dysmenorrhea

**Meaning:** Painful menstruation caused by uterine contractions or underlying disease.

**Common medicine / therapy examples:** NSAIDs; hormonal contraception in selected patients.

**RAG safety note:** Severe/new pain may indicate endometriosis or another condition.

### 157. Motion-related nausea

**Meaning:** Nausea associated with travel or movement.

**Common medicine / therapy examples:** Meclizine, dimenhydrinate or scopolamine in selected patients.

**RAG safety note:** Sedation and contraindications must be considered.

## Implementation notes for RAG

1. Keep one disease/condition as one logical chunk where possible. 2. Keep medicine names normalized to generic names and store brand names as aliases. 3. Preserve source, date and version metadata. 4. For production clinical use, validate every treatment statement against current authoritative guidelines and official drug information. 5. Do not allow the model to invent doses, contraindications, interactions or treatment recommendations when retrieved evidence does not support them.

Reference sources: World Health Organization, 2025 Model List of Essential Medicines and related 2025 Expert Committee report; U.S. National Library of Medicine MedlinePlus drug information. These sources provide authoritative frameworks, but this curated document is not a complete worldwide drug catalogue.