

Leandro Vendramin

# Groups, radical rings and the Yang–Baxter equation

A combinatorial approach to solutions

Tuesday 6<sup>th</sup> July, 2021



*A Dino*

The Yang–Baxter equation (YBE) arose from Yang’s work on statistic mechanics. In 1967 Yang tried to find the eigenfunctions of a one-dimensional fermion gas with delta function interaction. This was a rather difficult problem. He solved it and showed that a crucial identity in the intermediate steps was a matrix equation

$$A(u)B(u+v)A(v) = B(v)A(u+v)B(u).$$

Later, Baxter, in his solution of another problem in physics, the 8-vertex model again used the YBE. In 1980 Faddeev coined the term "Yang-Baxter Equation". A number of exciting developments in physics and mathematics have led to the conclusion that the YBE is a fundamental mathematical structure with connections to various subfields of mathematics such as knot theory, braid theory, operator theory, Hopf algebras, quantum groups, 3-manifolds, the monodromy of differential equations...

I got the feeling that the YBE is the next pervasive algebraic equation after the Jacobi identity.

Compiled: Tuesday 6<sup>th</sup> July, 2021, 20:04.

In Chapter 1 basic definitions and examples of solutions introduced. The main result of the chapter is Theorem 1.11, where the deep relationship between solutions and group actions is studied.

The first part of Chapter 2 provides an introduction to the theory of radical rings. After recalling basic definitions and stating basic properties, the theory of the Jacobson radical is explored. The second part of the chapter is devoted to involutive solutions. One of the main results of this chapter is Rump’s theorem, which states that radical rings produce solutions. In this chapter we also introduce cycle sets, which are structures that turn out to be equivalent to involutive solutions.

In Chapter 4 we introduce the theory of braces. One of the main results of this chapter is Theorem ??, which proves that braces produce arbitrary solutions. We introduce skew cycle sets and prove in Theorem... that skew cycle sets and arbitrary solutions are equivalent.

In Chapter 5 we study 1-cocycles. In Theorem... Sysak’s theorem states that...

Chapter 12.26 is about factorization of groups and braces. First we prove Itô’s theorem in the case of groups: Every group that admits a factorization through two abelian subgroups is meta-abelian.

Thanks: Jingpeng Shen

# Contents

<b>1</b>	<b>The Yang–Baxter equation</b>	<b>1</b>
<b>2</b>	<b>Radical rings</b>	<b>7</b>
<b>3</b>	<b>Racks</b>	<b>23</b>
<b>4</b>	<b>Braces</b>	<b>31</b>
<b>5</b>	<b>Complements</b>	<b>45</b>
<b>6</b>	<b>Nilpotent groups</b>	<b>55</b>
<b>7</b>	<b>Solvable groups</b>	<b>67</b>
<b>8</b>	<b>Factorizations</b>	<b>73</b>
<b>9</b>	<b>The structure brace of a solution</b>	<b>81</b>
<b>10</b>	<b>Bieberbach groups</b>	<b>83</b>
<b>11</b>	<b>Garside groups</b>	<b>85</b>
<b>12</b>	<b>Invariant subgroups</b>	<b>87</b>
<b>13</b>	<b>Multipermutation solutions</b>	<b>97</b>
<b>14</b>	<b>Ordered groups</b>	<b>107</b>
<b>15</b>	<b>Transitive groups</b>	<b>109</b>
<b>16</b>	<b>Regular subgroups</b>	<b>113</b>
<b>17</b>	<b>The transfer map</b>	<b>119</b>

<b>References</b> .....	125
<b>Index</b> .....	129

# Chapter 1

## The Yang–Baxter equation

YB

### 1A

In [26], Drinfeld briefly discuss set-theoretic solutions to the Yang–Baxter equation. He observed that it makes sense to consider the Yang–Baxter equation in the category of sets and that "maybe it would be interesting to study set-theoretical solutions".


**Definition 1.1.** A *set-theoretic solution* to the Yang–Baxter equation (YBE) is a pair  $(X, r)$ , where  $X$  is a set and  $r: X \times X \rightarrow X \times X$  is a bijective map that satisfies

$$(r \times \text{id})(\text{id} \times r)(r \times \text{id}) = (\text{id} \times r)(r \times \text{id})(\text{id} \times r),$$

where, if  $r(x, y) = (\sigma_x(y), \tau_y(x))$ , then

$$\begin{aligned} r \times \text{id}: X \times X \times X &\rightarrow X \times X \times X, & (r \times \text{id})(x, y, z) &= (\sigma_x(y), \tau_y(x), z), \\ \text{id} \times r: X \times X \times X &\rightarrow X \times X \times X, & (\text{id} \times r)(x, y, z) &= (x, \sigma_y(z), \tau_z(y)). \end{aligned}$$

The solution  $(X, r)$  is said to be *finite* if  $X$  is a finite set.



**Figure 1.1:** The Yang–Baxter equation.

fig:braid

For  $n \geq 2$ , the *braid group*  $\mathbb{B}_n$  is defined as the group with generators  $\sigma_1, \dots, \sigma_{n-1}$  and relations

$$\begin{aligned}\sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1} & \text{if } 1 \leq i \leq n-2, \\ \sigma_i \sigma_j &= \sigma_j \sigma_i & \text{if } |i-j| \geq 1.\end{aligned}$$

Let  $(X, r)$  be a set-theoretic solution to the YBE. Write  $X^n = X \times \cdots \times X$  ( $n$ -times). For  $i < n$  let  $r_{i,i+1} = \text{id}_{X^{i-1}} \times r \times \text{id}_{X^{n-i-1}} : X^n \rightarrow X^n$ . Then the map  $\sigma_i \mapsto r_{i,i+1}$  extends to an action of  $\mathbb{B}_n$  on  $X^n$ .

**Example 1.2.** Let  $X$  be a set. Then  $(X, r)$ , where  $r(x, y) = (y, x)$ , is a solution to the YBE. This solution is known as the *trivial solution* over the set  $X$ .

By convention, we write

$$r(x, y) = (\sigma_x(y), \tau_y(x)).$$

lem: YB

**Lemma 1.3.** *Let  $X$  be a non-empty set and  $r : X \times X \rightarrow X \times X$  be a bijective map. Then  $(X, r)$  is a set-theoretic solution to the YBE if and only if*

$$\sigma_x \circ \sigma_y = \sigma_{\sigma_x(y)} \circ \sigma_{\tau_y(x)}, \quad \sigma_{\tau_{\sigma_y(z)}(x)} \tau_z(y) = \tau_{\sigma_{\tau_y(x)}(z)} \sigma_x(y), \quad \tau_z \circ \tau_y = \tau_{\tau_z(y)} \circ \tau_{\sigma_y(z)}$$

for all  $x, y, z \in X$ .

*Proof.* We write  $r_1 = r \times \text{id}$  and  $r_2 = \text{id} \times r$ . We first compute

$$\begin{aligned}r_1 r_2 r_1(x, y, z) &= r_1 r_2(\sigma_x(y), \tau_y(x), z) = r_1(\sigma_x(y), \sigma_{\tau_y(x)}(z), \tau_z \sigma_x(y)) \\ &= (\sigma_{\sigma_x(y)} \sigma_{\tau_y(x)}(z), \tau_{\sigma_{\tau_y(x)}(z)} \sigma_x(y), \tau_z \tau_y(x)).\end{aligned}$$

Then we compute

$$\begin{aligned}r_2 r_1 r_2(x, y, z) &= r_2 r_1(x, \sigma_y(z), \tau_z(y)) = r_2(\sigma_x \sigma_y(z), \tau_{\sigma_y(z)}(x), \tau_z(y)) \\ &= (\sigma_x \sigma_y(z), \sigma_{\tau_{\sigma_y(z)}(x)} \tau_z(y), \tau_{\tau_z(y)} \tau_{\sigma_y(z)}(x))\end{aligned}$$

and the claim follows.  $\square$

If  $(X, r)$  is a solution, by definition the map  $r : X \times X \rightarrow X \times X$  is invertible. By convention, we write

$$r^{-1}(x, y) = (\hat{\sigma}_x(y), \hat{\tau}_y(x)).$$

Note that this implies that

$$x = \hat{\sigma}_{\sigma_x(y)} \tau_y(x), \quad y = \hat{\tau}_{\tau_y(x)} \sigma_x(y).$$

Since  $(X, r^{-1})$  is a solution, Lemma 1.3 implies that the following formulas hold:

$$\hat{\tau}_y \circ \hat{\tau}_x = \hat{\tau}_{\tau_y(x)} \circ \hat{\tau}_{\sigma_x(y)}, \quad \hat{\sigma}_x \circ \hat{\sigma}_y = \hat{\sigma}_{\sigma_x(y)} \circ \hat{\sigma}_{\tau_y(x)}.$$

**Definition 1.4.** A *homomorphism* between the set-theoretic solutions  $(X, r)$  and  $(Y, s)$  is a map  $f : X \rightarrow Y$  such that the diagram



$$\begin{array}{ccc}
X \times X & \xrightarrow{r} & X \times X \\
f \times f \downarrow & & \downarrow f \times f \\
Y \times Y & \xrightarrow{s} & Y \times Y
\end{array}$$

is commutative. An *isomorphism* of solutions is a bijective homomorphism of solutions.

Since we are interested in studying the combinatorics behind set-theoretic solutions to the YBE, it makes sense to study the following family of solutions.

**Definition 1.5.** We say that a solution  $(X, r)$  to the YBE is *non-degenerate* if the maps  $\sigma_x$  and  $\tau_x$  are permutations of  $X$ .

By convention, a *solution* we will mean a non-degenerate solution to the YBE.

lem:LYZ

**Lemma 1.6.** Let  $(X, r)$  be a solution.

1) Given  $x, u \in X$ , there exist unique  $y, v \in X$  such that  $r(x, y) = (u, v)$ .

2) Given  $y, v \in X$ , there exist unique  $x, u \in X$  such that  $r(x, y) = (u, v)$ .

*Proof.* For the first claim take  $y = \sigma_x^{-1}(u)$  and  $v = \tau_y(x)$ . For the second,  $x = \tau_y^{-1}(v)$  and  $u = \sigma_x(y)$ .  $\square$

The bijectivity of  $r$  means that any row determines the whole square. Lemma 1.6 means that any column also determines the whole square, see Figure 1.2.




fig:braid

**Figure 1.2:** Any row or column determines the whole square.

**Example 1.7.** If the map  $(x, y) \mapsto (\sigma_x(y), \tau_y(x))$  satisfies the Yang–Baxter equation, then so does  $(x, y) \mapsto (\tau_x(y), \sigma_y(x))$ .

exa:Lyubashenko

**Example 1.8.** Let  $X$  be a non-empty set and  $\sigma$  and  $\tau$  be bijections on  $X$  such that  $\sigma \circ \tau = \tau \circ \sigma$ . Then  $(X, r)$ , where  $r(x, y) = (\sigma(y), \tau(x))$ , is a non-degenerate solution. This is known as the *permutation solution* associated with permutations  $\sigma$  and  $\tau$ .

exa:Wada

**Example 1.9.** Let  $G$  be a group. Then  $(G, r)$ , where  $r(x, y) = (xy^{-1}x^{-1}, xy^2)$ , is a solution.

exa:Venkov

**Example 1.10.** Let  $G$  be a group. Then  $(G, r)$ , where  $r(x, y) = (xyx^{-1}, x)$ , is a solution.

The now prove the main theorem of the chapter. The result shows an intriguing connection between group actions and non-degenerate solutions. It was proved by Lu, Yan and Zhu.

thm:LYZ

**Theorem 1.11.** *Let  $G$  be a group and let  $\xi : G \times G \rightarrow G$ ,  $\xi(x, y) = x \triangleright y$ , be a left action of  $G$  on itself, and let  $\eta : G \times G \rightarrow G$ ,  $\eta(x, y) = x \triangleleft y$ , be a right action of  $G$  on itself. If the compatibility condition*

$$uv = (u \triangleright v)(u \triangleleft v)$$

*holds for all  $u, v \in G$ , then the pair  $(G, r)$ , where*

$$r : G \times G \rightarrow G \times G, \quad r(u, v) = (u \triangleright v, u \triangleleft v)$$

*is a bijective solution.*

*Proof.* We write  $r_1 = r \times \text{id}$  and  $r_2 = \text{id} \times r$ . Let

$$r_1 r_2 r_1(u, v, w) = (u_1, v_1, w_1), \quad r_2 r_1 r_2(u, v, w) = (u_2, v_2, w_2).$$

The compatibility condition implies that  $u_1 v_1 w_1 = u_2 v_2 w_2$ . So we need to prove that  $u_1 = u_2$  and  $v_1 = v_2$ . From Lemma 1.3 we note that

$$\begin{aligned} u_1 &= (u \triangleright v) \triangleright ((u \triangleleft v) \triangleright w), & v_1 &= (u \triangleleft v) \triangleleft w, \\ u_2 &= u \triangleright (v \triangleright w), & v_2 &= (u \triangleleft (v \triangleright w)) \triangleleft (v \triangleleft w). \end{aligned}$$

Using the compatibility condition and the fact that  $\xi$  is a left action,

$$u_1 = ((u \triangleright v)(u \triangleleft v)) \triangleright w = (uv) \triangleright w = u \triangleright (v \triangleright w) = u_2.$$

Similarly, since  $\eta$  is a right action,

$$v_2 = u \triangleleft ((v \triangleright w)(v \triangleleft w)) = u \triangleleft (vw) = (u \triangleleft v) \triangleleft w = v_1.$$

To prove that  $r$  is invertible we proceed as follows. Write  $r(u, v) = (x, y)$ , thus  $u \triangleright v = x$ ,  $u \triangleleft v = y$  and  $uv = xy$ . Since

$$(y \triangleright v^{-1})u = (y \triangleright v^{-1})(y \triangleleft v^{-1}) = yv^{-1} = x^{-1}u,$$

it follows that  $y \triangleright v^{-1} = x^{-1}$ , i.e.  $v^{-1} = y^{-1} \triangleright x^{-1}$ . Similarly,

$$v(u^{-1} \triangleleft x) = (u^{-1} \triangleright x)(u^{-1} \triangleleft x) = u^{-1}x = vy^{-1}$$

implies that  $u^{-1} = y^{-1} \triangleleft x^{-1}$ . Clearly  $r^{-1} = \zeta \circ (i \times i) \circ r \circ (i \times i) \circ \zeta$ , is the inverse of  $r$ , where  $\zeta(x, y) = (y, x)$  and  $i(x) = x^{-1}$ .  $\square$

**Proposition 1.12.** *Under the assumptions of Theorem 1.11, if  $r(x, y) = (u, v)$ , then*

$$r(x^{-1}, y^{-1}) = (u^{-1}, v^{-1}), \quad r(x^{-1}, u) = (y, v^{-1}), \quad r(v, y^{-1}) = (u^{-1}, x).$$

*Proof.* In the proof of Theorem 1.11 we found that the inverse of  $r$  is given by  $r^{-1} = \zeta \circ (i \times i) \circ r \circ (i \times i) \circ \zeta$ , where  $\zeta(x, y) = (y, x)$  and  $i(x) = x^{-1}$ , it follows that  $r(x^{-1}, y^{-1}) = (u^{-1}, v^{-1})$ . To prove the equality  $r(x^{-1}, u) = (y, v^{-1})$  we proceed as follows. Since  $r(x, y) = (u, v)$ , it follows that  $x \triangleright y = u$ . Then  $x^{-1} \triangleright u = y$  and hence  $r(x^{-1}, u) = (y, z)$  for some  $z \in G$ . Since  $xy = uv$  and  $x^{-1}u = yz$ , it follows that  $yt = yv^{-1}$ . Then  $z = v^{-1}$ . Similarly one proves  $r(v, y^{-1}) = (u^{-1}, x)$ .  $\square$

## Exercises

prob:Wada

**1.1.** Let  $G$  be a group. Prove that  $r(x, y) = (xy^{-1}x^{-1}, xy^2)$  is involutive if and only if  $x^2 = 1$  for all  $x \in G$ .

**1.2.** Let  $X$  be a finite non-empty set and  $r: X \times X \rightarrow X \times X, (x, y) \mapsto (\sigma_x(y), \tau_y(x))$ , be a map. Prove that  $(X, r)$  is a solution if and only if the maps  $\sigma_x: X \rightarrow X$  are bijective for all  $x \in X$ ,  $r^2 = \text{id}_{X \times X}$  and

$$\sigma_x \circ \sigma_{\sigma_x^{-1}(y)} = \sigma_y \circ \sigma_{\sigma_y^{-1}(x)}$$

for all  $x, y \in X$ .

**1.3.** Prove that if  $(X, r)$  be a solution...? FIXME

prob:perm\_group

**1.4.** Let  $(X, r)$  be a solution. Prove that  $\mathcal{G}(X, r) \simeq \langle (\sigma_x, \tau_x^{-1}) : x \in X \rangle$ .

## Notes

The first papers where set-theoretic solutions are studied are those of Etingof, Schedler and Soloviev [29] and Gateva–Ivanova and Van den Bergh [33]. Both papers deal with non-degenerate involutive solutions.

In [26], Drinfeld attributes Example 1.8 to Lyubashenko. Example 1.9 appears in the work of Wada [65]. Proposition 2.56 was proved by Rump [53].

Theorem 1.11 goes back to Lu, Yan and Zhu, see [45]. Similar results can be found in the work of Etingof, Schedler and Soloviev [29] for involutive solutions and in Soloviev’s paper [61].



## Chapter 2

### Radical rings

radical

#### 2A

We will consider rings possibly without identity. Thus a **ring** is an abelian group  $R$  with an associative multiplication  $(x, y) \mapsto xy$  such that  $(x + y)z = xz + yz$  and  $x(y + z) = xy + xz$  for all  $x, y, z \in R$ . If there is an element  $1 \in R$  such that  $x1 = 1x = x$  for all  $x \in R$ , we say that  $R$  is a ring (or a unitary ring). A **subring**  $S$  of  $R$  is an additive subgroup of  $R$  closed under multiplication.

**Example 2.1.**  $2\mathbb{Z} = \{2m : m \in \mathbb{Z}\}$  is a ring.

A **left ideal** (resp. **right ideal**) is a subring  $I$  of  $R$  such that  $rI \subseteq I$  (resp.  $Ir \subseteq I$ ) for all  $r \in R$ . An **ideal** (also two-sided ideal) of  $R$  is a subring  $I$  of  $R$  that is both a left and a right ideal of  $R$ .

**Example 2.2.** If  $I$  and  $J$  are both ideals of  $R$ , then the sum  $I + J = \{x + y : x \in I, y \in J\}$  and the intersection  $I \cap J$  are both ideals of  $R$ . The product  $IJ$ , defined as the additive subgroup of  $R$  generated by  $\{xy : x \in I, y \in J\}$ , is also an ideal of  $R$ .

**Example 2.3.** If  $R$  is a ring, the set  $Ra = \{xa : x \in R\}$  is a left ideal of  $R$ . Similarly, the set  $aR = \{ax : x \in R\}$  is a right ideal of  $R$ . The set  $RaR$ , which is defined as the additive subgroup of  $R$  generated by  $\{xay : x, y \in R\}$ , is an ideal of  $R$ .

**Example 2.4.** If  $R$  is a unitary ring, then  $Ra$  is the left ideal generated by  $a$ ,  $aR$  is the right ideal generated by  $a$  and  $RaR$  is the ideal generated by  $a$ . If  $R$  is not unitary, the left ideal generated by  $a$  is  $Ra + \mathbb{Z}a$ , the right ideal generated by  $a$  is  $aR + \mathbb{Z}a$  and the ideal generated by  $a$  is  $RaR + Ra + aR + \mathbb{Z}a$ .

A ring  $R$  is said to be **simple** if  $R^2 \neq \{0\}$  and the only ideals of  $R$  are  $0$  and  $R$ . The condition  $R^2 \neq \{0\}$  is trivially satisfied in the case of rings with identity, as  $1 \in R^2$ .

**Example 2.5.** Division rings are simple.

Let  $S$  be a unitary ring. Recall that  $M_n(S)$  is the ring of  $n \times n$  square matrices with entries in  $S$ . If  $A = (a_{ij}) \in M_n(S)$  y  $E_{ij}$  is the matrix such that  $(E_{ij})_{kl} = \delta_{ik}\delta_{jl}$ , then

$$E_{ij}AE_{kl} = a_{jk}E_{il} \quad (2.1) \quad \boxed{\text{eq:trick}}$$

for all  $i, j, k, l \in \{1, \dots, n\}$ .

**Exercise 2.6.** If  $D$  is a division ring, then  $M_n(D)$  is simple.

Let  $R$  be a ring. A left  $R$ -module (or module, for short) is an abelian group  $M$  together with a map  $R \times M \rightarrow M$ ,  $(r, m) \mapsto rm$ , such that

$$(r+s)m = rm + sm, \quad r(m+n) = rm + rs, \quad r(sm) = (rs)m$$

for all  $r, s \in R$ ,  $m, n \in M$ . If  $R$  has an identity  $1$  and  $1m = m$  holds for all  $m \in M$ , the module  $M$  is said to be **unitary**. If  $M$  is a unitary module, then  $M = RM \neq \{0\}$ .

The module  $M$  is said to be **simple** if  $RM \neq \{0\}$  and the only submodules of  $M$  are  $0$  and  $M$ . If  $M$  is a simple module, then  $M \neq \{0\}$ .

lemma:simple

**Lemma 2.7.** Let  $M$  be a non-zero module. Then  $M$  is simple if and only if  $M = Rm$  for all  $0 \neq m \in M$ .

*Proof.* Assume that  $M$  is simple. Let  $m \neq 0$ . Since  $Rm$  is a submodule of the simple module  $M$ , either  $Rm = 0$  or  $Rm = M$ . Let  $N = \{n \in M : Rn = 0\}$ . Since  $N$  is a submodule of  $M$  and  $RM \neq \{0\}$ ,  $N = \{0\}$ . Therefore  $Rm = Ms$ , as  $m \neq 0$ . Now assume that  $M = Rm$  for all  $m \neq 0$ . Let  $L$  be a non-zero submodule of  $M$  and let  $0 \neq x \in L$ . Then  $M = L$ , as  $M = Rx \subseteq L$ .  $\square$

**Example 2.8.** Let  $D$  be a division ring and let  $V$  be a non-zero vector space (over  $D$ ). If  $R = \text{End}_D(V)$ , then  $V$  is a simple  $R$ -módulo with  $fv = f(v)$ ,  $f \in R$ ,  $v \in V$ .

exa:I\_k

**Example 2.9.** Let  $n \geq 2$ . If  $D$  is a division ring and  $R = M_n(D)$ , then each

$$I_k = \{(a_{ij}) \in R : a_{ij} = 0 \text{ para } j \neq k\}$$

is an  $R$ -module isomorphic to  $D^n$ . Thus  $M_n(D)$  is a simple ring that is not a simple  $M_n(D)$ -module.

A left ideal  $L$  of a ring  $R$  is said to be **minimal** if  $L \neq \{0\}$  and  $L$  does not strictly contain other left ideals of  $R$ . Similarly one defines right minimal ideals and minimal ideals.

**Example 2.10.** Let  $D$  be a division ring and let  $R = M_n(D)$ . Then  $L = RE_{11}$  is a minimal left ideal.

**Example 2.11.** Let  $L$  be a non-zero left ideal. If  $RL \neq \{0\}$ , then  $L$  is minimal if and only if  $L$  is a simple  $R$ -module.

A left (resp. right) ideal  $L$  of  $R$  is said to be **regular** if there exists  $e \in R$  such that  $r - re \in L$  (resp.  $r - er \in L$ ) for all  $r \in R$ . If  $R$  is a ring with identity, every left (or right) ideal is regular. A left (resp. right) ideal  $I$  of  $R$  is said to be **maximal** if  $I \neq M$  and  $I$  is not properly contained in any other left (resp. right) ideal of  $R$ . A standard application of Zorn's lemma proves that every unitary ring contains a maximal left (or right) ideal. Similarly one defines maximal ideals.

proposition:R/I

**Proposition 2.12.** *Let  $R$  be a ring and  $M$  be a module. Then  $M$  is simple if and only if  $M \simeq R/I$  for some maximal regular left ideal  $I$ .*

*Proof.* Assume that  $M$  is simple. Then  $M = Rm$  for some  $m \neq 0$  by Lemma 2.7. The map  $\phi: R \rightarrow M, r \mapsto rm$ , is an epimorphism of  $R$ -modules, so the first isomorphism theorem implies that  $M \simeq R/\ker \phi$ .

We claim that  $I = \ker \phi$  is a maximal ideal. The correspondence theorem and the simplicity of  $M$  imply that  $I$  is a maximal ideal (because each left ideal  $J$  such that  $I \subseteq J$  yields a submodule of  $R/I$ ).

We claim that  $I$  is regular. Since  $M = Rm$ , there exists  $e \in R$  such that  $m = em$ . If  $r \in R$ , then  $r - re \in I$  since  $\phi(r - re) = \phi(r) - \phi(re) = rm - r(em) = 0$ .

Supongamos ahora que  $L$  es maximal y regular. Por el teorema de la correspondencia,  $R/L$  no tiene submódulos propios no nulos. Veamos entonces que  $R(R/L) \neq 0$ . Si  $R(R/L) = 0$  y  $r \in R$ , entonces, como  $L$  es regular,  $r - re \in L$  y luego  $r \in L$  pues

$$0 = r(e + I) = re + I = r + I,$$

una contradicción a la maximalidad de  $L$ . □

We will now discuss primitive rings.

Let  $R$  be a ring and  $M$  be a left  $R$ -module. For a subset  $N \subseteq M$  we define the **annihilator** of  $N$  as the subset

$$\text{Ann}_R(N) = \{r \in R : rn = 0 \ \forall n \in N\}.$$

**Example 2.13.**  $\text{Ann}_{\mathbb{Z}}(\mathbb{Z}/n) = n\mathbb{Z}$ .

The following exercise is standard.

**Exercise 2.14.** Let  $R$  be a ring and  $M$  be a module. If  $N \subseteq M$  is a subset, then  $\text{Ann}_R(N)$  is a left ideal of  $R$ . If  $N \subseteq M$  is a submodule of  $R$ , then  $\text{Ann}_R(N)$  is an ideal of  $R$ .

A module  $M$  is said to be **faithful** if  $\text{Ann}_R(M) = \{0\}$ .

**Example 2.15.** If  $K$  is a field, then  $K^n$  is a faithful unitary  $M_n(K)$ -module.

**Example 2.16.** If  $V$  is vector space over a field  $K$ , then  $V$  is faithful unitary  $\text{End}_K(V)$ -module.

A ring  $R$  is said to be **primitive** if there exists a faithful simple  $R$ -módulo. Since we are considering left modules, our definition of primitive rings is that of left primitive rings. By convention, a primitive ring will always mean a left primitive ring. The use of right modules yields to the notion of right primitive rings.

proposition:simple=>prim

**Proposition 2.17.** *If  $R$  is a simple unitary ring, then  $R$  is primitive.*

*Proof.* Since  $R$  is unitary, there exists a maximal left ideal  $I$  and, moreover,  $R$  is regular. By Proposition 2.12,  $R/I$  is a simple  $R$ -module. Since  $\text{Ann}_R(R/I)$  is an ideal of  $R$  and  $R$  is simple, either  $\text{Ann}_R(R/I) \in \{0\}$  or  $\text{Ann}_R(R/I) = R$ . Moreover, since  $1 \notin \text{Ann}_R(R/I)$ , it follows that  $\text{Ann}_R(R/I) = \{0\}$ .  $\square$

osition:prim+conm=cuerpo

**Proposition 2.18.** *If  $R$  is a commutative ring, then  $R$  is primitive if and only if  $R$  is a field.*

*Proof.* If  $R$  is a field, then  $R$  is primitive because it is a unitary simple ring, see Proposition 2.17. If  $R$  is a primitive commutative ring, Proposition 2.12 implies that there exists a maximal regular ideal  $I$  such that  $R/I$  is a faithful simple  $R$ -module. Since  $I \subseteq \text{Ann}_R(R/I) = \{0\}$  and  $I$  is regular, there exists  $e \in R$  such that  $r = re = er$ . Therefore  $R$  is a unitary commutative ring. Since  $I = \{0\}$  is a maximal ideal,  $R$  is a field.  $\square$

**Example 2.19.** The ring  $\mathbb{Z}$  is not primitive.

An ideal  $P$  of a ring  $R$  is said to be **primitive** if  $P = \text{Ann}_R(M)$  for some simple  $R$ -module  $M$ .

lemma:primitivo

**Lemma 2.20.** *Let  $R$  be a ring and  $P$  be an ideal of  $R$ . Then  $P$  is primitive if and only if  $R/P$  is a primitive ring.*

*Proof.* Assume that  $P = \text{Ann}_R(M)$  for some  $R$ -module  $M$ . Then  $M$  is a simple  $R/P$ -module with  $(r+P)m = rm$ ,  $r \in R$ ,  $m \in M$ . This is well-defined, as  $P = \text{Ann}_R(M)$ . Since  $M$  is a simple  $R$ -module, it follows that  $M$  is a simple  $R/P$ -module. Moreover,  $\text{Ann}_{R/P} M = \{0\}$ . Indeed, if  $(r+P)M = 0$ , then  $r \in \text{Ann}_R M = P$  and hence  $r+P = P$ .

Assume now that  $R/P$  is primitive. Let  $M$  be a faithful simple  $R/P$ -module. Then  $rm = (r+P)m$ ,  $r \in R$ ,  $m \in M$ , turns  $M$  into an  $R$ -module. It follows that  $M$  is simple and that  $P = \text{Ann}_R(M)$ .  $\square$

**Example 2.21.** Let  $R_1, \dots, R_n$  be primitive ring and  $R = R_1 \times \dots \times R_n$ . Then each  $P_i = R_1 \times \dots \times R_{i-1} \times \{0\} \times R_{i+1} \times \dots \times R_n$  is a primitive ideal of  $R$  since  $R/P_i \simeq R_i$ .

lemma:maxprim

**Lemma 2.22.** *Let  $R$  be a ring. Si  $P$  es un ideal primitivo, existe un ideal a izquierda  $L$  maximal tal que  $P = \{x \in R : xR \subseteq L\}$ . Recíprocamente, si  $L$  es un ideal a izquierda maximal y regular, entonces  $\{x \in R : xR \subseteq L\}$  es un ideal primitivo.*

*Proof.* Assume that  $P = \text{Ann}_R(M)$  for some simple  $R$ -module  $M$ . By Proposition 2.12, there exists a regular maximal left ideal  $L$  such that  $M \simeq R/L$ . Then  $P = \text{Ann}_R(R/L) = \{x \in R : xR \subseteq L\}$ .



Conversely, let  $L$  a regular maximal left ideal. By Proposition 2.12,  $R/L$  is a simple  $R$ -module simple. Then

$$\text{Ann}_R(R/L) = \{x \in R : xR \subseteq L\}$$

if a primitive ideal.  $\square$

**Proposition 2.23.** *Maximal ideals of unitary rings are primitive.*

*Proof.* Let  $R$  be a ring with identity and  $M$  be a maximal ideal of  $R$ . Then  $R/M$  is a simple unitary ring by Proposition 2.12. Then  $R/M$  is primitive by Proposition 2.17. By lemma 2.20,  $M$  is primitive.  $\square$

**Exercise 2.24.** Prove that every primitive ideal of a commutative ring is maximal.

**Exercise 2.25.** Prove that  $M_n(R)$  is primitive if and only if  $R$  is primitive.

Let us discuss the Jacobson radical and radical rings.

Let  $R$  be a ring. The **Jacobson radical**  $J(R)$  is the intersection of all the annihilators of simple left  $R$ -modules. If  $R$  does not have simple left  $R$ -modules, then  $J(R) = R$ . From the definition it follows that  $J(R)$  is an ideal. Moreover,

$$J(R) = \bigcap \{P : P \text{ left primitive ideal}\}.$$

If  $I$  is an ideal of  $R$  and  $n \in \mathbb{N}$ ,  $I^n$  is the additive subgroup of  $R$  generated by the set  $\{y_1 \dots y_n : y_j \in I\}$ . An ideal  $I$  of  $R$  is **nilpotent** if  $I^n = \{0\}$  for some  $n \in \mathbb{N}$ . Similarly one defines right or left nil ideals. Note that an ideal  $I$  is nilpotent if and only if there exists  $n \in \mathbb{N}$  such that  $x_1 x_2 \dots x_n = 0$  for all  $x_1, \dots, x_n \in I$ .

An element  $x$  of a ring is said to be **nil** (or nilpotent) if  $x^n = 0$  for some  $n \in \mathbb{N}$ . An ideal  $I$  of a ring is said to be nil if every element of  $I$  is nil. Every nilpotent ideal is nil, as  $I^n = 0$  implies  $x^n = 0$  for all  $x \in I$ .

**Example 2.26.** Let  $R = \mathbb{C}[x_1, x_2, \dots] / (x_1, x_2^2, x_3^3, \dots)$ . The ideal  $I = (x_1, x_2, x_3, \dots)$  is nil in  $R$ , as it is generated by nilpotent element. However, it is not nilpotente. Indeed, if  $I$  is nilpotent, then there exists  $k \in \mathbb{N}$  such that  $I^k = 0$  and hence  $x_i^k = 0$  for all  $i$ , a contradiction since  $x_{k+1}^k \neq 0$ .

pro:nilJ

**Proposition 2.27.** *Let  $R$  be a ring. Then every nil left ideal (resp. right ideal) is contained in  $J(R)$ .*

*Proof.* Assume that there is a nil left ideal (resp. right ideal)  $I$  such that  $I \not\subseteq J(R)$ . There exists a simple  $R$ -module  $M$  such that  $n = xm \neq 0$  for some  $x \in I$  and some  $m \in M$ . Since  $M$  is simple,  $Rn = M$  and hence there exists  $r \in R$  such that

$$(rx)m = r(xm) = rn = m \quad (\text{resp. } (xr)n = x(rn) = xm = n).$$

Thus  $(rx)^k m = m$  (resp.  $(xr)^k n = n$ ) for all  $k \geq 1$ , a contradiction since  $rx \in I$  (resp.  $xr \in I$ ) is a nilpotent element.  $\square$

Let  $R$  be a ring. An element  $a \in R$  is said to be **left quasi-regular** if there exists  $r \in R$  such that  $r + a + ra = 0$ . Similarly,  $a$  is said to be **right quasi-regular** if there exists  $r \in R$  such that  $a + r + ar = 0$ .

exercise:circ

**Exercise 2.28.** Let  $R$  be a ring. Prove that  $R \times R \rightarrow R$ ,  $(r, s) \mapsto r \circ s = r + s + rs$ , is an associative operation with neutral element 0.

**Exercise 2.29.** Let  $R = \mathbb{Z}/3 = \{0, 1, 2\}$ . Compute the multiplication table with respect to the circle operation given by the previous exercise.

If  $R$  is unitary, an element  $x \in R$  is left quasi-regular (resp. right quasi-regular) if and only if  $1 + x$  is left invertible (resp. right invertible). In fact, if  $r \in R$  is such that  $r + x + rx = 0$ , then  $(1 + r)(1 + x) = 1 + r + x + rx = 1$ . Conversely, if there exists  $y \in R$  such that  $y(1 + x) = 1$ , then

$$(y - 1) \circ x = y - 1 + x + (y - 1)x = 0.$$

**Example 2.30.** If  $x \in R$  is a nilpotent element, then  $y = \sum_{n \geq 1} x^n \in R$  is quasi-regular. En efecto, si existe  $N$  tal que  $x^N = 0$ , la suma que define al elemento  $y$  es finita y cumple que  $y + (-x) + y(-x) = 0$ .

A left ideal  $I$  of  $R$  is said to be **left quasi-regular** (resp. right quasi-regular) if every element of  $I$  is left quasi-regular (resp. right quasi-regular). A left ideal is said to be **quasi-regular** if it is left and right quasi-regular. Similarly one defines right quasi-regular ideals and quasi-regular ideals.

lemma:casiregular

**Lemma 2.31.** Let  $I$  be a left ideal of  $R$ . If  $I$  is left quasi-regular, then  $I$  is quasi-regular.

*Proof.* Let  $x \in I$ . Let us prove that  $x$  is right quasi-regular. Since  $I$  is left quasi-regular, there exists  $r \in R$  such that  $r \circ x = r + x + rx = 0$ . Since  $r = -x - rx \in I$ , there exists  $s \in R$  tal que  $s \circ r = s + r + sr = 0$ . Then  $s$  is right quasi-regular and

$$x = 0 \circ x = (s \circ r) \circ x = s \circ (r \circ x) = s \circ 0 = s. \quad \square$$

Let  $(A, \leq)$  be a partially order set, this means that  $A$  is a set together with a reflexive, transitive and anti-symmetric binary relation  $R$  en  $A \times A$ , where  $a \leq b$  if and only if  $(a, b) \in R$ . Recall that the relation is reflexive if  $a \leq a$  for all  $a \in A$ , the relation is transitive if  $a \leq b$  and  $b \leq c$  imply that  $a \leq c$  and the relation is anti-symmetric if  $a \leq b$  and  $b \leq a$  imply  $a = b$ .

The elements  $a, b \in A$  are said to be **comparable** if  $a \leq b$  or  $b \leq a$ . An element  $a \in A$  is said to be **maximal** if  $c \leq a$  for all  $c \in A$  that is comparable with  $a$ . An **upper bound** for a non-empty subset  $B \subseteq A$  is an element  $d \in A$  such that  $b \leq d$  for all  $b \in B$ . A **chain** in  $A$  is a subset  $B$  such that every pair of elements of  $B$  are comparable. **Zorn's lemma** states the following property:

If  $A$  is a non-empty partially ordered set such that every chain in  $A$  contains an upper bound in  $A$ , then  $A$  contains a maximal element.

Our application of Zorn's lemma:

lemma:maxreg

**Lemma 2.32.** *Let  $R$  be a ring and  $x \in R$  be an element that is not left quasi-regular. Then there exists a maximal left ideal  $M$  such that  $x \notin M$ . Moreover,  $R/M$  is a simple  $R$ -module and  $x \notin \text{Ann}_R(R/M)$ .*

*Proof.* Let  $T = \{r + rx : r \in R\}$ . A straightforward calculation shows that  $T$  is a left ideal of  $R$  such that  $x \notin T$  (if  $x \in T$ , then  $r + rx = -x$  for some  $r \in R$ , a contradiction since  $x$  is not left quasi-regular).

The only left ideal of  $R$  containing  $T \cup \{x\}$  is  $R$ . Indeed, if there exists a left ideal  $U$  containing  $T$ , then  $x \notin U$ , since otherwise every  $r \in R$  could be written as  $r = (r + rx) + r(-x) \in U$ .

Let  $\mathcal{S}$  be the set of proper left ideals of  $R$  containing  $T$  partially ordered by inclusion. If  $\{K_i : i \in I\}$  is a chain in  $\mathcal{S}$ , then  $K = \cup_{i \in I} K_i$  is an upper bound for the chain ( $K$  is a proper, as  $x \notin K$ ). Zorn's lemma implies that  $\mathcal{S}$  admits a maximal element  $M$ . Thus  $M$  is a maximal left ideal such that  $x \notin M$ . Moreover,  $M$  is regular since  $r + r(-x) \in T \subseteq M$  for all  $r \in R$ . Therefore  $R/M$  is a simple  $R$ -module by Proposition 2.12. Since  $x(x + M) \neq 0$  (if  $x^2 \in M$ , then  $x \in M$ , as  $x + x^2 \in T \subseteq M$ ), it follows that  $x \notin \text{Ann}_R(R/M)$ .  $\square$

If  $x \in R$  is not left quasi-regular, Lemma 2.32 implies that there exists a simple  $R$ -module  $M$  such  $x \notin \text{Ann}_R(M)$ . Thus  $x \notin J(R)$ .

thm:casireg\_eq

**Theorem 2.33.** *Let  $R$  be a ring and  $x \in R$ . The following statements are equivalent:*

- 1) *The left ideal generated by  $x$  is quasi-regular.*
- 2)  *$Rx$  is quasi-regular.*
- 3)  *$x \in J(R)$ .*

*Proof.* The implication (1)  $\implies$  (2) is trivial, as  $Rx$  is included in the left ideal generated by  $x$ .

We now prove (2)  $\implies$  (3). If  $x \notin J(R)$ , then Lemma 2.32 implies that there exists a simple  $R$ -module  $M$  such that  $xm \neq 0$  for some  $m \in M$ . The simplicity of  $M$  implies that  $R(xm) = M$ . Thus there exists  $r \in R$  such that  $rxm = -m$ . There is an element  $s \in R$  such that  $s + rx + s(rx) = 0$  and hence

$$-m = rxm = (-s - srx)m = -sm + sm = 0,$$

a contradiction.

Finally, to prove (3)  $\implies$  (1) it is enough to note that  $x$  is left quasi-regular. Thus the left ideal generated by  $x$  is quasi-regular by Lemma 2.31.  $\square$

The theorem immediately implies the following corollary.

**Corollary 2.34.** *If  $R$  is a ring, then  $J(R)$  is a quasi-regular ideal that contains every left quasi-regular ideal.*

The following result is somewhat what we all had in mind.

thm:J(R)

**Theorem 2.35.** *Let  $R$  be a ring such that  $J(R) \neq R$ . Then*

$$J(R) = \bigcap \{I : I \text{ regular maximal left ideal of } R\}.$$

*Proof.* We only prove the non-trivial inclusion. Let

$$K = \bigcap \{I : I \text{ regular maximal left ideal of } R\}.$$

By Proposition 2.12,

$$J(R) = \bigcap \{\text{Ann}_R(R/I) : I \text{ regular maximal left ideal of } R\}.$$

Let  $I$  be a regular maximal left ideal. If  $r \in J(R) \subseteq \text{Ann}_R(R/I)$ , then, since  $I$  is regular, there exists  $e \in R$  such that  $r - re \in I$ . Since

$$re + I = r(e + I) = 0,$$

$re \in I$  and hence  $r \in I$ . Thus  $J(R) \subseteq K$ .  $\square$

**Example 2.36.** Each maximal ideals of  $\mathbb{Z}$  is of the form  $p\mathbb{Z} = \{pm : m \in \mathbb{Z}\}$  for some prime number  $p$ . Thus  $J(\mathbb{Z}) = \bigcap_p p\mathbb{Z} = \{0\}$ .

We now review some basic results useful to compute radicals.

**Proposition 2.37.** *Let  $\{R_i : i \in I\}$  be a family of rings. Then*

$$J\left(\prod_{i \in I} R_i\right) = \prod_{i \in I} J(R_i).$$

*Proof.* Let  $R = \prod_{i \in I} R_i$  and  $x = (x_i)_{i \in I} \in R$ . The left ideal  $Rx$  is quasi-regular if and only if each left ideal  $R_i x_i$  is quasi-regular in  $R_i$ , as  $x$  is quasi-regular in  $R$  if and only if each  $x_i$  is quasi-regular in  $R_i$ . Thus  $x \in J(R)$  if and only if  $x_i \in J(R_i)$  for all  $i \in I$ .  $\square$

For the next result we shall need a lemma.

lemma:trickJ1

**Lemma 2.38.** *Let  $R$  be a ring and  $x \in R$ . If  $-x^2$  is a left quasi-regular element, then  $x$  también.*

*Proof.* Sea  $r \in R$  tal que  $r + (-x^2) + r(-x^2) = 0$  y sea  $s = r - x - rx$ . Entonces  $x$  es casi-regular a izquierda pues

$$\begin{aligned} s + x + sx &= (r - x - rx) + x + (r - x - rx)x \\ &= r - x - rx + x + rx - x^2 - rx^2 = r - x^2 - rx^2 = 0. \end{aligned} \quad \square$$

proposition:J(I)

**Proposition 2.39.** *If  $I$  is an ideal of  $R$ , then  $J(I) = I \cap J(R)$ .*

*Proof.* Since  $I \cap J(R)$  is an ideal of  $I$ , if  $x \in I \cap J(R)$ , then  $x$  is left quasi-regular in  $R$ . Let  $r \in R$  be such that  $r + x + rx = 0$ . Since  $r = -x - rx \in I$ ,  $x$  is left quasi-regular in  $I$ . Thus  $I \cap J(R) \subseteq J(I)$ .

Let  $x \in J(I)$  and  $r \in R$ . Since  $-(rx)^2 = (-rxr)x \in I(J(I)) \subseteq J(I)$ , the element  $-(rx)^2$  is left quasi-regular in  $I$ . Thus  $rx$  is left quasi-regular by Lemma 2.38.  $\square$

A ring  $R$  is said to be **radical** if  $J(R) = R$ .

**Example 2.40.** If  $R$  is a ring, then  $J(R)$  is a radical ring, by Proposition 2.39.

**Example 2.41.** The Jacobson radical of  $\mathbb{Z}/8$  is  $\{0, 2, 4, 6\}$ .

There are several characterizations of radical rings.

theorem:anillo\_radical

**Theorem 2.42.** Let  $R$  be ring. The following statements are equivalent:

- 1)  $R$  is radical.
- 2)  $R$  admits no simple  $R$ -modules.
- 3)  $R$  no tiene ideales a izquierda maximales y regulares.
- 4)  $R$  no tiene ideales a izquierda primitivos.
- 5) Every element of  $R$  is quasi-regular.
- 6)  $(R, \circ)$  is a group.

*Proof.* The equivalence (1)  $\iff$  (5) follows from Theorem 2.33.

The equivalence (5)  $\iff$  (6) is left as an exercise.

Let us prove that (1)  $\implies$  (2). Assume that there exists a simple  $R$ -module  $N$ . Since  $R = J(R) \subseteq \text{Ann}_R(N)$ ,  $R = \text{Ann}_R(N)$ . Hence  $RN = \{0\}$ , a contradiction to the simplicity of  $N$ .

To prove (2)  $\implies$  (3) we note that for each regular and maximal left ideal  $I$ , the quotient  $R/I$  is a simple  $R$ -module by Proposition 2.12.

To prove (3)  $\implies$  (4) assume that there is a primitive left ideal  $I = \text{Ann}_R(M)$ , where  $M$  is some simple  $R$ -module. Since  $R = J(R) \subseteq I$ , it follows that  $I = R$ , a contradiction to the simplicity of  $M$ .

Finally we prove (4)  $\implies$  (2). If  $M$  is a simple  $R$ -module, then  $\text{Ann}_R(M)$  is a primitive left ideal.  $\square$

**Example 2.43.** Let

$$A = \left\{ \frac{2x}{2y+1} : x, y \in \mathbb{Z} \right\}.$$

Then  $A$  is a radical ring, as the inverse of the element  $\frac{2x}{2y+1}$  with respect to the circle operation  $\circ$  is

$$\left( \frac{2x}{2y+1} \right)' = \frac{-2x}{2(x+y)+1}.$$

A ring  $R$  is said to be **nil** if for every  $x \in R$  there exists  $n = n(x)$  such that  $x^n = 0$ .

**Exercise 2.44.** Prove that a nil ring is a radical ring.

**Exercise 2.45.** Let  $\mathbb{R}[X]$  be the ring of power series with real coefficients. Prove that the ideal  $X\mathbb{R}[X]$  consisting of power series with zero constant term is a radical ring that is not nil.

The following problem is maybe the most important open problem in non-commutative ring theory.

conj:Koethe

**Conjecture 2.46 (Köthe).** Let  $R$  be a ring. The sum of two arbitrary nil left ideals of  $R$  is nil.

The conjecture is known to be true in several cases. Exercises?

thm:Jnilpotente

**Theorem 2.47.** If  $R$  is a left artinian ring, then  $J(R)$  is nilpotent.

*Proof.* Let  $J = J(R)$ . Since  $R$  is a left artinian ring, the sequence  $(J^m)_{m \in \mathbb{N}}$  of left ideals stabilizes. There exists  $k \in \mathbb{N}$  such that  $J^k = J^l$  for all  $l \geq k$ . We claim that  $J^k = \{0\}$ . If  $J^k \neq \{0\}$  let  $\mathcal{S}$  the set of left ideals  $I$  such that  $J^k I \neq \{0\}$ . Since

$$J^k J^k = J^{2k} = J^k \neq \{0\},$$

the set  $\mathcal{S}$  is non-empty. Since  $R$  is left artinian,  $\mathcal{S}$  has a minimal element  $I_0$ . Since  $J^k I_0 \neq \{0\}$ , let  $x \in I_0 \setminus \{0\}$  be such that  $J^k x \neq \{0\}$ . Moreover,  $J^k x$  is a left ideal of  $R$  contained in  $I_0$  and such that  $J^k x \in \mathcal{S}$ , as  $J^k(J^k x) = J^{2k} x = J^k x \neq \{0\}$ . The minimality of  $I_0$  implies that,  $J^k x = I_0$ . In particular, there exists  $r \in J^k \subseteq J(R)$  such that  $rx = x$ . Since  $-r \in J(R)$  is left quasi-regular, there exists  $s \in R$  such that  $s - r - sr = 0$ . Thus

$$x = rx = (s - sr)x = sx - s(rx) = sx - sx = 0,$$

a contradiction. □

**Corollary 2.48.** Let  $R$  be a left artinian ring. Each nil left ideal is nilpotent and  $J(R)$  is the unique maximal nilpotent ideal of  $R$ .

*Proof.* Let  $L$  be a nil left ideal of  $R$ . By Proposition 2.27,  $L$  is contained in  $J(R)$ . Thus  $L$  is nilpotent, as  $J(R)$  is nilpotent by Theorem 2.47. □

**Theorem 2.49.** Let  $R$  be a ring and  $n \in \mathbb{N}$ . Then  $J(M_n(R)) = M_n(J(R))$ .

*Proof.* We first prove that  $J(M_n(R)) \subseteq M_n(J(R))$ . If  $J(R) = R$ , the theorem is clear. Let us assume that  $J(R) \neq R$  and let  $J = J(R)$ . If  $M$  is a simple  $R$ -module, then  $M^n$  is a simple  $M_n(R)$ -module with the usual multiplication. Let  $x = (x_{ij}) \in J(M_n(R))$  and  $m_1, \dots, m_n \in M$ . Then

$$x \begin{pmatrix} m_1 \\ \vdots \\ m_n \end{pmatrix} = 0.$$

In particular,  $x_{ij} \in \text{Ann}_R(M)$  for all  $i, j \in \{1, \dots, n\}$ . Hence  $x \in M_n(J)$ .

We now prove that  $M_n(J) \subseteq J(M_n(R))$ . Let

$$J_1 = \begin{pmatrix} J & 0 & \cdots & 0 \\ J & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ J & 0 & \cdots & 0 \end{pmatrix} \quad \text{and} \quad x = \begin{pmatrix} x_1 & 0 & \cdots & 0 \\ x_2 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ x_n & 0 & \cdots & 0 \end{pmatrix} \in J_1.$$

Since  $x_1$  is quasi-regular, there exists  $y_1 \in R$  such that  $x_1 + y_1 + x_1 y_1 = 0$ . If

$$y = \begin{pmatrix} y_1 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix},$$

then  $u = x + y + xy$  is lower triangular, as

$$u = \begin{pmatrix} 0 & 0 & \cdots & 0 \\ x_2 y_1 & 0 & \cdots & 0 \\ x_3 y_1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ x_n y_1 & 0 & \cdots & 0 \end{pmatrix}.$$

Since  $u^n = 0$ , the element

$$v = -u + u^2 - u^3 + \cdots + (-1)^{n-1} u^{n-1}$$

is such that  $u + v + uv = 0$ . Thus  $x$  is right quasi-regular, as

$$x + (y + v + yv) + x(y + v + yv) = 0,$$

and therefore  $J_1$  is right quasi-regular. Similarly one proves that each  $J_i$  is right quasi-regular and hence  $J_i \subseteq J(M_n(R))$  for all  $i \in \{1, \dots, n\}$ . In conclusion,

$$J_1 + \cdots + J_n \subseteq J(M_n(R))$$

and therefore  $M_n(J) \subseteq J(M_n(R))$ . □

For completeness we recall basic results on the Jacobson radical in the case of unitary rings.

**Exercise 2.50.** Let  $R$  be a unitary ring. Then

$$J(R) = \bigcap \{M : M \text{ is a left maximal ideal}\}.$$

**Exercise 2.51.** Let  $R$  be a unitary ring. The following statements are equivalent:

- 1)  $x \in J(R)$ .
- 2)  $xM = 0$  for all simple  $R$ -module  $M$ .
- 3)  $x \in P$  for all primitive left ideal  $P$ .

- 4)  $1 + rx$  is invertible for all  $r \in R$ .
- 5)  $1 + \sum_{i=1}^n r_i x s_i$  is invertible for all  $n \in \mathbb{N}$  and all  $r_i, s_i \in R$ .
- 6)  $x$  belongs to every left maximal ideal maximal.

## 2B

We now go back to study solutions to the YBE and discuss the intriguing interplay between radical rings and involutive solutions.

**Definition 2.52.** A solution  $(X, r)$  is said to be *involutive* if  $r^2 = \text{id}$ .

For  $n \geq 2$ , the *symmetric group*  $\mathbb{S}_n$  can be presented as the group with generators  $\sigma_1, \dots, \sigma_{n-1}$  and relations

$$\begin{aligned} \sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1} & \text{if } 1 \leq i \leq n-2, \\ \sigma_i \sigma_j &= \sigma_j \sigma_i & \text{if } |i-j| \geq 1, \\ \sigma_i^2 &= 1 & \text{for all } i \in \{1, \dots, n-1\}. \end{aligned}$$

Let  $(X, r)$  be an involutive solution. Then the map  $\sigma_i \mapsto r_{i,i+1} = \text{id}_{X^{i-1}} \times r \times \text{id}_{X^{n-i-1}}$  extends to an action of  $\mathbb{S}_n$  on  $X^n$ .

**Example 2.53.** Let  $X$  be a non-empty set and  $\sigma$  be a bijection on  $X$ . Then  $(X, r)$ , where  $r(x, y) = (\sigma(y), \sigma^{-1}(x))$ , is an involutive solution.

We now present a very important family of involutive solutions. These examples show an intriguing connection between the YBE and the theory of non-commutative rings.

**Example 2.54.** Let  $p$  be a prime and let  $A = \mathbb{Z}/(p^2)$  be the cyclic additive group of order  $p^2$ . The operation  $x \circ y = x + y + pxy$  turns  $A$  into a radical ring.

**Example 2.55.** Let  $A = \left\{ \frac{2x}{2y+1} : x, y \in \mathbb{Z} \right\}$ . The operation  $a \circ b = a + b + ab$  turns  $A$  into a radical ring. A straightforward computation shows that

$$\left( \frac{x}{2y+1} \right)' = \frac{2(-x)}{2(x+y)+1}.$$

The following fundamental family of solutions appears in [53]. It turns out to be fundamental in the study of set-theoretic solutions to the YBE.

pro:Rump

**Proposition 2.56.** Let  $R$  be a radical ring. Then  $(R, r)$ , where

$$r(x, y) = (-x + x \circ y, (-x + x \circ y)' \circ x \circ y)$$

is an involutive solution.



The proposition can be demonstrated using Theorem 1.11, see Exercise 2.2. We will prove a stronger result in Theorem 4.22.

PRO:T

**Proposition 2.57.** *Let  $(X, r)$  be an involutive solution. Then the map  $T: X \rightarrow X$ ,  $x \mapsto \sigma_x^{-1}(x)$ , is invertible with inverse  $T^{-1}(y) = \tau_y^{-1}(y)$  and*

$$T^{-1} \circ \sigma_x^{-1} \circ T = \tau_x$$

for all  $x \in X$ .

*Proof.* Let  $U(x) = \tau_x^{-1}(x)$ . Since  $r$  is involutive,

$$(U(x), x) = r^2(U(x), x) = r(\sigma_{U(x)}(x), x) = (\sigma_{\sigma_{U(x)}(x)}(x), \tau_x \sigma_{U(x)}(x)).$$

The second coordinate can be written as  $U(x) = \sigma_{U(x)}(x)$ . This implies that

$$T(U(x)) = \sigma_{U(x)}^{-1}(U(x)) = x.$$

Similarly one obtains  $U(T(x)) = x$ .

Since  $(X, r)$  is a solution, Lemma 1.3 implies that  $\sigma_x \sigma_y = \sigma_{\sigma_x(y)} \sigma_{\tau_y(x)}$  holds for all  $x, y \in X$ . Then

$$\sigma_y^{-1} T(x) = \sigma_y^{-1} \sigma_x^{-1}(x) = \sigma_{\tau_y(x)}^{-1} \sigma_{\sigma_x(y)}^{-1}(x) = \sigma_{\tau_y(x)}^{-1} \tau_y(x) = T \tau_y(x)$$

for all  $y \in X$ , by Equality (2.2). □

Note that if  $(X, r)$  is a non-degenerate involutive solution, then

$$(x, y) = r^2(x, y) = r(\sigma_x(y), \tau_y(x)) = (\sigma_{\sigma_x(y)} \tau_y(x), \tau_{\tau_y(x)} \sigma_x(y)).$$

Hence

$$\tau_y(x) = \sigma_{\sigma_x(y)}^{-1}(x), \quad \sigma_x(y) = \tau_{\tau_y(x)}^{-1}(y) \tag{2.2}$$

eq:involutive

for all  $x, y \in X$ . Thus for involutive solutions it is enough to know  $\{\sigma_x : x \in X\}$ , as from this we obtain the set  $\{\tau_x : x \in X\}$ .

**Definition 2.58.** A *cycle set* is a pair  $(X, \cdot)$ , where  $X$  is a non-empty set provided with a binary operation  $X \times X \rightarrow X$ ,  $(x, y) \mapsto x \cdot y$ , such that

$$(x \cdot y) \cdot (x \cdot z) = (y \cdot x) \cdot (y \cdot z) \tag{2.3}$$

eq:cycle\_set

holds for all  $x, y, z \in X$  and each map  $\varphi_x: X \rightarrow X$ ,  $y \mapsto x \cdot y$ , is bijective. A cycle set  $(X, \cdot)$  is said to be *non-degenerate* if the map  $X \rightarrow X$ ,  $x \mapsto x \cdot x$ , is bijective.

**Definition 2.59.** Let  $X$  and  $Z$  be cycle sets. A *homomorphism* between the cycle sets  $X$  and  $Z$  is a map  $f: X \rightarrow Z$  such that  $f(x \cdot y) = f(x) \cdot f(y)$  for all  $x, y \in X$ . An *isomorphism* of cycle sets is a bijective homomorphism of cycle sets.

Cycle sets and cycle set homomorphisms form a category. It is possible to prove that the category of solutions is equivalent to the category of cycle sets, see Exercise 2.1.

thm:CS

**Theorem 2.60.** *There exists a bijective correspondence between non-isomorphic involutive solutions and non-isomorphic non-degenerate cycle sets.*

For the readers who are not familiar with the above-mentioned result, the bijective correspondence is given by

$$r(x, y) = (x * y, (x * y) \cdot x),$$

where  $x * y = z$  if and only if  $x \cdot z = y$ . We leave the proof for the reader, see Exercise 2.4. However, we will prove a more general result in Theorem 3.14.

Theorem 2.60 can be used to construct and enumerate small involutive solutions [2]. Table 2.1 shows the number of non-isomorphic involutive solutions of size  $\leq 10$ . For size  $\leq 7$  the numbers of Table 2.1 coincide with those in [29] but differ by two for  $n = 8$ , as two solutions of size eight are missing in [29].

**Table 2.1:** Involutive solutions of size  $\leq 10$ .

$n$	2	3	4	5	6	7	8	9	10
solutions	2	5	23	88	595	3456	34530	321931	4895272

tab:IYB

prob:cycle\_sets

**2.1.** Prove that the category of non-degenerate cycle sets and the category of solutions are equivalent.

prob:Rump

**2.2.** Prove Proposition 2.56.

**2.3.** If  $X$  is a cycle set, then  $x \cdot (y \cdot y) = ((y * x) \cdot y) \cdot ((y * x) \cdot y)$ , where  $y * x = z$  if and only if  $y \cdot z = x$ .

prob:CS

**2.4.** Prove Theorem 2.60.

## Open problems

**Problem 2.1.** Construct and enumerate involutive solutions of size 11.

**Problem 2.2.** Estimate the number of solutions of size  $n$  for  $n \rightarrow \infty$ .

## Notes

The material on non-commutative ring theory is standard, see for example [11]. Radical rings were introduced by Jacobson in [38]. Nil rings were used by Zelmanov in his solution to Burnside's problem, see for example [67].

Conjecture 2.46 is the well-known Köthe's conjecture. The conjecture was first formulated in 1930, see [41]. It is known to be true in several cases. In full generality, the problem is still open. In [42] Krempa proved that the following statements are equivalent:

- 1) Köthe's conjecture is true.
- 2) If  $R$  is a nil ring, then  $R[X]$  is a radical ring.
- 3) If  $R$  is a nil ring, then  $M_2(R)$  is a nil ring.
- 4) Let  $n \geq 2$ . If  $R$  is a nil ring, then  $M_n(R)$  is a nil ring.

In 1956 Amitsur formulated the following conjecture, see for example [4]: If  $R$  is a nil ring, then  $R[X]$  is a nil ring. In [55] Smoktunowicz found a counterexample to Amitsur's conjecture. This counterexample suggests that Köthe's conjecture might be false. A simplification of Smoktunowicz's example appears in [47]. See [56, 57] for more information on Köthe's conjecture and related topics.

Rump introduced cycle sets in [52]. The bijective correspondence of Theorem 2.60 was also proved by Rump in [52]. A similar result can be found in [29, Proposition 2.2].

The numbers of Table 2.1 were computed in [2] using a combination of [30] and constraint programming techniques. The algorithm is based on an idea of Plemmons [49], originally conceived to construct non-isomorphic semigroups.



## Chapter 3

### Racks

#### 3A

defn:rack

**Definition 3.1.** A *rack* is a pair  $(X, \triangleright)$ , where  $X$  is a non-empty set and  $X \times X \rightarrow X$ ,  $(x, y) \mapsto x \triangleright y$ , is a binary operation on  $X$  such that the maps  $\rho_y: X \rightarrow X$ ,  $x \mapsto x \triangleleft y$ , are bijective for all  $y \in X$ , and

$$(x \triangleleft y) \triangleleft z = (x \triangleleft z) \triangleleft (y \triangleleft z) \quad (3.1)$$

eq:rack

for all  $x, y, z \in X$ .

Racks are used in low-dimensional topology [27], singularities [12] and in the classification of finite-dimensional pointed Hopf algebras [5].

**Example 3.2.** Let  $X$  be a set. Then  $x \triangleleft y = x$  turns  $X$  into a rack. This is the *trivial rack* on  $X$ .

**Example 3.3.** Let  $X = \mathbb{Z}/n$ . Then  $x \triangleleft y = 2y - x$  turns  $X$  into a rack. This is the *dihedral rack* of size  $n$ .

**Example 3.4.** Let  $A$  be an abelian group and  $f \in \text{Aut}(A)$ . Then

$$x \triangleleft y = (\text{id} - f)(y) + f(x)$$

turns  $A$  into a rack. These racks are known as the *Alexander racks*.

**Definition 3.5.** Let  $X$  and  $Z$  be racks. A *rack homomorphism* between the racks  $X$  and  $Z$  is a map  $f: X \rightarrow Z$  such that  $f(x \triangleleft y) = f(x) \triangleleft f(y)$  for all  $x, y \in X$ . An *isomorphism* of racks is a bijective rack homomorphism.

For  $n \in \mathbb{N}$ , let  $r(n)$  be the number of isomorphism classes of racks of size  $n$ . Some values of  $r(n)$  appear in Table 3.1, see for example [64].

**Table 3.1:** Enumeration of non-isomorphic racks.

tab:racks

$n$	2	3	4	5	6	7	8	9	10	11	12	13
$r(n)$	2	6	19	74	353	2080	16023	159526	2093244	36265070	836395102	25794670618

pro:Venkov

**Proposition 3.6.** *Let  $X$  be a non-empty set and  $X \times X \rightarrow X$ ,  $(x, y) \mapsto x \triangleleft y$  be a binary operation on  $X$ . Then  $r(x, y) = (y, x \triangleleft y)$  is a solution if and only if  $(X, \triangleleft)$  is a rack.*

*Proof.* The map  $r$  satisfies  $(r \times \text{id})(\text{id} \times r)(r \times \text{id}) = (\text{id} \times r)(r \times \text{id})(\text{id} \times r)$  if and only if (3.1) holds for all  $x, y, z \in X$ . The solution  $(X, r)$  is non-degenerate if the maps  $X \rightarrow X$ ,  $x \mapsto x \triangleleft y$ , are bijective.  $\square$

The connection between racks and solutions goes deeper than the phenomenon appearing in Proposition 3.6.

pro:derived

**Proposition 3.7.** *Let  $(X, r)$  be a solution. Then*

$$x \triangleleft y = \sigma_y \tau_{\sigma_x^{-1}(y)}(x) = \sigma_y \widehat{\sigma}_y^{-1}(x) \quad (3.2)$$

eq:derived

*turns  $X$  into a rack and each  $\sigma_x$  is a rack homomorphism. Moreover,  $(X, r)$  is involutive if and only if the rack  $(X, \triangleleft)$  is trivial.*

*Proof.* Since  $r(x, \sigma_x^{-1}(y)) = (y, \tau_{\sigma_x^{-1}(y)}(x))$ , it follows that  $\widehat{\sigma}_y^{-1}(x) = \tau_{\sigma_x^{-1}(y)}(x)$  for all  $x, y \in X$ . Hence the second equality of (3.2) holds.

To prove that each  $\tau_z$  is a rack homomorphism it is enough to show that

$$\sigma_x(y) \triangleleft \sigma_x \sigma_y(z) = \sigma_x(y \triangleleft \sigma_y(z))$$

for all  $x, y \in X$ . Write  $r(x, y) = (u, v)$ . On the one hand, by Lemma 1.3,

$$\sigma_x(y) \triangleleft \sigma_x \sigma_y(z) = u \triangleleft \sigma_u \sigma_v(z) = \sigma_{\sigma_u \sigma_v(z)} \tau_{\sigma_{\sigma_y(x)}(z)} \sigma_x(y) = \sigma_{\sigma_x \sigma_y(z)} \sigma_{\tau_{\sigma_y(z)}(x)} \tau_z(y).$$

On the other hand,

$$\sigma_x(y \triangleleft \sigma_y(z)) = \sigma_x \sigma_{\sigma_y(z) \tau_z(y)} = \sigma_{\sigma_x \sigma_y(z)} \sigma_{\tau_{\sigma_y(z)}(x)} \tau_z(y).$$

By Proposition 3.6, in order to prove that  $(X, \triangleright)$  is a rack it is enough to show that  $r(x, y) = (y, x \triangleleft y)$  satisfies the YBE. For that purpose, we demonstrate that the map  $J: X^3 \rightarrow X^3$ ,  $J(x, y, z) = (x, \sigma_x(y), \sigma_x \sigma_y(z))$  is invertible and satisfies

$$(\text{id} \times s) \circ J = J \circ (\text{id} \times r), \quad (s \times \text{id}) \circ J = J \circ (r \times \text{id}).$$

The map  $(x, y, z) \mapsto (x, \sigma_x^{-1}(y), \sigma_{\sigma_x^{-1}(y)}^{-1} \sigma_x^{-1}(z))$  is the inverse of  $J$ .

Since  $\sigma_x$  is a rack homomorphism,

$$\sigma_x(y) \triangleleft \sigma_x \sigma_y(z) = \sigma_x(y \triangleleft \sigma_y(z)) = \sigma_x \sigma_{\sigma_y(z)} \tau_{\sigma_y^{-1} \sigma_y(z)}(y) = \sigma_x \sigma_{\sigma_y(z)} \tau_z(y)$$

Then it follows that

$$\begin{aligned}
 (\text{id} \times s)J(x, y, z) &= (\text{id} \times s)(x, \sigma_x(y), \sigma_x \sigma_y(z)) \\
 &= (x, \sigma_x \sigma_y(z), \sigma_x(y) \triangleleft \sigma_x \sigma_y(z)) \\
 &= (x, \sigma_x \sigma_y(z), \sigma_x \sigma_{\sigma_y(z)} \tau_z(y)) \\
 &= J(x, \sigma_y(z), \tau_z(y)) \\
 &= J(\text{id} \times r)(x, y, z).
 \end{aligned}$$

Similarly one proves that  $(s \times \text{id}) \circ J = J \circ (r \times \text{id})$ . This implies that  $(X, s)$  is a solution and hence  $(X, \triangleleft)$  is a rack by Proposition 3.6.

If  $(X, r)$  is involutive, then  $x \triangleleft \sigma_x(y) = \sigma_{\sigma_x(y)} \tau_y(x) = x$  by (2.2). Conversely, if  $x \triangleleft y = x$  for all  $x, y \in X$ , then  $r$  is involutive, as

$$r^2(x, \sigma_x^{-1}(y)) = r(y, \sigma_y^{-1}(x)) = (x, \sigma_x^{-1}(y)). \quad \square$$

The rack constructed in Proposition 3.7 is known as the *derived solution* of  $(X, r)$ . There is a dual version of the derived rack:

pro:derived\_dual

**Proposition 3.8.** *Let  $(X, r)$  be a solution. Then*

$$x \blacktriangleleft y = \tau_y \sigma_{\tau_x^{-1}(y)}(x) = \tau_y \widehat{\tau_y}^{-1}(x)$$

*turns  $X$  into a rack and each  $\tau_x$  is a rack homomorphism.*

*Proof.* Since  $(X, r)$  is a solution, then so is  $(X, r_0)$ , where  $r_0(x, y) = (\tau_x(y), \sigma_y(x))$ . Then the claim follows from Proposition 3.8 applied to the solution  $(X, r_0)$ .  $\square$

In general, the racks constructed in Propositions 3.7 and 3.8 are different:

**Example 3.9.** Let  $X = \{1, \dots, 5\}$  and  $(X, r)$  be the solution given by

$$\begin{aligned}
 \sigma_1 &= \text{id}, & \sigma_2 &= \text{id}, & \sigma_3 &= \text{id}, & \sigma_4 &= (13)(45), & \sigma_5 &= (12)(45), \\
 \tau_1 &= \text{id}, & \tau_2 &= \text{id}, & \tau_3 &= \text{id}, & \tau_4 &= (23)(45), & \tau_5 &= (23)(45).
 \end{aligned}$$

On the one hand the derived rack of  $(X, r)$  is given by the permutations

$$\sigma_1 \widehat{\sigma_1}^{-1} = \sigma_2 \widehat{\sigma_2}^{-1} = \sigma_3 \widehat{\sigma_3}^{-1} = \text{id}, \quad \sigma_4 \widehat{\sigma_4}^{-1} = (132), \quad \sigma_5 \widehat{\sigma_5}^{-1} = (123).$$

On the other hand, the dual derived rack by

$$\tau_1 \widehat{\tau_1}^{-1} = \tau_2 \widehat{\tau_2}^{-1} = \tau_3 \widehat{\tau_3}^{-1} = \text{id}, \quad \tau_4 \widehat{\tau_4}^{-1} = (123), \quad \tau_5 \widehat{\tau_5}^{-1} = (132).$$

We now prove that the racks of Propositions 3.7 and 3.8 are isomorphic. We shall need a lemma.

lem:T\_invertible

**Lemma 3.10.** *Let  $(X, r)$  be a solution. The map  $T: X \rightarrow X$ ,  $x \mapsto \sigma_x^{-1}(x)$ , is invertible with inverse  $U: X \rightarrow X$ ,  $x \mapsto \tau_x^{-1}(x \blacktriangleleft x)$ .*

*Proof.* Let  $x \in X$  and  $y = U(x) = \tau_x^{-1}(x \blacktriangleleft x)$ . Then  $\tau_x(y) = x \blacktriangleleft x = \tau_x \widehat{\tau}_x^{-1}(x)$  and hence  $y = \widehat{\tau}_x^{-1}(x)$ . Then  $\widehat{\tau}_x(y) = x$  and

$$r^{-1}(y, x) = (\widehat{\sigma}_y(x), x) = (z, x),$$

where  $z \in X$  is such that  $\sigma_z(x) = y$ . By Lemma 1.3,  $\sigma_y = \sigma_z$ . Then it follows that  $x = \sigma_y^{-1}(y) = T(y)$ . Therefore  $y = U(x) = U(T(y))$ .

To prove that  $T(U(x)) = x$ , first note that

$$r(\tau_x^{-1}(x), x) = (\sigma_{\tau_x^{-1}(x)}(x), x)$$

and Lemma 1.3 imply that  $\sigma_{\tau_x^{-1}(x)} = \sigma_{\sigma_{\tau_x^{-1}(x)}(x)}$ . Now

$$\begin{aligned} T(U(x)) &= T(\tau_x^{-1}(x \blacktriangleleft x)) = T(\sigma_{\tau_x^{-1}(x)}(x)) \\ &= \sigma_{\sigma_{\tau_x^{-1}(x)}(x)}^{-1} \sigma_{\tau_x^{-1}(x)}(x) = \sigma_{\tau_x^{-1}(x)}^{-1} \sigma_{\tau_x^{-1}(x)}(x) = x. \end{aligned} \quad \square$$

There is version of Proposition 2.57 for arbitrary solutions. A similar result appears in Exercise 3.3.

**Proposition 3.11.** *Let  $(X, r)$  be a solution. Then  $T : X \rightarrow X$ ,  $x \mapsto \tau_x^{-1}(x)$ , is a bijective map such that*

$$T \circ \tau_y = \widehat{\sigma}_y^{-1} \circ T, \quad T \circ \widehat{\tau}_y = \sigma_y^{-1} \circ T$$

and  $T(x \blacktriangleleft y) = T(x) \blacktriangleleft T(y)$  for all  $x, y \in X$ .

*Proof.* Lemma 3.10 proves that  $T$  is bijective. We now compute

$$\begin{aligned} T \tau_y(x) &= \sigma_{\tau_y(x)}^{-1} \tau_y(x) = \sigma_{\tau_y(x)}^{-1} \sigma_{\sigma_x(y)}^{-1} \sigma_{\sigma_x(y)} \tau_y(x) \\ &= \sigma_y^{-1} \sigma_x^{-1} \sigma_{\sigma_x(y)} \tau_y(x) = \sigma_y^{-1} \sigma_x^{-1} (x \blacktriangleleft \sigma_x(y)) = \sigma_y^{-1} (T(y) \blacktriangleleft y) = \widehat{\sigma}_y^{-1} T(x). \end{aligned}$$

Since  $\widehat{\tau}_y(x) = \sigma_{\widehat{\sigma}_x(y)}^{-1}(x)$ , Lemma 1.3 implies that

$$T \widehat{\tau}_y(x) = \sigma_{\widehat{\tau}_y(x)}^{-1} \widehat{\tau}_y(x) = \sigma_{\widehat{\tau}_y(x)}^{-1} \sigma_{\widehat{\sigma}_x(y)}^{-1} = \sigma_y^{-1} \sigma_x^{-1}(x) = \sigma_y^{-1} T(x).$$

These formulas imply that

$$T \circ \tau_y \circ \widehat{\tau}_y^{-1} = \widehat{\sigma}_y^{-1} \circ T \circ \widehat{\tau}_y^{-1} = \widehat{\sigma}_y^{-1} \circ \sigma_y \circ T. \quad (3.3) \quad \boxed{\text{eq:T\_rack}}$$

We evaluate Equality (3.3) on  $X$ . On the one hand,  $T(x \blacktriangleleft y) = T \sigma_x \widehat{\sigma}_x^{-1}(y)$ . On the other hand,

$$\widehat{\sigma}_y^{-1} \sigma_y T(x) = \sigma_y^{-1} \sigma_y \widehat{\sigma}_y^{-1} \sigma_y T(x) = \sigma_y^{-1} (\sigma_y T(x) \blacktriangleleft y) = T(x) \blacktriangleleft T(y). \quad \square$$

As it happens in the involutive case, there is a nice combinatorial structure that describes a solution.



defn:skewCS

**Definition 3.12.** A *skew cycle set* is a triple  $(X, \triangleleft, \cdot)$ , where  $X$  is a non-empty set,  $(X, \triangleleft)$  is a rack and  $X \times X \rightarrow X$ ,  $(x, y) \mapsto x \cdot y$ , is a binary operation on  $X$  such that the maps  $X \rightarrow X$ ,  $y \mapsto x \cdot y$ , are bijective rack homomorphisms, and

$$(x \cdot y) \cdot (x \cdot z) = (y \cdot (x \triangleleft y)) \cdot (y \cdot z) \quad (3.4)$$

eq:skew\_CS

for all  $x, y, z \in X$ . A skew cycle set  $(X, \triangleleft, \cdot)$  is said to be non-degenerate if the map  $X \times X$ ,  $x \mapsto x \cdot x$ , is bijective.

FIXME

**Definition 3.13.** Let  $X$  and  $Z$  be skew cycle sets. A *homomorphism* between the cycle sets  $X$  and  $Z$  is a map  $f: X \rightarrow Z$  such that  $f(x \cdot y) = f(x) \cdot f(y)$  for all  $x, y \in X$ . An *isomorphism* of cycle sets is a bijective homomorphism of cycle sets.

Cycle sets and cycle set homomorphisms form a category. It is possible to prove that the category of solutions is equivalent to the category of cycle sets, see Exercise 2.1.

Theorem 2.60 can be generalized to arbitrary solutions.

thm:skewCS

**Theorem 3.14.** *There exists a bijective correspondence between solutions and non-degenerate skew cycle sets.*

*Proof.* Let  $(X, r)$  be a solution and  $(X, \triangleleft)$  its derived rack. We will prove that the operation  $x \cdot y = \sigma_x^{-1}(y)$  turns  $(X, \triangleleft)$  into a skew cycle set. By Proposition 3.7, the maps  $X \rightarrow X$ ,  $y \mapsto x \cdot y$ , are bijective rack homomorphisms.

On the one hand, since  $r(x, \sigma_x^{-1}(y)) = (y, \tau_{\sigma_x^{-1}(y)}(x))$ ,

$$\begin{aligned} (x \cdot y) \cdot (x \cdot z) &= \sigma_x^{-1}(y) \cdot \sigma_x^{-1}(z) = \sigma_{\sigma_x^{-1}(y)}^{-1} \sigma_x^{-1}(z) \\ &= \left( \sigma_x \circ \sigma_{\sigma_x^{-1}(y)} \right)^{-1} (z) = \left( \sigma_y \circ \sigma_{\tau_{\sigma_x^{-1}(y)}(x)} \right)^{-1} (z). \end{aligned}$$

On the other hand,

$$\begin{aligned} (y \cdot (x \triangleleft y)) \cdot (y \cdot z) &= \sigma_y^{-1}(\sigma_y \tau_{\sigma_x^{-1}(y)}(x)) \cdot \sigma_y^{-1}(z) \\ &= \sigma_{\tau_{\sigma_x^{-1}(y)}(x)}^{-1} \sigma_y^{-1}(z) = \left( \sigma_y \circ \sigma_{\tau_{\sigma_x^{-1}(y)}(x)} \right)^{-1} (z). \end{aligned}$$

Now we prove the converse statement. For  $x, y \in X$  let

$$\sigma_x(y) = x * y, \quad \tau_y(x) = \sigma_{\sigma_x(y)}^{-1}(x \triangleleft \sigma_x(y)),$$

where  $x * y = z$  if and only if  $x \cdot z = y$ . Since  $X$  is a skew cycle set, each  $\sigma_x$  is bijective. Let us prove that the  $\tau_x$  are bijective. Equality (3.4) with  $y = \sigma_x(z)$  implies that

$$\sigma_z^{-1} \sigma_x^{-1} = \sigma_{\sigma_x^{-1}(y)}^{-1} \sigma_x^{-1} = \sigma_{\sigma_x^{-1}(x \triangleleft y)}^{-1} \sigma_y^{-1} = \sigma_{\sigma_x(z)}^{-1} \sigma_{\sigma_x(z)}^{-1} = \sigma_{\tau_z(x)}^{-1} \sigma_{\sigma_x(z)}^{-1}$$

for all  $x, z \in X$ . Since each  $\sigma_x$  is a rack homomorphism,

$$\tau_y(x) = \sigma_{\sigma_x(y)}^{-1}(x \triangleleft \sigma_x(y)) = \sigma_{\sigma_x(y)}^{-1} \sigma_x(\sigma_x^{-1}(x) \triangleleft y) = \sigma_{\tau_y(x)} \sigma_y^{-1}(\sigma_x^{-1}(x) \triangleleft y).$$

Therefore  $T \circ \tau_y = \sigma_y^{-1} \circ \rho_y \circ T$ , where  $T: X \rightarrow X$ ,  $T(x) = x \cdot x$  and  $\rho_y: X \rightarrow X$ ,  $\rho_y(x) = x \triangleleft y$  are bijective maps. In particular,  $\tau_y$  is bijective for all  $y \in X$ .

Now we prove that... solution?

invertible?

□

Theorem 3.14 can be used to construct small solutions, see Table 3.2.

**Table 3.2:** Enumeration of non-involutive solutions.

$n$	2	3	4	5	6	7	8
$s(n)$	2	21	253	3519	100071	4602720	422449480

tab:non\_involutive

## Exercises

prob:xx

**3.1.** Prove that  $x \triangleright x = x \blacktriangleright x$  for all  $x \in X$ .

prob:tau\_hat

**3.2.** Prove that  $\widehat{\tau}_x(y \triangleright z) = \widehat{\tau}_x(y) \triangleright \widehat{\tau}_x(z)$  for all  $x, y, z \in X$ .

prob:variationT

**3.3.** Let  $(X, r)$  be a solution and let  $(X, \triangleright)$  be its derived rack. Prove that

$$T \sigma_x(y) = \tau_x^{-1}(x \triangleright T(y))$$

for all  $x \in X$ , where  $T: X \rightarrow X$ ,  $T(y) = \tau_y^{-1}(y)$ .

prob:guitar

**3.4.** Let  $(X, r)$  be a solution and  $(X, \triangleright)$  its derived rack. Let  $T_2(x, y) = (\tau_y(x), y)$  and  $T_{n+1} = Q_n \circ (T_n \times \text{id})$  for  $n \geq 2$ , where

$$Q_n(x_1, \dots, x_{n+1}) = (\tau_{x_{n+1}}(x_1), \dots, \tau_{x_{n+1}}(x_n), x_{n+1}).$$

Prove that  $T_n \circ r_{i,i+1} = s_{i,i+1} \circ T_n$  for all  $n \geq 2$  and  $i \in \{1, \dots, n-1\}$ .

## Open problems

problem:racks14

**Problem 3.1.** Enumerate isomorphism classes of racks of size 14.

**Problem 3.2.** Enumerate non-involutive solutions of size  $\geq 9$ .

## Notes

A particular family of racks turns out to be useful in combinatorial knot theory. A quandle is a rack  $(X, \triangleleft)$  such that  $x \triangleleft x = x$  for all  $x \in X$ .

In [26], Drinfeld attributes Proposition 3.6 to Venkov.

There are several papers on the enumeration of isomorphic classes of finite racks [6, 10, 35]. Estimations on the number of finite racks of size  $n$  appear in [10].

The numbers of Table 3.2 were computed using Theorem 3.14 essentially with the same technique used to construct involutive solutions [2]. The construction of non-involutive solutions of size 9 seems to be feasible with these methods. However, it should be noted that a huge number of solutions is expected.

Exercises 3.1 and 3.2 appear in [44].

The map  $J$  of Exercise 3.4 is known as the *guitar map*. It was first considered by Etingof, Schedler and Soloviev in [29] for involutive solutions. The construction was extended to non-involutive solutions by Soloviev in [61] and Lu, Yan and Zhu in [45]. In [25] Dehornoy used the inverse of the guitar map to develop his right-cyclic calculus and to obtain short proofs for results on the structure group of involutive solutions. In [5] Andruskiewitsch and Graña use the guitar map to study certain isomorphisms of Nichols algebras. A particular case of the guitar map also appears in the work of Przytycki [51].

The derived rack of a solution was first defined in the work of Soloviev [61]. Most of the properties of the derived racks mentioned in this chapter were proved in [44].

Problem 3.1 appears in [64].



## Chapter 4

### Braces

braces

#### 4A

By convention, an additive group  $A$  will be a (not necessarily abelian) group with binary operation  $(a, b) \mapsto a + b$ . The identity of  $A$  will be denoted by  $0$  and the inverse of an element  $a$  will be denoted by  $-a$ .

def:brace

**Definition 4.1.** A *brace* is a triple  $(A, +, \circ)$ , where  $(A, +)$  and  $(A, \circ)$  are (not necessarily abelian) groups and

$$a \circ (b + c) = (a \circ b) - a + (a \circ c) \quad (4.1)$$

eq:compatibility

holds for all  $a, b, c \in A$ , where  $-a$  denotes the inverse of  $a$  with respect to the group structure given by  $(a, b) \mapsto a + b$ . The groups  $(A, +)$  and  $(A, \circ)$  are respectively the *additive* and *multiplicative* group of the brace  $A$ .

We write  $a'$  to denote the inverse of  $a$  with respect to the circle operation  $\circ$ .

Our definition is that of a left brace. Right braces are defined similarly, one needs to replace (4.1) by

$$(a + b) \circ c = a \circ c - c + b \circ c.$$

There is a bijective correspondence between left and right braces, see Exercise 4.1. For that reason, a brace will always mean a left brace.

**Definition 4.2.** Let  $\mathcal{X}$  be a property of groups. A brace  $A$  is said to be of  $\mathcal{X}$ -type if its additive group belongs to  $\mathcal{X}$ .

One particularly interesting families of braces is the family of *braces of abelian type*, that is braces with abelian additive group. Braces of abelian type were introduced by Rump in [53] to study involutive solutions to the Yang–Baxter equation. In the literature, braces of abelian type are called *left braces*.

exa:trivial

**Example 4.3.** Let  $A$  be an additive group. Then  $A$  is a brace with  $a \circ b = a + b$  for all  $a, b \in A$ . A brace  $(A, +, \circ)$  such that  $a \circ b = a + b$  for all  $a, b \in A$  is said to be *trivial*. Similarly, the operation  $a \circ b = b + a$  turns  $A$  into a brace.

exa:times

**Example 4.4.** Let  $A$  and  $B$  be braces. Then  $A \times B$  with

$$(a, b) + (a_1, b_1) = (a + a_1, b + b_1), \quad (a, b) \circ (a_1, b_1) = (a \circ a_1, b \circ b_1),$$

is a brace.

exa:sd

**Example 4.5.** Let  $A$  and  $M$  be additive groups and let  $\alpha: A \rightarrow \text{Aut}(M)$  be a group homomorphism. Then  $M \times A$  with

$$(x, a) + (y, b) = (x + y, a + b), \quad (x, a) \circ (y, b) = (x + \alpha_a(y), a + b)$$

is a brace. Similarly,  $M \times A$  with

$$(x, a) + (y, b) = (x + \alpha_a(y), a + b), \quad (x, a) \circ (y, b) = (x + y, b + a)$$

is a brace.

exa:s3c6

**Example 4.6.** Let  $A = \mathbb{S}_3$  be the symmetric group in three letters. Write  $A$  as an additive group. Let  $\lambda: A \rightarrow \mathbb{S}_A$  be the map given by

$$\begin{aligned} \lambda_{\text{id}} &= \lambda_{(123)} = \lambda_{(132)} = \text{id}, \\ \lambda_{(12)} &= \lambda_{(23)} = \lambda_{(13)} = \text{conjugation by } (23). \end{aligned}$$

It is easy to check that  $\lambda_{a+\lambda_a(b)} = \lambda_a \lambda_b$  for all  $a, b \in A$ . Hence  $A$  is a brace by Exercise 4.5. Since the transposition  $(12)$  has order six in the multiplicative group of  $A$ , it follows that the additive group of  $A$  is isomorphic to  $\mathbb{S}_3$  and the multiplicative group of  $A$  is isomorphic to the cyclic group of order six.

The following example is motivated by the paper [66].

exa:WX

**Example 4.7.** Let  $A$  be an additive group and  $B$  and  $C$  be subgroups of  $A$  such that  $A$  admits an *exact factorization* as  $A = B + C$ . Thus each  $a \in A$  can be written in a unique way as  $a = b + c$  for some  $b \in B$  and  $c \in C$ . The map

$$B \times C \rightarrow A, \quad (b, c) \mapsto b - c,$$

is bijective. Using this map we transport the group structure of the direct product  $B \times C$  into the set  $A$ . For  $a = b + c \in A$  and  $a_1 \in A$  let

$$a \circ a_1 = b + a_1 + c.$$

Then  $(A, \circ)$  is a group isomorphic to  $B \times C$ . Moreover, if  $x, y \in A$ , then

$$a \circ x - a + a \circ y = b + x + c - (b + c) + b + y + c = b + x + y + c = a \circ (x + y)$$

and therefore  $(A, +, \circ)$  is a brace.

We now give concrete examples of the previous construction.

exa:QR

**Example 4.8.** Let  $n \in \mathbb{N}$ . The group  $\mathbf{GL}_n(\mathbb{C})$  admits an exact factorization through the subgroups  $U(n)$  and  $T(n)$ , where  $U(n)$  is the unitary group and  $T(n)$  is the group of upper triangular matrices with positive diagonal entries. Therefore there exists a brace with additive group isomorphic to  $\mathbf{GL}_n(\mathbb{C})$  and multiplicative group isomorphic to  $U(n) \times T(n)$ .

The following examples appeared in the theory of Hopf–Galois extensions, see [14, Corollary 1.1].

exa:a5a4c5

**Example 4.9.** The alternating simple group  $\mathbb{A}_5$  admits an exact factorization through the subgroups  $A = \langle (123), (12)(34) \rangle \simeq \mathbb{A}_4$  and  $B = \langle (12345) \rangle \simeq C_5$ . There exists a brace with additive group isomorphic to  $\mathbb{A}_5$  and multiplicative group isomorphic to  $\mathbb{A}_4 \times C_5$ .

exa:PSL27S4C7

**Example 4.10.** The simple group  $\mathbf{PSL}_2(7)$  admits an exact factorization through the subgroups  $A \simeq \mathbb{S}_4$  and  $B \simeq C_7$ . There exists a brace with additive group isomorphic to  $\mathbf{PSL}_2(7)$  and multiplicative group isomorphic to  $\mathbb{S}_4 \times C_7$ .

lem:basic

**Lemma 4.11.** *Let  $A$  be a brace. Then the following properties hold:*

- 1)  $0 = 1$ .
- 2)  $a \circ (-b + c) = a - (a \circ b) + (a \circ c)$  for all  $a, b, c \in A$ .
- 3)  $a \circ (b - c) = (a \circ b) - (a \circ c) + a$  for all  $a, b, c \in A$ .

*Proof.* The first claim follows from the compatibility condition (4.1) with  $c = 1$ . To prove the second claim let  $d = b + c$ . Then (4.1) becomes

$$a \circ d = a \circ b - a + a \circ (-b + d)$$

and the claim follows. The third claim is proved similarly. □

pro:lambda

**Proposition 4.12.** *Let  $A$  be a brace. For each  $a \in A$ , the map*

$$\lambda_a: A \rightarrow A, \quad b \mapsto -a + (a \circ b),$$

*is bijective. Moreover, the map  $\lambda: (A, \circ) \rightarrow \text{Aut}(A, +)$ ,  $a \mapsto \lambda_a$ , is a group homomorphism.*

*Proof.* The inverse of  $\lambda$  is given by  $\lambda_a^{-1}: A \rightarrow A$ ,  $b \mapsto a' \circ (a + b)$ . To prove that  $\lambda_a \in \text{Aut}(A, +)$  we note that

$$\lambda_a(b + c) = -a + a \circ (b + c) = -a + a \circ b - a + a \circ c = \lambda_a(b) + \lambda_a(c).$$

To prove that  $\lambda$  is a group homomorphism, we use Lemma 4.11 to obtain

$$\begin{aligned} \lambda_a(\lambda_b(c)) &= -a + a \circ (-b + b \circ c) \\ &= -a + a \circ (-b) - a + a \circ (b \circ c) = -a \circ b + a \circ (b \circ c) = \lambda_{a \circ b}(c). \quad \square \end{aligned}$$

If  $A$  is a brace, the map  $\lambda$  is the previous proposition yields a left action from  $(A, \circ)$  on  $(A, +)$  by automorphisms. There is also a right action  $(A, \circ)$  on  $(A, +)$  by automorphisms:

pro:mu

**Proposition 4.13.** *Let  $A$  be a brace. For each  $a \in A$ , the map*

$$\mu_a: A \rightarrow A, \quad b \mapsto \lambda_a(b)' \circ a \circ b,$$

*is bijective. Moreover, the map  $\mu: (A, \circ) \rightarrow \mathbb{S}_A$ ,  $a \mapsto \mu_a$ , satisfies  $\mu_b \circ \mu_a = \mu_{a \circ b}$  for all  $a, b \in A$ .*

*Proof.* Let  $a, b, c \in A$ . To prove that  $\mu$  is a brace anti-homomorphism, we compute

$$\mu_{b \circ a}(c) = \lambda_c((b \circ a)' \circ c \circ b \circ a)$$

and

$$\mu_a \mu_b(c) = \mu_a(\lambda_c(b)' \circ c \circ b) = \lambda_{\lambda_c(b)' \circ c \circ b}(a)' \circ (\lambda_c((b \circ a)' \circ c \circ b)).$$

Using the formulas (4.2),

$$\begin{aligned} \lambda_c(b \circ a)' &= \lambda_c(b + \lambda_b(a)) = (\lambda_c(b) + \lambda_{c \circ b}(a))' \\ &= (\lambda_c(b) \circ \lambda_{\lambda_c(b)}^{-1} \lambda_{c \circ b}(a))' = \lambda_{\lambda_c(b)' \circ c \circ b}(a)' \circ \lambda_c(b)', \end{aligned}$$

which proves that  $\mu$  is an anti-homomorphism.

To compute the inverse of  $\mu_b$  we proceed as follows. Since  $a' \circ (-a) = 2a$  by Lemma 4.11,

$$\begin{aligned} (\lambda_a(b)' \circ a \circ b)' &= b' \circ (a' \circ \lambda_a(b)) \\ &= b' \circ (a' \circ (-a + a \circ b)) = b' \circ (a' + b) = b' \circ a' - b. \end{aligned}$$

From this one immediately obtains that  $\mu_b^{-1}(a) = (b \circ a' - b)'$ . □

Let  $A$  be a brace. The previous proposition implies that

$$a \circ b = a + \lambda_a(b), \quad a + b = a \circ \lambda_a^{-1}(b), \quad \lambda_a(a') = -a \quad (4.2)$$

eq:formulas

hold for  $a, b \in A$ . Moreover, if

$$a * b = \lambda_a(b) - b = -a + a \circ b - b,$$

then the following identities are easily verified:

$$a * (b + c) = a * b + b + a * c - b, \quad (4.3)$$

$$(a \circ b) * c = (a * (b * c)) + b * c + a * c. \quad (4.4)$$

These last two identities are similar to the usual *commutator identities*.

**Definition 4.14.** A *homomorphism* between two braces  $A$  and  $B$  is a group homomorphism  $f: A \rightarrow B$  such that  $f\lambda_a = \lambda_{f(a)}f$  for all  $a \in A$ . The *kernel* of  $f$  is



$$\ker f = \{a \in A : f(a) = 0\}.$$

Braces and brace homomorphisms form a category.

**Definition 4.15.** A brace  $A$  is said to be a *two-sided* if

$$(a+b) \circ c = a \circ c - c + b \circ c \quad (4.5) \quad \boxed{\text{eq:right\_compatibility}}$$

holds for all  $a, b, c \in A$ .

If  $A$  is a two-sided brace, then

$$a \circ (-b) = a - a \circ b + a, \quad (-a) \circ b = b - a \circ b + b \quad (4.6) \quad \boxed{\text{eq:2sided}}$$

hold for all  $a, b \in A$ . The first equality holds for every brace and follows from Lemma 4.11. The second equality follows from (4.5).

**Example 4.16.** Any brace with abelian multiplicative group is two-sided.

**Example 4.17.** Let  $n \in \mathbb{N}$  be such that  $n = p_1^{a_1} \cdots p_k^{a_k}$ , where the  $p_j$  are distinct primes, all  $a_j \in \{0, 1, 2\}$  and  $p_i^m \not\equiv 1 \pmod{p_j}$  for all  $i, j, m$  with  $1 \leq m \leq a_i$ . Then every brace of size  $n$  is a two-sided brace of abelian type, since every group of order  $n$  is abelian, see for example [48].

Two-sided braces of abelian type form an interesting family of rings without unit.

Braces are a far reaching generalizations of radical rings. The following result was proved by Rump in [53].

thm:radical

**Theorem 4.18.** A brace of abelian type is two-sided if and only if it is a radical ring.

*Proof.* Assume first that  $A$  is a two-sided brace of abelian type. Then  $(A, +)$  is an abelian group. Let us prove that the operation

$$ab = -a + a \circ b - b$$

turns  $A$  into a rng. Left distributivity follows from the compatibility condition:

$$a(b+c) = -a + a \circ (b+c) - (b+c) = -a + a \circ b - a + a \circ c - c - b = ab + ac.$$

Similarly, since  $A$  is two-sided one proves  $(a+b)c = ac + bc$ . It remains to show that the multiplication is associative. On the one hand, using the first equality of (4.6) and the brace compatibility condition, we write

$$\begin{aligned} a(bc) &= a(-b + b \circ c - c) \\ &= -a + a \circ (-b + b \circ c - c) - (-b + b \circ c - c) \\ &= -a + a \circ (-b) - a + a \circ (b \circ c) - a + a \circ (-c) + c - b \circ c + b \\ &= a \circ (b \circ c) - a \circ b - a \circ c - b \circ c + a + b + c, \end{aligned}$$

since the group  $(A, +)$  is abelian. On the other hand, the second equality of (4.6) and Equality (4.5) imply that

$$\begin{aligned} (ab)c &= (-a + a \circ b - b)c - (-a + a \circ b - b) + (-a + a \circ b - b) \circ c - c \\ &= b - a \circ b + a + (-a) \circ c - c + (a \circ b) \circ c - c + (-b) \circ c - c \\ &= (a \circ b) \circ c - a \circ b - a \circ c - b \circ c + a + b + c. \end{aligned}$$

It then follows that the multiplication is associative.

Conversely, if  $A$  is a radical ring, say with ring multiplication  $(a, b) \mapsto ab$ , then  $a \circ b = a + ab + b$  turns  $A$  into a two-sided brace of abelian type. In fact, since  $A$  is a radical ring, then  $(A, +)$  is an abelian group and  $(A, \circ)$  is a group. Moreover,

$$a \circ (b + c) = a + a(b + c) + (b + c) = a + ab + ac + b + c = a \circ b - a + a \circ c.$$

Similarly one proves  $(a + b) \circ c = a \circ c - c + b \circ c$ .  $\square$

A brace is said to be *associative* if the operation  $(x, y) \mapsto x * y = \lambda_x(y) - y$  is associative. In [18, Question 2.1(2)], Cedó, Gateva-Ivanova and Smoktunowicz asked if associative braces of abelian type are always radical rings. To answer this question, we need some lemmas.

**Lemma 4.19.** *If  $A$  is an associative brace of abelian type, then  $(-a) * b = -(a * b)$  holds for all  $a, b \in A$ . In particular,  $(-a) \circ b = 2b - (a \circ b)$  for all  $a, b \in A$ .*

*Proof.* The associativity implies that

$$\begin{aligned} (a * (-a)) * b &= (a * (-a) + a + (-a)) * b \\ &= a * ((-a) * b) + (-a) * b + a * b \\ &= (a * (-a)) * b + (-a) * b + a * b \end{aligned}$$

and therefore  $(-a) * b = -(a * b)$ . From this the claim follows.  $\square$

If  $A$  is a brace of abelian type, then one proves by induction that

$$a \circ \left( \sum_{i=1}^n b_i - \sum_{j=1}^m c_j \right) = \sum_{i=1}^n a \circ b_i - \sum_{j=1}^m a \circ c_j + (m - n + 1)a \quad (4.7) \quad \boxed{\text{eq:Lau}}$$

holds for all  $a, b, c \in A$ , see Exercise 4.7.

$\boxed{\text{thm:Lau}}$

**Theorem 4.20.** *If  $A$  is an associative brace of abelian type, then  $A$  is a radical ring.*

*Proof.* We need to prove that the right compatibility condition holds. Since  $A$  is associative,  $(a * b) * c = a * (b * c)$  for all  $a, b, c \in A$ . Write the associativity condition between  $a, b, c \in A$  as

$$(a \circ b - a - b) \circ c - (a \circ b - a - b) - c = a \circ (b \circ c - b - c) - a - (b \circ c - b - c),$$

which is equivalent to

$$a' \circ ((a \circ b - a - b) \circ c - a \circ b) = a' \circ (a \circ (b \circ c - b - c) - a - a - b \circ c + 2c).$$

By using the formula (4.7) with  $n = 1$  and  $m = 2$  in the right hand side and with  $n = m = 3$  in the left hand side,

$$a' \circ (a \circ b - a + (-b)) = b + a' \circ (-b)$$

Now (4.7) with  $n = 2$  and  $m = 1$  implies that the associativity of  $A$  is equivalent to

$$(b + a' \circ (-b)) \circ c + c = b \circ c + a' \circ (-b) \circ c. \quad (4.8)$$

eq:asociatividad

Let  $b, c \in A$ . If  $d \in A$ , then there exists  $a \in A$  such that  $d = a' \circ (-b)$ . Equality (4.8) implies that

$$(b + d) \circ c + c = b \circ c + d \circ c. \quad \square$$

The previous result is not true if the brace is not of abelian type.

Now we show a brace that is not two-sided:

#### Example 4.21.

In Proposition 2.56 we used radical rings to produce examples of solutions. A natural question arises: Does one need radical rings? Surprisingly, radical rings are just the tip of the iceberg.

thm:YB

**Theorem 4.22.** *Let  $A$  be a brace. Then  $(A, r)$ , where*

$$r: A \times A \rightarrow A \times A, \quad r(x, y) = (-x + x \circ y, (-x + x \circ y)' \circ x \circ y),$$

*is a solution.*

*Proof.* By Theorem 1.11, since  $x \circ y = (-x + x \circ y) \circ ((-x + x \circ y)' \circ x \circ y)$  for all  $x, y \in A$ , we only need to check that  $x \triangleright y = \lambda_x(y) = -x + x \circ y$  is a left action of  $(A, \circ)$  on the set  $A$  and that  $x \triangleleft y = \mu_y(x) = (-x + x \circ y)' \circ x \circ y$  is a right action of  $(A, \circ)$  on the set  $A$ . For the left action we use Proposition 4.12 and for the right action we use Proposition 4.13.  $\square$

In Theorem 4.22 it is possible to prove that the solution is involutive if and only if the additive group of the brace is abelian. The next result generalizes this fact. We shall need a lemma.

lem:|r|

**Lemma 4.23.** *Let  $A$  be a brace and  $r$  be its associated solution. Then*

$$\begin{aligned} r^{2n}(a, b) &= (-n(a \circ b) + a + n(a \circ b), \\ &\quad (-n(a \circ b) + a + n(a \circ b))' \circ a \circ b), \end{aligned} \quad (4.9)$$

eq:r^2n

$$\begin{aligned} r^{2n+1}(a, b) &= (-n(a \circ b) - a + (n+1)(a \circ b), \\ &\quad (-n(a \circ b) - a + (n+1)(a \circ b))' \circ a \circ b), \end{aligned} \quad (4.10)$$

eq:r^2n+1

for all  $n \geq 0$ . Moreover, the following statements hold:

- 1)  $r^{2n} = \text{id}$  if and only if  $a + nb = nb + a$  for all  $a, b \in A$ .  
 2)  $r^{2n+1} = \text{id}$  if and only if  $\lambda_a(b) = n(a \circ b) + a - n(a \circ b)$  for all  $a, b \in A$ .

*Proof.* It suffices to prove (4.9) and (4.10). We proceed by induction on  $n$ . The case  $n = 0$  is trivial for (4.9) and (4.10). Assume that the claim holds for some  $n > 0$ . If  $n$  is even, by applying the map  $r$  to Equation (4.9) we obtain that

$$\begin{aligned} r^{2n+1}(a, b) &= r(-n(a \circ b) + a + n(a \circ b), (-n(a \circ b) + a + n(a \circ b))' \circ a \circ b) \\ &= (-n(a \circ b) - a + (n+1), (-n(a \circ b) - a + (n+1)(a \circ b))' \circ a \circ b). \end{aligned}$$

Thus Equation (4.10) holds. If  $n$  is odd, a similar argument shows that (4.9) holds. The other claims follow easily from Equations (4.9) and (4.10).  $\square$

Recall that the (minimal) *exponent*  $\exp(G)$  of a finite group  $G$  is the minimal  $n$  such that  $g^n = 1$  for all  $g \in G$ .

thm: |r|

**Theorem 4.24.** *Let  $A$  be a finite brace with more than one element and let  $G$  be the additive group of  $A$ . If  $r$  is the solution associated with  $A$ , then  $r$  has order  $2\exp(G/Z(G))$ .*

*Proof.* Let  $n$  be such that  $r$  has odd order, say  $r^{2n+1} = \text{id}$ . By applying Lemma 4.23 one obtains that  $-a + (n+1)(a \circ b) = n(a \circ b) + a$  for all  $a, b \in A$ . In particular, if  $b = 0$ , then  $a = 0$ , a contradiction. Therefore we may assume that the order of the permutation  $r$  is  $2n$ , where

$$n = \min\{k : kb + a = a + kb \text{ for all } a, b \in A\}.$$

Now one computes

$$\begin{aligned} n &= \min\{k : kb \in Z(G) \text{ for all } b \in A\} \\ &= \min\{k : k(b + Z(G)) = Z(G) \text{ for all } b \in A\} = \exp(G/Z(G)). \end{aligned} \quad \square$$

An immediate consequence:

**Corollary 4.25.** *Let  $A$  be a finite brace and  $r$  be its associated solution. Then  $r$  is involutive if and only if  $A$  is of abelian type.*

## 4B

**Definition 4.26.** Let  $A$  be a brace. A *subbrace* of  $A$  is a non-empty subset  $B$  of  $A$  such that  $(B, +)$  is a subgroup of  $(A, +)$  and  $(B, \circ)$  is a subgroup of  $(A, \circ)$ .

**Definition 4.27.** Let  $A$  be a brace. A *left ideal* of  $A$  is a subgroup  $(I, +)$  of  $(A, +)$  such that  $\lambda_a(I) \subseteq I$  for all  $a \in A$ , i.e.  $\lambda_a(x) \in I$  for all  $a \in A$  and  $x \in I$ . A *strong left ideal* of  $A$  is a left ideal  $I$  of  $A$  such that  $(I, +)$  is a normal subgroup of  $(A, +)$ .

**Proposition 4.28.** *A left ideal  $I$  of a brace  $A$  is a subbrace of  $A$ .*

*Proof.* We need to prove that  $(I, \circ)$  is a subgroup of  $(A, \circ)$ . Clearly  $I$  is non-empty, as it is an additive subgroup of  $A$ . If  $x, y \in I$ , then  $x \circ y = x + \lambda_x(y) \in I + I \subseteq I$  and  $x' = -\lambda_x(x) \in I$ .  $\square$

**Example 4.29.** Let  $A$  be a brace. Then

$$\text{Fix}(A) = \{a \in A : \lambda_x(a) = a \text{ for all } x \in A\}$$

is a left ideal of  $A$ .

**Definition 4.30.** An *ideal* of  $A$  is a strong left ideal  $I$  of  $A$  such that  $(I, \circ)$  is a normal subgroup of  $(A, \circ)$ .

In general

$$\{\text{subbraces}\} \subsetneq \{\text{left ideals}\} \subsetneq \{\text{strong left ideals}\} \subsetneq \{\text{ideals}\}.$$

For example,  $\text{Fix}(A)$  is not a strong left ideal of  $A$ .

**Example 4.31.** Consider the semidirect product  $A = \mathbb{Z}/(3) \rtimes \mathbb{Z}/(2)$  of the trivial braces  $\mathbb{Z}/(3)$  and  $\mathbb{Z}/(2)$  via the non-trivial action of  $\mathbb{Z}/(2)$  over  $\mathbb{Z}/(3)$ . Then

$$\lambda_{(x,y)}(a,b) = (x,y)(a,b) - (x,y) = (x + (-1)^y a, y + b) - (x,y) = ((-1)^y a, b).$$

Then  $\text{Fix}(A) = \{(0,0), (0,1)\}$  is not a normal subgroup of  $(A, \circ)$  and hence  $\text{Fix}(A)$  is not a strong left ideal of  $A$ .

**Example 4.32.** Let  $f: A \rightarrow B$  be a brace homomorphism. Then  $\ker f$  is an ideal of  $A$ .

Let  $I$  and  $J$  be ideals of a  $A$ . Then  $I \cap J$  is an ideal of  $A$ , see Exercise 4.9. The sum  $I + J$  of  $I$  and  $J$  is defined as the additive subgroup of  $A$  generated by all the elements of the form  $u + v$ ,  $u \in I$  and  $v \in J$ .

**Proposition 4.33.** *Let  $A$  be a brace and let  $I$  and  $J$  be ideals of  $A$ . Then  $I + J$  is an ideal of  $A$ .*

*Proof.* Let  $a \in A$ ,  $u \in I$  and  $v \in J$ . Then  $\lambda_a(u + v) \in I + J$  and hence it follows that  $\lambda_a(I + J) \subseteq I + J$ . Moreover,

$$(u + v) * a = (u \circ \lambda_u^{-1}(v)) * a = u * (\lambda_u^{-1}(v) * a) + \lambda_u^{-1}(v) * a + u * a \in I + J.$$

This formula implies that

$$a \circ (u + v) \circ a' = a + \lambda_a((u + v) + (u + v) * a') - a \in I + J.$$

Thus it follows that  $a \circ (I + J) \circ a' \subseteq I + J$ .

Finally  $I + J$  is a normal subgroup of  $(A, +)$  since

$$a + \sum_k (u_k + v_k) - a = \sum_k ((a + u_k - a) + (a + v_k - a)) \in I + J$$

whenever  $u_k \in I$  and  $v_k \in J$  for all  $k$ .  $\square$

**Definition 4.34.** Let  $A$  be a brace. The subset  $\text{Soc}(A) = \ker \lambda \cap Z(A, +)$  is the *socle* of  $A$ .

lem:socle

**Lemma 4.35.** Let  $A$  be a brace and  $a \in \text{Soc}(A)$ . Then

$$b + b \circ a = b \circ a + b \quad \text{and} \quad \lambda_b(a) = b \circ a \circ b'$$

both hold for all  $b \in A$ .

*Proof.* Let  $b \in A$ . Since  $b' \circ (b \circ a + b) = a - b'$  and  $b' \circ (b + b \circ a) = -b' + a$ , the first claim follows since  $a \in Z(A, +)$ . Now we prove the second claim:

$$b \circ a \circ b' = b \circ (a \circ b') = b \circ (a + b') = b \circ a - b = -b + b \circ a = \lambda_b(a). \quad \square$$

pro:socle

**Proposition 4.36.** Let  $A$  be a brace. Then  $\text{Soc}(A)$  is an ideal of  $A$ .

*Proof.* Clearly  $0 \in \text{Soc}(A)$ , since  $\lambda$  is a group homomorphism. Let  $a, b \in \text{Soc}(A)$  and  $c \in A$ . Since  $b \circ (-b) = b + (-b) = 0$ , it follows that  $b' = -b \in \text{Soc}(A)$ . The calculation

$$\lambda_{a-b}(c) = \lambda_{a \circ b'}(c) = \lambda_a \lambda_b^{-1}(c) = c,$$

implies that  $a - b \in \ker \lambda$ . Since  $a - b \in Z(A, +)$ , it follows that  $(\text{Soc}(A), +)$  is a normal subgroup of  $(A, +)$ .

For each  $d \in A$ ,  $a + c' \circ d = c' \circ d + a$  by Lemma 4.35. Then

$$\begin{aligned} d + \lambda_c(a) &= d - c + c \circ a = c \circ (c' \circ d + a) \\ &= c \circ (a + c' \circ d) = c \circ a - c + d = -c + c \circ a + d = \lambda_c(a) + d, \end{aligned}$$

that is  $\lambda_c(a)$  is central in  $(A, +)$ . Moreover,

$$\begin{aligned} \lambda_c(a) + d &= -c + c \circ a + d = c \circ a - c + d \\ &= c \circ (a + (c' \circ d)) = c \circ a \circ c' \circ d = \lambda_c(a) \circ d \end{aligned}$$

and hence

$$\lambda_{\lambda_c(a)}(d) = -\lambda_c(a) + \lambda_c(a) \circ d = -\lambda_c(a) + \lambda_c(a) + d = d.$$

Therefore  $\text{Soc}(A)$  is a strong left ideal of  $A$ . In fact,  $\text{Soc}(A)$  is an ideal of  $A$ , as  $c \circ a \circ c' = \lambda_c(a) \in \text{Soc}(A)$ .  $\square$

As a corollary we obtain that the socle of a brace  $A$  is a trivial brace of abelian type. In particular, if  $a \in \text{Soc}(A)$ , then  $a$  is a central element such that  $a \circ b = a + b$  for all  $b \in B$ .

pro:soc\_kernels

**Proposition 4.37.** Let  $A$  be a brace. Then  $\text{Soc}(A) = \ker \lambda \cap \ker \mu$ .

*Proof.* Let  $a \in \text{Soc}(A)$ . Then  $\lambda_a = \text{id}$  and  $a \in Z(A, +)$ . Let  $c = \mu_a(b) = \lambda_b(a)' \circ b \circ a$ . Then  $b \circ a = \lambda_b(a) \circ c = (-b + b \circ a) \circ c$ . Since this is equivalent to

$$a \circ c' = b' \circ (-b + b \circ a) = b' \circ (-b) - b' + a = b' + a = a + b' = a \circ b',$$

it follows that  $c' = b'$  and therefore  $c = b$ . Thus  $a \in \ker \lambda \cap \ker \mu$ .

Conversely, let  $a \in \ker \lambda \cap \ker \mu$  and  $b \in A$ . Then  $b' = \mu_a(b') = \lambda_{b'}(a)' \circ b' \circ a$ , so  $\lambda_{b'}(a) = b' \circ a \circ b$ . Now

$$b + a = b \circ \lambda_b^{-1}(a) = b \circ \lambda_{b'}(a) = b \circ b' \circ a \circ b = a \circ b = a + \lambda_a(b) = a + b$$

implies that  $a \in \text{Soc}(A)$ .  $\square$

Another important ideal was defined in [15].

**Definition 4.38.** Let  $A$  be a brace. The *annihilator* of  $A$  is defined as the set  $\text{Ann}(A) = \text{Soc}(A) \cap Z(A, \circ)$ .

Note that  $\text{Ann}(A) \subseteq \text{Fix}(A)$ .

**Proposition 4.39.** The annihilator of a brace  $A$  is an ideal of  $A$ .

*Proof.* Let  $a \in A$  and  $x \in \text{Ann}(A)$ . Since  $\text{Ann}(A) \subseteq Z(A, +) \cap Z(A, \circ)$ , we only need to note that  $\lambda_a(x) = x \in \text{Ann}(A)$ .  $\square$

If  $X$  and  $Y$  are subsets of a brace  $A$ ,  $X * Y$  is defined as the subgroup of  $(A, +)$  generated by elements of the form  $x * y$ ,  $x \in X$  and  $y \in Y$ , i.e.

$$X * Y = \langle x * y : x \in X, y \in Y \rangle_+.$$

pro:A\*I

**Proposition 4.40.** Let  $A$  be a brace. A subgroup  $I$  of  $(A, +)$  is a left ideal of  $A$  if and only if  $A * I \subseteq I$ .

*Proof.* Let  $a \in A$  and  $x \in I$ . If  $I$  is a left ideal, then  $a * x = \lambda_a(x) - x \in I$ . Conversely, if  $A * I \subseteq I$ , then  $\lambda_a(x) = a * x + x \in I$ .  $\square$

pro:I\*A

**Proposition 4.41.** Let  $A$  be a brace. A normal subgroup  $I$  of  $(A, +)$  is an ideal of  $A$  if and only if  $\lambda_a(I) \subseteq I$  for all  $a \in A$  and  $I * A \subseteq I$ .

*Proof.* Let  $x \in I$  and  $a \in A$ . Assume first that  $I$  is invariant under the action of  $\lambda$  and that  $I * A \subseteq I$ . Then

$$\begin{aligned} a \circ x \circ a' &= a + \lambda_a(x \circ a') \\ &= a + \lambda_a(x + \lambda_x(a')) = a + \lambda_a(x) + \lambda_a \lambda_x(a') + a - a \\ &= a + \lambda_a(x + \lambda_x(a') - a') - a = a + \lambda_a(x + x * a') - a \end{aligned} \quad (4.11)$$

eq:trick:I\*A

and hence  $I$  is an ideal.

Conversely, assume that  $I$  is an ideal. Then  $I * A \subseteq I$  since

$$\begin{aligned} x * a &= -x + x \circ a - a \\ &= -x + a \circ (a' \circ x \circ a) - a = -x + a + \lambda_a(a' \circ x \circ a) - a \in I. \end{aligned} \quad \square$$

If  $A$  is a brace and  $I$  is an ideal of  $A$ , then  $a + I = a \circ I$  for all  $a \in A$ . Indeed,  $a \circ x = a + \lambda_a(x) \in a + I$  and  $a + x = a \circ \lambda_a^{-1}(x) = a \circ \lambda_{a'}(x) \in a \circ I$  for all  $a \in A$  and  $x \in I$ . This allows us to prove that there exists a unique brace structure over  $A/I$  such that the map

$$\pi: A \rightarrow A/I, \quad a \mapsto a + I = a \circ I,$$

is a brace homomorphism. The brace  $A/I$  is the *quotient brace* of  $A$  modulo  $I$ . It is possible to prove the isomorphism theorems for braces, see Exercises 4.14, 4.15, 4.16 and 4.17.

## Exercises

prob:left\_right

**4.1.** Prove that there exists a bijective correspondence between left and right braces.

**4.2.** Let  $p$  be a prime number. Prove that  $\mathbb{Z}/(p^2)$  is a brace of abelian type with the operation  $x \circ y = x + y + pxy$ .

**4.3.** Let  $A$  be a brace. Prove that

$$\mu_b(a) = \lambda_{\lambda_a(b)}^{-1}(-a \circ b + a + a \circ b).$$

prob:star

**4.4.** Let  $A$  be an additive (not necessarily abelian) group. Prove that a brace structure over  $A$  is equivalent to an operation  $A \times A \rightarrow A$ ,  $(a, b) \mapsto a * b$ , such that

$$a * (b + c) = a * b + b + a * c - b$$

holds for all  $a, b, c \in A$ , and the operation  $a \circ b = a + a * b + b$  turns  $A$  into a group.

prob:equivalences

**4.5.** Let  $(A, +, \circ)$  be a triple, where  $(A, +)$  and  $(A, \circ)$  are groups, and let  $\lambda: A \rightarrow \mathbb{S}_A$ ,  $a \mapsto \lambda_a$ ,  $\lambda_a(b) = -a + a \circ b$ . Prove that the following statements are equivalent:

- 1)  $A$  is a brace.
- 2)  $\lambda_{a \circ b}(c) = \lambda_a \lambda_b(c)$  for all  $a, b, c \in A$ .
- 3)  $\lambda_a(b + c) = \lambda_a(b) + \lambda_a(c)$  for all  $a, b, c \in A$ .

prob:2sided

**4.6.** Let  $A$  be a brace such that  $\lambda_a(a) = a$  for all  $a \in A$ . Then  $A$  is two-sided.

prob:Lau

**4.7.** Prove Equality (4.7).

prob:radical

**4.8.** Recall that two-sided braces are equivalent to radical rings. Prove that under this equivalence, (left) ideals of the radical ring correspond to (left) ideals of the associated brace.

prob:sum\_ideals

**4.9.** Prove that the intersection of ideals is an ideal.

**4.10.** Let  $A$  be a brace and  $I$  be a characteristic subgroup of the additive group of  $A$ . Prove that  $I$  is a left ideal of  $A$ .



**4.11.** Let  $A = \dots$ . Prove that  $A$  has only three ideals: ... Let  $I$  be the ideal of  $A$  of size four. Prove that  $A * I$  has size two and hence it is not an ideal of  $A$ .

prob:Bachiller1

**4.12.** Prove that the socle of a brace  $A$  is the kernel of the group homomorphism  $(A, \circ) \rightarrow \text{Aut}(A, +) \times \mathbb{S}_A$ ,  $a \mapsto (\lambda_a, \mu_a^{-1})$ .

prob:Bachiller2

**4.13.** Prove that the socle of a brace  $A$  is the kernel of the group homomorphism  $(A, \circ) \rightarrow \text{Aut}(A, +) \times \text{Aut}(A, +)$ ,  $a \mapsto (\lambda_a, \xi_a)$ , where  $\xi_a(b) = a + \lambda_a(b) - a$ .

prob:iso1

**4.14.** Let  $f: A \rightarrow B$  be a brace homomorphism. Prove that  $A/\ker f \simeq f(A)$ .

prob:iso2

**4.15.** Let  $A$  be a brace and  $B$  be a subbrace of  $A$ . If  $I$  is an ideal of  $B$ , then  $B \circ I$  is a subbrace of  $B$ ,  $B \cap I$  is an ideal of  $B$  and  $(B \circ I)/I \simeq B/(B \cap I)$ .

prob:iso3

**4.16.** Let  $A$  be a brace and  $I$  and  $J$  be ideals of  $A$ . If  $I \subseteq J$ , then  $A/J \simeq (A/I)/(J/I)$ .

prob:correspondence

**4.17.** Let  $A$  be a brace and  $I$  be an ideal of  $A$ . There is a bijective correspondence between (left) ideals of  $A$  containing  $I$  and (left) ideals of  $A/I$ .

## Notes

Braces of abelian type were introduced by Rump in [53] for studying involutive solutions to the YBE. Rump's definition was reformulated by Cedó, Jespers and Okniński in [20]. With this definition at hand, Guarnieri and Vendramin introduced arbitrary braces in [34].

Exercise 4.5 combines results of Bachiller, Rump [53] and Gateva-Ivanova [32]. Exercise 4.6 comes from [20].

Theorem 4.18 was proved by Rump in [53].

Theorem 4.20 was proved by Lau [43] and independently by Kinyon (unpublished). It answers a question of Cedó, Gateva-Ivanova and Smoktunowicz, see [18].

Theorem 4.22 was proved for braces of abelian type appears implicit in the work [53] of Rump, see also [20]. The general case was proved by Guarnieri and Vendramin in [34].

Theorem 4.24 was proved by Smoktunowicz and Vendramin in [60].

Exercise 4.6 comes from [20]. Exercises 4.12 and 4.13 appear in [8].

The socle was defined by Rump in [53]. The annihilator first appeared in the work [15] of Catino, Colazzo and Stefanell.



## Chapter 5

### Complements

cocycles

**A**

An **extension** of  $K$  by  $Q$  is a short exact sequence

$$1 \longrightarrow K \longrightarrow G \longrightarrow Q \longrightarrow 1$$

This means that  $G$  is a group with a normal subgroup  $N$  isomorphic to  $K$  such that  $G/N \simeq Q$ .

**Example 5.1.**  $C_6$  and  $\mathbb{S}_3$  are both extensions of  $C_3$  by  $C_2$ .

**Example 5.2.**  $C_6$  is an extension of  $C_2$  by  $C_3$ .

**Example 5.3.** The direct product  $K \times Q$  of the groups  $K$  and  $Q$  is an extension of  $K$  by  $Q$  and an extension of  $Q$  by  $K$ .

**Example 5.4.** Let  $G$  be an extension of  $K$  by  $Q$ . If  $L$  is a subgroup of  $G$  containing  $K$ , then  $L$  is an extension of  $K$  by  $L/K$ .

Let  $E : 1 \longrightarrow K \longrightarrow G \longrightarrow Q \longrightarrow 1$  be an extension. A **lifting** of  $E$  is a map  $\ell : Q \rightarrow G$  such that  $p(\ell(x)) = x$  for all  $x \in Q$ .

xca:lifting

**Exercise 5.5.** Let  $E : 1 \longrightarrow K \longrightarrow G \xrightarrow{p} Q \longrightarrow 1$  be an extension.

- 1) If  $\ell : Q \rightarrow G$  is a lifting, then  $\ell(Q)$  is a transversal of  $\ker p$  in  $G$ .
- 2) Each transversal of  $\ker p$  in  $G$  induces a lifting  $\ell : Q \rightarrow G$ .
- 3) If  $\ell : Q \rightarrow G$  is a lifting, then  $\ell(xy) \ker p = \ell(x)\ell(y) \ker p$ .

An extension  $E$  **splits** if there is a lifting of  $E$  that is a group homomorphism.

Let  $Q$  and  $K$  be groups. Assume that  $Q$  acts by automorphism on  $K$ . A map  $\varphi : Q \rightarrow K$  is said to be a 1-**cocycle** (or a derivation) if

$$\varphi(xy) = \varphi(x)(x \cdot \varphi(y))$$

for all  $x, y \in Q$ . The set of 1-cocycles  $Q \rightarrow K$  is defined as

$$\text{Der}(Q, K) = Z^1(Q, K) = \{\delta: Q \rightarrow K : \delta \text{ is 1-cocycle}\}.$$

**Example 5.6.** Let  $Q$  acts on  $K$  by automorphisms. For each  $k \in K$ , the map  $Q \rightarrow K$ ,  $x \mapsto [k, x] = kxk^{-1}x^{-1}$ , is a derivation.

xca:1cocycle

**Exercise 5.7.** Let  $\varphi: Q \rightarrow K$  be a 1-cocycle.

- 1)  $\varphi(1) = 1$ .
- 2)  $\varphi(y^{-1}) = (y^{-1} \cdot \varphi(y))^{-1} = y^{-1} \cdot \varphi(y)^{-1}$ .
- 3) The set  $\ker \varphi = \{x \in Q : \varphi(x) = 1\}$  is a subgroup of  $Q$ .

A subgroup  $K$  of  $G$  admits a **complement**  $Q$  if  $G$  admits an exact factorization through  $K$  and  $Q$ , i.e.  $G = KQ$  with  $K \cap Q = \{1\}$ . A classical example is the semidirect product  $G = K \rtimes Q$ , where  $K$  is a normal subgroup of  $G$  and  $Q$  is a subgroup of  $G$  such that  $K \cap Q = \{1\}$ .

thm:complements

**Theorem 5.8.** Let  $Q$  acts by automorphism on  $K$ . Then there exists a bijective correspondence between the set  $\mathcal{C}$  of complements  $K$  in  $K \rtimes Q$  and the set  $\text{Der}(Q, K)$  of 1-cocycles  $Q \rightarrow K$ .

*Proof.* Since  $Q$  acts by conjugation on  $K$ , it follows that  $\delta \in \text{Der}(Q, K)$  if and only if  $\delta(xy) = \delta(x)x\delta(y)x^{-1}$  for all  $x, y \in Q$ . In this case, one obtains that  $\delta(1) = 1$  and  $\delta(x^{-1}) = x^{-1}\delta(x)^{-1}x$ .

Let  $C \in \mathcal{C}$ . If  $x \in Q$ , then there exist unique elements  $k \in K$  and  $c \in C$  such that  $x = k^{-1}c$ . Hence the map  $\delta_C: Q \rightarrow K$ ,  $x \mapsto k$ , is well-defined. Moreover,  $\delta_C(x)x = c \in C$ .

We claim that  $\delta_C \in \text{Der}(Q, K)$ . If  $x, x_1 \in Q$ , we write  $x = k^{-1}c$  and  $x_1 = k_1^{-1}c_1$  for  $k, k_1 \in K$  and  $c, c_1 \in C$ . Since  $K$  is a normal subgroup of the semidirect product  $K \rtimes Q$ , we can write  $xx_1$  as  $xx_1 = k_2c_2$ , where  $k_2 = k^{-1}(ck_1^{-1}c^{-1}) \in K$ ,  $c_2 = cc_1 \in C$ . Thus  $\delta_C(xx_1)xx_1 = cc_1 = \delta_C(x)x\delta_C(x_1)x_1$  implies that  $\delta_C(xx_1) = \delta_C(x)x\delta_C(x_1)x^{-1}$ . So there is a map  $F: \mathcal{C} \rightarrow \text{Der}(Q, K)$ ,  $F(C) = \delta_C$ .

We now construct a map  $G: \text{Der}(Q, K) \rightarrow \mathcal{C}$ . For each  $\delta \in \text{Der}(Q, K)$  we find a complement  $\Delta$  of  $K$  in  $K \rtimes Q$ . Let  $\Delta = \{\delta(x)x : x \in Q\}$ . We claim that  $\Delta$  is a subgroup of  $K \rtimes Q$ . Since  $\delta(1) = 1$ ,  $1 \in \Delta$ . If  $x, y \in Q$ , then  $\delta(x)x\delta(y)y = \delta(x)x\delta(y)x^{-1}xy = \delta(xy)xy \in \Delta$ . Finally, if  $x \in Q$ , then

$$(\delta(x)x)^{-1} = x^{-1}\delta(x)^{-1}xx^{-1} = \delta(x^{-1})x^{-1}.$$

Let us prove that  $\Delta \cap K = \{1\}$ . If  $x \in Q$  is such that  $\delta(x)x \in K$ , then since  $\delta(x) \in K$ , it follows that  $x \in K \cap Q = 1$ . If  $g \in G$ , then there are unique  $k \in K$  and  $x \in Q$  such that  $g = kx$ . We write  $g = k\delta(x)^{-1}\delta(x)x$ . Since  $k\delta(x)^{-1} \in K$  and  $\delta(x)x \in \Delta$ , we conclude that  $G = K\Delta$ . Thus there is a well-defined map  $G: \text{Der}(Q, K) \rightarrow \mathcal{C}$ ,  $G(\delta) = \Delta$ .

We claim that  $G \circ F = \text{id}_{\mathcal{C}}$ . Let  $C \in \mathcal{C}$ . Then

$$G(F(C)) = G(\delta_C) = \{\delta_C(x)x : x \in Q\} = C,$$

by construction. (We know that  $\delta_C(x)x \in C$ . Conversely, if  $c \in C$ , we write  $c = kx$  for unique elements  $k \in K$  and  $x \in Q$ . Thus  $x = k^{-1}c$  and hence  $c = \delta_C(x)x$ .)

Finally, we prove that  $F \circ G = \text{id}_{\text{Der}(Q,K)}$ . Let  $\delta \in \text{Der}(Q,K)$ . Then

$$F(G(\delta)) = F(\Delta) = \delta_\Delta.$$

Finally, we need to show that  $\delta_\Delta = \delta$ . Let  $x \in Q$ . There exists  $\delta(y)y \in \Delta$  for some  $y \in Q$  such that  $x = k^{-1}\delta(y)y$ . Thus  $\delta_\Delta(x)x = \delta(y)y$  and hence  $\delta(x) = \delta(y)$  by the uniqueness.  $\square$

Let the group  $Q$  acts by automorphism on  $K$ . A derivation  $\delta \in \text{Der}(Q,K)$  is said to be **inner** if there exists  $k \in K$  such that  $\delta(x) = [k, x]$  for all  $x \in Q$ . The set of **inner derivations** will be denoted by

$$\text{Inn}(Q, K) = B^1(Q, K) = \{\delta \in \text{Der}(Q, K) : \delta \text{ is inner}\}.$$

An inner derivation is also called a **1-coboundary**.

theorem:Sysak

**Theorem 5.9 (Sysak).** *Sean  $Q$  y  $K$  grupos tales que  $Q$  actúa por automorfismos en  $K$ . Sea  $\delta \in \text{Der}(Q, K)$ .*

- 1)  $\Delta = \{\delta(x)x : x \in Q\}$  es un complemento para  $K$  en  $K \rtimes Q$ .
- 2)  $\delta \in \text{Inn}(Q, K)$  si y sólo si  $Q$  y  $\Delta$  son conjugados en  $K$ .
- 3)  $\ker \delta = Q \cap \Delta$ .
- 4)  $\delta$  es sobreyectiva si y sólo si  $K \rtimes Q = \Delta Q$ .

*Proof.* In the proof of Theorem 5.8 we found that  $\Delta$  is a complement of  $K$  in  $K \rtimes Q$ .

Let us prove the second statement. If  $\delta$  is inner, then there exists  $k \in K$  such that  $\delta(x) = [k, x] = kxk^{-1}x^{-1}$  for all  $x \in Q$ . Since  $\delta(x)x = kxk^{-1}$  for all  $x \in Q$ ,  $\Delta = kQk^{-1}$ . Conversely, if there exists  $k \in K$  such that  $\Delta = kQk^{-1}$ , for each  $x \in Q$  there exists  $y \in Q$  such that  $\delta(x)x = kyk^{-1}$ . Since  $[k, y] = kyk^{-1}y^{-1} \in K$ ,  $\delta(x) \in K$  and  $\delta(x)x = [k, y]y \in KQ$ , we conclude that  $x = y$  and hence  $\delta(x) = [k, x]$ .

Let us prove the third statement. If  $x \in Q$  is such that  $\delta(x)x = y \in Q$ , then

$$\delta(x) = yx^{-1} \in K \cap Q = \{1\}.$$

Conversely, if  $x \in Q$  is such that  $\delta(x) = 1$ , then  $x = \delta(x)x \in Q \cap \Delta$ .

Finally we prove the fourth statement. If  $\delta$  is surjective, then for each  $k \in K$  there exists  $y \in Q$  such that  $\delta(y) = k$ . Thus  $K \rtimes Q \subseteq \Delta Q$ , as

$$kx = \delta(y)x = (\delta(y)y)y^{-1}x \in \Delta Q.$$

Moreover,  $\Delta Q \subseteq K \rtimes Q$ , as  $\delta(x) \in K$  for all  $x \in Q$ . Conversely, if  $k \in K$  y  $x \in Q$  there exist  $y, z \in Q$  such that  $kx = \delta(y)yz$ . Then it follows that  $k = \delta(y)$ .  $\square$

A group  $G$  admits a **triple factorization** if there are subgroups  $A$ ,  $B$  and  $M$  such that  $G = MA = MB = AB$  y  $A \cap M = B \cap M = \{1\}$ . The following result is An immediate consequence of Sysak's theorem:

**Corollary 5.10.** *If the group  $Q$  acts by automorphisms on  $K$  and  $\delta \in \text{Der}(Q, K)$  is surjective, then  $G = K \rtimes Q$  admits a triple factorization.*

Another consequence:

xca:kerlcocycle

**Exercise 5.11.** Let  $\delta \in \text{Der}(Q, K)$ .

- 1) Prove that  $\delta$  is injective if and only if  $\ker \delta = \{1\}$ .
- 2) Prove that if  $\delta$  is bijective, then  $K$  admits a complement  $\Delta$  in  $K \rtimes Q$  such that  $K \rtimes Q = K \rtimes \Delta = \Delta Q$  and  $Q \cap \Delta = \{1\}$ .

## B

lemma:lcocycle

**Lemma 5.12.** *Let  $G$  be a group and  $N$  be a normal subgroup of  $G$ . If  $G$  acts on  $N$  by conjugation and  $\varphi: G \rightarrow N$  is a 1-cocycle with kernel  $K$ , then  $\varphi(x) = \varphi(y)$  if and only if  $xK = yK$ . In particular,  $(G : K) = |\varphi(G)|$ .*

*Proof.* If  $\varphi(x) = \varphi(y)$ , then, since

$$\varphi(x^{-1}y) = \varphi(x^{-1})(x^{-1} \cdot \varphi(y)) = \varphi(x^{-1})(x^{-1} \cdot \varphi(x)) = \varphi(x^{-1}x) = \varphi(1) = 1,$$

we obtain that  $xK = yK$ . Conversely, if  $x^{-1}y \in K$ , then, since

$$1 = \varphi(x^{-1}y) = \varphi(x^{-1})(x^{-1} \cdot \varphi(y)),$$

we conclude that  $\varphi(y) = x \cdot \varphi(x^{-1})^{-1}$ . Thus  $\varphi(x) = \varphi(y)$ .

The second claim is now trivial, as  $\varphi$  is constant in each coset of  $K$  and there are  $(G : K)$  different possible values.  $\square$

lemma:d

**Lemma 5.13.** *Let  $G$  be a finite group,  $N$  be a normal abelian subgroup of  $G$  and  $S, T$  and  $U$  be transversal of  $N$  in  $G$ . Let*

$$d(S, T) = \prod st^{-1} \in N,$$

where the product is taken over all  $s \in S$  and  $t \in T$  such that  $sN = tN$ . The following statements hold:

- 1)  $d(S, T)d(T, U) = d(S, U)$ .
- 2)  $d(gS, gT) = gd(S, T)g^{-1}$  for all  $g \in G$ .
- 3)  $d(nS, S) = n^{(G:N)}$  for all  $n \in N$ .

*Proof.* If  $s \in S, t \in T$  and  $u \in U$  are such that  $sN = tN = uN$ , then, since  $N$  is abelian and  $(st^{-1})(tu^{-1}) = su^{-1}$ ,

$$d(S, T)d(T, U) = \prod (st^{-1})(tu^{-1}) = \prod su^{-1} = d(S, U).$$

Since  $sN = tN$  if and only if  $gsN = gtN$  for all  $g \in G$ ,

$$g \left( \prod st^{-1} \right) g^{-1} = \prod g s t^{-1} g^{-1} = \prod (gs)(gt)^{-1} = d(gS, gT).$$

Finally, since  $N$  is normal,  $nsN = sN$  for all  $n \in N$ . Thus

$$d(nS, S) = \prod (ns)s^{-1} = n^{(G:N)}. \quad \square$$

We now prove the first version of Schur–Zassenhaus’ theorem.

SchurZassenhaus:abelian

**Theorem 5.14 (Schur–Zassenhaus).** *Sea  $G$  un grupo finito y sea  $N$  un subgrupo normal abeliano de  $G$ . Si  $|N|$  y  $(G : N)$  son coprimos,  $N$  se complementa en  $G$ . Si  $N$  se complementa en  $G$ , todos los complementos son conjugados.*

*Proof.* Let  $T$  be a transversal of  $N$  in  $G$ . Let  $\theta: G \rightarrow N$ ,  $\theta(g) = d(gT, T)$ . Since  $N$  is abelian, Lemma 5.13 implies that  $\theta$  is a 1-cocycle, where  $G$  acts on  $N$  by conjugation:

$$\begin{aligned} \theta(xy) &= d(xyT, T) = d(xyT, xT)d(xT, T) \\ &= (xd(yT, T)x^{-1})d(xT, T) = (x \cdot \theta(y))\theta(x). \end{aligned}$$

*Claim.*  $\theta|_N: N \rightarrow N$  is surjective.

If  $n \in N$ , Lemma 5.13 implies that  $\theta(n) = d(nT, T) = n^{(G:N)}$ . Since  $|N|$  and  $(G : N)$  are coprime, there exist  $r, s \in \mathbb{Z}$  such that  $r|N| + s(G : N) = 1$ . Thus

$$n = n^{r|N| + s(G:N)} = (n^s)^{(G:N)} = \theta(n^s).$$

Let  $H = \ker \theta$ . We claim that  $H$  is a complement for  $N$ . We know that  $H$  is a subgroup of  $G$ . Since

$$|N| = |\theta(G)| = (G : H) = \frac{|G|}{|H|}$$

Lemma 5.12, FIXME Como  $N \cap H$  es un subgrupo de  $N$  y de  $H$ , entonces  $N \cap H = \{1\}$  pues  $|N|$  y  $(G : N) = |H|$  son coprimos. Como  $|NH| = |N||H| = |G|$ , se concluye que  $G = NH$  y entonces  $H$  es un complemento para  $N$ .

Veamos ahora que dos complementos para  $N$  son conjugados. Sea  $K$  un complemento de  $N$  en  $G$ . Como  $NK = G$  y  $N \cap K = \{1\}$ ,  $K$  es un transversal para  $N$ . Sea  $m = d(T, K) \in N$ . Como  $\theta|_N$  es sobreyectiva, existe  $n \in N$  tal que  $\theta(n) = m$ . Por el lema 5.13, para todo  $k \in K$  se tiene

$$kmk^{-1} = kd(T, K)k^{-1} = d(kT, kK) = d(kT, K) = d(kT, T)d(T, K) = \theta(k)m.$$

Entonces, como  $N$  es abeliano,  $\theta(n^{-1}) = m^{-1}$  y luego

$$\begin{aligned} \theta(nkn^{-1}) &= \theta(n)n\theta(kn^{-1})n^{-1} = m\theta(kn^{-1}) \\ &= m\theta(k)k\theta(n^{-1})k^{-1} = m\theta(k)km^{-1}k^{-1} = 1. \end{aligned}$$

Queda demostrado entonces que  $nKn^{-1} \subseteq H = \ker \theta$ . Como  $|K| = (G : N) = |H|$ , se concluye que  $nKn^{-1} = H$ .  $\square$

En el siguiente teorema vemos que no es necesario suponer que el subgrupo normal  $N$  es abeliano.

thm:SchurZassenhaus

**Theorem 5.15 (Schur–Zassenhaus).** *Sea  $G$  un grupo finito y sea  $N$  un subgrupo normal de  $G$ . Si  $|N|$  y  $(G : N)$  son coprimos entonces  $N$  se complementa en  $G$ .*

*Proof.* Procederemos por inducción en  $|G|$ . Si existe un subgrupo propio  $K$  de  $G$  tal que  $NK = G$  entonces, como  $(K : K \cap N) = (G : N)$  es coprimo con  $|N|$ , es también coprimo con  $|K \cap N|$ . Como además  $K \cap N$  es normal en  $K$ , por hipótesis inductiva,  $K \cap N$  se complementa en  $K$ , y luego existe un subgrupo  $H$  de  $K$  tal que  $|H| = (K : K \cap N) = (G : N)$ .

Supongamos entonces que no existe un subgrupo propio  $K$  de  $G$  tal que  $NK = G$ . Podemos suponer que  $N \neq 1$  (de lo contrario, basta tomar  $G$  como complemento de  $N$  en  $G$ ). Como  $N$  está contenido en todo subgrupo maximal de  $G$  (pues si existe un maximal  $M \subsetneq G$  tal que  $N \not\subseteq M$  entonces  $NM = G$ ), se tiene que  $N \subseteq \Phi(G)$ . Por el teorema de Frattini ??,  $\Phi(G)$  es nilpotente y luego  $N$  es nilpotente; en particular,  $Z(N) \neq \{1\}$ . Sea  $\pi: G \rightarrow G/Z(N)$  el morfismo canónico. Como  $N$  es normal en  $G$  y  $Z(N)$  es característico en  $N$ ,  $Z(N)$  es normal en  $G$ . Además

$$(\pi(G) : \pi(N)) = \frac{|\pi(G)|}{|\pi(N)|} = \frac{|G/Z(N)|}{|N/N \cap Z(N)|} = (G : N)$$

es coprimo con  $|N|$ , y entonces es también coprimo con  $|\pi(N)|$ . Por hipótesis inductiva,  $\pi(N)$  admite un complemento en  $G/Z(N)$ , digamos  $\pi(K)$  para algún subgrupo  $K$  de  $G$ . Luego  $G = NK$  pues  $\pi(G) = \pi(N)\pi(K) = \pi(NK)$ . Como entonces  $K = G$  (pues sabíamos que no existe  $K$  tal que  $G = NK$ ),  $\pi(N)$  es abeliano pues

$$\pi(Z(N)) = \pi(N) \cap \pi(K) = \pi(N) \cap \pi(G) = \pi(N).$$

Luego  $N \subseteq Z(N)$  es abeliano y entonces, por el teorema ??, el subgrupo  $N$  admite un complemento.  $\square$

urZassenhaus:conjugacion

**Theorem 5.16.** *Let  $G$  be a finite group and  $N$  be a normal subgroup of  $G$  such that  $|N|$  and  $(G : N)$  are coprime. If either  $N$  or  $G/N$  is solvable, then all complements of  $N$  in  $G$  are conjugate.*

*Proof.* Let  $G$  be a minimal counterexample, so there are complements  $K_1$  and  $K_2$  of  $N$  in  $G$  such that  $K_1$  and  $K_2$  are not conjugate and  $|G|$  is minimal with this property.

*Claim.* Each subgroup  $U$  of  $G$  satisfies the hypotheses of the theorem with respect to the normal subgroup  $U \cap N$ .

Since  $N$  is normal in  $G$ , the subgroup  $U \cap N$  is normal in  $U$ . Moreover,  $|U \cap N|$  and  $(U : U \cap N)$  are coprime, as  $|U \cap N|$  divides  $|N|$  and  $(U : U \cap N) = (UN : N)$  divides  $(G : N)$ . If  $G/N$  is solvable, then  $U/U \cap N$  is solvable since  $U/U \cap N$  is isomorphic to a subgroup of  $G/N$ . If  $N$  is solvable, the subgroup  $U \cap N$  is solvable.



*Claim.* If there is a normal subgroup  $L$  of  $G$  such that  $\pi(N)$  is normal in  $\pi(G)$ , where  $\pi: G \rightarrow G/L$  is the canonical map, then  $\pi(G)$  satisfies the hypotheses of the theorem with respect to  $\pi(N)$ . In this case, if  $H$  is a complement of  $N$  in  $G$ , then  $\pi(H)$  is a complement of  $\pi(N)$  in  $\pi(G)$ .

If  $N$  is solvable, then  $\pi(N)$  is solvable. If  $G/N$  is solvable, then the group  $\pi(G)/\pi(N) \simeq G/NL$  is solvable. Moreover,  $(\pi(G) : \pi(N)) = \frac{|G/L|}{|N/N \cap L|}$  divides the index  $(G : N)$  of  $N$  in  $G$ .

If  $H$  is a complement of  $N$  in  $G$ ,  $|\pi(H)|$  and  $|\pi(N)|$  are coprime. Thus  $\pi(H)$  is a complement of  $\pi(N)$ , as  $\pi(G) = \pi(N)\pi(H) = \pi(NH)$  and  $\pi(N) \cap \pi(H) = \{1\}$ .

*Claim.*  $N$  is minimal normal in  $G$ .

Let  $M \neq \{1\}$  be a normal subgroup of  $G$  such that  $M \subseteq N$ . Let  $\pi: G \rightarrow G/M$  be the canonical map. The group  $\pi(G)$  satisfies the hypotheses of the theorem with respect to the normal subgroup  $\pi(N)$ . By the minimality of  $|G|$ , there exists  $x \in G$  such that  $\pi(xK_1x^{-1}) = \pi(K_2)$ . The subgroup  $U = MK_2$  satisfies the hypotheses of the theorem with respect to the normal subgroup  $U \cap N$ . Since  $xK_1x^{-1} \cup K_2 \subseteq U$ , we conclude that  $xK_1x^{-1}$  and  $K_2$  are complements of  $U \cap N$  in  $U$ . Thus  $MK_2 = G$ , as  $xK_1x^{-1}$  and  $K_2$  are not conjugate and  $|G|$  is minimal. Therefore  $M = N$ , as

$$\frac{|K_2|}{|M \cap K_2|} = (MK_2 : M) = (G : M) = \frac{|NK_2|}{|M|} = (N : M)|K_2|.$$

*Claim.*  $N$  is not solvable and  $G/N$  is solvable.

Otherwise, by Lemma 7.5,  $N$  is minimal normal and hence abelian. This yields a contradiction, as the previous version of the Schur–Zassenhaus’s theorem implies that  $K_1$  and  $K_2$  are conjugate.

Let  $p: G \rightarrow G/N$  be the canonical map and  $S$  be such that  $p(S)$  is minimal normal in  $p(G) = G/N$ . By Lemma 7.5,  $p(S)$  is a  $p$ -group for some prime number  $p$ . Since  $G = NK_1 = NK_2$  and  $N \subseteq S$ , Dedekind’s lemma implies that

$$S = N(S \cap K_1) = N(S \cap K_2).$$

Thus  $S \cap K_1$  and  $S \cap K_2$  are complements of  $N$  in  $S$ . Since

$$p(S) = p(S \cap K_1) = p(S \cap K_2)$$

is a  $p$ -group,  $p$  divides  $|S|$ . The group  $S$  satisfies the hypotheses of the theorem with respect to the normal subgroup  $N$ , so  $|N|$  and  $(S : N)$  are coprime. If  $p \mid |N|$ , then  $p \nmid (S : N) = |p(S)|$ , a contradiction. Therefore  $p \nmid |N|$ . This implies that  $S \cap K_1$  and  $S \cap K_2$  are Sylow  $p$ -subgroups of  $S$ , as

$$|S \cap K_1| = (S : N) = |S \cap K_2|.$$

By the second Sylow’s theorem, there exists  $s \in S$  such that

$$S \cap sK_1s^{-1} = S \cap K_2.$$

In particular,  $S \neq G$  by the minimality of  $|G|$ . Let

$$L = S \cap K_2 = S \cap sK_1s^{-1} \neq \{1\}.$$

Since  $S$  is normal in  $G$ , it follows that  $sK_1s^{-1} \cup K_2 \subseteq N_G(L)$  (because  $L$  is normal both in  $sK_1s^{-1}$  and in  $K_2$ ). The subgroups  $sK_1s^{-1} \subseteq N_G(L)$  and  $K_2 \subseteq N_G(L)$  are complements of  $N \cap N_G(L)$  in  $N_G(L)$ . Thus  $N_G(L) = G$  by the minimality of  $|G|$  (if  $N_G(L) \neq G$ , then both  $sK_1s^{-1}$  and  $K_2$  are conjugate in  $G$  because they are conjugate in  $N_G(L)$ ). Therefore  $L$  is normal in  $G$ .

Let  $\pi_L: G \rightarrow G/L$  be the canonical map. Since both  $\pi_L(K_1)$  and  $\pi_L(K_2)$  are complements of  $\pi_L(N)$  in  $\pi_L(G)$ , the minimality of  $|G|$  implies that there exists  $g \in G$  such that  $\pi_L(gK_1g^{-1}) = \pi_L(K_2)$ , so there exists  $g \in G$  such that  $(gK_1g^{-1})L = K_2L$ . Thus  $gK_1g^{-1} \cup K_2 \subseteq \langle K_2, L \rangle = K_2$ , as  $L \subseteq K_2$ . In conclusion,  $gK_1g^{-1} = K_2$ , a contradiction to the minimality of  $|G|$ .  $\square$

By Feit–Thompson’s theorem, we do not need to assume that  $N$  or  $G/N$  is solvable. Indeed, since every group of odd order is solvable and  $|N|$  and  $(G : N)$  are coprime, it follows that either  $|N|$  or  $(G : N)$  is odd.

## C

Let  $A$  be an additive group and  $G$  be a group and let  $G \times A \rightarrow A$ ,  $(g, a) \mapsto g \cdot a$ , is a left action of  $G$  on  $A$  by automorphisms. This means that the action of  $G$  on  $A$  satisfies  $g \cdot (a + b) = g \cdot a + g \cdot b$  for all  $g \in G$  and  $a, b \in A$ . A *bijective 1-cocycle* is a bijective map  $\pi: G \rightarrow A$  such that

$$\pi(gh) = \pi(g) + g \cdot \pi(h) \tag{5.1}$$

eq:1cocycle

for all  $g, h \in G$ . We now prove the equivalence between braces and bijective 1-cocycles.

thm:1cocycle

**Theorem 5.17.** *Over any additive group  $A$  the following data are equivalent:*

- 1) *A group  $G$  and a bijective 1-cocycle  $\pi: G \rightarrow A$ .*
- 2) *A brace structure over  $A$ .*

*Proof.* Consider on  $A$  a second group structure given by

$$a \circ b = \pi(\pi^{-1}(a)\pi^{-1}(b)) = a + \pi^{-1}(a) \cdot b$$

for all  $a, b \in A$ . Since  $G$  acts on  $A$  by automorphisms,

$$\begin{aligned} a \circ (b + c) &= \pi(\pi^{-1}(a)\pi^{-1}(b+c)) = a + \pi^{-1}(a) \cdot (b+c) \\ &= a + \pi^{-1}(a) \cdot b + \pi^{-1}(a) \cdot c = a \circ b + a \circ c \end{aligned}$$

holds for all  $a, b, c \in A$ .

Conversely, assume that the additive group  $A$  has a brace structure. Let  $G$  be the multiplicative group of  $A$  and  $\pi = \text{id}$ . By Proposition 4.12,  $a \mapsto \lambda_a$ , is a group homomorphism and hence  $G$  acts on  $A$  by automorphisms. Then (5.1) holds and therefore  $\pi: G \rightarrow A$  is a bijective 1-cocycle.  $\square$

The construction of Theorem 5.17 is functorial, see Exercise 5.1.

exa:d8q8

**Example 5.18.** Let

$$D_4 = \langle r, s : r^4 = s^2 = 1, srs = r^{-1} \rangle$$

be the dihedral group of eight elements and let

$$Q_8 = \{1, -1, i, -i, j, -j, k, -k\}$$

be the quaternion group of eight elements. Let  $\pi: Q_8 \rightarrow D_4$  be given by

$$\begin{array}{llll} 1 \mapsto 1, & -1 \mapsto r^2, & -k \mapsto r^3 s, & k \mapsto rs, \\ i \mapsto s, & -i \mapsto r^2 s, & j \mapsto r^3, & -j \mapsto r. \end{array}$$

Since  $\pi$  is bijective, a straightforward calculation shows that  $D_4$  with

$$x + y = xy, \quad x \circ y = \pi(\pi^{-1}(x)\pi^{-1}(y))$$

is a two-sided brace with additive group isomorphic to  $D_4$  and multiplicative group isomorphic to  $Q_8$ .

## Exercises

prob:1cocycle

**5.1.** Let  $\pi: G \rightarrow A$  and  $\eta: H \rightarrow B$  be bijective 1-cocycles. A *homomorphism* between these bijective 1-cocycles is a pair  $(f, g)$  of group homomorphisms  $f: G \rightarrow H$ ,  $g: A \rightarrow B$  such that

$$\begin{aligned} \eta \circ f &= g \circ \pi, \\ g(h \cdot a) &= f(h) \cdot g(a), \end{aligned} \quad a \in A, h \in G.$$

Bijjective 1-cocycles and homomorphisms form a category. For a given additive group  $A$  the full subcategory of the category of bijective 1-cocycles with objects  $\pi: G \rightarrow A$  is equivalent to the full subcategory of the category of braces with additive group  $A$ .

## Open problems

## Notes

In the case of braces of abelian type, Theorem 5.17 is implicit in the work of Rump, see [53, 54] or [20]. Similar results appear in the work of Etingof, Schedler and Soloviev [29], Lu, Yan and Zhu [45] and Soloviev [61]. In [28] Etingof and Gelaki give a method of constructing finite-dimensional complex semisimple triangular Hopf algebras. They show how any non-abelian group which admits a bijective 1-cocycle gives rise to a semisimple minimal triangular Hopf algebra which is not a group algebra.

## Chapter 6

### Nilpotent groups

#### A

If  $G$  is a group and  $x, y, z \in G$ , the conjugation (as a left action) will be denoted by  ${}^x y = xyx^{-1}$ . The commutator between  $x$  and  $y$  is then

$$[x, y] = xyx^{-1}y^{-1} = ({}^x y)y^{-1}.$$

We also write  $[x, y, z] = [x, [y, z]]$ . If  $X, Y$  and  $Z$  are subgroups of  $G$ , we write  $[X, Y, Z] = [X, [Y, Z]]$ . Note that  $[X, Y] = [Y, X]$ .

xca:HallWitt

**Exercise 6.1 (Hall–Witt).** Let  $G$  be a group and  $x, y, z \in G$ . Then

$$({}^y [x, y^{-1}, z]) ({}^z [y, z^{-1}, x]) ({}^x [z, x^{-1}, y]) = 1. \quad (6.1)$$

eq:HallWitt

Note that if  $G$  is such that  $[G, G]$  is central, then Jacobi’s identity turns out to be Jacobi’s identity.

lemma:3subgrupos

**Lemma 6.2 (three subgroups lemma).** Let  $X, Y$  and  $Z$  be subgroups of  $G$  such that  $[X, Y, Z] = [Y, Z, X] = \{1\}$ . Then  $[Z, X, Y] = \{1\}$ .

*Proof.* Since  $[x, y] \in C_G(z)$  implies that  $[X, Y] \subseteq C_G(Z)$ , it is enough to prove that  $[z, x^{-1}, y] = 1$  for all  $x \in X, y \in Y$  and  $z \in Z$ . Since  $[y^{-1}, z] \in [Y, Z]$ , it follows that  $[x, y^{-1}, z] \in [X, Y, Z] = \{1\}$ . Thus  ${}^y [x, y^{-1}, z] = 1$ . Similarly,  ${}^z [y, z^{-1}, x] = 1$ . Hence the Hall–Witt identity yields  $[z, x^{-1}, y] = 1$ .  $\square$

lemma:3subgrupos\_general

**Lemma 6.3.** Let  $N$  be a normal subgroup of  $G$  and let  $X, Y$  and  $Z$  be subgroups of  $G$ . If  $[X, Y, Z] \subseteq N$  and  $[Y, Z, X] \subseteq N$ , then  $[Z, X, Y] \subseteq N$ .

*Proof.* Let  $\pi: G \rightarrow G/N$  be the canonical map. Since  $[X, Y, Z] \subseteq N$ ,

$$\begin{aligned} \{1\} &= \pi([X, Y, Z]) = \pi([X, [Y, Z]]) \\ &= [\pi(X), \pi([Y, Z])] = [\pi(X), [\pi(Y), \pi(Z)]] = [\pi(X), \pi(Y), \pi(Z)]. \end{aligned}$$

Similarly one proves that  $[\pi(Y), \pi(Z), \pi(X)] = \{1\}$ . By the three subgroups lemma,  $[\pi(Z), \pi(X), \pi(Y)] = \{1\}$ , so  $[Z, X, Y] \subseteq N$ .  $\square$

The **lower central series** of a group  $G$  is the sequence  $\gamma_k(G)$ ,  $k \in \mathbb{N}$ , defined recursively as

$$\gamma_1(G) = G, \quad \gamma_{i+1}(G) = [G, \gamma_i(G)] \quad i \geq 1.$$

A group  $G$  is said to be **nilpotent** if there exists positive integer  $c$  such that  $\gamma_{c+1}(G) = \{1\}$ . The smallest  $c$  such that  $\gamma_{c+1}(G) = \{1\}$  is the **nilpotency index** (or **nilpotency class**) of  $G$ .

**Example 6.4.** A group is nilpotent of class one if and only if it is abelian.

**Example 6.5.** The group  $G = \mathbb{A}_4$  is not nilpotent, as

$$\gamma_1(G) = G, \quad \gamma_j(G) = \{\text{id}, (12)(34), (13)(24), (14)(23)\} \simeq C_2 \times C_2$$

for all  $j \geq 2$ .

xca:gamma

**Exercise 6.6.** Let  $G$  be group. Prove the following statements:

- 1) Each  $\gamma_i(G)$  is a characteristic group of  $G$ .
- 2)  $\gamma_i(G) \supseteq \gamma_{i+1}(G)$  for all  $i \geq 1$ .
- 3) If  $f: G \rightarrow H$  is a surjective group homomorphism, then  $f(\gamma_i(G)) = \gamma_i(H)$  for all  $i \geq 1$ .

theorem:nilpotent

**Theorem 6.7.** Let  $G$  be a nilpotent group.

- 1) If  $H$  is a subgroup of  $G$ , then  $H$  is nilpotent.
- 2) If  $f: G \rightarrow H$  is a surjective group homomorphism, then  $H$  is nilpotent.

*Proof.* For the first statement note that  $\gamma_i(H) \subseteq \gamma_i(G)$  for all  $i \geq 1$ . Let us prove the second claim, if there exists  $c$  such that  $\gamma_{c+1}(G) = \{1\}$ , then

$$\gamma_{c+1}(H) = f(\gamma_{c+1}(G)) = f(\{1\}) = \{1\}. \quad \square$$

**Example 6.8.** The group  $\mathbf{SL}_2(3)$  is not nilpotent, as  $\mathbb{A}_4$  is a quotient of  $\mathbf{SL}_2(3)$ .

There exist a non-nilpotent group  $G$  with a normal subgroup  $K$  such that  $K$  and  $G/K$  are both nilpotent. For example,  $G = \mathbb{S}_3$  and  $K = \mathbb{A}_3$ .

**Exercise 6.9.** Let  $p$  be a prime number. Prove that finite  $p$ -groups are nilpotent.

theorem:gamma

**Theorem 6.10.** Let  $G$  be a group. Then  $[\gamma_i(G), \gamma_j(G)] \subseteq \gamma_{i+j}(G)$  for all  $i, j \geq 1$ .

*Proof.* We proceed by induction on  $j$ . The case where  $j = 1$  is trivial, as by definition one has  $[G, \gamma_j(G)] = \gamma_{j+1}(G)$ . Assume now that the result holds for some  $j \geq 1$  and all  $i \geq 1$ . We first note that

$$[G, \gamma_i(G), \gamma_j(G)] = [\gamma_i(G), G, \gamma_j(G)] = [\gamma_{i+1}(G), \gamma_j(G)] \subseteq \gamma_{i+j+1}(G)$$

by the inductive hypothesis. Moreover, again using the inductive hypothesis,

$$[\gamma_i(G), \gamma_j(G), G] \subseteq [\gamma_{i+j}(G), G] = \gamma_{i+j+1}(G).$$

Lemma 6.3 implies that  $[\gamma_j(G), G, \gamma_i(G)] \subseteq \gamma_{i+j+1}(G)$ . Thus

$$[\gamma_i(G), \gamma_{j+1}(G)] = [\gamma_{j+1}(G), \gamma_i(G)] = [\gamma_j(G), G, \gamma_i(G)] \subseteq \gamma_{i+j+1}(G). \quad \square$$

Podríamos considerar conmutadores arbitrarios donde no necesariamente se asocia siempre hacia la izquierda. Por ejemplo  $[G, G, G] = [[G, G], G]$  y  $[G, [G, G]]$  son ambos conmutadores de peso tres.

**Corollary 6.11.** *Sea  $G$  un grupo. Entonces todo conmutador de peso  $n$  está contenido en  $\gamma_n(G)$ .*

*Proof.* Procederemos por inducción en  $n$ . El caso  $n = 1$  es trivial. Supongamos entonces que el resultado es válido para algún  $n \geq 1$ . Tenemos entonces un conmutador de la forma  $[A, B]$ , donde  $A$  es un conmutador de peso  $k$ ,  $B$  es un conmutador de peso  $l$  y  $n + 1 = k + l$ . Como  $k < n$  y  $l < n$ , la hipótesis inductiva implica que  $A \subseteq \gamma_k(G)$  y  $B \subseteq \gamma_l(G)$ . Luego  $[A, B] \subseteq [\gamma_k(G), \gamma_l(G)] \subseteq \gamma_{k+l}(G)$  por el teorema anterior.  $\square$

En el siguiente lema veremos que los grupos nilpotentes satisfacen la **condición normalizadora**.

lemma:normalizadora

**Lemma 6.12 (condición normalizadora).** *Sea  $G$  un grupo nilpotente. Si  $H$  es un subgrupo propio de  $G$  entonces  $H \subsetneq N_G(H)$ .*

*Proof.* Sabemos que existe  $c$  tal que  $G = \gamma_1(G) \supseteq \cdots \supseteq \gamma_{c+1}(G) = \{1\}$ . Como  $\{1\} = \gamma_{c+1}(G) \subseteq H$  y  $\gamma_1(G) \not\subseteq H$ , sea el mínimo  $k$  tal que  $\gamma_k(G) \subseteq H$ . Como

$$[\gamma_{k-1}(G), H] \subseteq [\gamma_{k-1}(G), G] = \gamma_k(G) \subseteq H,$$

se tiene que  $xHx^{-1} \subseteq H$  para todo  $x \in \gamma_{k-1}(G)$ , es decir  $\gamma_{k-1}(G) \subseteq N_G(H)$ . Si  $N_G(H) = H$  entonces  $\gamma_{k-1}(G) \subseteq H$ , que contradice la minimalidad de  $k$ .  $\square$

Si  $G$  es un grupo se define sucesión  $\zeta_0(G), \zeta_1(G), \dots$  inductivamente de la siguiente forma:

$$\zeta_0(G) = \{1\}, \quad \zeta_{i+1}(G) = \{g \in G : [g, x] \in \zeta_i(G) \text{ para todo } x \in G\}, \quad i \geq 0.$$

Por ejemplo:  $\zeta_1(G) = Z(G)$ .

lemma:central\_ascendente

**Lemma 6.13.** *Sea  $G$  un grupo. Para todo  $i \geq 0$  el conjunto  $\zeta_i(G)$  es un subgrupo normal de  $G$ .*

*Proof.* Procederemos por inducción en  $i$ . El caso  $i = 0$  es trivial pues  $\zeta_0(G) = 1$ . Supongamos entonces que el resultado es válido para  $i$ . Veamos primero que  $\zeta_{i+1}(G)$  es un subgrupo de  $G$ . Sean  $g, h \in \zeta_{i+1}(G)$  y sea  $x \in G$ . Por hipótesis inductiva,

$$\begin{aligned} [g^{-1}, x] &= (xg^{-1})[g, x^{-1}](xg^{-1})^{-1} \in (xg^{-1})\zeta_i(G)(xg^{-1})^{-1} = \zeta_i(G), \\ [gh, x] &= [g, h x h^{-1}][h, x] \in \zeta_i(G). \end{aligned}$$

Como  $1 \in \zeta_{i+1}(G)$ , se concluye que todos los  $\zeta_i(G)$  son subgrupos de  $G$ . La normalidad también se demuestra por inducción en  $i$ : si  $g \in \zeta_{i+1}(G)$ ,  $x \in G$  entonces  $xgx^{-1} \in \zeta_{i+1}(G)$  pues

$$[xgx^{-1}, y] = x[g, x^{-1}yx]x^{-1} \in \zeta_i(G)$$

para todo  $y \in G$ . □

Sea  $G$  un grupo. Se define la **serie central ascendente** de  $G$  como la sucesión

$$1 = \zeta_0(G) \subseteq \zeta_1(G) \subseteq \zeta_2(G) \subseteq \cdots$$

Recordemos que un grupo  $G$  se dice **perfecto** si  $[G, G] = G$ .

theorem:Grun

**Theorem 6.14 (Grün).** Si  $G$  es un grupo perfecto, entonces  $Z(G/Z(G)) = \{1\}$ .

*Proof.* Si usamos el lema de los tres subgrupos con  $X = Y = G$  y  $Z = \zeta_2(G)$ ,  $\{1\} = [\zeta_2(G), G, G] = [\zeta_2(G), [G, G]] = [\zeta_2(G), G]$ . Luego  $\zeta_2(G) \subseteq Z(G)$  y entonces  $\zeta_2(G) = Z(G/Z(G)) = \{1\}$ . □

Si  $H$  y  $K$  son subgrupos de  $G$  se define

$$[H, K] = \langle [h, k] : h \in H, k \in K \rangle.$$

Sea  $G$  un grupo. Se dice que un subgrupo  $K$  de  $G$  **normaliza** a  $H$  si  $K \subseteq N_G(H)$ . Se dice que un subgrupo  $K$  de  $G$  **centraliza** a  $H$  si  $K \subseteq C_G(H)$ , es decir si y sólo si  $[H, K] = \{1\}$ .

**Exercise 6.15.** Sean  $K$  y  $H$  subgrupos de  $G$  con  $K \subseteq H$  y  $K$  normal en  $G$ . Demuestre que  $[H, G] \subseteq K$  si y sólo si  $H/K \subseteq Z(G/K)$ .

lemma:gamma\_zeta

**Lemma 6.16.** Sea  $G$  un grupo. Existe  $c$  tal que  $\zeta_c(G) = G$  si y sólo si  $\gamma_{c+1}(G) = \{1\}$ . Más aún, en ese caso

$$\gamma_{i+1}(G) \subseteq \zeta_{c-i}(G)$$

para todo  $i \in \{0, 1, \dots, c\}$ .

*Proof.* Supongamos primero que  $\zeta_c(G) = G$ . Por inducción vamos a demostrar que  $\gamma_{i+1}(G) \subseteq \zeta_{c-i}(G)$ . Como el caso  $i = 0$  es trivial, supongamos que el resultado es válido para un cierto  $i \geq 0$ . Si  $g \in \gamma_{i+2}(G) = [\gamma_{i+1}(G), G]$ , podemos escribir

$$g = \prod_{k=1}^N [g_k, x_k],$$

donde  $g_1, \dots, g_N \in \gamma_{i+1}(G)$  y  $x_1, \dots, x_N \in G$ . Por hipótesis inductiva



$$g_j \in \gamma_j(G) \subseteq \zeta_{c-j}(G)$$

y entonces  $[g_j, x_j] \in \zeta_{c-i-1}(G)$  para todo  $j$ . Luego  $g \in \zeta_{c-(i+1)}(G)$ . La implicación que queremos queda demostrada al tomar  $i = c$ .

Supongamos ahora que  $\gamma_{c+1}(G) = 1$ . Demostremos por inducción en  $c - i$  que  $\gamma_{c+1-i}(G) \subseteq \zeta_{c-i}(G)$ . El caso  $c - i = 0$  es trivial. Si el resultado es válido para algún  $c - i \geq 0$ , sea  $g \in \gamma_{c+2-i}(G) = [\gamma_{c+1-i}(G), G]$ . Escribimos

$$g = \prod_{k=1}^N [g_k, x_k]$$

con  $g_1, \dots, g_N \in \gamma_{c+1-i}(G) \subseteq \zeta_i(G)$  por hipótesis inductiva. Luego  $g \in \zeta_{c-(i+1)}(G)$  pues cada  $[g_j, x_j] \in \zeta_{i-1}(G)$ . Al tomar  $i = 0$  se obtiene la implicación buscada.  $\square$

**Example 6.17.** Sea  $G = \mathbb{S}_3$ . Entonces  $\zeta_j(G) = \{1\}$  para todo  $j \geq 0$ :

**Definition 6.18.** Sea  $G$  un grupo. Una **serie central** para  $G$  es una sucesión

$$G = G_0 \supseteq G_1 \supseteq \dots \supseteq G_n = \{1\}$$

de subgrupos normales de  $G$  tal que para cada  $i \in \{1, \dots, n\}$ ,  $\pi_i(G_{i-1})$  es un subgrupo de  $Z(G/G_i)$ , donde  $\pi_i: G \rightarrow G/G_i$  es el morfismo canónico.

lemma:serie\_central

**Lemma 6.19.** Sea  $G$  un grupo y sea  $G = G_0 \supseteq G_1 \supseteq \dots \supseteq G_n = \{1\}$  una serie central para  $G$ . Entonces  $\gamma_{i+1}(G) \subseteq G_i$  para todo  $i$ .

*Proof.* Procederemos por inducción en  $i$ . El caso  $i = 0$  es trivial. Supongamos que el resultado es válido para algún  $i \geq 0$ . Entonces

$$\gamma_{i+1}(G) = [G, \gamma_i(G)] \subseteq [G, G_{i-1}] \subseteq G_i$$

pues, como  $\pi_i(G_{i-1}) \subseteq Z(G/G_i)$ , entonces  $\pi([G, G_{i-1}]) = [\pi(G), \pi(G_{i-1})] = \{1\}$  y luego  $[G, G_{i-1}] \subseteq G_i$ .  $\square$

**Theorem 6.20.** Un grupo es nilpotente si y sólo si admite una serie central.

*Proof.* Si el grupo  $G$  es nilpotente, entonces los  $\gamma_j(G)$  forman una serie central para  $G$ . Recíprocamente, si  $G = G_0 \supseteq G_1 \supseteq \dots \supseteq G_n = \{1\}$  es una serie central para  $G$ , entonces, por el lema anterior,  $G$  es nilpotente pues

$$\gamma_{n+1}(G) \subseteq G_n = \{1\}. \quad \square$$

xca:nilpotente\_central

**Exercise 6.21.** Sea  $G$  un grupo. Demuestre que si  $K$  es un subgrupo de  $Z(G)$  tal que  $G/K$  es nilpotente, entonces  $G$  es nilpotente.

theorem:Z(nilpotent)

**Theorem 6.22 (Hirsch).** Sea  $G$  un grupo nilpotente. Si  $H$  es un subgrupo normal no trivial de  $G$  entonces  $H \cap Z(G) \neq \{1\}$ . En particular,  $Z(G) \neq \{1\}$ .

*Proof.* Como  $\zeta_0(G) = \{1\}$  y existe  $c$  tal que  $\zeta_c(G) = G$ , existe

$$m = \min\{k : H \cap \zeta_k(G) \neq \{1\}\}.$$

Como  $H$  es normal,

$$[H \cap \zeta_m(G), G] \subseteq H \cap [\zeta_m(G), G] \subseteq H \cap \zeta_{m-1}(G) = \{1\}.$$

Luego  $\{1\} \neq H \cap \zeta_m(G) \subseteq H \cap Z(G)$ . Si  $H = G$  entonces  $Z(G) \neq \{1\}$ .  $\square$

nilpotente\_minimalnormal

**Exercise 6.23.** Sea  $G$  un grupo nilpotente y sea  $M$  un subgrupo minimal-normal de  $G$ . Demuestre que  $M \subseteq Z(G)$ .

**Corollary 6.24.** Sea  $G$  un grupo nilpotente no abeliano y sea  $A$  un subgrupo maximal-normal y abeliano de  $G$ . Entonces  $A = C_G(A)$ .

*Proof.* Como  $A$  es abeliano,  $A \subseteq C_G(A)$ . Supongamos que  $A \neq C_G(A)$ . El centralizador  $C_G(A)$  es normal en  $G$  pues, como  $A$  es normal en  $G$ ,

$$gC_G(A)g^{-1} = C_G(gAg^{-1}) = C_G(A).$$

para todo  $g \in G$ . Sea  $\pi: G \rightarrow G/A$  el morfismo canónico. Entonces  $\pi(C_G(A))$  es un subgrupo normal no trivial de  $\pi(G)$ . Como  $G$  es nilpotente,  $\pi(G)$  es nilpotente y, por el teorema de Hirsch,  $\pi(C_G(A)) \cap Z(\pi(G)) \neq \{1\}$ . Sea  $x \in C_G(A) \setminus A$  tal que  $\pi(x)$  es central en  $\pi(G)$ . El grupo  $\langle A, x \rangle$  es abeliano pues  $x \in C_G(A)$ . Además  $\langle A, x \rangle$  es normal en  $G$  pues  $A$  es normal en  $G$  y si  $g \in G$  entonces  $gxg^{-1}x^{-1} \in A$  porque  $\pi(x)$  es central y luego  $gxg^{-1} \in \langle A, x \rangle$ . Luego  $A \subsetneq \langle A, x \rangle \subsetneq G$ , una contradicción.  $\square$

**Theorem 6.25.** Sea  $G$  un grupo nilpotente. Valen las siguientes afirmaciones:

- 1) Todo subgrupo minimal-normal tiene orden primo y es central.
- 2) Todo subgrupo maximal es normal de índice primo y contiene a  $[G, G]$ .

*Proof.* Demostremos la primera afirmación. Sea  $N$  un subgrupo minimal-normal. Como  $N \cap Z(G) \neq \{1\}$  por el teorema de Hirsch,  $N \cap Z(G)$  es un subgrupo normal de  $G$  contenido en  $N$ . Luego  $N = N \cap Z(G) \subseteq Z(G)$  por la minimalidad de  $N$ . En particular,  $N$  es abeliano. Además, como todo subgrupo de  $N$  es normal en  $G$ ,  $N$  es simple y luego  $N \simeq C_p$  para algún primo  $p$ .

Demostremos ahora la segunda afirmación. Si  $M$  es un subgrupo maximal,  $M$  es normal en  $G$  gracias a la condición normalizadora. La maximalidad de  $M$  implica que  $G/M$  no contiene subgrupos propios no triviales. Luego  $G/M \simeq C_p$  para algún primo  $p$ . Como en particular  $G/M$  es abeliano,  $[G, G] \subseteq M$ .  $\square$

Es importante remarcar que el teorema anterior no garantiza, por ejemplo, la existencia de subgrupos maximales. Recordemos que  $\mathbb{Q}$  es un grupo abeliano (por lo tanto, nilpotente) que no tiene subgrupos maximales.

proposition:g^n

**Proposition 6.26.** Sea  $G$  un grupo nilpotente y sea  $H$  un subgrupo de  $G$  de índice  $n$ . Si  $g \in G$  entonces  $g^n \in H$ .

*Proof.* El resultado es obvio en el caso en que  $H$  sea un subgrupo normal. Sea  $H_0 = H$  y  $H_{i+1} = N_G(H_i)$  para  $i \geq 0$ . Por definición,  $H_i$  es normal en  $H_{i+1}$  y además, como  $G$  es nilpotente, si  $H_i \neq G$  entonces  $H_i \subsetneq H_{i+1}$  por la condición normalizadora. Como  $(G : H)$  es finito, existe  $k$  tal que  $H_k = G$ . Veamos que

$$g^{(G:H)} = g^{(H_k:H_{k-1})(H_{k-1}:H_{k-2})\cdots(H_1:H_0)} \in H.$$

Observemos que  $g^{(H_k:H_{k-1})} \in H_{k-1}$  pues  $H_{k-1}$  es normal en  $H_k = G$ , y que, como  $g^{(H_k:H_{k-1})} \in H_k$ , entonces

$$g^{(H_k:H_{k-2})} = g^{(H_k:H_{k-1})(H_{k-1}:H_{k-2})} = \left(g^{(H_k:H_{k-1})}\right)^{(H_{k-1}:H_{k-2})} \in H_{k-2}$$

pues  $H_{k-2}$  es normal en  $H_{k-1}$ . Al repetir este argumento,  $g^{(G:H)} \in H$ .  $\square$

**Example 6.27.** La proposición anterior no vale si el grupo  $G$  no es nilpotente. Sea  $G = \mathbb{S}_3$  y sea  $H = \{\text{id}, (12)\}$  de índice tres. Si  $g = (13)$  entonces  $g^3 = (13) \notin H$ .

El lema que daremos a continuación es una herramienta útil para hacer demostraciones por inducción en grupos nilpotentes.

lemma : a [GG]

**Lemma 6.28.** Sea  $G$  un grupo nilpotente de clase  $c \geq 2$ . Si  $x \in G$  entonces el subgrupo  $\langle x, [G, G] \rangle$  es nilpotente de clase  $< c$ .

*Proof.* Sea  $H = \langle x, [G, G] \rangle$ . Si  $x \in [G, G]$ , el resultado es cierto. Supongamos entonces que  $x \notin [G, G]$ . Observemos que

$$H = \{x^n c : n \in \mathbb{Z}, c \in [G, G]\},$$

pues  $[G, G]$  es normal en  $G$ . Para demostrar el lema alcanza entonces con probar que  $[H, H] \subseteq \gamma_3(G)$ . Sean  $h = x^n c, k = x^m d \in H$  con  $c, d \in [G, G]$ . Como

$$[h, x^m] = [x^n, [c, x^m]][c, x^m] \in \gamma_4(G)\gamma_3(G) \subseteq \gamma_3(G),$$

entonces

$$\begin{aligned} [h, k] &= [h, x^m][x^m, [h, d]][h, d] \\ &= [x^n, [c, x^m]][c, x^m][x^m, [h, d]][h, d] \in \gamma_3(G). \end{aligned} \quad \square$$

Veamos un ejemplo, está hecho con la computadora.

**Example 6.29.** Sea  $G = \mathbb{D}_8 = \langle r, s : r^8 = s^2 = 1, srs = r^{-1} \rangle$  el grupo diedral de orden 16. El grupo  $G$  es nilpotente de clase tres y  $[G, G] = \{1, r^2, r^4, r^6\} \simeq C_4$ . El subgrupo  $\langle s, [G, G] \rangle \simeq \mathbb{D}_4$  es nilpotente de clase dos.

Ahora una aplicación del lema.

theorem : T (nilpotent)

**Theorem 6.30.** Si  $G$  es un grupo nilpotente, entonces

$$T(G) = \{g \in G : g^n = 1 \text{ para algún } n \in \mathbb{N}\}$$

es un subgrupo de  $G$ .

*Proof.* Sean  $a, b \in T(G)$  y sean

$$A = \langle a, [G, G] \rangle, \quad B = \langle b, [G, G] \rangle.$$

Como  $A$  y  $B$  son nilpotentes por el lema anterior, por hipótesis inductiva,  $T(A)$  es un subgrupo de  $A$  y  $T(B)$  es un subgrupo de  $B$ . Como  $T(A)$  es característico en  $A$  y  $A$  es normal en  $G$ ,  $T(A)$  es normal en  $G$ . Similarmente se demuestra que  $T(B)$  es normal en  $B$ . Veamos ahora que todo elemento de  $T(A)T(B)$  tiene orden finito: si  $x \in T(A)T(B)$ , digamos  $x = a_1 b_1$  con  $a_1$  de orden  $m$ , entonces  $x$  tiene orden finito pues

$$x^m = (a_1 b_1)^m = (a_1 b_1 a_1^{-1})(a_1^2 b_1 a_1^{-2}) \cdots (a_1^{m-1} b_1 a_1^{-m+1}) b_1 \in T(B).$$

Para entender mejor el truco hagamos un caso concreto, digamos  $m = 3$ . La fórmula en este caso queda así:

$$\begin{aligned} (a_1 b_1)^3 &= (a_1 b_1)(a_1 b_1)(a_1 b_1) \\ &= (a_1 b_1 a_1^{-1})(a_1^2 b_1 a_1^{-2}) a_1^3 b_1 = (a_1 b_1 a_1^{-1})(a_1^2 b_1 a_1^{-2}) b_1, \end{aligned}$$

pues  $a_1^3 = 1$ .

El truco nos permite entonces demostrar que  $ab$  y  $a^{-1}$  tienen ambos orden finito. Luego  $T(G)$  es un subgrupo de  $G$ .  $\square$

Otra aplicación.

theorem:a=b

**Theorem 6.31.** Sea  $G$  un grupo nilpotente y sin torsión y sean  $a, b \in G$ . Si existe  $n \neq 0$  tal que  $a^n = b^n$  entonces  $a = b$ .

*Proof.* Procederemos por inducción en el orden de nilpotencia  $c$  de  $G$ . El resultado es trivialmente cierto si  $G$  es abeliano. Supongamos entonces que  $G$  es nilpotente de índice  $c \geq 1$ . Como  $\langle a, [G, G] \rangle$  es un subgrupo de  $G$  nilpotente de índice  $< c$ , y  $bab^{-1} = [b, a]a \in \langle a, [G, G] \rangle$ , por hipótesis inductiva,  $ba = ab$  pues

$$a^n = (bab^{-1})^n = b^n.$$

Luego  $(ab^{-1})^n = a^n b^{-n} = 1$  y por lo tanto, como  $G$  no tiene torsión, se concluye que  $a = b$ .  $\square$

**Corollary 6.32.** Sea  $G$  un grupo nilpotente sin torsión. Sean  $x, y \in G$  tales que  $x^n y^m = y^m x^n$  para algún  $n, m \neq 0$ , entonces  $xy = yx$ .

*Proof.* Sean  $a = x$  y  $b = y^n x y^{-n}$ . Como  $a^m = b^m$ , el teorema anterior implica que  $a = b$  y luego  $xy^n = y^n x$ . Al usar nuevamente ese teorema, esta vez con  $a = y$  y  $b = xyx^{-1}$ , se concluye que  $xy = yx$ .  $\square$

lemma:fg

**Lemma 6.33.** Sea  $G$  un grupo finitamente generado y sea  $H$  un subgrupo de índice finito. Entonces  $H$  es finitamente generado.

*Proof.* Supongamos que  $G$  está generado por  $\{g_1, \dots, g_m\}$ . Podemos suponer, sin pérdida de generalidad, que para cada  $i$  existe  $k$  tal que  $g_i^{-1} = g_k$ .

Sea  $\{1 = t_1, \dots, t_n\}$  un transversal de  $H$  en  $G$ , es decir un conjunto de representantes de  $G/H$ . Para  $i \in \{1, \dots, n\}$ ,  $j \in \{1, \dots, m\}$ , escribimos

$$t_i g_j = h(i, j) t_{k(i, j)}.$$

Vamos a demostrar que  $H$  está generado por los  $h(i, j)$ . Sea  $x \in H$ . Escribamos

$$\begin{aligned} x &= g_{i_1} \cdots g_{i_s} \\ &= (t_1 g_{i_1}) g_{i_2} \cdots g_{i_s} \\ &= h(1, i_1) t_{k_1} g_{i_2} \cdots g_{i_s} \\ &= h(1, i_1) h(k_1, i_2) t_{k_2} g_{i_3} \cdots g_{i_s} \\ &= h(1, i_1) h(k_1, i_2) \cdots h(k_{s-1}, i_s) t_{k_s}, \end{aligned}$$

donde  $k_1, \dots, k_{s-1} \in \{1, \dots, n\}$ . Como  $t_{k_s} \in H$  pues  $x \in H$ , entonces  $t_{k_s} = 1 \in H$  y luego  $x$  está generado por los  $h(i, j)$ .  $\square$

theorem:T(G) finito

**Theorem 6.34.** Sea  $G$  un grupo finitamente generado, de torsión y nilpotente. Entonces  $G$  es finito.

*Proof.* Procederemos por inducción en la clase de nilpotencia  $c$ . El caso  $c = 1$  es verdadero pues  $G$  es abeliano. Supongamos entonces que el resultado es válido para  $c \geq 1$ . Como  $[G, G]$  y  $G/[G, G]$  son nilpotentes de clase  $< c$ , finitamente generados por el lema anterior y de torsión, por hipótesis inductiva se tiene que  $[G, G]$  y  $G/[G, G]$  son finitos. Luego  $G$  es también finito, de hecho puede probarse que  $G$  es de orden  $|[G, G]|(G : [G, G])$ .  $\square$

El siguiente lema también resultará muy útil, especialmente en caso de trabajar con grupos finitos nilpotentes.

lemma:normalizador

**Lemma 6.35.** Sean  $G$  un grupo finito,  $p$  un primo que divide a  $|G|$  y  $P \in \text{Syl}_p(G)$ . Entonces

$$N_G(N_G(P)) = N_G(P).$$

*Proof.* Sea  $H = N_G(P)$ . Como  $P$  es normal en  $H$ ,  $P$  es el único  $p$ -subgrupo de Sylow de  $H$ . Para ver que  $N_G(H) = H$  basta demostrar que  $N_G(H) \subseteq H$ . Sea  $g \in N_G(H)$ . Como

$$gPg^{-1} \subseteq gHg^{-1} = H,$$

$gPg^{-1} \in \text{Syl}_p(H)$  y  $H$  tiene un único  $p$ -subgrupo de Sylow,  $P = gPg^{-1}$ . Luego  $g \in N_G(P) = H$ .  $\square$

theorem:nilpotente:eq

**Theorem 6.36.** Sea  $G$  un grupo finito. Son equivalentes:

- 1)  $G$  es nilpotente.
- 2) Todo subgrupo de Sylow de  $G$  es normal.
- 3)  $G$  es producto directo de sus subgrupos de Sylow.

*Proof.* Veamos que (1)  $\implies$  (2). Sea  $P \in \text{Syl}_p(G)$ . Queremos ver que  $P$  es normal en  $G$ , es decir  $N_G(P) = G$ . Por el lema anterior,  $N_G(N_G(P)) = N_G(P)$ . La condición normalizadora implica entonces que  $N_G(P) = G$ .

Veamos ahora que (2)  $\implies$  (3). Sean  $p_1, \dots, p_k$  los factores primos de  $|G|$  y para cada  $i \in \{1, \dots, k\}$  sea  $P_i \in \text{Syl}_{p_i}(G)$ . Por hipótesis, cada  $P_j$  es normal en  $G$ .

Vamos a demostrar que  $P_1 \cdots P_j \simeq P_1 \times \cdots \times P_j$  para todo  $j$ . El caso  $j = 1$  es trivial. Supongamos entonces que el resultado vale para algún  $j \geq 1$ . Como

$$N = P_1 \cdots P_j \simeq P_1 \times \cdots \times P_j$$

es normal en  $G$  y tiene orden coprimo con  $|P_{j+1}|$ ,  $N \cap P_{j+1} = \{1\}$ . Luego

$$NP_{j+1} \simeq N \times P_{j+1} \simeq P_1 \times \cdots \times P_j \times P_{j+1}$$

pues  $P_{j+1}$  es también normal en  $G$ .

Ahora que sabemos que  $P_1 \cdots P_k \simeq P_1 \times \cdots \times P_k$  es un subgrupo de orden  $|G|$ , se concluye que  $G = P_1 \times \cdots \times P_k$ .

Para ver que (3)  $\implies$  (1) simplemente hace falta observar que todo  $p$ -grupo es nilpotente (proposición ??) y que el producto directo de finitos nilpotentes es nilpotente.  $\square$

xca:truco

**Exercise 6.37.** Sea  $G$  un grupo finito. Demuestre que si  $P \in \text{Syl}_p(G)$  y  $M$  es un subgrupo de  $G$  tal que  $N_G(P) \subseteq M$  entonces  $M = N_G(M)$ .

xca:normalizadora

**Exercise 6.38.** Sea  $G$  un grupo finito. Son equivalentes:

- 1)  $G$  es nilpotente.
- 2) Si  $H \subsetneq G$  es un subgrupo entonces  $H \subsetneq N_G(H)$ .
- 3) Todo subgrupo maximal de  $G$  es normal en  $G$ .

**Theorem 6.39.** Sea  $G$  un grupo finito nilpotente. Si  $p$  es un primo que divide al orden de  $G$ , existe un subgrupo minimal-normal de orden  $p$  y existe un subgrupo maximal de índice  $p$ .

*Proof.* Supongamos que  $|G| = p^\alpha m$ , donde  $p$  es un primo coprimo con  $m$ . Escribamos  $G = P \times H$ , donde  $P \in \text{Syl}_p(G)$ . Como  $Z(P)$  es un subgrupo normal no trivial de  $P$ , todo subgrupo de  $Z(P)$  minimal-normal en  $G$  tiene orden  $p$  (y esos subgrupos existen porque  $G$  es un grupo finito). Por otro lado, como  $P$  contiene un subgrupo de índice  $p$ , que resulta maximal. Luego  $P \times H$  también contiene un subgrupo maximal de índice  $p$ .  $\square$

xca:pgrupos

**Exercise 6.40.** Sea  $p$  un primo y sea  $G$  un grupo no trivial de orden  $p^n$ . Demuestre las siguientes afirmaciones:

- 1)  $G$  tiene un subgrupo normal de orden  $p$ .

2) Para todo  $j \in \{0, \dots, n\}$  existe un subgrupo normal de  $G$  de orden  $p^j$ .

nilpotente\_equivalencia

**Exercise 6.41.** Sea  $G$  un grupo finito. Demuestre que las siguientes afirmaciones son equivalentes:

- 1)  $G$  es nilpotente.
- 2) Cualesquiera dos elementos de órdenes coprimos conmutan.
- 3) Todo cociente no trivial de  $G$  tiene centro no trivial.
- 4) Si  $d$  divide al orden de  $G$ , existe un subgrupo normal de  $G$  de orden  $d$ .

El siguiente resultado, que puede demostrarse en forma completamente elemental, fue descubierto en 2014.

**Theorem 6.42 (Baumslag–Wiegold).** Sea  $G$  un grupo finito tal que  $|xy| = |x||y|$  si  $x$  e  $y$  son elementos de órdenes coprimos. Entonces  $G$  es nilpotente.

*Proof.* Sean  $p_1, \dots, p_n$  los distintos primos que dividen al orden de  $G$ . Para cada  $i \in \{1, \dots, n\}$  sea  $P_i \in \text{Syl}_{p_i}(G)$ . Primero vamos a demostrar que  $G = P_1 \cdots P_n$ . La inclusión no trivial equivale a demostrar que la función

$$\psi: P_1 \times \cdots \times P_n \rightarrow G, \quad (x_1, \dots, x_n) \mapsto x_1 \cdots x_n$$

es sobreyectiva. Procederemos de la siguiente forma. Primero vemos que la función  $\psi$  es inyectiva. En efecto, si  $\psi(x_1, \dots, x_n) = \psi(y_1, \dots, y_n)$ , entonces

$$x_1 \cdots x_n = y_1 \cdots y_n.$$

Si  $y_n \neq x_n$ , entonces  $x_1 \cdots x_{n-1} = (y_1 \cdots y_{n-1})y_n x_n^{-1}$ . Pero  $x_1 \cdots x_{n-1}$  es un elemento de orden coprimo con  $p_n$  y  $y_1 \cdots y_{n-1}y_n x_n^{-1}$  es un elemento de orden múltiplo de  $p_n$ , una contradicción. Entonces  $x_n = y_n$  y luego, el mismo argumento, prueba que  $\psi$  es inyectiva. Como  $|P_1 \times \cdots \times P_n| = |G|$ , se concluye que  $\psi$  es biyectiva. En particular,  $\psi$  es sobreyectiva y luego  $G = P_1 \cdots P_n$ .

Veamos ahora que cada  $P_j$  es normal en  $G$ . Sea  $j \in \{1, \dots, n\}$  y sea  $x_j \in P_j$ . Sea  $g \in G$  y sea  $y_j = gx_jg^{-1}$ . Como  $y_j \in G$ , podemos escribir  $y_j = z_1 \cdots z_n$  con  $z_k \in P_k$  para todo  $k$ . Como el orden de  $y_j$  es una potencia del primo  $p_j$ , el elemento  $z_1 \cdots z_n$  tiene orden una potencia de  $p_j$  y luego  $z_k = 1$  para todo  $k \neq j$  y además  $y_j = z_j \in P_j$ . Como cada subgrupo de Sylow es normal en  $G$ , se concluye que  $G$  es nilpotente.  $\square$

lemma:commutador

**Lemma 6.43.** Si  $x, y \in G$  son tales que  $[x, y] \in C_G(x) \cap C_G(y)$ , entonces

$$[x, y]^n = [x^n, y] = [x, y^n]$$

para todo  $n \in \mathbb{Z}$ .

*Proof.* Procederemos por inducción en  $n \geq 0$ . El caso  $n = 0$  es trivial. Supongamos entonces que el resultado vale para algún  $n \geq 0$ . Entonces, como  $[x, y] \in C_G(x)$ ,

$$[x, y]^{n+1} = [x, y]^n [x, y] = [x^n, y][x, y] = [x^n, y]xyx^{-1}y^{-1} = x[x^n, y]yx^{-1}y^{-1} = [x^{n+1}, y].$$

Para demostrar el lema en el caso  $n < 0$  basta observar que, como  $[x, y]^{-1} = [x^{-1}, y]$ ,  $[x, y]^{-n} = [x^{-1}, y]^n = [x^{-n}, y]$ .  $\square$

lemma:Hall

**Lemma 6.44 (Hall).** *Sea  $G$  un grupo nilpotente de clase dos y  $x, y \in G$ . Entonces*

$$(xy)^n = [y, x]^{n(n-1)/2} x^n y^n$$

para todo  $n \in \mathbb{N}$ .

*Proof.* Procederemos por inducción en  $n$ . Como el caso  $n = 1$  es trivial, supongamos que el resultado es válido para algún  $n \geq 1$ . Entonces, gracias al lema anterior,

$$\begin{aligned} (xy)^{n+1} &= (xy)^n(xy) = [y, x]^{n(n-1)/2} x^n y^{n-1} (yx)y \\ &= [y, x]^{n(n-1)/2} x^n [y^n, x] xy^{n+1} = [y, x]^{n(n-1)/2} [y, x]^n x^{n+1} y^{n+1}. \end{aligned} \quad \square$$

lemma:class2

**Lemma 6.45.** *Sea  $p > 2$  un número primo y sea  $P$  un  $p$ -grupo de clase de nilpotencia  $\leq 2$ . Si  $[y, x]^p = 1$  para todo  $x, y \in P$  entonces  $P \rightarrow [P, P]$ ,  $x \mapsto x^p$ , es un morfismo de grupos.*

*Proof.* Por lema de Hall,  $(xy)^p = [y, x]^{p(p-1)/2} x^p y^p = x^p y^p$ .  $\square$

thm:class2

**Theorem 6.46.** *Sea  $p > 2$  un número primo y sea  $P$  un  $p$ -grupo de clase de nilpotencia  $\leq 2$ . Entonces  $\{x \in P : x^p = 1\}$  es un subgrupo de  $P$ .*

*Proof.* Como  $P$  tiene clase de nilpotencia dos, los conmutadores son centrales. Para cada  $x \in G$ , la función  $g \mapsto [g, x]$  es un morfismo de grupos pues

$$[gh, x] = ghxh^{-1}g^{-1}x^{-1} = g[h, x]xg^{-1}x^{-1} = [g, x][h, x].$$

En particular, si  $x, y \in P$  con  $x^p = y^p = 1$ , entonces

$$[x, y]^p = [x^p, y] = [1, y] = 1.$$

Luego, al usar el lema de Hall, se concluye que  $(xy)^p = [y, x]^{p(p-1)/2} x^p y^p = 1$ .  $\square$



## Chapter 7

### Solvable groups

#### A

For a group  $G$  we define

$$G^{(0)} = G, \quad G^{(i+1)} = [G^{(i)}, G^{(i)}] \quad i \geq 0.$$

The **derived series** of  $G$  is the sequence

$$G = G^{(0)} \supseteq G^{(1)} \supseteq G^{(2)} \supseteq \dots$$

Each  $G^{(i)}$  is a characteristic subgroup of  $G$ . We say that  $G$  is **solvable** if  $G^{(n)} = \{1\}$  for some  $n \in \mathbb{N}$ . Clearly every abelian group is solvable. A non-abelian simple group cannot be solvable. Nilpotent groups are solvable.

**Exercise 7.1.** The group  $\mathbf{SL}_2(3)$  is solvable.

Let  $p$  be a prime number. An **elementary abelian**  $p$ -group is a group  $P$  such that  $x^p = 1$  for all  $x \in P$ . A subgroup  $M$  of a group  $G$  is said to be **minimal normal** if  $M \neq \{1\}$ ,  $M$  is normal in  $G$  and the unique normal subgroup of  $G$  strictly contained in  $M$  is the trivial subgroup. Every finite group contains a minimal normal subgroup.

**Example 7.2.** If a normal subgroup  $M$  is minimal (with respect to the inclusion), then it is minimal normal. The converse statement is not true. The subgroup of  $\mathbb{A}_4$  generated by  $(12)(34)$ ,  $(13)(24)$  and  $(14)(23)$  is minimal normal in  $\mathbb{A}_4$  but it is not minimal.

**Example 7.3.** Let  $G = \mathbb{D}_6 = \langle r, s : r^6 = s^2 = 1, srs = r^{-1} \rangle$  be the dihedral group of size twelve. The subgroups  $S = \langle r^2 \rangle$  and  $T = \langle r^3 \rangle$  are minimal normal subgroups.

**Exercise 7.4.** Let  $G = \mathbf{SL}_2(3)$ . The unique minimal normal subgroup of  $G$  is  $Z(\mathbf{SL}_2(3)) \simeq C_2$ :

A subgroup  $H$  of a group  $G$  is said to be **characteristic** if  $f(H) \subseteq H$  for all  $f \in \text{Aut}(G)$ . The center  $Z(G)$  and the commutator subgroup  $[G, G]$  of  $G$  are both characteristic subgroups of  $G$ . Every characteristic subgroup of  $G$  is normal in  $G$ . If  $H$  is a characteristic subgroup of  $K$  and  $K$  is normal in  $G$ , then  $H$  is normal in  $G$ .

lemma:minimal\_normal

**Lemma 7.5.** *Let  $M$  be a minimal normal subgroup of  $G$ . If  $M$  is solvable and finite, then  $M$  is an elementary abelian  $p$ -group for some prime number  $p$ .*

*Proof.* Since  $M$  is solvable,  $[M, M] \subsetneq M$ . Moreover,  $[M, M]$  is normal in  $G$ , as  $[M, M]$  is characteristic in  $M$  and  $M$  is normal in  $G$ . Since  $M$  is minimal normal, it follows that  $[M, M] = \{1\}$  and hence  $M$  is abelian. Now if  $M$  is finite, there exists a prime number  $p$  such that  $P = \{x \in M : x^p = 1\}$  is a non-trivial subgroup of  $M$ . Since  $P$  is characteristic in  $M$ , the subgroup  $P$  is normal in  $G$ . Thus  $P = M$ .  $\square$

theorem:resoluble

**Theorem 7.6.** *Let  $G$  be a group.*

- 1) *Each subgroup  $H$  of  $G$  is solvable.*
- 2) *Let  $K$  be a normal subgroup of  $G$ . Then  $G$  is solvable if and only if  $K$  and  $G/K$  are both solvable.*

*Proof.* By induction,  $H^{(i)} \subseteq G^{(i)}$  for all  $i \geq 0$ . Let us prove the second claim. Let  $Q = G/K$  and  $\pi: G \rightarrow Q$  be the canonical map. By induction we prove that  $\pi(G^{(i)}) = Q^{(i)}$  for all  $i \geq 0$ . The case  $i = 0$  is trivial, as  $\pi$  is surjective. Now assume that the result holds for some  $i \geq 0$ . Then

$$\pi(G^{(i+1)}) = \pi([G^{(i)}, G^{(i)}]) = [\pi(G^{(i)}), \pi(G^{(i)})] = [Q^{(i)}, Q^{(i)}] = Q^{(i+1)}.$$

Assume that  $Q$  and  $K$  are both solvable. Since  $Q$  is solvable, there exists  $n$  such that  $Q^{(n)} = \{1\}$ . Since  $\pi(G^{(n)}) = Q^{(n)} = \{1\}$ , it follows that  $G^{(n)} \subseteq K$ . Since  $K$  is solvable, there exists  $m$  such that

$$G^{(n+m)} \subseteq (G^{(n)})^{(m)} \subseteq K^{(m)} = \{1\},$$

and hence  $G$  is solvable.

Let us now assume that  $G$  is solvable. There exists  $n \in \mathbb{N}$  such that  $G^{(n)} = \{1\}$ . Thus  $Q$  is solvable, as  $Q^n = f(G^{(n)}) = f(\{1\}) = \{1\}$ . The group  $K$  is also solvable, as it is a subgroup of  $G$ .  $\square$

**Example 7.7.** Let  $n \geq 5$ . The group  $\mathbb{S}_n$  is not solvable.

**Exercise 7.8.** Let  $p$  be a prime number and  $G$  be a finite  $p$ -group. Prove that  $G$  is solvable.

To prove Wielandt's theorem on solvable groups we need the following lemma.

lemma:4Wielandt

**Lemma 7.9.** *Let  $G$  be a finite group. If  $H$  and  $K$  are subgroups of  $G$  of coprime indices, then  $G = HK$  and  $(H : H \cap K) = (G : K)$ .*

*Proof.* Let  $D = H \cap K$ . Since

$$(G : D) = \frac{|G|}{|H \cap K|} = (G : H)(H : H \cap K),$$

$(G : H)$  divides  $(G : D)$ . Similarly,  $(G : K)$  divides  $(G : D)$ . Since  $(G : H)$  and  $(G : K)$  are coprime,  $(G : H)(G : K)$  divides  $(G : D)$ . In particular,

$$\frac{|G|}{|H|} \frac{|G|}{|K|} = (G : H)(G : K) \leq (G : D) = \frac{|G|}{|H \cap K|}$$

and hence  $|G| = |HK|$ . Since

$$|G| = |HK| = |H||K|/|H \cap K|,$$

it follows that  $(G : K) = (H : H \cap K)$ . □

The **normal closure**  $H^G$  of a subgroup  $H$  of  $G$  is the subgroup

$$H^G = \langle xHx^{-1} : x \in G \rangle$$

generated by all conjugates of  $H$ . The subgroup  $H^G$  is the smallest normal subgroup of  $G$  containing  $H$ .

**Example 7.10.** Let  $G = \mathbb{A}_4$  and  $H = \{\text{id}, (12)(34)\}$ . Then

$$H^G = \{\text{id}, (12)(34), (13)(24), (14)(23)\} \simeq C_2 \times C_2.$$

theorem:Wielandt:solvable

**Theorem 7.11 (Wielandt).** *Let  $G$  be a finite group and  $H$ ,  $K$  and  $L$  be subgroups of  $G$  with pair-wise coprime indices. If  $H$ ,  $K$  and  $L$  are solvable, then  $G$  is solvable.*

*Proof.* Assume the theorem is not valid and let  $G$  be a minimal counterexample. Then  $G$  is not trivial. Let  $N$  be a minimal-normal subgroup of  $G$  and  $\pi: G \rightarrow G/N$ ,  $g \mapsto gN$ , be the canonical map. Since by definition  $N$  is non-trivial, it follows that  $|G/N| < |G|$ . The subgroups  $\pi(H) = \pi(HN)$ ,  $\pi(K) = \pi(KN)$  and  $\pi(L) = \pi(LN)$  of  $\pi(G) = G/N$  are solvable. The correspondence theorem implies that the indices of  $\pi(H)$ ,  $\pi(K)$  and  $\pi(L)$  in  $\pi(G)$  are pair-wise coprime. By the minimality of  $G$ , the group  $\pi(G)$  is solvable. If  $H = \{1\}$ , then  $|G| = (G : H)$  is coprime with  $(G : K)$  and hence  $G = K$  is solvable. So we may assume that  $H \neq \{1\}$ . Let  $M$  be a minimal normal subgroup of  $H$ . By Lemma 7.5,  $M$  is a  $p$ -group for some prime number  $p$ . We may assume that  $p$  does not divide  $(G : K)$  (if  $p$  divides  $(G : K)$ , then  $p$  does not divide  $(G : L)$  and hence it is enough to replace  $K$  by  $L$ ). There exists  $P \in \text{Syl}_p(G)$  such that  $P \subseteq K$ . By Sylow's theorem, there exists  $g \in G$  such that  $M \subseteq gKg^{-1}$ . Since  $(G : gKg^{-1}) = (G : K)$  and  $(G : H)$  are coprime, Lemma 7.9 implies that  $G = (gKg^{-1})H$ .

We claim that all conjugate of  $M$  are included in  $gKg^{-1}$ . If  $x \in G$ , then  $x = uv$  for some  $u \in gKg^{-1}$  and  $v \in H$ . Since  $M$  is normal in  $H$ ,

$$xMx^{-1} = (uv)M(uv)^{-1} = uMu^{-1} \subseteq gKg^{-1}.$$

In particular,  $\{1\} \neq M^G \subseteq gKg^{-1}$  is solvable, as  $gKg^{-1}$  is solvable. The minimality of  $G$  implies that  $G/M^G$  is solvable. Hence  $G$  is solvable by Theorem 7.6.  $\square$

Let  $G$  be a finite group of order  $p^\alpha m$  with  $p$  a prime number coprime with  $m$ . A subgroup  $H$  of  $G$  is a  **$p$ -complement** if  $|H| = m$ .

**Example 7.12.** Sea  $G = \mathbb{S}_3$ . Then  $H = \langle (123) \rangle$  is a 2-complement and  $K = \langle (12) \rangle$  is a 3-complement.

A famous theorem of Burnside states that finite groups whose order are divisible by exactly two primes are solvable.

**Theorem 7.13 (Burnside).** *Let  $p$  and  $q$  be prime numbers and let  $G$  be a group of order  $p^\alpha q^\beta$ . Then  $G$  is solvable.*

There is a quite easy proof that uses basic character theory, see for example. A non-character-theoretic proof is known but it is harder, see [36].

theorem:Hall:solvable

**Theorem 7.14 (Hall).** *Let  $G$  be a finite group that admits a  $p$ -complement for all primes  $p$  dividing the order of  $G$ . Then  $G$  is solvable.*

*Proof.* Let  $|G| = p_1^{\alpha_1} \cdots p_k^{\alpha_k}$  with the  $p_j$  being distinct primes. We proceed by induction on  $k$ . If  $k = 1$ , then  $G$  is a  $p$ -group and the result is clear. If  $k = 2$ , then Burnside's theorem implies the claim. Assume now that  $k \geq 3$ . For each  $j \in \{1, 2, 3\}$  let  $H_j$  be  $p_j$ -complement in  $G$ . Since  $|H_j| = |G|/p_j^{\alpha_j}$ , the subgroups  $H_j$  have coprime indices.

We claim that  $H_1$  is solvable. Note that  $|H_1| = p_2^{\alpha_2} \cdots p_k^{\alpha_k}$ . Let  $p$  be a prime number that divides  $|H_1|$  and let  $Q$  be a  $p$ -complement in  $G$ . Since  $(G : H_1)$  and  $(G : Q)$  are coprime, Lemma 7.9 implies that

$$(H_1 : H_1 \cap Q) = (G : Q).$$

Thus  $H_1 \cap Q$  is a  $p$ -complement in  $H_1$ . Hence  $H_1$  is solvable by the inductive hypothesis. Similarly,  $H_2$  and  $H_3$  are both solvable.

Since  $H_1, H_2$  and  $H_3$  are solvable and have coprime indices, Wielandt's theorem implies the claim.  $\square$

## B

We now use Hall's theorem to obtain information related to the structure of finite braces.

thm:add\_nilpotent

**Theorem 7.15.** *Let  $A$  be a finite brace of nilpotent type. Then the multiplicative group of  $A$  is solvable.*

*Proof.* Let  $K$  be the additive group of  $A$  and  $G$  be the multiplicative group of  $A$ . Assume that  $|A| = p_1^{\alpha_1} \cdots p_n^{\alpha_n}$  for different primes numbers  $p_1, \dots, p_n$ . Since  $K$  is nilpotent, each  $K_i \in \text{Syl}_{p_i}(K)$  is normal in  $K$ , so each  $K_i$  is a left ideal of  $A$ . It follows that for each  $i \in \{1, \dots, n\}$  both  $K_i$  and  $\prod_{j \neq i} K_j$  are braces of coprime order. In particular, for each  $i \in \{1, \dots, n\}$  there exists a subgroup of  $G$  of order coprime with  $p_i$ . Then  $G$  is solvable by Hall's theorem.  $\square$

## Exercises

prob:G(X,r) solvable

**7.1.** Let  $(X, r)$  be a finite involutive solution. Prove that  $G(X, r)$  is solvable.

## Notes

Solvable groups...

In [29], Etingof, Schedler and Soloviev proved that the structure group of a finite involutive solution is always solvable.



## Chapter 8

### Factorizations

factorizations

#### A

In Chapter 4 we found that groups with an exact factorization produce braces. In this chapter we will study a different relationship between factorizations and braces.

A group  $G$  is said to be **factorized through subgroups**  $A$  and  $B$  if  $G = AB$ . We remark that we do not assume that  $A \cap B = \{1\}$ .

**Definition 8.1.** A group  $G$  is **metabelian** if  $[G, G]$  is abelian.

A group  $G$  is metabelian if and only if there is a normal subgroup  $K$  of  $G$  such that  $K$  and  $G/K$  are abelian. The groups  $\mathbb{S}_3$  and  $\mathbb{A}_4$  are metabelian.

**Exercise 8.2.** Let  $G$  be a metabelian group.

- 1) If  $H$  is a subgroup of  $G$ , then  $H$  is metabelian.
- 2) If  $f: G \rightarrow H$  is a group homomorphism, then  $f(H)$  is metabelian.

**Exercise 8.3.** Prove that  $\mathbf{SL}_2(3)$  is metabelian.

A straightforward calculation shows that the following formulas hold:

$$\begin{aligned} [a, bc] &= [a, b]b[a, c]b^{-1}, \\ [ab, c] &= a[b, c]a^{-1}[a, c]. \end{aligned}$$

The following theorem is considered the most satisfying result about group factorization. The proof is based on a surprisingly short and smart calculation with commutators.

theorem:Ito

**Theorem 8.4 (Itô).** *Let  $G = AB$  be a factorization of  $G$  through the abelian subgroups  $A$  and  $B$ . Then  $G$  is metabelian.*

*Proof.* Since  $G = AB$ , it follows that  $AB = BA$ . Let us prove that  $[A, B]$  is a normal subgroup of  $G$ . Let  $a, a_1, \alpha, \alpha_1 \in A$  and  $b, b_1, \beta, \beta_1 \in B$  be such that  $\alpha b \alpha^{-1} = b_1 a_1$ ,  $\beta a \beta^{-1} = a_2 b_2$ . Since

$$\begin{aligned}\alpha[a, b]\alpha^{-1} &= a(\alpha b \alpha^{-1})a^{-1}(\alpha b^{-1}\alpha^{-1}) = ab_1a_1a^{-1}a_1^{-1}b_1^{-1} = [a, b_1] \in [A, B] \\ \beta[a, b]\beta^{-1} &= (\beta a \beta^{-1})\beta b \beta^{-1}(\beta a^{-1}\beta^{-1})b^{-1} = a_2b_2bb_2^{-1}a_2^{-1}b^{-1} = [a_2, b] \in [A, B],\end{aligned}$$

it follows that  $[A, B]$  is a normal subgroup of  $G$ .

Now we prove that  $[A, B]$  is abelian. Since

$$\begin{aligned}\beta\alpha[a, b]\alpha^{-1}\beta^{-1} &= \beta[a, b_1]\beta^{-1} = (\beta a \beta^{-1})b_1(\beta a^{-1}\beta^{-1})b_1^{-1} = [a_2, b_1], \\ \alpha\beta[a, b]\beta^{-1}\alpha^{-1} &= \alpha[a_2, b]\alpha^{-1} = a_2(\alpha b \alpha^{-1})a_2^{-1}(\alpha b^{-1}\alpha^{-1}) = [a_2, b_1],\end{aligned}$$

a direct calculation shows that

$$[\alpha^{-1}, \beta^{-1}][a, b][\alpha^{-1}, \beta^{-1}]^{-1} = [a, b].$$

Since two arbitrary generators of  $[A, B]$  commute, the group  $[A, B]$  is abelian.

Finally we note that  $[G, G] = [A, B]$ . Since  $[A, B]$  is normal in  $G$ ,

$$[a_1b_1, a_2b_2] = a_1[a_2, b_1]^{-1}a_1^{-1}a_2[a_1, b_2]a_2^{-1} \subseteq [A, B]. \quad \square$$

Sysak found the following generalization of Itô's theorem:

**Theorem 8.5 (Sysak).** *If  $A$  and  $B$  are abelian subgroups of  $G$  and  $H$  is a subgroup of  $G$  contained in the set  $AB$ , then  $H$  is metabelian.*

The proof appears in [62].

There are several other interesting results in the theory of factorizable groups. Another important result that is worth mentioning is the following theorem.

**Theorem 8.6 (Kegel–Wielandt).** *Let  $G$  be a finite group. If there are nilpotent subgroups  $A$  and  $B$  of  $G$  such that  $G = AB$ , then  $G$  is solvable.*

The proof appears for example in [3, Theorem 2.4.3].

The theorem of Kegel–Wielandt turns out to be the main tool in the proof of the following result on the structure of finite braces. This proves a conjecture of Byott, see [14].

thm:mul\_nilpotent

**Theorem 8.7.** *Let  $A$  be a finite brace with nilpotent multiplicative group. Then the additive group of  $A$  is solvable.*

*Proof.* Let  $K$  be the additive group of  $A$  and  $G$  be the multiplicative group of  $A$ . The group  $\Gamma = K \rtimes G$  has multiplication

$$(g, \lambda_g)(h, \lambda_h) = (g + \lambda_g(h), \lambda_g \lambda_h) = (g \circ h, \lambda_{g \circ h}).$$

Let  $f: G \rightarrow \Gamma$ ,  $g \mapsto (g, \lambda_g)$ . Then  $f$  is a group homomorphism and  $f(G)$  is nilpotent. Since  $\lambda(G)$  is nilpotent, the finite group  $K \rtimes \lambda(G) = f(G)\lambda(G)$  is a product of nilpotent groups. By the theorem of Kegel–Wielandt,  $K \rtimes \lambda(G)$  is solvable. Hence  $K$  is solvable.  $\square$



## B

It turns out to be interesting to study factorization of braces.

**Definition 8.8.** Let  $A$  be a brace and let  $B$  and  $C$  be left ideals of  $A$ . We say that  $A$  admits a *factorization* through  $B$  and  $C$  if  $A = B + C$ .

Note that if a brace  $A$  admits a factorization through  $B$  and  $C$ , then it follows that

$$A = B + C = C + B = B \circ C = C \circ B.$$

Now we prove an analog of Itô's theorem in the context of braces. It turns out that one needs to consider factorizations through strong left ideals. We also need the following definition:

**Definition 8.9.** A brace  $A$  is said to be *meta-trivial* if  $A^{(2)}$  is a trivial brace.

Clearly a brace  $A$  is meta-trivial if and only there is an ideal  $I$  of  $A$  such that  $I$  and  $A/I$  are trivial as braces.

lem:calcbraces

**Lemma 8.10.** Let  $A$  be a brace. For any  $x, y, z \in A$  the following statements hold:

- 1)  $x * (y + z) = x * y + y + x * z - y$ ,
- 2)  $(x \circ y) * z = x * (y * z) + y * z + x * z$ .

lem:calculations

**Lemma 8.11.** Let  $A$  be a brace such that  $A = B + C$ , where  $B$  and  $C$  are left ideals. If  $B$  and  $C$  are trivial as braces then, for any  $b, \beta \in B$  and  $c, \gamma \in C$ , the following statements hold:

- 1)  $\lambda_{\beta \circ \gamma} = \lambda_{\gamma \circ \beta}$ ,
- 2)  $(c + b) \circ \beta - \beta = c + b + c * \beta$ ,
- 3)  $b \circ c \circ b' \circ c' = b \circ c - c \circ b = b + \lambda_b(c) - \lambda_c(b) - c \in \ker \lambda$ .

*Proof.* To prove (1) put  $c_1 = \lambda_\beta(c) \in C$  and  $b_1 = \lambda_\gamma(b) \in B$ . As  $B$  and  $C$  are trivial braces,  $\lambda_\beta(b + c) = \lambda_\beta(b) + \lambda_\beta(c) = b + c_1$  and similarly  $\lambda_\gamma(b + c) = b_1 + c$ . Then  $\lambda_{\beta \circ \gamma}(b + c) = b_1 + c_1 = \lambda_{\gamma \circ \beta}(b + c)$ .

Let us prove (2). As  $B$  is a trivial brace, it follows from (4.2) that

$$\begin{aligned} (c + b) \circ \beta - \beta &= (c \circ \lambda_{c'}(b)) \circ \beta - \beta \\ &= c \circ (\lambda_{c'}(b) \circ \beta) - \beta \\ &= c \circ (\lambda_{c'}(b) + \beta) - \beta \\ &= c \circ \lambda_{c'}(b) - c + c \circ \beta - \beta \\ &= c + b + c * \beta. \end{aligned}$$

Part (3) follows from the following computation

$$\begin{aligned}
b \circ c \circ b' \circ c' &= (b \circ c) + \lambda_{b \circ c}(b' + \lambda_{b'}(c')) \\
&= b + \lambda_b(c) + \lambda_{b \circ c}(b') + \lambda_{b \circ c \circ b'}(c') \\
&= b + \lambda_b(c) + \lambda_c(b') + \lambda_{b \circ b'}(c') \\
&= b + \lambda_b(c) + \lambda_c(-b) - c \\
&= b \circ c - c \circ b.
\end{aligned}$$

Moreover, by (1) it follows that  $b \circ c \circ b' \circ c' \in \ker \lambda$ .  $\square$

**Lemma 8.12.** *Let  $A$  be a brace such that  $A = B + C$  is a factorization through left ideals  $B$  and  $C$ . If  $B$  and  $C$  are trivial braces, then:*

- 1)  $B * C$  and  $C * B$  are strong left ideals of  $A$ ,
- 2)  $B * C$  and  $C * B$  are trivial braces, and
- 3)  $A^{(2)} = C * B + B * C = B * C + C * B$ .

*Proof.* Since  $C$  is a left ideal, it follows that  $B * C \subseteq C$ . Let  $b, \beta \in B$  and  $c, \gamma \in C$ . As  $C$  is trivial, it follows that

$$\begin{aligned}
\lambda_{b \circ c}(\beta * \gamma) &= \lambda_b(\beta * \gamma) \\
&= \lambda_b \lambda_\beta(\gamma) - \lambda_b(\gamma) \\
&= \lambda_{b \circ \beta \circ b'} \lambda_b(\gamma) - \lambda_b(\gamma) \\
&= (b \circ \beta \circ b') * \lambda_b(\gamma) \in B * C.
\end{aligned}$$

Hence  $B * C$  is a left ideal and trivial as a brace.

Let  $a \in A$ ,  $b \in B$  and  $c \in C$ . Write  $a = b_1 + c_1$ , with  $b_1 \in B$  and  $c_1 \in C$ . Then

$$\begin{aligned}
a + (b * c) - a &= a + \lambda_b(c) - c - a \\
&= -(b * a) + b * (a + c) \\
&= -(b * (b_1 + c_1)) + b * (b_1 + c_1 + c).
\end{aligned} \tag{8.1}$$

As  $B + C = C + B$ , it follows that for any  $\beta \in B$  and  $\gamma \in C$ , there exist  $\beta_1 \in B$  and  $\gamma_1 \in C$  such that  $\beta + \gamma = \gamma_1 + \beta_1$ . Hence, for any  $b \in B$  it holds that

$$b * (\beta + \gamma) = b * (\gamma_1 + \beta_1) = b * \gamma_1 + \gamma_1 + b * \beta_1 - \gamma_1 = b * \gamma_1,$$

as  $B$  is trivial. Applying this on (8.1) it follows that  $B * C$  is a normal subgroup of  $(A, +)$ . This proves (1) and (2) for  $B * C$ . The proof for  $C * B$  is similar.

Now we show that  $A^{(2)} \subseteq C * B + B * C$ . Let  $b, b_1 \in B$  and  $c, c_1 \in C$ . Then

$$\begin{aligned}
(b \circ c) * (b_1 + c_1) &= (b \circ c) * b_1 + b_1 + (b \circ c) * c_1 - b_1 \\
&= \lambda_{b \circ c}(b_1) - b_1 + b_1 + b * (c * c_1) + c * c_1 + b * c_1 - b_1 \\
&= \lambda_c(b_1) - b_1 + b_1 + b * c_1 - b_1 \\
&= c * b_1 + b_1 + b * c_1 - b_1 \in C * B + B * C.
\end{aligned}$$

Clearly  $C * B + B * C \subseteq A^{(2)}$  and thus  $A^{(2)} = C * B + B * C = B * C + C * B$ .  $\square$

thm:Ito\_braces

**Theorem 8.13.** *Let  $A$  be a brace. If  $A = B + C$  is a factorization through strong left ideals  $B$  and  $C$  that are trivial as braces, then  $A$  is right nilpotent of class at most three. In particular,  $A$  is meta-trivial.*

*Proof.* By Lemma 8.12,  $B * C$  and  $C * B$  are strong left ideals of  $A$ , and both are trivial as braces. Furthermore,

$$A^{(2)} = B * C + C * B = (B * C) \circ (C * B).$$

It rests to show that  $A^{(2)}$  acts trivially on  $A$ . We first show that  $B * C$  acts trivially on  $A$ . For that purpose, let  $b \in B$ ,  $c \in C$  and  $a \in A$ . Write  $a = \beta + \gamma$ , where  $\beta \in B$  and  $\gamma \in C$ . Then

$$(b * c) * (\beta + \gamma) = (b * c) * \beta + \beta + (b * c) * \gamma - \beta = (b * c) * \beta,$$

as  $C$  is a trivial brace. By Lemma 8.11(3),

$$(b \circ c - c \circ b) + \beta = (b \circ c - c \circ b) \circ \beta = (b + \lambda_b(c) - \lambda_c(b) - c) \circ \beta.$$

Since  $(B, +)$  is a normal subgroup of  $(A, +)$ ,

$$b \circ c - c \circ b = b + \lambda_b(c) - \lambda_c(b) - c = \lambda_b(c) - c + b_1$$

for some  $b_1 \in B$ . By Lemma 8.11(2),

$$\begin{aligned} (b \circ c - c \circ b) + \beta &= (\lambda_b(c) - c + b_1) \circ \beta \\ &= \lambda_b(c) - c + b_1 + (b * c) * \beta + \beta \end{aligned}$$

and therefore  $(b * c) * \beta = 0$ . Thus  $B * C$  acts trivially on  $A$ . As  $(C, +)$  also is a normal subgroup of  $(A, +)$ , it follows by symmetry that  $C * B$  acts trivially on  $A$ . Hence  $A^{(2)}$  acts trivially on  $A$ .  $\square$

**Corollary 8.14.** *Let  $A$  be a brace. Assume that  $A = B + C$ , where  $B$  and  $C$  are (not necessarily strong) left ideals, which are trivial as braces. Then  $A$  has a meta-trivial ideal  $I$  such that  $A/I$  is a trivial brace.*

*Proof.* By Lemma 8.12, the ideal  $A^{(2)}$  has a factorization through the strong left ideals  $B * C$  and  $C * B$ , which are trivial braces. By Theorem 8.13,  $A^{(2)}$  is meta-trivial and hence the claim follows.  $\square$

Theorem 8.13 has application to involutive solutions.

thm:MP

**Theorem 8.15.** *Let  $(X, r)$  be an involutive non-degenerate (not necessarily finite) solution of the Yang–Baxter equation with  $|X| \geq 2$ . If the brace of abelian type  $\mathcal{G}(X, r)$  admits a factorization through left ideals, which are trivial as left braces, then  $(X, r)$  is a multipermutation solution of level at most three.*

*Proof.* Let  $A = \mathcal{G}(X, r)$  and  $G = G(X, r)$ . Then Theorem 8.13 yields  $A^{(m)} = 0$  for some  $m \leq 3$ . Because  $G/\text{Soc}(G) \cong A$  as left braces, we get  $G^{(m)} \subseteq \text{Soc}(G)$ , and

thus  $G^{(m+1)} = 0$ . Hence  $G$  is a right nilpotent left brace of class at most four and, by [?, Proposition 6],  $(G, r_G)$  is a multipermutation solution of level at most three. Therefore, by [32, Theorem 5.15],  $(X, r)$  is a multipermutation solution of level at most three.  $\square$

This shows that properties of the involutive non-degenerate set-theoretic solution  $(X, r)$  are not completely determined by the group theory of the additive and multiplicative groups of the left brace  $\mathcal{G}(X, r)$ .

exa:B(8,27)

**Example 8.16.** Let  $X = \{1, 2, 3, 4\}$  and  $r(x, y) = (\sigma_x(y), \tau_y(x))$  be the irretractable involutive non-degenerate solution given by

$$\begin{aligned} \sigma_1 &= (34), & \sigma_2 &= (1324), & \sigma_3 &= (1423), & \sigma_4 &= (12), \\ \tau_1 &= (24), & \tau_2 &= (1432), & \tau_3 &= (1234), & \tau_4 &= (13). \end{aligned}$$

The associated left brace  $\mathcal{G}(X, r)$  has additive group  $C_2^3$  and multiplicative group  $D_8$ . Furthermore,  $\mathcal{G}(X, r)$  is not right nilpotent. Hence it is impossible to decompose the left brace  $\mathcal{G}(X, r)$  as in Theorem 8.15.

**Example 8.17.** The left brace  $B(8, 26)$  has the same additive and multiplicative groups as the brace  $\mathcal{G}(X, r)$  of Example 8.16 but it has a factorization as in Theorem 8.15. This shows that  $B(8, 26)$  is right nilpotent.

## Exercises

prob:decomposable

**8.1.** Let  $A$  be a brace. If there exists a proper strong left ideal  $I$ , then  $(A, r_A)$  is decomposable as  $A = I \cup A \setminus I$ .

prob:Ito\_relaxed

**8.2.** Prove that the assumptions of Theorem 8.13 cannot be relaxed.

prob:Ito\_version2

**8.3.** Let  $A$  be a non-zero brace that has a factorization  $A = B + C$  through left ideals  $B$  and  $C$ , where both are trivial as braces. If  $B$  is a strong left ideal of  $A$ , then  $B$  or  $C$  contains a non-zero ideal  $I$  of  $A$  that acts trivially on  $A$ .

prob:mul\_abelian

**8.4.** Let  $A$  be a brace with abelian multiplicative group. Prove that the additive group of  $A$  is meta-abelian.

prob:mul\_cyclic

**8.5.** Let  $A$  be a finite brace with cyclic multiplicative group. Prove that the additive group of  $A$  is supersolvable.

## Open problems

problem:Byott

**Problem 8.1.** Let  $A$  be a brace with solvable additive group. Is the multiplicative group of  $A$  solvable?

**Notes**

Theorem 8.13 was proved by Jespers, Kubat, Antwerpen and Vendramin in [39]. Exercises 8.2 and 8.3 also appear in there. One cannot expect a naive result similar to that of Kegel–Wielandt in the context of braces.

Theorem 8.7 was proved by Tsang and Qin in [63]. Exercises 8.4 and 8.5 also appear in [63].

Problem 8.1 was formulated by Byott in [14].



## Chapter 9

### The structure brace of a solution

structure\_brace

To prove that the structure group  $G(X, r)$  of a solution  $(X, r)$  is a brace, we follow the proof of Lu, Yan and Zhu. They use the language of braided groups.

**Definition 9.1.** A *braided group* is a pair  $(G, r)$ , where  $G$  is a group with operation  $m: G \times G \rightarrow G$ ,  $m(x, y) = xy$ , and  $r: G \times G \rightarrow G \times G$  is a bijective map such that

- 1)  $r(xy, z) = (\text{id} \times m)r_1r_2(x, y, z)$  for all  $x, y, z \in G$ ,
- 2)  $r(x, yz) = (m \times \text{id})r_2r_1(x, y, z)$  for all  $x, y, z \in G$ ,
- 3)  $r(1, x) = (x, 1)$  and  $r(x, 1) = (1, x)$  for all  $x \in G$ , and
- 4)  $m \circ r = m$ .

The map  $r$  is called a *braiding operator* on  $G$ .

*Proof.*

□

**Theorem 9.2.** Let  $G$  be a group. Then  $G$  admits a braiding operator if and only if there is a brace structure on the set  $G$  with multiplicative group  $G$ .

*Proof.*

□

### Exercises

**9.1.** Prove that a braiding operator is a solution.





## **Chapter 10**

### **Bieberbach groups**

**A**

**B**



## Chapter 11

### Garside groups

In this chapter prove that the structure group of a finite involutive solution is a Garside group.

A *monoid* is a non-empty set  $M$  provided with an associative binary operation  $M \times M \rightarrow M$ ,  $(x, y) \mapsto xy$ , and an identity element. A monoid  $M$  is said to be *cancellative* if

$$xy = xz \implies y = z \quad \text{and} \quad xy = zy \implies x = z$$

for all  $x, y, z \in M$ .

**Definition 11.1.** A *Garside monoid* is a pair  $(M, \Delta)$ , where  $M$  is a cancellative monoid such that

- 1) There exists a map  $d: M \rightarrow \mathbb{N}$  such that  $d(xy) \geq d(x) + d(y)$  and  $d(x) \neq 0$  if  $x \neq 1$ .
- 2)
- 3)  $\Delta$  is a Garside element of  $M$ ...
- 4) The family of all divisors of  $\Delta$  is finite.

**Definition 11.2.** A group  $G$  is said to be a *Garside group* if...

Structure groups of involutive solutions are Garside groups.

thm:Chouraqui

**Theorem 11.3.** Let  $(X, r)$  be an involutive solution. Then  $G(X, r)$  is a Garside group.

*Proof.*

□

At this point it is easy to prove the following important result of Gateva–Ivanova and Van den Bergh.

thm:torsion\_free

**Theorem 11.4.** Let  $(X, r)$  be an involutive solution. Then  $G(X, r)$  has no torsion. In particular,  $G(X, r)$  is a Bieberbach group.

*Proof.*

□

As a consequence we obtain the following result on linear representations of the structure group of an involutive solution.

thm:ESS

**Theorem 11.5.**

*Proof.*

□

thm:D

**Theorem 11.6.**

*Proof.*

□

## Exercises

## Open problems

## Notes

Theorem 11.3 was proved by Chouraqui in [22]. Our proof is based on the work of Dehornoy [25] and the presentation of Cedó's survey [17].

Theorem 11.4 was proved by Gateva–Ivanova and Van den Bergh in [33] using somewhat different methods.

## Chapter 12

### Invariant subgroups

#### A

We say that a group  $G$  acts on a group  $K$  by automorphism if the (left) action

$$G \times K \rightarrow K, \quad (g, x) \mapsto g \cdot x,$$

satisfies  $g \cdot (xy) = (g \cdot x)(g \cdot y)$  for all  $g \in G$  and  $x, y \in K$ . The group

$$C_K(G) = \{x \in K : g \cdot x = x \text{ for all } g \in G\}$$

acts on the set of  $G$ -orbits by left multiplication. Indeed, if  $x \in K$  and  $c \in C_K(G)$ , then  $g \cdot c = c$  for all  $g \in G$ . Thus

$$\begin{aligned} c(G \cdot x) &= \{c(g \cdot x) : g \in G\} \\ &= \{(g \cdot c)(g \cdot x) : g \in G\} = \{g \cdot (cx) : g \in G\} = G \cdot (cx). \end{aligned}$$

The following theorem goes back to Deaconescu and Walls [24]. Our proof is that of Isaacs, see [37].

thm:DeaconescuWalls

**Theorem 12.1 (Deaconescu–Walls).** *Let the group  $G$  acts by automorphism on a finite group  $K$ . Let  $C = C_K(G)$  and  $N = C \cap [G, K]$ , where  $[G, K]$  is the subgroup of  $K$  generated by  $[g, x] = (g \cdot x)x^{-1}$  for all  $g \in G$  and  $x \in K$ . Then the index  $(C : N)$  divides the number of  $G$ -orbits of  $K$ .*

*Proof.* The group  $C$  acts by left multiplication on the set  $\Omega$  of  $G$ -orbits on  $K$ . Let  $X = G \cdot x \in \Omega$  be an orbit and  $C_X$  be the stabilizer of  $X$  in  $C$  de  $X$ . If  $c \in C_X$ , then  $cX = X$ . In particular, if  $c \in C_X$ , then  $cx = g \cdot x$  for some  $g \in G$ . Thus

$$c = (g \cdot x)x^{-1} = [g, x] \in [G, K]$$

and hence  $C_X \subseteq N$ .

To prove that  $(C : N)$  divides the size of  $\Omega$ , decompose  $\Omega$  as a disjoint union of  $C$ -orbits. Then it is enough to show that  $(C : N)$  divides the size of each  $C$ -orbit. If

$X \in \Omega$ , then  $C \cdot X$  has size

$$(C : C_X) = (C : N)(N : C_X).$$

Thus  $(C : N)$  divides the size of  $C \cdot X$ .  $\square$

cor:  $Z(K)$  subset  $[K, K]$

**Corollary 12.2.** *Let  $K$  be a non-trivial finite group with  $k$  conjugacy classes. If  $|Z(K)|$  and  $k$  are coprime, then  $Z(K) \subseteq [K, K]$ .*

*Proof.* Let the group  $K$  acts on  $K$  by conjugation, which is an action by automorphism. Deaconescu–Walls’ theorem implies that  $(Z(K) : Z(K) \cap [K, K])$  divides  $k$ . Since  $k$  and  $|Z(K)|$  are coprime, it follows that  $Z(K) = Z(K) \cap [K, K] \subseteq [K, K]$ .  $\square$

Let  $K$  be a group and  $f \in \text{Aut}(K)$ . Then  $f$  is **central** if  $f(x)x^{-1} \in Z(K)$  for all  $x \in K$ . Note that  $f \in \text{Aut}(K)$  is central if and only if  $f \in C_{\text{Aut}(K)}(\text{Inn}(K))$ .

**Corollary 12.3.** *Let  $K$  be a finite group with  $k$  conjugacy classes and  $c$  central automorphisms. If  $\gcd(|K|, kc) = 1$ , then  $[K, K] = Z(K)$ .*

*Proof.* By Corollary 12.2,  $Z(K) \subseteq [K, K]$ .

Let us prove that  $Z(K) \supseteq [K, K]$ . Let  $G = C_{\text{Aut}(K)}(\text{Inn}(K))$ . Since  $\gcd(|K|, kc) = 1$  and  $(C_K(G) : C_K(G) \cap [G, K])$  divides  $c$ , Deaconescu–Walls’ theorem, it follows that  $C_K(G) = C_K(G) \cap [G, K]$ . Since  $[K, K] \subseteq C_K(G)$ , as

$$a \cdot [x, y] = [(a \cdot x)x^{-1}x, (a \cdot y)y^{-1}y] = [x, y]$$

for all  $a \in G$ ,  $x, y \in K$  and  $[G, K] \subseteq Z(K)$ , we conclude that

$$[K, K] \subseteq C_K(G) = C_K(G) \cap [G, K] \subseteq [G, K] \subseteq Z(K). \quad \square$$

**Corollary 12.4.** *Let  $p$  be a prime number. If  $K$  is a group with  $p$  conjugacy classes, then  $Z(K) \subseteq [K, K]$  or  $|K| = p$ .*

*Proof.* Let  $K$  acts on  $K$  by conjugation. Since every element of  $C = Z(K)$  form a conjugacy class,  $|C| \leq p$ . If  $|C| = p$ , then  $K = C = Z(K)$  has  $p$  elements. Otherwise,  $\gcd(|C|, p) = 1$  and hence  $C \subseteq N = [K, K]$ .  $\square$

## B

## C

**Definition 12.5.** Let  $A$  be a brace. One defines  $A^1 = A$  and for  $n \geq 1$

$$A^{n+1} = A * A^n = \langle a * x : a \in A, x \in A^n \rangle_+.$$

The sequence  $A^1 \supseteq A^2 \supseteq A^3 \supseteq \cdots \supseteq A^n \supseteq \cdots$  is the *left series* of  $A$ .

pro:left\_series

**Proposition 12.6.** *Let  $A$  be a brace. Each  $A^n$  is a left ideal of  $A$ .*

*Proof.* We proceed by induction on  $n$ . The case  $n = 1$  is trivial, so we may assume that the result is true for some  $n \geq 1$ . Let  $a, b \in A$  and  $x \in A^n$ . By the inductive hypothesis,  $\lambda_a(x) \in A^n$  and hence

$$\lambda_a(b * x) = (a \circ b \circ a') * \lambda_a(x) \in A^{n+1},$$

where the equality follows by (??). This implies that  $\lambda_a(A^{n+1}) \subseteq A^{n+1}$ . Thus the result follows.  $\square$

**Definition 12.7.** A brace  $A$  is said to be *left nilpotent* if  $A^m = \{0\}$  for some  $m \geq 1$ .

Some basic properties of left nilpotent braces appear in Exercises 12.1–12.3.

pro:IcapFix

**Proposition 12.8.** *Let  $A$  be a left nilpotent brace and  $I$  be a non-zero left ideal of  $A$ . Then  $I \cap \text{Fix}(A) \neq \{0\}$ . In particular,  $\text{Fix}(A) \neq \{0\}$ .*

*Proof.* Let  $m = \max\{k : I \cap A^k \neq \{0\}\}$ . Since  $A * (I \cap A^m) \subseteq I \cap A^{m+1} = \{0\}$ , it follows that there exists a non-zero  $x \in I \cap A^m$  such that  $a * x = 0$  for all  $a \in A$ . Thus  $0 \neq x \in \text{Fix}(A) \cap I$ . For the second claim, apply the first case with  $I = A$ .  $\square$

Let  $A$  be a brace. Let  $A^{[1]} = A$  and for  $n \geq 1$  let  $A^{[n+1]}$  be the additive subgroup of  $A$  generated by elements from  $\{A^{[i]} * A^{[n+1-i]} : 1 \leq i \leq n\}$ . One easily proves by induction that  $A^{[k]} \supseteq A^{[k+1]}$  for all  $k \geq 1$ .

pro:Smoktunowicz

**Proposition 12.9.** *Let  $A$  be a brace. Each  $A^{[n]}$  is a left ideal of  $A$ .*

*Proof.* Each  $A^{[n]}$  is a subgroup of  $(A, +)$ . Since  $A * A^{[n]} \subseteq A^{[n+1]} \subseteq A^{[n]}$ , the claim follows from Proposition 4.40.  $\square$

There exists a brace  $A$  such that  $A^{[n]} = A^{[n+1]} \neq \{0\}$  for some positive integer  $n$  and  $A^{[n+2]} = \{0\}$ .

exa:funny

**Example 12.10.** Let

$$G = \langle r, s : r^8 = s^2 = 1, srs = r^7 \rangle \simeq \mathbb{D}_{16},$$

$$K = \langle a, b : 8a = 2b = 0, a + b = b + a \rangle \simeq \mathbb{Z}/(8) \times \mathbb{Z}/(2).$$

The group  $G$  acts by automorphisms on  $K$  via

$$r \cdot a = a + b, \quad r \cdot b = 4a + b, \quad s \cdot a = 3a, \quad s \cdot b = 4a + b.$$

A direct calculation shows that the map  $\pi : G \rightarrow K$  given by

$$\begin{array}{llll} 1 \mapsto 0, & r \mapsto a, & r^2 \mapsto 2a + b, & r^3 \mapsto 7a + b, \\ r^4 \mapsto 4a, & r^5 \mapsto 5a, & r^6 \mapsto 6a + b, & r^7 \mapsto 3a + b, \\ rs \mapsto 6a, & r^2s \mapsto 7a, & r^3s \mapsto b, & r^4s \mapsto 5a + b, \\ r^5s \mapsto 2a, & r^6s \mapsto 3a, & r^7s \mapsto 4a + b, & s \mapsto a + b, \end{array}$$

is a bijective 1-cocycle. Therefore there exists a brace  $A$  with additive group isomorphic to  $K$  and multiplicative group isomorphic to  $G$ . The addition of  $A$  is that of  $K$  and the multiplication is given by

$$x \circ y = \pi(\pi^{-1}(x)\pi^{-1}(y)), \quad x, y \in K.$$

Since

$$\begin{aligned} a * a &= -a + a \circ a - a = -a + (2a + b) - a = b, \\ (5a + b) * a &= -(5a + b) + (5a + b) \circ a - a = -(5a + b) + b - a = 2a, \end{aligned}$$

it follows that  $A^{[2]}$  contains  $\langle 2a, b \rangle_+ = \{0, 2a, 4a, 6a, b, 2a + b, 4a + b, 6a + b\}$ , the additive subgroup of  $(A, +)$  generated by  $2a$  and  $b$ . Therefore  $A^{[2]} = \langle 2a, b \rangle_+$  since  $A^{[2]} \neq A$ . Routine calculations prove that

$$A^{[3]} = \{0, 2a + b, 4a, 6a + b\}, \quad A^{[4]} = A^{[5]} = \{0, 4a\}, \quad A^{[6]} = \{0\}.$$

**Definition 12.11.** For a brace  $A$  let  $\ell_1(a) = a$  and  $\ell_{k+1}(a) = a * \ell_k(a)$  for  $n \geq 1$ . The brace  $A$  is said to be *left nil* if there exists a positive integer  $n$  such that  $\ell_n(a) = 0$  for all  $a \in A$ .

**Definition 12.12.** For a brace  $A$  let  $\rho_1(a) = a$  and  $\rho_{k+1}(a) = \rho_k(a) * a$  for  $n \geq 1$ . The brace  $A$  is said to be *right nil* if there exists a positive integer  $n$  such that  $\rho_n(a) = 0$  for all  $a \in A$ .

**Definition 12.13.** A brace  $A$  is said to be *strongly nilpotent* if there is a positive integer  $n$  such that  $A^{[n]} = 0$ .

**Definition 12.14.** A brace  $A$  is said to be *strongly nil* if for every  $a \in A$  there is a positive integer  $n = n(a)$  such that any  $*$ -product of  $n$  copies of  $a$  is zero.

We first prove that if both groups of a finite brace  $A$  are nilpotent, then  $A$  can be decomposed as a direct product of braces of prime-power size.

sum

**Lemma 12.15.** Let  $A$  be a brace such that the additive group is a direct sum of ideals  $I_1, I_2$ , that is  $A = I_1 + I_2$  and  $I_1 \cap I_2 = \{0\}$ . Then the map  $f : A \rightarrow I_1 \times I_2$  defined by  $f(a_1 + a_2) = (a_1, a_2)$ , for all  $a_1 \in I_1$  and  $a_2 \in I_2$ , is an isomorphism of braces.

*Proof.* The operations of the brace  $I_1 \times I_2$  are defined component-wise. Clearly  $f$  is an isomorphism of the additive groups of  $A$  and  $I_1 \times I_2$ . Let  $a_1 \in I_1$  and  $a_2 \in I_2$ . Since  $I_1$  and  $I_2$  are ideals we have that

$$a_1 + a_2 - a_1 - a_2, a_1 * a_2, a_2 * a_1 \in I_1 \cap I_2 = \{0\},$$

thus  $a_1 + a_2 = a_2 + a_1$  and  $a_1 \circ a_2 = a_1 + a_2 = a_2 \circ a_1$ . Hence

$$\begin{aligned} f((a_1 + a_2) \circ (b_1 + b_2)) &= f(a_1 \circ a_2 \circ b_1 \circ b_2) = f(a_1 \circ b_1 \circ a_2 \circ b_2) \\ &= f(a_1 \circ b_1 + a_2 \circ b_2) = (a_1 \circ b_1, a_2 \circ b_2) \\ &= (a_1, a_2) \circ (b_1, b_2) = f(a_1 + a_2) \circ f(b_1 + b_2), \end{aligned}$$



for all  $a_1, b_1 \in I_1$  and  $a_2, b_2 \in I_2$ .  $\square$

thm:direct

**Theorem 12.16.** *Let  $n$  be a positive integer. Let  $A$  be a brace such that the additive group is a direct sum of ideals  $I_1, \dots, I_n$ , that is every element  $a \in A$  is uniquely written as  $a = a_1 + \dots + a_n$ , with  $a_j \in I_j$  for all  $j$ . Then the map*

$$f : A \rightarrow I_1 \times \dots \times I_n, \quad f(a_1 + \dots + a_n) = (a_1, \dots, a_n),$$

for all  $a_j \in I_j$ , is an isomorphism of braces.

*Proof.* We shall prove the result by induction on  $n$ . For  $n = 1$ , it is clear. Suppose that  $n > 1$  and that the result is true for  $n - 1$ . Let  $A_1 = I_1 + \dots + I_{n-1}$ . Then  $A_1$  is an ideal of  $A$  and  $A$  is the direct sum of the ideals  $A_1$  and  $I_n$ . By Lemma 12.15, the map  $f_1 : A \rightarrow A_1 \times I_n$  defined by  $f(a + a_n) = (a, a_n)$ , for all  $a \in A_1$  and  $a_n \in I_n$ , is an isomorphism of braces. By the induction hypothesis, the map

$$f_2 : A_1 \rightarrow I_1 \times \dots \times I_{n-1}, \quad f_2(a_1 + \dots + a_{n-1}) = (a_1, \dots, a_{n-1}),$$

is an isomorphism of braces. Therefore  $f = (f_2 \times \text{id}) \circ f_1 : A \rightarrow I_1 \times \dots \times I_n$  is an isomorphism of braces and  $f(a_1 + \dots + a_n) = (a_1, \dots, a_n)$ , for all  $a_j \in I_j$ . The result then follows.  $\square$

cor:product

**Corollary 12.17.** *Let  $A$  be a finite brace such that  $(A, +)$  and  $(A, \circ)$  are nilpotent. Let  $I_1, \dots, I_n$  be the distinct Sylow subgroups of the additive group of  $A$ . Then  $I_1, \dots, I_n$  are ideals of  $A$  and the map*

$$f : A \rightarrow I_1 \times \dots \times I_n, \quad f(a_1 + \dots + a_n) = (a_1, \dots, a_n),$$

for all  $a_j \in I_j$ , is an isomorphism of braces.

*Proof.* Since  $(A, +)$  is nilpotent, for every prime divisor  $p$  of the order of  $A$ , there is a unique Sylow  $p$ -subgroup  $I$  of  $(A, +)$ . Hence  $I$  is a normal subgroup of  $(A, +)$ , and  $\lambda_a(b) \in I$  for all  $a \in A$  and  $b \in I$ . Thus  $I$  is a left ideal of  $A$  and thus it is a Sylow  $p$ -subgroup of  $(A, \circ)$ . Since  $(A, \circ)$  is nilpotent,  $I$  is the unique Sylow  $p$ -subgroup of  $(A, \circ)$  and, thus, it is normal in  $(A, \circ)$ . Therefore  $I$  is an ideal of  $A$ . Hence  $I_1, \dots, I_n$  are ideals of  $A$  and clearly the additive group of  $A$  is the direct sum of  $I_1, \dots, I_n$ . The result follows by Theorem 12.16.  $\square$

Let  $A$  be a brace. Let  $G$  be the multiplicative group of  $A$  and  $K$  be the additive group of  $A$ . The group  $G$  acts on  $K$  by automorphisms. Let  $\Gamma$  be the semidirect product  $\Gamma = K \rtimes G$ . The operation of  $\Gamma$  is

$$(x, g)(y, h) = (x + \lambda_g(y), g \circ h).$$

Identifying each  $g \in G$  with  $(0, g) \in \Gamma$  and each  $x \in K$  with  $(x, 0) \in \Gamma$ ,

$$\begin{aligned} [g, x] &= gxg^{-1}x^{-1} = (0, g)(x, 0)(0, g')(-x, 0) \\ &= (\lambda_g(x), g)(-\lambda_g^{-1}(x), g') = (\lambda_g(x) - x, 0) = \lambda_g(x) - x = g * x. \end{aligned}$$

Let  $K_0 = K = A^1$  and  $K_{n+1} = [G, K_n] = A^{n+2}$  for  $n \geq 0$ . The elements of the left series of  $A$  are iterated commutators of the group  $\Gamma$ .

prop:pgroups

**Proposition 12.18.** *Let  $p$  be a prime and  $A$  be brace of size  $p^m$ . Then  $A$  is left nilpotent.*

*Proof.* Let  $G$  be the multiplicative group of  $A$  and  $K$  be the additive group of  $A$ . Since the semidirect product  $\Gamma = K \rtimes G$  is a  $p$ -group, it is nilpotent. Thus there exists  $k$  such that the  $k$ -repeated commutator  $[\Gamma, \Gamma, \dots, \Gamma]$ , where  $\Gamma$  appears  $k$ -times, is trivial. Since  $A^k = [G, \dots, G, K] \subseteq [\Gamma, \dots, \Gamma]$ , it follows that  $A$  is left nilpotent.  $\square$

HERE The following results follow immediately from theorems of P. Hall:

**Lemma 12.19.** *Let  $A$  be finite brace such that  $A^3 = 0$ . Then the additive group of  $A^2$  is abelian. In fact,  $A^2$  is a trivial brace.*

*Proof.* The first part follows by [?, Theorem 6]. Note that  $(A^2)^2 \subseteq A^3 = 0$ , hence  $a \circ b = a + b$  for all  $a, b \in A^2$ , and the result follows.  $\square$

thm:A2

**Theorem 12.20.** *Let  $A$  be left nilpotent skew left brace. Then the following statements hold:*

- 1) *The additive group of  $A^2$  is locally nilpotent.*
- 2) *The multiplicative group of  $A/\ker \lambda$  is locally nilpotent.*

*Proof.* Since each element of the left series of  $A$  is a repeated commutator, the first claim follows from Hall's theorem [?, Theorem 4]. To prove the second claim, we use the notation above Proposition 12.18. Let  $K = [G, X]G \subseteq \Gamma$  and  $H = [G, X]X$ . Let  $C$  be the centralizer of  $H$  in  $K$ . Then by [?, Theorem 4],  $K/C$  is locally nilpotent. Note that, since  $X$  is normal in  $\Gamma$ ,  $H = X$ . Hence  $G \cap C$  is the centralizer of  $X$  in  $G$ , that is

$$\begin{aligned} G \cap C &= \{g \in G \mid gxg^{-1} = x, \text{ for all } x \in X\} \\ &= \{g \in A \mid \lambda_g(x) = x, \text{ for all } x \in A\} = \ker \lambda. \end{aligned}$$

Thus  $(GC)/C \cong G/(G \cap C) = G/\ker \lambda$  is locally nilpotent.  $\square$

The assumption on the nilpotency of the additive group in Theorem ?? is needed (see Example ??).

**Corollary 12.21.** *Let  $A$  be a finite brace of size  $p^n$  for some prime number  $p$  and some positive integer  $n$ . Then either  $A$  is the trivial brace of order  $p$  or it is not simple.*

*Proof.* By Theorem ??,  $A$  is left nilpotent. In particular, if  $A \neq 0$ , then  $A^2 \neq A$ . Since  $A^2$  is an ideal either  $A$  is not simple or  $A^2 = 0$ . Assume that  $A^2 = 0$ . In this case,  $a \circ b = a + b$  for all  $a, b \in A$ . Therefore  $[A, A]$  is a proper ideal of  $A$ . Hence, either  $A$  is not simple or  $[A, A] = 0$ . Assume that  $A^2 = [A, A] = 0$ . In this case  $A$  is a trivial brace and the result follows.  $\square$

lem:sylow\_leftideals

**Lemma 12.22.** *Let  $A$  be a finite skew left brace with nilpotent additive group. Let  $p$  and  $q$  distinct prime numbers and let  $P$  and  $Q$  be Sylow subgroups of  $(A, +)$  of sizes  $p^n$  and  $q^m$ , respectively. Then  $P$ ,  $Q$  and  $P + Q$  are left ideals of  $A$ .*

*Proof.* Let us first prove that  $P$  is a left ideal. Since  $(A, +)$  is nilpotent,  $P$  is a normal subgroup of  $(A, +)$ . Let  $a \in A$  and  $x \in P$ . Then  $\lambda_a(x) \in P$  since  $\lambda_a$  is a group homomorphism. Similarly one proves that  $Q$  is a left ideal. From this it follows that  $P + Q$  is a left ideal.  $\square$

The following is based on [58, Theorem 5(1)]. However, the proof is completely different.

thm:P\*Q=0

**Theorem 12.23.** *Let  $A$  be a finite skew left brace with nilpotent additive group. Let  $p$  and  $q$  distinct prime numbers and let  $A_p$  and  $A_q$  be Sylow subgroups of  $(A, +)$  of sizes  $p^n$  and  $q^m$ , respectively. If  $p$  does not divide  $q^t - 1$  for all  $t \in \{1, \dots, m\}$ , then  $A_p * A_q = 0$ . In particular,  $\lambda_x(y) = y$  for all  $x \in A_p$  and  $y \in A_q$ .*

*Proof.* By Lemma 12.22  $A_p$ ,  $A_q$  and  $A_p + A_q$  are left ideals of  $A$ . In particular,  $A_p + A_q$  is a skew subbrace of  $A$  and  $A_p$  and  $A_q$  are Sylow subgroups of  $(A_p + A_q, \circ)$ . By Sylow's theorem, the number  $n_p$  of Sylow  $p$ -subgroups of the multiplicative group of  $A_p + A_q$  is

$$n_p = [A_p + A_q : N] \equiv 1 \pmod{p},$$

where  $N = \{g \in A_p + A_q : g \circ A_p \circ g' = A_p\}$  is the normalizer of  $A_p$  in the multiplicative group of  $A_p + A_q$ . Since  $[A_p + A_q : N] = q^s$  for some  $s \in \{0, \dots, m\}$  and  $p$  does not divide  $q^t - 1$  for all  $t \in \{1, \dots, m\}$ , it follows that  $s = 0$  and hence  $A_p$  is a normal subgroup of the multiplicative group of  $A_p + A_q$ . Thus  $A_p$  is an ideal of the skew left brace  $A_p + A_q$ . Since  $A_p$  is an ideal of  $A_p + A_q$  and  $A_q$  is a left ideal, we have that  $A_p * A_q \subseteq A_p \cap A_q = 0$ , and the result follows.  $\square$

**Corollary 12.24.** *Let  $A$  be a skew left brace of size  $p_1^{\alpha_1} \cdots p_k^{\alpha_k}$ , where  $p_1 < p_2 < \cdots < p_k$  are prime numbers and  $\alpha_1, \dots, \alpha_k$  are positive integers. Assume that the additive group of  $A$  is nilpotent. Let  $A_j$  be the Sylow  $p_j$ -subgroups of the additive group of  $A$ . Assume that, for some  $j \leq k$ ,  $p_j$  does not divide  $p_i^{t_i} - 1$  for all  $t_i \in \{1, \dots, \alpha_i\}$  for all  $i \neq j$ . Then  $\text{Soc}(A_j) \subseteq \text{Soc}(A)$ .*

*Proof.* Write  $A = A_1 + \cdots + A_k$ . Let  $a \in \text{Soc}(A_j)$  and  $b \in A$ . Hence there exist elements  $b_k \in A_k$  such that  $b = b_1 + \cdots + b_k$ . By Theorem 12.23,  $\lambda_a(b_i) = b_i$ , for all  $i \neq j$ . Then  $\lambda_a(b) = \lambda_a(b_1) + \cdots + \lambda_a(b_k) = b_1 + \cdots + b_k = b$  and hence  $a \in \text{Soc}(A)$ . Thus the result follows.  $\square$

Let  $A$  be a skew left brace. For subsets  $X$  and  $Y$  of  $A$  we define inductively  $L_0(X, Y) = Y$  and  $L_{n+1}(X, Y) = X * L_n(X, Y)$  for  $n \geq 0$ .

**Definition 12.25.** Let  $p$  be a prime number. A finite skew left brace  $A$  of nilpotent type is said to be *left  $p$ -nilpotent* if there exists  $n \geq 1$  such that  $L_n(A, A_p) = 0$ , where  $A_p$  is the Sylow  $p$ -subgroup of  $(A, +)$ .

lem:factorization

**Lemma 12.26.** *Let  $A$  be a skew left brace such that its additive group is the direct product of the left ideals  $B$  and  $C$ . Then  $A * (B + C) = A * B + A * C$ . Moreover, if  $A = \oplus_{i=1}^n B_i$  where the  $B_i$  are left ideals, then*

$$A * \sum_{i=1}^n B_i = \sum_{i=1}^n A * B_i.$$

*Proof.* Let  $a \in A$ ,  $b \in B$  and  $c \in C$ . Then

$$a * (b + c) = a * b + b + a * c - b = a * b + a * c$$

holds for all  $a \in A$ ,  $b \in B$  and  $c \in C$ . The second part follows by induction.  $\square$

pro:left\_p

**Proposition 12.27.** *Let  $A$  be a finite skew left brace of nilpotent type. Then  $A$  is left nilpotent if and only if  $A$  is left  $p$ -nilpotent for all  $p \in \pi(A)$ .*

*Proof.* For each  $p \in \pi(A)$  there exists  $n(p) \in \mathbb{N}$  such that  $L_{n(p)}(A, A_p) = \{0\}$ . Let  $n = \max\{n(p) : p \in \pi(A)\}$ . Then  $L_n(A, A_p) = \{0\}$  for all  $p \in \pi(A)$ . Since  $A$  is of nilpotent type, the additive group  $(A, +)$  is isomorphic to the direct sum of the  $A_p$  for  $p \in \pi(A)$ . Then Lemma 12.26 implies that

$$L_n(A, A) = \sum_{p \in \pi(A)} L_n(A, A_p) = \{0\}.$$

The other implication is trivial.  $\square$

We now recall some notation about commutators. Given a brace  $A$ , the group  $(A, \circ)$  acts on  $(A, +)$  by automorphisms. If in the semidirect product  $(A, +) \rtimes (A, \circ)$  we identify  $a$  with  $(0, a)$  and  $b$  with  $(b, 0)$ , then

$$\begin{aligned} [a, b] &= (0, a)(b, 0)(0, a)^{-1}(b, 0)^{-1} = (0, a)(b, 0)(0, a')(-b, 0) \\ &= (\lambda_a(b), a)(-\lambda_{a'}(b), a') = (\lambda_a(b) - b, 0) \\ &= (a * b, 0) \end{aligned}$$

Under this identification, we write  $[X, Y] = X * Y$  for any pair of subsets  $X, Y \subseteq A$ . Then the iterated commutator satisfies

$$[X, \dots, X, Y] = [X, [X, \dots, [X, Y] \dots]] = L_n(X, Y),$$

where the subset  $X$  appears  $n$  times.

thm:left\_p

**Theorem 12.28.** *Let  $A$  be a finite brace of nilpotent type. The following statements are equivalent:*

- 1)  $A$  is left  $p$ -nilpotent.
- 2)  $A_{p'} * A_p = \{0\}$ .
- 3) The group  $(A, \circ)$  is  $p$ -nilpotent.

*Proof.* We first prove that (1) implies (2). Since  $A$  is left  $p$ -nilpotent, there exists  $n \in \mathbb{N}$  such that  $L_n(A_{p'}, A_p) \subseteq L_n(A, A_p) = \{0\}$ . Since  $(A_{p'}, \circ)$  acts by automorphisms on  $(A_p, +)$  and this is a coprime action, it follows from [36, Lemma 4.29] that

$$L_1(A_{p'}, A_p) = A_{p'} * A_p = A_{p'} * (A_{p'} * A_p) = L_2(A_{p'}, A_p).$$

By induction one then proves that  $A_{p'} * A_p = L_n(A_{p'}, A_p) = \{0\}$ .

We now prove that (2) implies (3). It is enough to prove that  $(A_{p'}, \circ)$  is a normal subgroup of  $(A, \circ)$ . By using Lemma 12.26,

$$A_{p'} * A = A_{p'} * (A_p + A_{p'}) = (A_{p'} * A_p) + (A_{p'} * A_{p'}) \subseteq A_{p'}.$$

since  $A_{p'}$  is a left ideal of  $A$  and  $A_{p'} * A_p = \{0\}$ . Then  $A_{p'}$  is an ideal of  $A$  by Lemma ?? and [?, Lemma 1.9]. In particular,  $(A_{p'}, \circ)$  is a normal subgroup of  $(A, \circ)$ .

Finally we prove that (3) implies (1). We need to prove that  $L_n(A_p, A_p) = 0$  for some  $n$ . Since  $(A, \circ)$  is  $p$ -nilpotent, there exists a normal  $p$ -complement that is a characteristic subgroup of  $(A, \circ)$ . This group is  $A_{p'}$  and hence  $A_{p'}$  is an ideal of  $A$ . Then  $A_{p'} * A_p \subseteq A_{p'} \cap A_p = 0$ . We now prove that  $L_n(A, A_p) = L_n(A_p, A_p)$  for all  $n \geq 0$ . The case where  $n = 0$  is trivial, so assume that the result holds for some  $n \geq 0$ . By the inductive hypothesis,

$$L_{n+1}(A, A_p) = A * L_n(A, A_p) = A * L_n(A_p, A_p).$$

Thus it is enough to prove that  $A * L_n(A_p, A_p) \subseteq A_p * L_n(A_p, A_p)$ . Let  $a \in A$  and  $b \in L_n(A_p, A_p)$ . Write  $a = x \circ y$  for  $x \in A_p$  and  $y \in A_{p'}$ . Then

$$a * b = (x \circ y) * b = x * (y * b) + y * b + x * b = x * b \in A_p * L_n(A_p, A_p)$$

since  $A_{p'} * A_p = 0$ . The skew left brace  $A_p$  is left nilpotent by [?, Proposition 4.4], so there exists  $n \in \mathbb{N}$  such that  $L_n(A_p, A_p) = 0$ .  $\square$

## Exercises

prob:LN\_direct

**12.1.** Let  $A_1, \dots, A_k$  be left nilpotent braces. Prove that  $A_1 \times \dots \times A_k$  is left nilpotent.

prob:LN\_surj

**12.2.** Let  $f: A \rightarrow B$  be a surjective homomorphism of braces. Prove that if  $A$  is left nilpotent, then  $B$  is left nilpotent.

prob:LN\_sub

**12.3.** Let  $A$  be a left nilpotent brace and  $B \subseteq A$  be a sub brace. Prove that  $B$  is left nilpotent.

prob:nil=>leftnilpotent

**12.4.** Prove that nil braces of abelian/nilpotent type? left nilpotent.

## Open problems

tion:rightnil=>rightnilp

**Problem 12.1.** Let  $A$  be a finite right nil brace. Is  $A$  right nilpotent?

stronglynil=>stronglynilp

**Problem 12.2.** Let  $A$  be a finite strongly nil brace. Is  $A$  strongly nilpotent?

## Notes

The left series of a brace was defined by Rump [53] in the context of braces of abelian type. Precisely in that paper he proved Proposition 12.18 by a different method in the case of braces of abelian type.

Strongly nilpotent braces of abelian type were defined by Smoktunowicz in [59]. These definitions extend to skew left braces, see [21].

Theorem 12.16 was proved by Byott in the context of Hopf–Galois extensions [13].

Theorem ?? was proved by Smoktunowicz in [59] for braces of abelian type and it was extended to nilpotent type in [21].

Theorem ?? was proved By Smoktunowicz in [59, Theorem 1.1] for braces of abelian type. The generalization to braces of nilpotent type appeared in [21, Theorem 4.8]. The proof presented in this chapter appeared in [1] and it is heavily based on the ideas of Ballester–Bolinches, Meng and Romero [46].

Theorem 12.28 was proved by Ballester–Bolinches, Meng and Romero for braces of abelian type.

Exercise 12.4 was proved in the case of braces of abelian type by Smoktunowicz [58].

## Chapter 13

### Multipermutation solutions

#### A

If  $X$  is a solution, we consider over  $X$  the relation

$$x \sim y \iff \sigma_x = \sigma_y \text{ and } \tau_x = \tau_y.$$

Then  $\sim$  is an equivalence relation. Let  $\bar{X}$  be the set of equivalence classes and  $[x]$  denote the equivalence class of  $x$ .

**Proposition 13.1.** *Let  $(X, r)$  be a solution. Then  $(\bar{X}, \bar{r})$ , where*

$$\bar{r}([x], [y]) = ([\sigma_x(y)], [\tau_y(x)]),$$

*is a solution.*

*Proof.* We first prove that  $\bar{r}$  is well-defined. Let  $x, y \in X$  be such that  $x \sim y$  and let  $z \in X$ . Since  $(X, r)$  is a solution, Lemma 1.3 implies that

$$\sigma_{\sigma_x(z)} \circ \sigma_{\sigma_z(x)} = \sigma_x \circ \sigma_z = \sigma_y \circ \sigma_z = \sigma_{\sigma_y(z)} \circ \sigma_{\sigma_z(y)},$$

it follows that  $\sigma_{\sigma_z(x)} = \sigma_{\sigma_z(y)}$  and hence  $\sigma_z(x) \sim \sigma_z(y)$ . Similarly  $\tau_{\tau_z(x)} = \tau_{\tau_z(y)}$  and therefore  $\bar{r}$  is well-defined.

We now prove that  $\bar{r}$  is invertible.

...

□

In the case of involutive solutions, it follows from Proposition 2.57 that  $\sigma_x = \sigma_y$  if and only if  $\tau_x = \tau_y$ .

**Definition 13.2.** Let  $(X, r)$  be a solution. The solution  $\text{Ret}(X, r) = (\bar{X}, \bar{r})$  induced by the equivalence relation  $\sim$  is the *retraction* of  $(X, r)$ .

We define inductively  $\text{Ret}^0(X, r) = (X, r)$ ,  $\text{Ret}^1(X, r) = \text{Ret}(X, r)$  and

$$\text{Ret}^{n+1}(X, r) = \text{Ret}(\text{Ret}^n(X, r)) \quad n \geq 1.$$

**Definition 13.3.** A solution  $(X, r)$  is said to be of *multipermutation level*  $n$  if  $n$  is the smallest non-negative integer such that  $|\text{Ret}^n(X, r)| = 1$ . The solution  $(X, r)$  is said to be *irretractable* if  $\text{Ret}(X, r) = (X, r)$ .

The trivial solution over the set with one element is a multipermutation of level zero. Permutation solutions are multipermutation solutions of level one.

**Example 13.4.**

**Example 13.5.**

**Table 13.1:** Involutive solutions of size  $\leq 10$ .

$n$	2	3	4	5	6	7	8	9	10
solutions	2	5	23	88	595	3456	34530	321931	4895272
multipermutation	2	5	21	84	554	3295	32155	305916	4606440
irretractable	0	0	2	4	9	13	191	685	3590

tab:INV\_mp

For size  $\leq 7$  the numbers of Table 13.1 coincide with those in [29] but there are some differences for solutions of size eight.

**Table 13.2:** Non-involutive solutions of size  $\leq 8$ .

$n$	2	3	4	5	6	7	8
solutions	2	21	230	3519	100071	4602720	422449480
multipermutation	15	206	3165	95517	4461805	416725250	
irretractable	6	24	98	514	2659	17370	

tab:mp

thm:CJKAV

**Theorem 13.6.** Let  $(X, r)$  be a finite multipermutation solution. If  $|X| > 1$ , then  $r$  has even order.

*Proof.* Since  $(X, r) \rightarrow \text{Ret}(X, r), x \mapsto [x]$  is a homomorphism of solutions, it follows that the order of the solution  $\bar{r}$  divides the order of  $r$ . Assume that  $(X, r)$  has multipermutation level  $n$ . There exists a homomorphism of solutions  $(X, r) \rightarrow \text{Ret}^{n-1}(X, r)$ , thus it is enough to prove the theorem in the case where  $r(x, y) = (\sigma(y), \tau(x))$  for commuting permutations  $\sigma$  and  $\tau$ , i.e. multipermutation solutions of level one. If  $r$  has order  $2k + 1$ , then

$$(x, y) = r^{2k+1}(x, y) = (\sigma^{k+1}\tau^k(y), \sigma^k\tau^{k+1}(x)).$$

This implies that  $\sigma^{k+1}\tau^k(y) = x$  for all  $x, y \in X$ . This equality in particular implies that  $x = y$  because  $\sigma^{k+1}\tau^k$  is a permutation, a contradiction.  $\square$

The connection between the socle of a brace and the retract of a solution was discovered by Rump in the case of involutive solutions and braces of abelian type, see [53].



pro:add\_cyclic

**Proposition 13.7.** *Let  $A$  be a brace and  $(A, r)$  be its associated solution. Then the retraction  $\text{Ret}(A, r)$  is the canonical solution associated with the quotient brace  $A/\text{Soc}(A)$ .*

*Proof.* The equivalence relation  $\sim$  on  $A$  is defined as  $a \sim b$  if and only if  $\lambda_a = \lambda_b$  and  $\mu_a = \mu_b$ . Let  $\bar{A}$  be the set of equivalence classes. The equivalence class of an element  $a$  is then

$$\begin{aligned} [a] &= \{b \in A : a \sim b\} = \{b \in A : \lambda_a = \lambda_b, \mu_a = \mu_b\} \\ &= \{b \in A : a' \circ b \in \ker \lambda \cap \ker \mu\} = \{b \in A : a' \circ b \in \text{Soc}(A)\}, \end{aligned}$$

by Proposition 4.37. This means that  $[a] = [b]$  if and only if  $\pi(a) = \pi(b)$ , where  $\pi: A \rightarrow A/\text{Soc}(A)$ ,  $x \mapsto x \circ \text{Soc}(A)$ , is the canonical brace homomorphism. Moreover,  $A/\text{Soc}(A) = \bar{A}$  as sets. Now we compute the retraction of  $(A, r)$ :

$$\begin{aligned} \bar{r}([a], [b]) &= ([\lambda_a(b)], [\mu_b(a)]) = (\pi(\lambda_a(b)), \pi(\mu_b(a))) \\ &= (\lambda_{\pi(a)}(\pi(b)), \mu_{\pi(b)}(\pi(a))) = (\lambda_{[a]}([b]), \mu_{[b]}([a])). \end{aligned}$$

Therefore  $\text{Ret}(A, r) = (A/\text{Soc}(A), \bar{r})$ .  $\square$

Now...

pro:impl

**Proposition 13.8.** *Let  $(X, r)$  and  $(Y, s)$  be solutions. Each surjective homomorphism of solutions  $f: (X, r) \rightarrow (Y, s)$  induces a surjective homomorphism of solutions  $\text{Ret}(X, r) \rightarrow \text{Ret}(Y, s)$ .*

*Proof.* Write  $r(x, y) = (\sigma_x(y), \tau_y(x))$  and  $s(x, y) = (\lambda_x(y), \mu_y(x))$ . Let  $x, x_1 \in X$  be such that  $x \sim x_1$ . If  $z \in X$ , then

$$\lambda_{f(x)}f(z) = f(\sigma_x(z)) = f(\sigma_{x_1}(z)) = \lambda_{f(x_1)}f(z).$$

Since  $f$  is surjective, it follows that  $\lambda_{f(x)} = \lambda_{f(x_1)}$ . A similar calculation proves that  $\mu_{f(x)} = \mu_{f(x_1)}$ . If  $\pi: (Y, s) \rightarrow \text{Ret}(Y, r)$ ,  $y \mapsto [y]$ , is the canonical map, the composition  $\pi \circ f: (X, r) \rightarrow \text{Ret}(Y, s)$  is a surjective homomorphism of solutions. Therefore the map  $\text{Ret}(X, r) \rightarrow \text{Ret}(Y, s)$ ,  $[x] \mapsto \pi(f(x))$ , is then a well-defined surjective homomorphism of solutions.  $\square$

pro:impl\_subsol

**Proposition 13.9.** *Let  $(X, r)$  be a solution of finite multipermutation level  $m$  and  $Y \subseteq X$  be such that  $r(Y \times Y) \subseteq Y \times Y$ . Then the subsolution  $(Y, r|_{Y \times Y})$  is of finite multipermutation level  $\leq m$ .*

*Proof.*  $\square$

**Theorem 13.10.** *Let  $(X, r)$  be a solution. The following statements are equivalent:*

- 1)  $(X, r)$  has finite multipermutation level.
- 2)  $(\mathcal{G}(X, r), r_{\mathcal{G}(X, r)})$  has finite multipermutation level.
- 3)  $(G(X, r), r_{G(X, r)})$  has finite multipermutation level.

*Proof.* Let us first prove that (2) implies (1). The map  $X \rightarrow \mathcal{G}(X, r)$ ,  $x \mapsto (\lambda_x, \mu_x^{-1})$ , is a homomorphism of solutions that induces an injective homomorphism of solutions  $\text{Ret}(X, r) \rightarrow (\mathcal{G}(X, r), r_{\mathcal{G}(X, r)})$ . Since  $(\mathcal{G}(X, r), r_{\mathcal{G}(X, r)})$  has finite multipermutation level,  $(X, r)$  has finite multipermutation level by Proposition 13.9.

Let us now prove that (3) implies (2). The canonical map  $G(X, r) \rightarrow \mathcal{G}(X, r)$  yields a surjective homomorphism of solutions. Then Proposition 13.8 applies.  $\square$

...

The following result appeared in [29].

**Proposition 13.11.** *Let  $(X, r)$  be a finite involutive solution. If the additive group of the brace  $\mathcal{G}(X, r)$  is cyclic, then  $(X, r)$  is multipermutation.*

*Proof.* Let  $(X, r)$  be a counterexample of minimal cardinality. If  $K$  is the additive group of  $\mathcal{G}(X, r)$ , then  $K$  is finite and cyclic. Write  $G$  for the multiplicative group of  $\mathcal{G}(X, r)$ . Since  $|\text{Aut}(K)| = \varphi(|K|) < |K|$ , where  $\varphi$  is the Euler function, the group homomorphism  $\lambda : G \rightarrow \text{Aut}(K)$  has a non-trivial kernel, so  $\text{Soc}(\mathcal{G}(X, r))$  is non-zero. This implies that  $(X, r)$  is retractable. Since  $\mathcal{G}(X, r)/\text{Soc}(\mathcal{G}(X, r))$  is a brace with cyclic additive group and  $\text{Ret}(X, r)$  is an involutive solution, the minimality of  $|X|$  implies that  $\text{Ret}(X, r)$  is a multipermutation solution, and hence so is  $(X, r)$ , a contradiction.  $\square$

The converse of the previous proposition does not hold.

**Example 13.12.** Let  $X = \{1, 2, 3, 4\}$  and  $r(x, y) = (\varphi_x(y), \varphi_y(x))$ , where

$$\varphi_1 = \varphi_2 = \text{id}, \quad \varphi_3 = (34), \quad \varphi_4 = (12)(34).$$

Then  $(X, r)$  is an involutive multipermutation solution. One easily checks that  $\mathcal{G}(X, r) \simeq C_2 \times C_2$ .

A similar idea proves the following result:

thm:mul\_cyclic

**Theorem 13.13.** *Let  $(X, r)$  be a finite involutive solution. If the multiplicative group of the brace  $\mathcal{G}(X, r)$  is cyclic, then  $(X, r)$  is multipermutation.*

*Proof.* Let  $(X, r)$  be a counterexample of minimal cardinality. Write  $K$  for the additive group of  $\mathcal{G}(X, r)$  and  $G = \langle g \rangle$  for the multiplicative group of  $\mathcal{G}(X, r)$ . Since the image of the group homomorphism  $\lambda : G \rightarrow \text{Aut}(K)$  is cyclic generated by  $\lambda_g$  and  $|\lambda_g| < |G|$  by Horosevskii's theorem, see [36, Corollary 3.3], it follows that  $\lambda$  has a non-trivial kernel, so  $\text{Soc}(\mathcal{G}(X, r))$  is non-zero. This implies that  $(X, r)$  is retractable. Since  $\mathcal{G}(X, r)/\text{Soc}(\mathcal{G}(X, r))$  is a brace with cyclic additive group and  $\text{Ret}(X, r)$  is an involutive solution, the minimality of  $|X|$  implies that  $\text{Ret}(X, r)$  is a multipermutation solution, and hence so is  $(X, r)$ , a contradiction.  $\square$

The previous result does not hold in the case of arbitrary solutions.

**Example 13.14.** Let  $X = \{1, 2, 3, 4, 5, 6\}$  and  $r(x, y) = (\sigma_x(y), \tau_y(x))$ , where

$$\begin{array}{lll} \sigma_1 = \text{id}, & \sigma_2 = \text{id}, & \sigma_3 = \text{id}, \\ \sigma_4 = (23)(56), & \sigma_5 = (23)(56), & \sigma_6 = (23)(56), \\ \tau_1 = \text{id}, & \tau_2 = (456), & \tau_3 = (465), \\ \tau_4 = \text{id}, & \tau_5 = (465), & \tau_6 = (456). \end{array}$$

The brace  $\mathcal{G}(X, r)$  has multiplicative group isomorphic to  $\mathbb{S}_3$  and additive group isomorphic to the cyclic group of order six.

We will see later that Theorem 13.13 is true for braces of nilpotent type. The following example appears in the work of Rump [53].

pro:radical

**Proposition 13.15.** *Let  $A$  be a finite non-trivial radical ring. Then  $\text{Soc}(A) \neq \{0\}$  and  $(A, r_A)$  is an involutive multipermutation solution.*

*Proof.* Let  $A$  be a counterexample of minimal size. This means that  $\text{Soc}(A) = \{0\}$  and all two-sided braces of abelian type of size  $< |A|$  have non-trivial socle. Since  $A$  is finite, there exists a non-zero minimal left ideal  $I$  of  $A$ . Recall that  $A$  is a radical ring with product  $a * b = \lambda_a(b) - b$ . Since  $A$  is a radical ring,  $A$  is a nil ring, which implies by Nakayama's lemma that  $I * A = \{0\}$ . This means that if  $x \in I$ , then  $x \in \text{Soc}(A)$ , as  $0 = x * a = \lambda_x(a) - a$  for all  $a \in A$ . In particular,  $\text{Soc}(A) \neq \{0\}$ , a contradiction.  $\square$

Proposition 13.15 has a nice application. The results appeared first in [19]. The proof presented here is from [20].

thm:CJO\_abelian

**Theorem 13.16.** *Let  $(X, r)$  be a finite involutive solution. If the multiplicative group of the brace  $\mathcal{G}(X, r)$  is abelian, then  $(X, r)$  is multipermutation.*

*Proof.*  $\square$

In [31], Gateva–Ivanova conjectured that finite involutive square-free solutions are retractable.

In [32] Gateva–Ivanova asked when...

Right nilpotency...

The following theorem characterizes multipermutation involutive solutions in terms of left orderability of groups. A group  $G$  is said to be *left ordered* if it admits a total ordering  $<$  such that

$$x < y \implies zx < zy$$

for all  $x, y, z \in G$ . Torsion-free abelian groups, free groups and braid groups are left ordered groups. See [23] for more information on ordered groups.

thm:BCV

**Theorem 13.17.** *Let  $(X, r)$  be a finite involutive solution. The following statements are equivalent:*

- 1)  $(X, r)$  is a multipermutation solution.
- 2)  $G(X, r)$  is poly- $\mathbb{Z}$ .
- 3)  $G(X, r)$  is left orderable.
- 4)  $G(X, r)$  is diffuse.

*Proof.*

□

Recall that a group  $G$  has the *unique product property* if for all finite non-empty subsets  $A$  and  $B$  of  $G$  there exists  $x \in G$  that can be written uniquely as  $x = ab$  with  $a \in A$  and  $b \in B$ .

It is natural to ask when  $G(X, r)$  has the unique product property. By Theorem 13.17, if  $(X, r)$  is an involutive multipermutation solution, then  $G(X, r)$  has the unique product property since  $G(X, r)$  is left orderable.

pro:4-19

**Example 13.18.** Let  $X = \{1, 2, 3, 4\}$  and  $r(x, y) = (\sigma_x(y), \tau_y(x))$  be the irretractable involutive solution given by

$$\begin{array}{llll} \sigma_1 = (12), & \sigma_2 = (1324), & \sigma_3 = (34), & \sigma_4 = (1423), \\ \tau_1 = (14), & \tau_2 = (1243), & \tau_3 = (23), & \tau_4 = (1342). \end{array}$$

We claim that the group  $G(X, r)$  with generators  $x_1, x_2, x_3, x_4$  and relations

$$\begin{array}{lll} x_1^2 = x_2x_4, & x_1x_3 = x_3x_1, & x_1x_4 = x_4x_3, \\ x_2x_1 = x_3x_2, & x_2^2 = x_4^2, & x_3^2 = x_4x_2. \end{array}$$

does not have the unique product property. Let  $x = x_1x_2^{-1}$  and  $y = x_1x_3^{-1}$  and

$$S = \{x^2y, y^2x, xyx^{-1}, (y^2x)^{-1}, (xy)^{-2}, y, (xy)^2x, (xy)^2, (xyx)^{-1}, yxy, y^{-1}, x, xyx, x^{-1}\}. \quad (13.1)$$

eq:Promislow

To prove that  $G(X, r)$  does not have the unique product property it is enough to prove that each  $s \in S^2 = \{s_1s_2 : s_1, s_2 \in S\}$  admits at least two different decompositions of the form  $s = ab = uv$  for  $a, b, u, v \in S$ . To perform these calculations we use the injective group homomorphism  $G \rightarrow \mathbf{GL}(5, \mathbb{Z})$  given by

$$\begin{array}{ll} x_1 \mapsto \begin{pmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, & x_2 \mapsto \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \\ x_3 \mapsto \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, & x_4 \mapsto \begin{pmatrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \end{array}$$

This faithful representation of  $G(X, r)$  allows us to compute all possible products of the form  $s_1s_2$  for all  $s_1, s_2 \in S$ . By inspection, each element of  $S^2$  admits at least two different representations.

thm:CJO\_mp

**Theorem 13.19.** *Let  $A$  be a finite brace of abelian type with multiplicative group  $G$ . Then there exists a finite solution  $(X, r)$  such that  $\text{Ret}(X, r)$  is isomorphic to  $(A, r_A)$  and  $\mathcal{G}(X, r) \simeq G$ .*

*Proof.* Let  $X = A \times \mathbb{Z}/(2)$ . For  $a, b \in A$ , let

$$\begin{aligned}\varphi_{(a,0)}(b, 0) &= (b, 0), & \varphi_{(a,0)}(b, 1) &= (b, 1), \\ \varphi_{(a,1)}(b, 0) &= (a' \circ b, 0), & \varphi_{(a,1)}(b, 1) &= (\lambda_a^{-1}(b), 1).\end{aligned}$$

The maps  $\varphi_{(a,\varepsilon)}$  are invertible for all  $a \in A$  and  $\varepsilon \in \mathbb{Z}/(2)$ . In fact,

$$\varphi_{(a,0)}^{-1} = \text{id}, \quad \varphi_{(a,1)}^{-1}(b, \varepsilon) = \begin{cases} a \circ b & \text{if } \varepsilon = 0, \\ \lambda_a(b) & \text{if } \varepsilon = 1. \end{cases}$$

So we need to check that

$$((a, \varepsilon_1) \cdot (b, \varepsilon_2)) \cdot ((a, \varepsilon_1) \cdot (c, \varepsilon_3)) = ((b, \varepsilon_2) \cdot (a, \varepsilon_1)) \cdot ((b, \varepsilon_2) \cdot (c, \varepsilon_3)) \quad (13.2)$$

eq:CJO\_tocheck

holds for all  $a, b, c \in A$  and  $\varepsilon_1, \varepsilon_2, \varepsilon_3 \in \mathbb{Z}/(2)$ . There are several cases to consider. Let us assume first that  $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, 1, 0)$ . Then (13.2) turns out to be

$$(\lambda_a^{-1}(b))' \circ a' \circ c, 0) = (\lambda_b^{-1}(a))' \circ b' \circ c, 0),$$

which holds for all  $a, b \in A$ , as

$$\lambda_a^{-1}(b)' \circ a' \circ c = (a \circ \lambda_a^{-1}(b))' \circ c = (a + b)' \circ c = (b + a)' \circ c = \lambda_b^{-1}(a)' \circ b' \circ c$$

because  $A$  is of abelian type. Let us now deal with the case  $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, 1, 1)$ . In this case Equality (13.2) turns out to be equivalent to

$$(\lambda_a^{-1}(b), 1) \cdot (\lambda_a^{-1}(c), 1) = (\lambda_b^{-1}(a), 1) \cdot (\lambda_b^{-1}(c), 1).$$

Since  $A$  is of abelian type,

$$\begin{aligned}(\lambda_a^{-1}(b), 1) \cdot (\lambda_a^{-1}(c), 1) &= \lambda_{\lambda_a^{-1}(b)}^{-1} \lambda_a^{-1}(c) = \lambda_{a \circ \lambda_a^{-1}(b)}^{-1}(c) = \lambda_{a+b}^{-1}(c) \\ &= \lambda_{b+a}^{-1}(c) = \lambda_{\lambda_b^{-1}(a)}^{-1} \lambda_b^{-1}(c) = (\lambda_b^{-1}(a), 1) \cdot (\lambda_b^{-1}(c), 1).\end{aligned}$$

The other cases are easier and require straightforward calculations.

Let  $\psi: G \rightarrow \mathbb{S}_X$ ,  $\psi(g) = \varphi_{(g',1)}$ . Since

$$\psi(g)(0, 0) = \varphi_{(g',1)}(0, 0) = (g \circ 0, 0) = (g, 0),$$

it follows that  $\psi$  is injective. Moreover,  $\psi$  is a group homomorphism, as

$$\begin{aligned}\psi(a)\psi(b)(c, 0) &= \psi(a)\varphi_{(b,1)}(c, 0) = \psi(a)(b \circ c, 0) \\ &= \varphi_{(a,1)}(b \circ c, 0) = (a \circ b \circ c, 0) = \varphi_{(a \circ b,1)}(c, 0) = \psi_{(a \circ b)}(c, 0)\end{aligned}$$

and

$$\begin{aligned}\psi(a)\psi(b)(c, 1) &= \varphi_{(a',1)}\varphi_{(b',1)}(c, 1) = \varphi_{(a',1)}(\lambda_{b'}^{-1}(c), 1) \\ &= (\lambda_{a'}^{-1}\lambda_{b'}^{-1}(c), 1) = (\lambda_{a \circ b}(c), 1) = \varphi_{(a \circ b,1)}(c, 1) = \psi(a \circ b)(c, 1).\end{aligned}$$

Since  $\psi$  is an injective group homomorphism,

$$G \simeq \psi(G) \simeq \langle \psi(a) : a \in A \rangle = \langle \varphi_{(a,1)} : a \in A \rangle \simeq \mathcal{G}(X, r).$$

Consider the equivalence relation on  $X$  given by  $x \sim y$  if and only if  $\varphi_x = \varphi_y$ . As usual  $[x]$  denotes the equivalence class of the element  $x \in X$  and  $\bar{X}$  is the set of equivalence classes. A straightforward computation shows that  $(a, 0) \sim (0, 1)$  for all  $a \in A$ . This implies that

$$\bar{p}: \bar{X} \rightarrow A, \quad \bar{p}([(a, \varepsilon)]) = \begin{cases} 0 & \text{if } \varepsilon = 0, \\ a & \text{if } \varepsilon = 1, \end{cases}$$

is a well-defined surjective map. We claim that  $\bar{p}$  is injective. Let  $(a, \varepsilon_1) \in \bar{X}$  and  $(b, \varepsilon_2) \in \bar{X}$  be such that  $\bar{p}([(a, \varepsilon_1)]) = \bar{p}([(b, \varepsilon_2)])$ . Since  $[(a, 0)] = [(0, 1)]$  for all  $a \in A$ , we only need to consider the case where  $\varepsilon_1 = \varepsilon_2 = 1$ . In this case,

$$a = \bar{p}([(a, \varepsilon_1)]) = \bar{p}([(b, \varepsilon_2)]) = b.$$

Thus  $\bar{p}$  is bijective. Now

$$\begin{aligned}r_A(\bar{p}([(a, 1)]), \bar{p}([(b, 1)])) &= r_A(a, b) = (\lambda_a(b), \mu_b(a)) = (\bar{p}[\lambda_a(b), 1], \bar{p}[\mu_b(a), 1]) \\ &= (\bar{p}\varphi_{(a,1)}(b, 1), \bar{p}\dots).\end{aligned}$$

□

**Theorem 13.20.** *Let  $A$  be...*

*Proof.*

□

## Exercises

prob:bounded\_mpl

**13.1.** Let  $(X, r)$  be a solution of finite multipermutation level  $m$ . Prove that any homomorphic image of  $(X, r)$  is a solution of finite multipermutation level  $\leq m$ .

prob:4-13

**13.2.** Let  $X = \{1, 2, 3, 4\}$  and  $r(x, y) = (\sigma_x(y), \tau_y(x))$  be the irretractable involutive solution given by

$$\begin{array}{llll}\sigma_1 = (34), & \sigma_2 = (1324), & \sigma_3 = (1423), & \sigma_4 = (12), \\ \tau_1 = (24), & \tau_2 = (1432), & \tau_3 = (1234), & \tau_4 = (13).\end{array}$$

Prove that  $G(X, r)$  does not have the unique product property.

## Open problems

**Problem 13.1.** Does the group  $G(X, r)$ ... A linear representation of this group is...

## Notes

Multipermutation involutive solutions were introduced in [29]. The notion was extended to the non-involutive case in [44].

Theorem 13.6 was proved in...

Proposition 13.8 was proved in [20] for involutive solutions. The general case goes back to...

Theorem 13.17 combines several results. The implication...

Theorem 13.19 appears in [20].

Non-involutive multipermutation solutions...

The set (13.1) appears in the work of Promislow [50]. Exercise 13.2 appears in the book of Jespers and Okniński, see [40, Example 8.2.14].





## Chapter 14

### Ordered groups

#### A

A group  $G$  is **left ordered** if there is a total ordering  $<$  on  $G$  such that  $x < y$  implies  $zx < zy$  for all  $x, y, z \in G$ . Similarly one defines right ordered groups.

**Example 14.1.** The group  $\mathbb{Z}$  is left ordered.

**Example 14.2.** If  $G$  is a left ordered group and  $H$  is a subgroup of  $G$ , then  $H$  is left ordered.

**Example 14.3.** Let  $G = \mathbb{Z}^2$  and  $v \in \mathbb{R}^2$  with irrational slope. Then...

**Proposition 14.4.** *Let ... be a short exact sequence of groups. If both  $K$  and  $Q$  are both left ordered, then  $G$  is left ordered.*

*Proof.*

□

#### 14B

#### 14C



## Chapter 15

### Transitive groups

**Definition 15.1.** A finite solution  $(X, r)$  is said to be **decomposable** if there is a decomposition  $X = X_1 \cup X_2$  of  $X$  into a disjoint union of non-empty subsets  $X_1$  and  $X_2$  such that  $r(X_1 \times X_1) \subseteq X_1 \times X_2$  and  $r(X_2 \times X_2) \subseteq X_2 \times X_2$ . A solution  $(X, r)$  is then **indecomposable** if it is not decomposable.

If  $(X, r)$  is a finite decomposable solution and  $X = X_1 \cup X_2$  is a decomposition, then the restrictions  $r|_{X_1 \times X_1}$  and  $r|_{X_2 \times X_2}$  are solutions. Moreover, it follows that  $r(X_1 \times X_2) \subseteq X_2 \times X_1$  and  $r(X_2 \times X_1) \subseteq X_1 \times X_2$ , see Exercise 15.1.

**Proposition 15.2.** A finite solution  $(X, r)$  is indecomposable if and only if the group

$$\langle \sigma_x, \tau_x : x, y \in X \rangle$$

acts transitively on  $X$ .

*Proof.* Let us assume that  $X = X_1 \cup X_2$  is a decomposition of  $X$  into non-empty orbits...  $\square$

Note that this group is in general not isomorphic to the permutation group of the solution.

**Definition 15.3.** A finite solution  $(X, r)$  is said to be **simple** if  $|X| > 1$  and for every surjective homomorphism  $f: (X, r) \rightarrow (Y, s)$  of solutions either  $f$  is an isomorphism or  $|Y| = 1$ .

**Example 15.4.**

**Example 15.5.**

**Example 15.6.**

**Proposition 15.7.** Let  $(X, r)$  be a finite simple solution. If  $|X| > 2$ , then  $(X, r)$  is indecomposable. involutive?

o:simple=>indecomposable

*Proof.* Let us assume that  $(X, r)$  is decomposable. Decompose  $X = X_1 \cup X_2$  for non-empty disjoint subsets  $X_1$  and  $X_2$  of  $X$  such that  $r(X_i \times X_i) \subseteq X_i \times X_i$  for  $i \in \{1, 2\}$ . Let  $Y = \{1, 2\}$  and  $s: Y \times Y \rightarrow Y \times Y$ ,  $s(x, y) = (y, x)$ . Since  $X = X_1 \cup X_2$  is a decomposition, it follows that  $r(X_i \times X_j) \subseteq X_j \times X_i$  for all  $i, j \in \{1, 2\}$ . Why? Let  $f: X \rightarrow Y$ ,  $f(x) = i$  if  $x \in X_i$ . Since  $f$  is then a surjective homomorphism of solutions and  $f$  is not an isomorphism (because  $|X| > 2$ ), the simplicity of  $(X, r)$  implies that  $|Y| = 1$ , a contradiction.  $\square$

**Proposition 15.8.** *Let  $(X, r)$  be a finite simple involutive solution. If  $|X|$  is not a prime number, then  $(X, r)$  is irretractable.*

*Proof.* Let us assume that  $(X, r)$  is retractable. Let  $(X, r) \rightarrow \text{Ret}(X, r)$ ,  $x \mapsto [x]$ , be the canonical map. Since it is a surjective homomorphism of solutions and  $(X, r)$  is retractable, the simplicity of  $(X, r)$  implies that  $|\text{Ret}(X, r)| = 1$ . Therefore  $(X, r)$  is a permutation solution, say  $r(x, y) = (\sigma(y), \tau(x))$  for some commuting permutations  $\sigma: X \rightarrow X$  and  $\tau: X \rightarrow X$ . Since  $|X| > 2$ , the solution  $(X, r)$  is indecomposable by Proposition 15.7. This implies that  $\sigma$  is a cycle of length  $|X|$  and  $\tau = \sigma^k$  for some  $k \in \mathbb{Z}$ . Let us assume that  $\sigma = (x_1 \cdots x_n)$ , where  $n = |X|$ . Since  $n$  is not a prime number,  $n = dm$  for some  $1 < d < n$ . Let  $Y = \mathbb{Z}/(d)$  and  $s: Y \times Y \rightarrow Y \times Y$ ,  $s(i, j) = (j + 1, i + k)$ . Then  $(Y, s)$  is a solution. The map  $f: X \rightarrow Y$ ,  $f(x_i) = i \bmod d$  satisfies  $f(\tau_{x_j}(x_i)) = i + k$  and

$$f(\sigma_{x_i}(x_j)) = \begin{cases} f(x_{j+1}) & \text{if } j < n, \\ 1 & \text{if } j = n. \end{cases}$$

Then a straightforward computation shows that  $f$  is a surjective homomorphism of solutions, a contradiction.  $\square$

The previous proposition cannot be extended to the non-involutive case.

**Example 15.9.** Let  $X = \{1, \dots, 6\}$ . The permutation solution with permutations  $\sigma = (153)(264)$  and  $\tau = (12)(34)(56)$  is indecomposable.

## Exercises

prob:decomposition

**15.1.** Let  $(X, r)$  be a finite decomposable solution and  $X = X_1 \cup X_2$  be a decomposition. Prove that  $r(X_1 \times X_2) \subseteq X_2 \times X_1$  and  $r(X_2 \times X_1) \subseteq X_1 \times X_2$ . Solution?

**15.2.** Let  $(X, r)$  be a finite involutive permutation solution. Prove that  $(X, r)$  is indecomposable if and only if  $\sigma$  is a cycle of length  $|X|$ .

**Open problems**

**Notes**



## Chapter 16

### Regular subgroups

For an additive group  $A$ , the **holomorph** of  $A$  is the semidirect product

$$\text{Hol}(A) = A \rtimes \text{Aut}(A).$$

This means that the operation is

$$(a, f)(b, g) = (a + f(b), f \circ g), \quad a, b \in A, \quad f, g \in \text{Aut}(A).$$

Every subgroup  $G$  of  $\text{Hol}(A)$  acts on  $A$  by

$$(x, f) \cdot a = \pi_1((x, f)(a, \text{id})) = \pi_1(x + f(a), f) = x + f(a), \quad a, x \in A, \quad f \in \text{Aut}(A),$$

where  $\pi_1 : \text{Hol}(A) \rightarrow A, (a, f) \mapsto a$ .

**Exercise 16.1.** The group  $\text{Hol}(A)$  acts transitively on  $A$  and the stabilizer  $a \in A$  is isomorphic to  $\text{Aut}(A)$ .

A subgroup  $G$  of  $\text{Hol}(A)$  is said to be *regular* if it acts regularly on  $A$ , this means that given  $a, b \in A$  there exists a unique  $(x, f) \in G$  such that

$$b = (x, f) \cdot a = x + f(a).$$

lem:bijective

**Lemma 16.2.** If  $G$  is a regular subgroup of  $\text{Hol}(A)$ , then  $\pi_1 : G \rightarrow A$  is bijective.

*Proof.* We first prove that restriction of  $\pi_1$  onto  $G$  is injective. Let  $(a, f) \in G$  and  $(b, g) \in G$  be such that  $\pi_1(a, f) = \pi_1(b, g)$ . Then  $a = b$ . Since  $G$  is a subgroup,

$$(-f^{-1}(a), f^{-1}) = (a, f)^{-1} \in G, \quad (-g^{-1}(a), g^{-1}) = (a, g)^{-1} \in G,$$

and hence  $f = g$  since

$$(-f^{-1}(a), f^{-1}) \cdot a = 0 = (-g^{-1}(a), g^{-1}) \cdot a$$

and  $G$  is a regular subgroup. Now we prove that  $\pi_1|_G$  is surjective. Let  $a \in A$ . Since  $G$  is regular, there exists  $(x, f) \in G$  such that  $x + f(a) = (x, f) \cdot a = 0$ , so  $(-f(a), f) \in G$  for some  $f \in \text{Aut}(A)$ . Then  $(a, f^{-1}) = (-f(a), f)^{-1} \in G$  and  $\pi_2|_G(a, f^{-1}) = a$ .  $\square$

Now we establish an important connection between braces and regular subgroups.

thm:regular

**Theorem 16.3.** *If  $A$  is a brace, then  $\Delta = \{(a, \lambda_a) : a \in A\}$  is a regular subgroup of  $\text{Hol}(A, +)$ . Conversely, if  $A$  is an additive group and  $G$  is a regular subgroup of  $\text{Hol}(A)$ , then  $A$  is a brace with*

$$a \circ b = a + f(b),$$

where  $(\pi_1|_G)^{-1}(a) = (a, f) \in G$ .

*Proof.* Assume first that  $A$  is a brace. Using (4.2) and that  $\lambda$  is a group homomorphism, it follows that  $\Delta = \{(a, \lambda_a) : a \in A\}$  is a subgroup of  $\text{Hol}(A, +)$ , as

$$\begin{aligned} (a, \lambda_a)^{-1} &= (\lambda_a^{-1}(-a), \lambda_a^{-1}) = (a', \lambda_{a'}) \in \Delta, \\ (a, \lambda_a)(b, \lambda_b) &= (a + \lambda_a(b), \lambda_a \circ \lambda_b) = (a \circ b, \lambda_{a \circ b}) \in \Delta. \end{aligned}$$

To see that  $\Delta$  is a regular subgroup, note that  $(c, \lambda_c) \cdot a = b$  implies that  $c = b \circ a'$ , as  $(A, \circ)$  is a group.

Assume now that  $A$  is an additive group and that  $G$  is a regular subgroup of  $\text{Hol}(A)$ . By Lemma 16.2, the restriction  $\pi_1|_G$  is bijective. Use the bijection  $\pi_1|_G$  to transport the operation of  $G$  into  $A$ :

$$a \circ b = \pi_1|_G((\pi_1|_G)^{-1}(a)(\pi_1|_G)^{-1}(b)) = a + f(b),$$

where  $a, b \in A$  and  $(\pi_1|_G)^{-1}(a) = (a, f) \in G$ . Then  $(A, \circ)$  is a group isomorphic to  $G$  and moreover  $A$  is a brace, as

$$\begin{aligned} a \circ (b + c) &= a + f(b + c) = a + f(b) + f(c) \\ &= a + f(b) - a + a + f(c) = a \circ b - a + a \circ c \end{aligned}$$

holds for all  $a, b, c \in A$ .  $\square$

The following lemma is from [9].

lem:BNY

**Lemma 16.4.** *Let  $A$  be a group. If  $H$  and  $K$  are conjugate regular subgroups of  $\text{Hol}(A)$ , then  $H$  and  $K$  are conjugate by an automorphism of  $A$ .*

*Proof.* Assume that  $H$  and  $K$  are conjugate in  $\text{Hol}(A)$ . Let  $(b, g) \in \text{Hol}(A)$  be such that  $(b, g)^{-1}H(b, g) = K$ . Since  $b \in A$ , the regularity of  $H$  implies that there exists  $(a, f) \in H$  such that  $a + f(b) = 0$ . Since  $(a, f) \in H$ ,

$$\begin{aligned} K &= (b, g)^{-1}H(b, g) = (b, g)^{-1}(a, f)^{-1}H(a, f)(b, g) \\ &= (0, f \circ g)^{-1}H(0, f \circ g) = (f \circ g)^{-1}H(f \circ g). \end{aligned} \quad \square$$



pro:regular

**Proposition 16.5.** *Let  $A$  be an additive group. There exists a bijective correspondence between isomorphism classes of brace structures with additive group  $A$  and conjugacy classes of regular subgroups of  $\text{Hol}(A)$ .*

*Proof.* Assume that the additive group  $A$  has two isomorphic brace structures given by  $(a, b) \mapsto a \circ b$  and  $(a, b) \mapsto a \times b$ . Let  $f: A \rightarrow A$  be a bijective map such that  $f(a+b) = f(a) + f(b)$  and  $f(a \circ b) = f(a) \times f(b)$  for all  $a, b \in A$ . We claim that the regular subgroups  $\{(a, \lambda_a) : a \in A\}$  and  $\{(a, \mu_a) : a \in A\}$ , where  $\lambda_a(b) = -a + a \circ b$  and  $\mu_a(b) = -a + a \times b$ , are conjugate. Since  $f$  is an isomorphism of braces,

$$f \circ \lambda_a \circ f^{-1} = \mu_{f(a)}$$

for all  $a \in A$ . This implies that  $(0, f)(a, \lambda_a)(0, f)^{-1} = (f(a), \mu_{f(a)})$  for all  $a \in A$  and hence the first claim follows.

Conversely, let  $H$  and  $K$  be conjugate regular subgroups of  $\text{Hol}(A)$ . Since  $H$  and  $K$  are conjugate in  $\text{Hol}(A)$ , by Lemma 16.4 there exists  $\varphi \in \text{Aut}(A)$  such that  $\varphi H \varphi^{-1} = K$ . The brace structure on  $A$  corresponding to the subgroup  $H$  is given by  $a \circ b = a + f(b)$ , where  $(a, f) = (\pi_1|_H)^{-1}(a) \in H$ , see Lemma 16.2. Since

$$\varphi(f, a)\varphi^{-1} = (\varphi(a), \varphi \circ f \circ \varphi^{-1}) \in K,$$

it follows that  $(\pi_1|_K)^{-1}(\varphi(a)) = (\varphi(a), \varphi \circ f \circ \varphi^{-1})$ . Since  $\varphi \in \text{Aut}(A)$ ,

$$\begin{aligned} \varphi(a) \times \varphi(b) &= \varphi(a) + (\varphi \circ f \circ \varphi^{-1})(\varphi(b)) \\ &= \varphi(a) + \varphi(f(b)) = \varphi(a + f(b)) = \varphi(a \circ b) \end{aligned}$$

and hence the braces corresponding to  $H$  and  $K$  are isomorphic.  $\square$

Now we present algorithm used to enumerate braces. It is based on Theorem 16.3. The use of Lemma 16.4 in Proposition 16.5 significantly improves the performance.

alg:regular

**Algorithm 16.6.** Let  $A$  be a finite group. To construct all braces with additive group  $A$  we proceed as follows:

- 1) Compute the holomorph  $\text{Hol}(A)$  of  $A$ .
- 2) Compute the list of regular subgroups of  $\text{Hol}(A)$  of order  $|A|$  up to conjugation.
- 3) For each representative  $G$  of regular subgroups of  $\text{Hol}(A)$  construct the map  $p: A \rightarrow G$  given by  $a \mapsto (a, f) \in G$ . Then the set  $A$  is a brace with additive group  $A$  and multiplication given by  $a \circ b = p^{-1}(p(a)p(b))$  for all  $a, b \in A$ .

To enumerate all isomorphism classes of braces structures with a fixed additive group the third step of Algorithm 16.6 is not needed. Algorithm 16.6 can be used to compute the number  $s(n)$  of non-isomorphic braces of size  $n$ . With small modifications it could be used to compute the number  $a(n)$  of non-isomorphic braces of abelian type of size  $n$ , or the number of non-isomorphic radical rings, or the number of non-isomorphic braces of nilpotent type. Some values for  $s(n)$  and  $a(n)$  appear in Table 16.1.

**Table 16.1:** The number of non-isomorphic braces.

$n$	1	2	3	4	5	6	7	8	9	10	11	12
$a(n)$	1	1	1	4	1	2	1	27	4	2	1	10
$s(n)$	1	1	1	4	1	6	1	47	4	6	1	38
$n$	13	14	15	16	17	18	19	20	21	22	23	24
$a(n)$	1	2	1	357	1	8	1	11	2	2	1	96
$s(n)$	1	6	1	1605	1	49	1	43	8	6	1	855
$n$	25	26	27	28	29	30	31	32	33	34	35	36
$a(n)$	4	2	37	9	1	4	1	25281	1	2	1	46
$s(n)$	4	6	101	29	1	36	1	1223061	1	6	1	400
$n$	37	38	39	40	41	42	43	44	45	46	47	48
$a(n)$	1	2	2	106	1	6	1	9	4	2	1	1708
$s(n)$	1	6	8	944	1	78	1	29	4	6	1	66209
$n$	49	50	51	52	53	54	55	56	57	58	59	60
$a(n)$	4	8	1	11	1	80	2	91	2	2	1	28
$s(n)$	4	51	1	43	1	1028	12	815	8	6	1	418
$n$	61	62	63	64	65	66	67	68	69	70	71	72
$a(n)$	1	2	11	?	1	4	1	11	1	4	1	489
$s(n)$	1	6	11	?	1	36	1	43	1	36	1	17790
$n$	73	74	75	76	77	78	79	80	81	82	83	84
$a(n)$	1	2	5	9	1	6	1	1985	804	2	1	34
$s(n)$	1	6	14	29	1	78	1	74120	8436	6	1	606
$n$	85	86	87	88	89	90	91	92	93	94	95	96
$a(n)$	1	2	1	90	1	16	1	9	2	2	1	195971
$s(n)$	1	6	1	800	1	294	1	29	8	6	1	?
$n$	97	98	99	100	101	102	103	104	105	106	107	108
$a(n)$	1	8	4	51	1	4	1	106	2	2	1	494
$s(n)$	1	53	4	711	1	36	1	944	8	6	1	11223
$n$	109	110	111	112	113	114	115	116	117	118	119	120
$a(n)$	1	6	2	1671	1	6	1	11	11	2	1	395
$s(n)$	1	94	8	65485	1	78	1	43	47	6	1	22711
$n$	121	122	123	124	125	126	127	128	129	130	131	132
$a(n)$	4	2	1	9	49	36	1	?	2	4	1	24
$s(n)$	4	6	1	29	213	990	1	?	8	36	1	324
$n$	133	134	135	136	137	138	139	140	141	142	143	144
$a(n)$	1	2	37	108	1	4	1	27	1	2	1	10215
$s(n)$	1	6	101	986	1	36	1	395	1	6	1	3013486
$n$	145	146	147	148	149	150	151	152	153	154	155	156
$a(n)$	1	2	9	11	1	19	1	90	4	4	2	40
$s(n)$	1	6	123	43	1	401	1	800	4	36	12	782
$n$	157	158	159	160	161	162	163	164	165	166	167	168
$a(n)$	1	2	1	209513	1	1374	1	11	2	2	1	443
$s(n)$	1	6	1	?	1	45472	1	43	12	6	1	28505

tab:braces

## Open problems

**Problem 16.1.** Estimate  $s(n)$  and  $a(n)$  for  $n \rightarrow \infty$ .

**Problem 16.2.** Construct and enumerate braces of size 64, 96, 128 and 160.

## Notes

Theorem 16.3 was first observed by Catino and Rizzo [16] and Bachiller [7].

Algorithm 16.6 and most of the numbers of Table 16.1 appeared in [34]. It should be noted that the number of braces of size 57 of [34] is incorrect; the correct value is  $s(57) = 8$ , as Table 16.1 shows. Lemma 16.4 appears in [9] and it is needed to compute the number  $a(n)$  of isomorphism classes of braces of abelian type of size  $n \in \{32, 81, 96, 144, 160, 162\}$  and the number  $s(n)$  of isomorphism classes of braces of size  $n \in \{32, 54, 80, 81, 108, 112, 120, 136, 144, 147, 150, 152, 162, 168\}$ .



## Chapter 17

### The transfer map

#### A

Let  $G$  be a group and  $H$  be a finite index subgroup. We will define a group homomorphism  $G \rightarrow H/[H, H]$ , known as the **transfer map** of  $G$  on  $H$ . Fix a **left transversal**  $T$  of  $H$  in  $G$ .

lem:sigma

**Lemma 17.1.** *Let  $G$  be a group and  $H$  be a subgroup of finite index  $n = (G : H)$ . Let  $S = \{s_1, \dots, s_n\}$  and  $T = \{t_1, \dots, t_n\}$  be transversals of  $H$  in  $G$ . If  $g \in G$ , there exist unique  $h_1, \dots, h_n \in H$  and a permutation  $\sigma \in \mathbb{S}_n$  such that*

$$gt_i = s_{\sigma(i)}h_i, \quad i \in \{1, \dots, n\}.$$

*Proof.* If  $i \in \{1, \dots, n\}$ , then there exists a unique  $j \in \{1, \dots, n\}$  such that  $gt_i \in s_jH$ . Thus there exists a unique  $h_i \in H$  such that  $gt_i = s_jh_i$ . Take  $\sigma(i) = j$  and thus there is a well-defined map  $\sigma: \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ . To prove that  $\sigma \in \mathbb{S}_n$  it is enough to check that  $\sigma$  is injective. If  $\sigma(i) = \sigma(k) = j$ , since  $gt_i = s_jh_i$  and  $gt_k = s_jh_k$ , it follows that  $t_i^{-1}t_k = h_i^{-1}h_k \in H$ . Hence  $i = k$ , as  $t_iH = t_kH$ .  $\square$

Let  $G$  be a group and  $H$  be a subgroup of  $G$  of finite index  $n$ . If  $T = \{t_1, \dots, t_n\}$  is a transversal of  $H$  in  $G$ , we define the map

$$v_T: G \rightarrow H/[H, H], \quad v_T(g) = \prod_{i=1}^n h_i$$

where  $gt_i = t_jh_i$ . Note that the product is well-defined since  $H/[H, H]$  is an abelian group. We now prove that the map does not depend on the transversal.

lem:nu\_T

**Lemma 17.2.** *Let  $G$  be a group and  $H$  be a subgroup of  $G$  of finite index. if  $T$  and  $S$  are transversals of  $H$  in  $G$ , then  $v_T = v_S$ .*

*Proof.* Assume that  $gs_i = s_{\sigma(i)}h_i$  for all  $i$ . Write  $s_i = t_ik_i$ ,  $k_i \in H$ . If  $l_i = k_{\sigma(i)}h_ik_i^{-1}$ , then

$$gt_i = gs_i k_i^{-1} = s_{\sigma(i)} h_i k_i^{-1} = t_{\sigma(i)} k_{\sigma(i)} h_i k_i^{-1} = t_{\sigma(i)} l_i$$

for all  $i \in \{1, \dots, n\}$ . Moreover,

$$s_{\sigma(i)}^{-1} g s_i = k_{\sigma(i)}^{-1} t_{\sigma(i)}^{-1} g t_i k_i.$$

Since  $H/[H, H]$  is abelian,

$$\begin{aligned} v_S(g) &= \prod_{i=1}^n s_{\sigma(i)}^{-1} g s_i = \prod_{i=1}^n k_{\sigma(i)}^{-1} t_{\sigma(i)}^{-1} g t_i k_i \\ &= \prod_{i=1}^n k_{\sigma(i)}^{-1} \prod_{i=1}^n k_i \prod_{i=1}^n t_{\sigma(i)}^{-1} g t_i = \prod_{i=1}^n t_{\sigma(i)}^{-1} g t_i = v_T(g). \end{aligned} \quad \square$$

By Lemma 17.2, if  $H$  is a finite-index subgroup of  $G$ , the map

$$v: G \rightarrow H/[H, H], \quad v(g) = v_T(g),$$

where  $T$  is some transversal of  $H$  in  $G$ , is well-defined.

theorem:transfer

**Theorem 17.3.** *Let  $G$  be a group and  $H$  be a finite-index subgroup of  $G$ . Then  $v(xy) = v(x)v(y)$  for all  $x, y \in G$ .*

*Proof.* Let  $T = \{t_1, \dots, t_n\}$  be a transversal of  $H$  in  $G$ . Let  $x, y \in G$ . By Lemma 17.1, there exist unique elements  $h_1, \dots, h_n, k_1, \dots, k_n \in H$  and there are permutations  $\sigma, \tau \in \mathbb{S}_n$  such that  $xt_i = t_{\sigma(i)} h_i$  and  $yt_i = t_{\tau(i)} k_i$ . Since

$$xyt_i = xt_{\tau(i)} k_i = t_{\sigma\tau(i)} h_{\tau(i)} k_i$$

and  $H/[H, H]$  is abelian,

$$v(xy) = \prod_{i=1}^n h_{\tau(i)} k_i = \prod_{i=1}^n h_{\tau(i)} \prod_{i=1}^n k_i = v(x)v(y). \quad \square$$

If  $G$  is a group and  $H$  is a finite-index subgroup of  $G$ , the **transfer homomorphism** is the group homomorphism  $v: G \rightarrow H/[H, H]$ ,  $v(g) = v_T(g)$ , for some transversal  $T$  of  $H$  in  $G$ .

lem:evaluation

**Lemma 17.4.** *Let  $G$  be a group and  $H$  be a subgroup of  $G$  with  $(G : H) = n$ . Let  $T = \{t_1, \dots, t_n\}$  be a transversal of  $H$  in  $G$ . For each  $g \in G$  there are elements  $s_1, \dots, s_m \in T$  and positive integers  $n_1, \dots, n_m$  (depending on  $g$ ) such that*

$$s_i^{-1} g^{n_i} s_i \in H, \quad n_1 + \dots + n_m = n \quad \text{and} \quad v(g) = \prod_{i=1}^m s_i^{-1} g^{n_i} s_i.$$

*Proof.* For each  $i$  there exist  $h_1, \dots, h_n \in H$  and  $\sigma \in \mathbb{S}_n$  such that  $gt_i = t_{\sigma(i)} h_i$ . Write  $\sigma$  as a product

$$\sigma = \alpha_1 \cdots \alpha_m$$

of disjoint cycles.

For each  $i \in \{1, \dots, n\}$ , write  $\alpha_i = (j_1 \cdots j_{n_i})$ . Since

$$gt_{j_k} = t_{\sigma(j_k)}h_{j_k} = \begin{cases} t_{j_1}h_{n_i} & \text{if } i = n_i, \\ t_{j_{k+1}}h_k & \text{otherwise,} \end{cases}$$

it follows that

$$t_{j_1}^{-1}g^{n_i}t_{j_1} = t_{j_1}^{-1}gg^{n_i-1}t_{j_1} = t_{j_1}^{-1}gt_{j_r}h_{j_{r-1}} \cdots h_{j_1} = h_{j_r} \cdots h_{j_1} \in H,$$

and we let  $s_i = t_{j_1}$ . Now the claim follows, since  $v(g) = h_1 \cdots h_n$ .  $\square$

prop:  $v(g) = g^n$

**Proposition 17.5.** *Let  $G$  be a group and  $H$  be a central subgroup of index  $n$ . Then  $v(g) = g^n$  for all  $g \in G$ .*

*Proof.* Let  $g \in G$ . By Lemma 17.4, there exist  $s_1, \dots, s_m \in H$  such that  $s_i^{-1}g^{n_i}s_i \in H$  and  $v(g) = \prod_{i=1}^m s_i^{-1}g^{n_i}s_i$ . Since  $H$  is central in  $G$ , then it is normal in  $G$ . Thus

$$g^{n_i} = s_i(s_i^{-1}g^{n_i}s_i)s_i^{-1} \in H \subseteq Z(G)$$

and hence

$$v(g) = \prod_{i=1}^m s_i^{-1}g^{n_i}s_i = \prod_{i=1}^m g^{n_i} = g^{\sum_{i=1}^m n_i} = g^n. \quad \square$$

**Exercise 17.6.** Let  $G$  be a group with a central subgroup  $H$  of index  $n$ . Then  $g \mapsto g^n$  is a group homomorphism.

corollary:  $[x, y]^n = 1$

**Exercise 17.7.** Let  $G$  be a group such that  $(G : Z(G)) = n$ . If  $x, y \in G$ , then  $[x, y]^n = 1$ .

Another application.

prop: semidirecto

**Proposition 17.8.** *Sea  $G$  un grupo finito y sea  $H$  un subgrupo abeliano de índice  $n$ , donde  $n$  es coprimo con  $|H|$ . Sea  $N = \ker(v : G \rightarrow H)$ . Entonces  $G \simeq N \rtimes H$ .*

*Proof.* Since  $H$  is abelian,  $H = H/[H, H]$  and the transfer map is  $v : G \rightarrow H$ . By Lemma 17.4,

$$v(h) = \prod_{i=1}^m s_i^{-1}h^{n_i}s_i = \prod_{i=1}^m h^{n_i} = h^{\sum_{i=1}^m n_i} = h^n.$$

The composition  $H \hookrightarrow G \xrightarrow{v} H$  is a group homomorphism.

We claim that it is an isomorphism. It is injective: If  $h^n = 1$ , then  $|h|$  divides  $|H|$  and divides  $n$ . Since  $n$  and  $|H|$  are coprime,  $h = 1$ . It is surjective: Since  $n$  and  $|H|$  are coprime, there exists  $m \in \mathbb{Z}$  such that  $nm \equiv 1 \pmod{|H|}$ . If  $h \in H$ , then  $h^m \in H$  and  $v(h^m) = h^{nm} = h$ .

Therefore  $G \simeq N \rtimes H$ , as  $N$  is normal in  $G$ ,  $N \cap H = \{1\}$  and  $G = NH$  (because  $|NH| = |N||H|$  and  $G/N \simeq H$ ).  $\square$

**Exercise 17.9.** Let  $H$  be a central subgroup of a finite group  $G$ . If  $|H|$  and  $|G/H|$  are coprime, then  $G \simeq H \times G/H$ .

## B

lemma:[s, t]

**Lemma 17.10.** *Let  $G$  be a group and  $T$  be a transversal of  $Z(G)$  in  $G$ . Then each commutator of  $G$  is of the form  $[s, t]$  for  $s, t \in T$ . In particular,  $G$  has a finite number of commutators if  $Z(G)$  is of finite index.*

*Proof.* Every element of  $G$  is of the form  $sx$  for  $s \in T$  and  $x \in Z(G)$ . To prove the first claim note that  $[sx, ty] = [s, t]$ , as  $x, y \in Z(G)$ . The second claim now follows from  $|T| = (G : Z(G))$ .  $\square$

theorem:Dietzmann

**Theorem 17.11 (Dietzmann).** *Let  $G$  be a group and  $X \subseteq G$  be a finite subset of  $G$  closed under conjugation. If there exists  $n \in \mathbb{N}$  such that  $x^n = 1$  for all  $x \in X$ , then  $\langle X \rangle$  is a finite subgroup of  $G$ .*

*Proof.* Let  $S = \langle X \rangle$  be the subgroup generated by  $S$ . Since  $x^{-1} = x^{n-1}$ , every element of  $S$  can be written as a finite product of elements of  $X$ .

Fix  $x \in X$ . We claim that if  $x \in X$  appears  $k \geq 1$  times in the representation of  $s$ , then  $s$  is a product of  $m$  elements of  $X$  where the first  $k$  elements are equal to  $x$ . Assume that

$$s = x_1 x_2 \cdots x_{t-1} x x_{t+1} \cdots x_m,$$

where each  $x_j \neq x$  for all  $j \in \{1, \dots, t-1\}$ . Then

$$s = x(x^{-1}x_1x)(x^{-1}x_2x) \cdots (x^{-1}x_{t-1}x)x_{t+1} \cdots x_m$$

is a product of  $m$  elements of  $X$  since  $X$  is closed under conjugation and the first element is our  $x$ . The same argument implies that  $s$  can be written as

$$s = x^k y_{k+1} \cdots y_m,$$

where the  $y_j$  are elements of  $X \setminus \{x\}$ .

Let  $s \in S$ . Write  $s$  as a product of  $m$  elements of  $X$ , where  $m$  is minimal. To see that  $S$  is finite it is enough to show that  $m \leq (n-1)|X|$ .

If  $m > (n-1)|X|$ , then at least one  $x \in X$  appears  $n$  times in the representation of  $s$ . Without loss of generality, write

$$s = x^n x_{n+1} \cdots x_m = x_{n+1} \cdots x_m,$$

a contradiction to the minimality of  $m$ .  $\square$

theorem:Schur\_commutador

**Theorem 17.12 (Schur).** *If  $Z(G)$  has finite index in  $G$ , then  $[G, G]$  is finite.*

*Proof.* Let  $X = \{[x, y] : x, y \in G\}$ . By Lemma 17.10,  $X$  is finite. Moreover,  $X$  is closed under conjugation,

$$g[x, y]g^{-1} = [gxg^{-1}, gyg^{-1}]$$

for all  $g, x, y \in G$ . If  $n = (G : Z(G))$ , then  $x^n = 1$  for all  $x \in X$  by Corollary 17.7. Thus the claim follows from Dietzmann's theorem (Theorem 17.11).  $\square$



**Corollary 17.13 (Sury).** *If the set of commutators of a group  $G$  is finite, then  $[G, G]$  is finite.*

*Proof.* Let  $C$  be the set of commutators of  $G$  and let  $H$  be the subgroup of  $G$  generated by  $C$ . The group  $H$  is finitely generated, say by the elements  $h_1, \dots, h_n$ . Since  $h \in Z(H)$  if and only if  $h \in C_H(H_i)$  for all  $i \in \{1, \dots, n\}$ , we conclude that  $Z(H) = \cap_{i=1}^n C_H(h_i)$ . Moreover, if  $h \in H$ , then  $hh_ih^{-1} = ch_i$  for some  $c \in C$ . Thus the conjugacy class of each  $h_i$  contains at most as many elements as  $C$ . This implies that

$$|H/Z(H)| = |H/\cap_{i=1}^n C_H(H_i)| \leq \prod_{i=1}^n (H : C_H(h_i)) \leq |C|^n.$$

Since  $H/Z(H)$  is finite,  $[H, H]$  is finite. Hence  $[G, G] = \langle C \rangle \subseteq [H, H]$  is a finite group.  $\square$

The corollary can be used to prove another proof of the following result.

**Theorem 17.14 (Hilton–Niroomand).** *Let  $G$  be a finitely generated group. If  $[G, G]$  is finite and  $G/Z(G)$  is generated by  $n$  elements, then*

$$|G/Z(G)| \leq |[G, G]|^n.$$

*Proof.* Assume that  $G/Z(G) = \langle x_1Z(G), \dots, x_nZ(G) \rangle$ . Let

$$f: G/Z(G) \rightarrow [G, G] \times \cdots \times [G, G], \quad y \mapsto ([x_1, y], \dots, [x_n, y]).$$

Note that  $f$  is well-defined: If  $y \in G$   $yz \in Z(G)$ , then

$$f(yz) = [x_i, yz] = [x_i, y] = f(y).$$

We claim that  $f$  is injective. Assume that  $f(xZ(G)) = f(yZ(G))$ . Then  $[x_i, x] = [x_i, y]$  for all  $i \in \{1, \dots, n\}$ . For each  $i$  we compute

$$\begin{aligned} [x^{-1}y, x_i] &= x^{-1}[y, x_i]x[x^{-1}, x_i] \\ &= x^{-1}[y, x_i][x_i, x]x = x^{-1}[x_i, y]^{-1}[x_i, x]x = x^{-1}[x_i, y]^{-1}[x_i, y]x = 1. \end{aligned}$$

This implies that  $x^{-1}y \in Z(G)$ . Indeed, since every  $g \in G$  can be written as  $g = x_kz$  for some  $k \in \{1, \dots, n\}$  and some  $z \in Z(G)$ , it follows that

$$[x^{-1}y, g] = [x^{-1}y, x_kz] = [x^{-1}y, x_k] = 1.$$

Thus  $f$  is injective and hence  $|G/Z(G)| \leq |[G, G]|^n$ .  $\square$

An application to infinite groups.

**Theorem 17.15.** *Let  $G$  be a torsion-free group that contains a finite-index subgroup isomorphic to  $\mathbb{Z}$ . Then  $G \simeq \mathbb{Z}$ .*

*Proof.* We may assume that  $G$  contains a finite-index normal subgroup isomorphic to  $\mathbb{Z}$ . Indeed, if  $H$  is a finite-index subgroup of  $G$  such that  $H \simeq \mathbb{Z}$ , then  $K = \bigcap_{x \in G} xHx^{-1}$  is a non-trivial normal subgroup of  $G$  (because  $K = \text{Core}_G(H)$  and  $G$  has no torsion) and hence  $K \simeq \mathbb{Z}$  (because  $K \subseteq H$ ) and  $(G : K) = (G : H)(H : K)$  is finite. The action of  $G$  on  $K$  by conjugation induces a group homomorphism  $\varepsilon : G \rightarrow \text{Aut}(K)$ . Since  $\text{Aut}(K) \simeq \text{Aut}(\mathbb{Z}) = \{-1, 1\}$ , there are two cases to consider.

Assume first that  $\varepsilon = \text{id}$ . Since  $K \subseteq Z(G)$ , let  $\nu : G \rightarrow K$  be the transfer homomorphism. By Proposition 17.5,  $\nu(g) = g^n$ , where  $n = (G : K)$ . Since  $G$  has no torsion,  $\nu$  is injective. Thus  $G \simeq \mathbb{Z}$  because it is isomorphic to a subgroup of  $K$ .

Assume now that  $\varepsilon \neq \text{id}$ . Let  $N = \ker \varepsilon \neq G$ . Since  $K \simeq \mathbb{Z}$  is abelian,  $K \subseteq N$ . The result proved in the previous paragraph applied to  $\varepsilon|_N = 1$  implies that  $N \simeq \mathbb{Z}$ , as  $N$  contains a finite-index subgroup isomorphic to  $\mathbb{Z}$ . Let  $g \in G \setminus N$ . Since  $N$  is normal in  $G$ ,  $G$  acts by conjugation on  $N$  and hence there exists a group homomorphism  $c_g \in \text{Aut}(N) \simeq \{-1, 1\}$ . Since  $K \subseteq N$   $g$  acts non-trivially on  $K$ ,

$$c_g(n) = gng^{-1} = n^{-1}$$

for all  $n \in N$ . Since  $g^2 \in N$ ,

$$g^2 = gg^2g^{-1} = g^{-2}.$$

Therefore  $g^4 = 1$ , a contradiction since  $g \neq 1$  and  $G$  has no torsion.  $\square$

## C

## References

1. E. Acri, R. Lutowski, and L. Vendramin. Retractability of solutions to the Yang-Baxter equation and  $p$ -nilpotency of skew braces. *Internat. J. Algebra Comput.*, 30(1):91–115, 2020.
2. O. Akgün, M. Mereb, and L. Vendramin. Enumeration of set-theoretic solutions to the Yang-Baxter equation, 2020.
3. B. Amberg, S. Franciosi, and F. de Giovanni. *Products of groups*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1992. Oxford Science Publications.
4. S. A. Amitsur. Nil radicals. Historical notes and some new results. In *Rings, modules and radicals (Proc. Internat. Colloq., Keszthely, 1971)*, pages 47–65. Colloq. Math. Soc. János Bolyai, Vol. 6, 1973.
5. N. Andruskiewitsch and M. Graña. From racks to pointed Hopf algebras. *Adv. Math.*, 178(2):177–243, 2003.
6. M. Ashford and O. Riordan. Counting racks of order  $n$ . *Electron. J. Combin.*, 24(2):Paper No. 2.32, 20, 2017.
7. D. Bachiller. Counterexample to a conjecture about braces. *J. Algebra*, 453:160–176, 2016.
8. D. Bachiller. Solutions of the Yang-Baxter equation associated to skew left braces, with applications to racks. *J. Knot Theory Ramifications*, 27(8):1850055, 36, 2018.
9. V. G. Bardakov, M. V. Neshchadim, and M. K. Yadav. Computing skew left braces of small orders. *Internat. J. Algebra Comput.*, 30(4):839–851, 2020.
10. S. R. Blackburn. Enumerating finite racks, quandles and kei. *Electron. J. Combin.*, 20(3):Paper 43, 9, 2013.
11. M. Brešar. *Introduction to noncommutative algebra*. Universitext. Springer, Cham, 2014.
12. E. Brieskorn. Automorphic sets and braids and singularities. In *Braids (Santa Cruz, CA, 1986)*, volume 78 of *Contemp. Math.*, pages 45–115. Amer. Math. Soc., Providence, RI, 1988.
13. N. P. Byott. Nilpotent and abelian Hopf-Galois structures on field extensions. *J. Algebra*, 381:131–139, 2013.
14. N. P. Byott. Solubility criteria for Hopf-Galois structures. *New York J. Math.*, 21:883–903, 2015.
15. F. Catino, I. Colazzo, and P. Stefanelli. Skew left braces with non-trivial annihilator. *J. Algebra Appl.*, 18(2):1950033, 23, 2019.
16. F. Catino and R. Rizzo. Regular subgroups of the affine group and radical circle algebras. *Bull. Aust. Math. Soc.*, 79(1):103–107, 2009.
17. F. Cedó. Left braces: solutions of the Yang-Baxter equation. *Adv. Group Theory Appl.*, 5:33–90, 2018.
18. F. Cedó, T. Gateva-Ivanova, and A. Smoktunowicz. Braces and symmetric groups with special conditions. *J. Pure Appl. Algebra*, 222(12):3877–3890, 2018.
19. F. Cedó, E. Jespers, and J. Okniński. Retractability of set theoretic solutions of the Yang-Baxter equation. *Adv. Math.*, 224(6):2472–2484, 2010.

20. F. Cedó, E. Jespers, and J. Okniński. Braces and the Yang-Baxter equation. *Comm. Math. Phys.*, 327(1):101–116, 2014.
21. F. Cedó, A. Smoktunowicz, and L. Vendramin. Skew left braces of nilpotent type. *Proc. Lond. Math. Soc.* (3), 118(6):1367–1392, 2019.
22. F. Chouraqui. Garside groups and Yang-Baxter equation. *Comm. Algebra*, 38(12):4441–4460, 2010.
23. A. Clay and D. Rolfsen. *Ordered groups and topology*, volume 176 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2016.
24. M. Deakonesku and G. L. Uolls. On the orbits of automorphism groups. *Sibirsk. Mat. Zh.*, 46(3):533–537, 2005.
25. P. Dehornoy. Set-theoretic solutions of the Yang-Baxter equation, RC-calculus, and Garside germs. *Adv. Math.*, 282:93–127, 2015.
26. V. G. Drinfel'd. On some unsolved problems in quantum group theory. In *Quantum groups (Leningrad, 1990)*, volume 1510 of *Lecture Notes in Math.*, pages 1–8. Springer, Berlin, 1992.
27. M. Elhamdadi and S. Nelson. *Quandles—an introduction to the algebra of knots*, volume 74 of *Student Mathematical Library*. American Mathematical Society, Providence, RI, 2015.
28. P. Etingof and S. Gelaki. A method of construction of finite-dimensional triangular semisimple Hopf algebras. *Math. Res. Lett.*, 5(4):551–561, 1998.
29. P. Etingof, T. Schedler, and A. Soloviev. Set-theoretical solutions to the quantum Yang-Baxter equation. *Duke Math. J.*, 100(2):169–209, 1999.
30. The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.11.1*, 2021.
31. T. Gateva-Ivanova. A combinatorial approach to the set-theoretic solutions of the Yang-Baxter equation. *J. Math. Phys.*, 45(10):3828–3858, 2004.
32. T. Gateva-Ivanova. Set-theoretic solutions of the Yang-Baxter equation, braces and symmetric groups. *Adv. Math.*, 338:649–701, 2018.
33. T. Gateva-Ivanova and M. Van den Bergh. Semigroups of  $I$ -type. *J. Algebra*, 206(1):97–112, 1998.
34. L. Guarnieri and L. Vendramin. Skew braces and the Yang-Baxter equation. *Math. Comp.*, 86(307):2519–2534, 2017.
35. J. Hoste and P. D. Shanahan. An enumeration process for racks. *Math. Comp.*, 88(317):1427–1448, 2019.
36. I. M. Isaacs. *Finite group theory*, volume 92 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2008.
37. I. M. Isaacs. Group actions and orbits. *Arch. Math. (Basel)*, 98(5):399–401, 2012.
38. N. Jacobson. The radical and semi-simplicity for arbitrary rings. *Amer. J. Math.*, 67:300–320, 1945.
39. E. Jespers, L. Kubat, A. Van Antwerpen, and L. Vendramin. Factorizations of skew braces. *Math. Ann.*, 375(3-4):1649–1663, 2019.
40. E. Jespers and J. Okniński. *Noetherian semigroup algebras*, volume 7 of *Algebra and Applications*. Springer, Dordrecht, 2007.
41. G. Köthe. Die Struktur der Ringe, deren Restklassenring nach dem Radikal vollständig reduzibel ist. *Math. Z.*, 32(1):161–186, 1930.
42. J. Krempa. Logical connections between some open problems concerning nil rings. *Fund. Math.*, 76(2):121–130, 1972.
43. I. Lau. An associative left brace is a ring. *J. Algebra Appl.*, 19(9):2050179, 6, 2020.
44. V. Lebed and L. Vendramin. On structure groups of set-theoretic solutions to the Yang-Baxter equation. *Proc. Edinb. Math. Soc.* (2), 62(3):683–717, 2019.
45. J.-H. Lu, M. Yan, and Y.-C. Zhu. On the set-theoretical Yang-Baxter equation. *Duke Math. J.*, 104(1):1–18, 2000.
46. H. Meng, A. Ballester-Bolínches, and R. Esteban-Romero. Left braces and the quantum Yang-Baxter equation. *Proc. Edinb. Math. Soc.* (2), 62(2):595–608, 2019.
47. P. P. Nielsen. Simplifying Smoktunowicz's extraordinary example. *Comm. Algebra*, 41(11):4339–4350, 2013.
48. J. Pakianathan and K. Shankar. Nilpotent numbers. *Amer. Math. Monthly*, 107(7):631–634, 2000.

49. R. Plemmons. Construction and analysis of non-equivalent finite semigroups. In *Computational Problems in Abstract Algebra (Proc. Conf., Oxford, 1967)*, pages 223–228. Pergamon, Oxford, 1970.
50. S. D. Promislow. A simple example of a torsion-free, nonunique product group. *Bull. London Math. Soc.*, 20(4):302–304, 1988.
51. J. H. Przytycki. Distributivity versus associativity in the homology theory of algebraic structures. *Demonstratio Math.*, 44(4):823–869, 2011.
52. W. Rump. A decomposition theorem for square-free unitary solutions of the quantum Yang-Baxter equation. *Adv. Math.*, 193(1):40–55, 2005.
53. W. Rump. Braces, radical rings, and the quantum Yang-Baxter equation. *J. Algebra*, 307(1):153–170, 2007.
54. W. Rump. The brace of a classical group. *Note Mat.*, 34(1):115–144, 2014.
55. A. Smoktunowicz. Polynomial rings over nil rings need not be nil. *J. Algebra*, 233(2):427–436, 2000.
56. A. Smoktunowicz. On some results related to Köthe’s conjecture. *Serdica Math. J.*, 27(2):159–170, 2001.
57. A. Smoktunowicz. Some results in noncommutative ring theory. In *International Congress of Mathematicians. Vol. II*, pages 259–269. Eur. Math. Soc., Zürich, 2006.
58. A. Smoktunowicz. A note on set-theoretic solutions of the Yang-Baxter equation. *J. Algebra*, 500:3–18, 2018.
59. A. Smoktunowicz. On Engel groups, nilpotent groups, rings, braces and the Yang-Baxter equation. *Trans. Amer. Math. Soc.*, 370(9):6535–6564, 2018.
60. A. Smoktunowicz and L. Vendramin. On skew braces (with an appendix by N. Byott and L. Vendramin). *J. Comb. Algebra*, 2(1):47–86, 2018.
61. A. Soloviev. Non-unitary set-theoretical solutions to the quantum Yang-Baxter equation. *Math. Res. Lett.*, 7(5-6):577–596, 2000.
62. Y. P. Sysak. Products of almost abelian groups. In *Investigations of groups with restrictions for subgroups (Russian)*, pages 81–85, iii. Akad. Nauk Ukrain. SSR, Inst. Mat., Kiev, 1988.
63. C. Tsang and C. Qin. On the solvability of regular subgroups in the holomorph of a finite solvable group. *Internat. J. Algebra Comput.*, 30(2):253–265, 2020.
64. P. Vojtěchovský and S. Y. Yang. Enumeration of racks and quandles up to isomorphism. *Math. Comp.*, 88(319):2523–2540, 2019.
65. M. Wada. Group invariants of links. *Topology*, 31(2):399–406, 1992.
66. A. Weinstein and P. Xu. Classical solutions of the quantum Yang-Baxter equation. *Comm. Math. Phys.*, 148(2):309–343, 1992.
67. E. Zelmanov. *Nil rings and periodic groups*. KMS Lecture Notes in Mathematics. Korean Mathematical Society, Seoul, 1992. With a preface by Jongsik Kim.



# Index

- 1-coborde, 47
- 1-cocyclo, 45
- $p$ -complemento, 70
  
- Annihilator, 41
- Automorfism
  - central, 88
  
- Bachiller, D., 43, 117
- Brace, 31
  - additive group, 31
  - associative, 36
  - left nil, 90
  - left nilpotent, 89
  - left series, 88
  - meta-trivial, 75
  - multiplicative group, 31
  - right nil, 90
  - strongly nil, 90
  - strongly nilpotent, 90
  - trivial, 31
  - two sided, 35
- Braid group, 1
- Braided group, 81
- Burnside
  - Theorem, 70
  
- Catino, F., 117
- Cedó, F., 43, 86
- Centralizador, 58
- Chouraqui, F., 86
- Clausura normal, 69
- Commutator identities
  - for braces, 34
- Condición normalizadora, 57
- Cycle set, 19
  - non-degenerate, 19
  
- Deaconescu–Walls
  - theorem, 87
- Dehornoy, P., 86
- Derivación
  - interior, 47
- Derivation, 45
- Derived
  - series, 67
- Dietzmann
  - theorem, 122
- Direct product
  - of braces, 32
- Drinfeld, V., 5
  
- Etingof, P., 5
  
- Garside
  - monoid, 85
- Gateva–Ivanova, T., 5, 43, 86
- Group
  - left ordered, 107
  - metabelian, 73
  - nilpotent, 56
  - with the unique product property, 102
- Grupo
  - perfecto, 58
- Guarnieri, L., 43
  
- Hall, P., 55
- Hall–Witt
  - identity, 55
- Holomorph, 113
- Homomorphism
  - of braces, 34
  - of cycle sets, 19
  - of racks, 23
  - of skew cycle sets, 27

- Ideal, 39
  - primitive, 10
- Jacobi
  - identity, 55
- Jacobi, G., 55
- Jacobson
  - radical ring, 18, 35
- Jespers, E., 43
- Kernel, 39
- Kinyon, M., 43
- Lau, I., 43
- Left
  - ideal, 38
- Lema
  - de Hall, 66
- Lemma
  - three subgroups, 55
  - Zorn, 12
- Lifting, 45
- Lower central series, 56
- Lu, J-H., 5
- Lyubashenko, V., 5
- Monoid, 85
  - Garside, 85
- Nilpotency index, 56
- Normalizador, 58
- Okniński, J., 43
- Quotient brace, 42
- Rack, 23
  - Alexander, 23
  - dihedral, 23
  - homomorphism, 23
  - isomorphism, 23
  - trivial, 23
- Radical ring, 18, 35
- Regular subgroup, 113
- Retraction
  - of a solution, 97
- Ring
  - nil, 15
  - primitive, 10
  - radical, 15
- Rizzo, ???, 117
- Rump, W., 5, 43
- Schedler, T., 5
- Schur
  - theorem, 122
- Schur-Zassenhaus
  - theorem, 49, 50
- Serie
  - central, 59
  - central ascendente, 58
- Skew cycle set, 27
  - non-degenerate, 27
- Smoktunowicz, A., 43
- Socle, 40
- Soloviev, A., 5
- Solution, 1
  - derived rack, 25
  - finite, 1
  - indecomposable, 109
  - involutive, 18
  - multipermutation, 98
  - non-degenerate, 3
  - simple, 109
  - trivial, 2
- Split
  - extension, 45
- Strong
  - left ideal, 38
- Subbrace, 38
- Subgroup
  - characteristic, 67
  - minimal normal, 67
- Symmetric group, 18
- Sysak, Y., 47
- Teorema
  - de Grün, 58
  - de Hirsch, 59
  - de Sysak, 47
- Teorema de
  - Baumslag-Wiegold, 65
  - Hilton-Niroomand, 123
- Theorem
  - of Itô, 73
  - of Kegel-Wielandt, 74
  - of Sysak, 74
- Transfer map, 119
- Van den Bergh, M., 5, 86
- Vendramin, L., 43
- Wada, M., 5
- Witt, E., 55
- Yan, M., 5
- Zhu, Y-C., 5